



https://helda.helsinki.fi

Forskolin attenuates the NLRP3 inflammasome activation and IL-1 beta secretion in human macrophages

Chen, Yan

2019-12

Chen, Y, Wen, JG, Feng, JJ, Wang, YH, Li, TF, Nurmi, K, Eklund, KK & Xing, D 2019, 'Forskolin attenuates the NLRP3 inflammasome activation and IL-1 beta secretion in human macrophages ', Pediatric Research, vol. 86, no. 6, pp. 692-698. https://doi.org/10.1038/s41390-019-0418-

http://hdl.handle.net/10138/341301 https://doi.org/10.1038/s41390-019-0418-4

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository. This is an electronic reprint of the original article. This reprint may differ from the original in pagination and typographic detail. Please cite the original version.

Forskolin attenuates the NLRP3 inflammasome activation and IL-1β secretion in human macrophages

Cite this article as: Yan Chen, Jian Guo Wen, Jin Jin Feng, Yi He Wang, Tian Fang Li, Katariina Nurmi, Kari K. Eklund and Dong Xing, Forskolin attenuates the NLRP3 inflammasome activation and IL-1 β secretion in human macrophages, *Pediatric Research* doi:10.1038/s41390-019-0544-z

This Author Accepted Manuscript is a PDF file of an unedited peer-reviewed manuscript that has been accepted for publication but has not been copyedited or corrected. The official version of record that is published in the journal is kept up to date and so may therefore differ from this version.

r til Terms of use and reuse: academic research for non-commercial purposes, see here for full terms. https://www.nature.com/authors/policies/license.html#AAMtermsV1

© 2019 Macmillan Publishers Limited, part of Springer Nature.

AUTHOR ACCEPTED MANUSCRIPT

Forskolin attenuates the NLRP3 inflammasome activation and IL-1β secretion in human macrophages

Yan Chen^{1, 3#}, Jian Guo Wen¹, Jin Jin Feng¹, Yi He Wang¹, Tian Fang Li², Katariina Nurmi³, Kari K. Eklund^{3, 4}, Dong Xing¹

¹Pediatric urodynamic Center and Department of Urology, Institute of Clinical Medicine, the First Affiliated Hospital of Zhengzhou University, 450052 Zhengzhou, China

²Helsinki Rheumatic Diseases and Inflammation Research Group, Institute of Clinicum, University of Helsinki, Haartmaninkatu 8,00290 Helsinki, Finland

³Department of Medicine, D of Rheumatology, the First Affiliated Hospital of Zhengzhou University, 450052 Zhengzhou, China

⁴Department of Medicine, Division of Rheumatology, Helsinki University Central Hospital, Haartmaninkatu 2,000290 Helsinki, Finland

Corresponding Author:

Yan Chen, M.D., Ph.D., 1) Pediatric urodynamic Center and Department of Urology, Institute of Clinical Medicine, the First Affiliated Hospital of Zhengzhou University, 450052 Zhengzhou, China. Fax: +86 371 66970906, Phone: +86 18537157126, Email: <u>chenyan.abby@163.com</u>

AUTHOR ACCEPTED MANUSCRIPT

Figure Legend:

•Nigericin activates the NLRP3 inflammasome by stimulating an intracellular K+ efflux through pannexin-1 dependent pathway that induces the assembly of the NLRP3 inflammasome (NLRP3-ASC-Caspase 1) and its subsequent activation of the secretion of IL-1 β in PMA-differentiated THP-1 macrophages.

•Forskolin increases the level of intracellular cAMP which then binds to NLRP3. Binding of cAMP to NLRP3 promotes the ubiquitination of NLRP3, thus leading into degradation of both cAMP and NLRP3 protein in proteosomes. The degradation of NLRP3 then interrupts the assembly of inflammasome (NLRP3-ASC-Caspase 1) and thus the activation of pro-caspase 1 autocatalysis . it .n of IL .n of I into caspase 1 which subsequently interrupting the maturation and secretion of IL-1β.

© 2019 Macmillan Publishers Limited, part of Springer Nature.

