

<https://helda.helsinki.fi>

---

## Antibacterial Activity of 2-Hydroxyisocaproic Acid (HICA) Against Obligate Anaerobic Bacterial Species Associated With Periodontal Disease

Sakko, Marjut

2021-01-01

---

Sakko , M , Rautemaa-Richardson , R , Sakko , S , Richardson , M & Sorsa , T 2021 , '  
Antibacterial Activity of 2-Hydroxyisocaproic Acid (HICA) Against Obligate Anaerobic  
Bacterial Species Associated With Periodontal Disease ' , Microbiology Insights , vol. 14 . <https://doi.org/10.1177/11771177211050086>

---

<http://hdl.handle.net/10138/342560>

<https://doi.org/10.1177/11771177211050086>

---

cc\_by

publishedVersion

---

*Downloaded from Helda, University of Helsinki institutional repository.*

*This is an electronic reprint of the original article.*

*This reprint may differ from the original in pagination and typographic detail.*

*Please cite the original version.*

# Antibacterial Activity of 2-Hydroxyisocaproic Acid (HICA) Against Obligate Anaerobic Bacterial Species Associated With Periodontal Disease

Microbiology Insights  
Volume 14: 1–7  
© The Author(s) 2021  
DOI: 10.1177/11786361211050086



Marjut Sakko<sup>1</sup> , Riina Rautemaa-Richardson<sup>2,3</sup>, Samuli Sakko<sup>1</sup>, Malcolm Richardson<sup>4,5</sup> and Timo Sorsa<sup>1,6</sup>

<sup>1</sup>Department of Oral and Maxillofacial Diseases, Clinicum, University of Helsinki, and Helsinki University Hospital, Helsinki, Finland. <sup>2</sup>Division of Infection, Inflammation and Respiratory Medicine, Faculty of Biology, Medicine and Health, The University of Manchester, Manchester, UK. <sup>3</sup>Department of Infectious Diseases, Manchester Academic Health Science Centre, Wythenshawe Hospital, Manchester University NHS Foundation Trust, Manchester, UK. <sup>4</sup>Mycology Reference Centre Manchester, ECMM Excellence Centre of Medical Mycology, Manchester University NHS Foundation Trust, Manchester, UK. <sup>5</sup>Manchester Fungal Infection Group, Division of Infection, Immunity and Respiratory Medicine, University of Manchester, Manchester, UK. <sup>6</sup>Department of Oral Diseases, Karolinska Institutet, Huddinge, Sweden.

**ABSTRACT:** Topical antiseptics are used to assist and further increase the effect of mechanical biofilm eradication and to potentially prevent new biofilm formation in periodontal treatment. This is of importance in treatment-resistant infections with 10% prevalence of all periodontitis cases to avoid the need for antibiotic therapy. The purpose of this study was to evaluate the antimicrobial activity of DL-2-hydroxyisocaproic acid on human pathogenic obligate anaerobic bacteria related to periodontitis. In this study antimicrobial activity of 2-hydroxyisocaproic acid was observed against 14 bacterial reference strains and clinical isolates of obligate anaerobic bacterial species using a microdilution method in 1.25 to 160 mg/mL concentrations of 2-hydroxyisocaproic acid. The 11 strains of bacteria included in this study are typically associated with periodontal disease; *Porphyromonas gingivalis*, *Fusobacterium nucleatum*, *Tannerella forsythia*, *Aggregatibacter actinomycetemcomitans*, and *Parvimonas micra*. Three strains of *Cutibacterium acnes*, normally associated with skin diseases, were tested for comparison. 90% inhibitory concentration was determined at 48 hours and minimum bactericidal concentration was determined after 72 hours incubation. The 2-hydroxyisocaproic acid was bactericidal at  $\geq 160$  mg/mL for all isolates tested. The reference strain of *T. forsythia*, and the reference strain and the clinical isolates of *C. acnes* were the most tolerant ones. The reference strains and clinical isolates of *F. nucleatum* and *A. actinomycetemcomitans* were killed at  $\geq 40$  mg/mL concentration. In conclusion, topical use of 2-hydroxyisocaproic acid could eventually be a well-tolerated and useful method in the therapy of patients with difficult-to-treat periodontal disease or other superficial infections to avoid unnecessary antibiotic use and the emergence of antibiotic resistance.

**KEYWORDS:** Anaerobic bacteria, bactericidal, 2-hydroxyisocaproic acid, 2-hydroxy-4-methyl pentanoic acid, 2-hydroxy-4-methyl valeric acid, leucic acid, new topical antimicrobial

RECEIVED: February 6, 2021. ACCEPTED: September 10, 2021.

TYPE: Original Research

**FUNDING:** The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by grants from the Finnish Association of Female Dentists, the Helsinki University Central Hospital Research Foundation (Y1014SL012, TYH 2016251, 2017251, 2018229, and 2019319) and the Karolinska Institutet, Stockholm, Sweden. Open access funded by Helsinki University Library.

**DECLARATION OF CONFLICTING INTERESTS:** The author(s) declared the following potential conflicts of interest with respect to the research, authorship, and/or publication of this article: Dr. Marjut Sakko, Dr. Riina Rautemaa-Richardson, Medical student Samuli Sakko, and Prof. Malcolm Richardson have nothing to declare. Prof. Timo Sorsa is one of the inventors of patent EP 0871438 B1: Use of alpha-hydroxy acids in the manufacture of medicament for the treatment of inflammation.

**CORRESPONDING AUTHOR:** Marjut Sakko, Department of Oral and Maxillofacial Diseases, Clinicum, University of Helsinki, PO Box 63, (Haartmaninkatu 8), Helsinki FIN-00014, Finland. Email: marjut.sakko@helsinki.fi

## Introduction

Periodontitis is an inflammatory disease of periodontal tissues, which is induced by dysbiotic polymicrobial biofilms.<sup>1</sup> The degree of inflammation and symptoms depends on the quantity of microbial cells and their virulence as well as host responses. Various cell surface proteins and carbohydrates, toxins, and hydrolytic enzymes are considered to be microbial virulence factors because they may by enhancing microbial attachment and protection from the host induce the microbial burden. The ability of microbes to form especially the dysbiotic biofilm is potentially an additional virulence factor.<sup>2</sup> The ability to find synergistic microbial species, the ability to evade or resist host responses and to release bacterial mediators such as lipopolysaccharides (LPS), and the release of tissue damaging

proteolytic enzymes have an impact on microbial survival in multispecies infections.<sup>3</sup>

Bacterial species associated with severe periodontitis are mainly obligate anaerobic rods, such as *Porphyromonas gingivalis*, *Fusobacterium nucleatum*, *Aggregatibacter actinomycetemcomitans*, *Prevotella* spp., *Campylobacter rectus* or some cocci such as *Parvimonas* spp. They are able to live and multiply in the anaerobic areas such as the gingival sulcus. *Treponemas* and viruses can also be typically found in periodontitis. It has been suggested that in the polymicrobial synergy and especially the dysbiosis theory of the pathogenesis of periodontitis some corner-stone pathogens have increased ability to form bacterial communities found in the presence of periodontal disease. Accordingly, commensal microbial species are suppressed.<sup>1</sup>



Topical antimicrobials potentially increase the effect of mechanical biofilm eradication. They may also prevent new biofilm formation as a part of the anti-infective treatment of periodontitis.<sup>4,5</sup> Antiseptics can be used as an oral rinse by the patient or injected into the gingival sulcus by a dentist. The use of these agents as an adjuvant to scaling and root planning has been suggested to potentially improve the outcome of chronic periodontitis.<sup>6,7</sup> This improved efficacy would be particularly helpful in the management of treatment-resistant infections representing some 10% of all periodontitis cases in order to reduce the use of systemic antibiotics in this patient group.<sup>8</sup> Overall, topical antiseptics are potential devices in the treatment of any kind of superficial infections for avoiding the potential spreading of infection.<sup>9</sup>

Hietala et al<sup>10</sup> identified 2-hydroxyisocaproic acid (HICA) as a bacterial amino acid with antimicrobial potential from a fermentation product of *Lactobacillus plantarum*. It is a by-product of the leucine-acetyl-CoA pathway in the leucine degradation pathway. For HICA elimination hydroxyisocaproic acid dehydrogenase (HicDH) enzyme is needed. Both of them are mainly produced by lactobacilli.<sup>11-23</sup> The reduction reaction of 2-ketoisocaproic acid (KICA) to HICA is catalyzed by hydroxyisocaproic acid dehydrogenase (HicDH). An NADH-dependent oxygenation reaction is needed for HICA elimination back to KICA.<sup>22</sup> The broad-spectrum antimicrobial activity of HICA has been shown against facultative anaerobic gram-positive and gram-negative bacteria,<sup>10,24</sup> obligate anaerobic *F. nucleatum*,<sup>24</sup> several fungal species,<sup>10,25-27</sup> and against *Candida albicans* biofilms.<sup>28</sup> HICA exerts anti-inflammatory properties as well.<sup>29</sup>

The European Food Safety Authority (EFSA)<sup>30</sup> has approved HICA as a flavoring substance in nutritional products. The maximum survey-delivered daily intake is 0.0012 µg/capita/day and theoretical maximum daily intake is 3800 µg/capita/day. The dose of the threshold of concern is 1800 µg/capita/day. HICA is also found to be present as a natural compound in human breast milk in minimal amounts, 17 µg/L.<sup>31</sup> Selis et al<sup>32</sup> showed that DL-HICA is not cytotoxic or genotoxic for human periodontal fibroblasts at <10 mg/mL after 24 hours incubation. They considered that HICA could be tested as a root canal medication for regenerative endodontics. Cytotoxicity of the widely used antiseptic chlorhexidine on human cells is well-reported in the literature. Verma et al<sup>33</sup> showed that chlorhexidine was toxic to periodontal fibroblasts beyond 10 mg/mL even after 1, 5, and 15 minutes incubation. Since HICA can be metabolized by human tissues<sup>34,35</sup> and is found naturally in nutrition it should be well-tolerated at least in the doses recommended by EFSA. HICA could eventually be a useful new and safe topical antimicrobial agent.

The hypothesis of this study was that HICA could be active also against obligate anaerobic periodontopathogenic bacteria, because it affects the microbial metabolism. The purpose of this study was to evaluate the microbial activity and spectrum

of HICA against obligately anaerobic bacterial in an environment that is favorable for their growth.

## Materials and Methods

### Bacterial strains

Fourteen obligate anaerobic strains were included in the study: 7 reference strains and 7 clinical isolates of *Porphyromonas gingivalis*, *Fusobacterium nucleatum*, *Tannerella forsythia*, *Aggregatibacter actinomycetemcomitans*, *Parvimonas micra*, and *Cutibacterium acnes*. Nine isolates belonged to the group of gram-negative rods, 2 of them were gram-positive cocci and 3 others were gram-positive rods (Table 1). In addition to gingival micro-organisms 3 strains of *C. acnes*, which can be found often in skin diseases acted as a control. Clinical isolates were obtained from the Clinical Microbiology Laboratory of Helsinki University Central Hospital HUSLAB (PD and TS isolates; Finland), and from the Department of Oral and Maxillofacial diseases, University of Helsinki (A, T, and R isolates; Finland). The reference strains were obtained from the American Type Culture Collection (ATCC; USA). The bacterial strains and isolates were identified by standard biochemical methods.

### Antimicrobial susceptibility testing

The susceptibility of obligate anaerobic bacteria for DL-HICA was tested by a modification of the standard microdilution method of the Clinical and Laboratory Standards Institute (CLSI)<sup>36</sup>; Methods for antimicrobial susceptibility testing of anaerobic bacteria, document M11-A5. Culture medium, inoculum concentration, and incubation volume were used according to the manual instructions. Bacterial cell densities were prepared at a concentration of 10<sup>7</sup> cfu/mL in broth, estimated by using McFarland standard 3.0 and diluted in serial 1:100. This was confirmed also by quantitative plate counts. 90% inhibitory concentration (IC<sub>90</sub>) was determined at 48 hours, which is the MIC (minimum inhibitory concentration) breakpoint according to manual instructions of the CLSI. The minimum bactericidal concentration (MBC) was determined after 72 hours incubation. The concentrations of HICA and the test pH of 6.5 were selected following on from a series of the pilot studies. Increasing concentrations of 1.38, 2.75, 5.5, 11, 22, 44, 88, and 176 mg/mL of HICA were prepared in thioglycollate broth. Controls free of antimicrobial agents were included in all experiments (negative control). Chlorhexidine digluconate (pH 6.5; 0.6 mg/mL) was incorporated as a positive control. To check for sterility, pure thioglycollate broth without bacteria was incubated. Microbial suspensions (20 µL) were added to the test solutions (180 µL) in 96-well cell-culture microtiter plates (F96 MicroWell Plates, NUNC, ThermoFisher Scientific, Leicestershire, UK) resulting in a final bacterial density of 10<sup>6</sup> cfu/mL in the wells. Final test concentrations of HICA resulted in 1.25 to 160 mg/

**Table 1.** Obligate anaerobic bacterial strains included in the study (n = 14) are listed in this table. Their designations, sources, minimum bactericidal concentrations (MBC) of HICA at 72 hours are shown.

ORGANISM	DESIGNATION	SOURCE	MBC AT 72 H (MG/ML)
Gram-negative rods			
<i>Porphyromonas gingivalis</i>	ATCC 33277	Gingival sulcus	≥80
<i>Porphyromonas gingivalis</i>	T 75048	Gingival sulcus	≥20
<i>Fusobacterium nucleatum</i>	ATCC 25586	Cervico-facial lesion	≥40
<i>Fusobacterium nucleatum</i>	PD 788	Gingival sulcus	≥40
<i>Tannerella forsythia</i>	ATCC 43037	Gingival sulcus	≥160
<i>Tannerella forsythia</i>	R 1015	Oral abscesses	≥80
<i>Aggregatibacter actinomycetemcomitans, serotype a</i>	ATCC 29523	Blood	≥10
<i>Aggregatibacter actinomycetemcomitans, serotype b</i>	ATCC 43718	Gingival sulcus	≥40
<i>Aggregatibacter actinomycetemcomitans, non-serotyped</i>	R 58	Oral abscesses	≥10
Gram-positive cocci			
<i>Parvimonas micra</i>	ATCC 33270	Purulent pleurisy	≥80
<i>Parvimonas micra</i>	A 299	Gingival sulcus	≥80
Gram-positive rods			
<i>Cutibacterium acnes</i>	ATCC 6919	Facial acne	≥160
<i>Cutibacterium acnes</i>	TS 9498	Breast skin	≥160
<i>Cutibacterium acnes</i>	TS 9512	Subarachnoidal lesion	≥160

Clinical bacterial isolates were obtained from the Clinical Microbiology Laboratory Helsinki University Central Hospital HUSLAB (PD and TS), and from the Department of Oral and Maxillofacial diseases, University of Helsinki (A, R and T). The reference strains were obtained from the American Type Culture Collection (ATCC; USA).


mL in the wells of the microtiter plates. Bacterial cultures were grown in an anaerobic atmosphere at 35°C for up to 72 hours. Oxygen was eliminated from the anaerobic jars with the Anoxomat® III Anaerobic Culture System (Advanced Instruments, Norwood, Massachusetts, USA). Anaerobic indicators were used to monitor the anaerobic test conditions during incubation (Resazurin Anaerobic Indicator, Oxoid, ThermoFisher Scientific, Leicestershire, UK). The suspensions in the wells were mixed using a sterile instrument to ensure homogenous cultures before measurement. To determine the optical density at 0, 24, 48, and 72 hours a spectrophotometer (450 nm; Victor X4, 2030 Multilabel reader, PerkinElmer, Massachusetts, USA) was used. Then, the microbicidal activity of HICA and the controls were determined by culture on Brucella Blood agar (BD BBL™ Brucella agar with 5% sheep blood, hemin and vitamin K1 prepared media, Anaerobe Systems, ThermoFisher Scientific, Leicestershire, UK) after the last measurement with spectrophotometer. About 20 µL of test suspension from each well was cultured on agar for another 72 hours. The experiments were performed twice in quadruplicate.

### Analysis of bacterial growth

Bacterial growth was determined in each test solution by changes in optical density. Bacterial growth inhibition was determined in comparison to maximum bacterial growth observed in the negative control for each bacterial strain. The results of the spectrophotometrical analyses are presented as means and standard error of mean. The results of bacterial growth inhibition over 90% (IC<sub>≥90</sub>) for each reference strain and clinical isolate were read at 48 hours. The results of microbicidal activity which were determined by test solution cultures on agar are shown as minimum bactericidal concentration (MBC) at 72 hours.

### Results

HICA was bactericidal at ≥160 mg/mL for all reference strains (ATCC) and clinical isolates included in the study, at 72 hours (Table 1). The reference strain and the clinical isolate of *P. gingivalis* and *P. micra*, and the clinical isolate of *T. forsythia* were killed at ≥80 mg/mL HICA. A concentration of ≥40 mg/mL HICA was bactericidal for the reference strain and the clinical isolate of *F. nucleatum*, and 2 reference strains of



**Figure 1.** Growth inhibition of gram-negative rods in the presence of increasing concentration of HICA. The activity of 5 to 80 mg/mL of HICA in thioglycollate broth on the clinical isolates (a, c, and e) and the reference strains (b, d, and f) of *P. gingivalis*, *F. nucleatum* and 10 to 160 mg/mL of HICA on *T. forsythia* in anaerobic test conditions at pH 6.5 up to 72 hours is shown. Chlorhexidine digluconate 0.6 mg/mL was used as a positive control and thioglycollate broth as a negative control.

*A. actinomycetemcomitans* including serotype a and b and 1 non-serotyped clinical isolate, at 72 hours.


At 48 hours, greater than 90% of bacterial growth of the reference strain, the clinical isolate of *P. gingivalis* and the clinical isolate of *F. nucleatum* were inhibited by  $\geq 20$  mg/mL concentration of HICA (Figure 1a–c). However, over 90% of bacterial growth of the reference strain of *F. nucleatum* was inhibited even by  $\geq 10$  mg/mL concentration (Figure 1d). Bacterial growth inhibition for the reference strain and the clinical isolate of *T. forsythia* was over 90% in the presence of  $\geq 80$  mg/mL HICA at 48 hours (Figure 1e and f). Dose-dependent inhibition was shown by HICA for all reference strains and clinical isolates of *P. gingivalis*, *F. nucleatum*, and *T. forsythia*.

Over 90% of the bacterial growth of the non-serotyped clinical isolate and the serotype a reference strain (ATCC 29523) of *A. actinomycetemcomitans* was inhibited in  $\geq 10$  mg/mL concentration of HICA at 48 hours (Figure 2a and b).


Over 90% of the growth of the reference strain of *A. actinomycetemcomitans* serotype b (ATCC 43718) was inhibited even in the presence of  $\geq 5$  mg/mL of HICA (Figure 2c). The concentration, in which over 50% but under 90% of the growth of the *A. actinomycetemcomitans* strains was variable, but dose-dependent inhibition could be seen up to 48 hours.

At 48 hours, the growth inhibition of the reference strain and the clinical isolate of *P. micra* was evident in the presence of  $\geq 40$  mg/mL HICA around 50% or higher (Figure 3a and b). Incubation for 72 hours was required to demonstrate 90% growth inhibition in the presence of  $\geq 40$  mg/mL HICA. The bacterial growth inhibition caused by HICA was mainly dose-dependent and especially time-dependent for the clinical isolate of *P. micra* up to 48 hours. The inhibition results of the reference strain were more variable.

*C. acnes* (formerly *Propionibacterium acnes*), was tested as a control in this study. At 48 hours, over 90% of the bacterial



**Figure 2.** Growth inhibition of gram-negative rods in 1.25 to 20 mg/mL of HICA. The activity of increasing concentrations of HICA in thioglycollate broth on the clinical isolate of *A. actinomycetemcomitans* non-serotyped (a), and the reference strains of serotype a (b) and serotype b (c) at pH 6.5 up to 72 hours in anaerobiosis is shown. Chlorhexidine digluconate 0.6 mg/mL was used as a positive control and thioglycollate broth as a negative control.



**Figure 3.** Growth inhibition of gram-positive cocci in 5 to 80 mg/mL of HICA. The activity of increasing concentrations of HICA in thioglycollate broth on the clinical isolate (a) and the reference strain (b) of *P. micra* in anaerobic test conditions at pH 6.5 up to 72 hours is shown. Chlorhexidine digluconate 0.6 mg/mL was used as a positive control and thioglycollate broth as a negative control.


growth of 2 clinical isolates of *C. acnes* was inhibited by  $\geq 80$  mg/mL of HICA (Figure 4a and b), but the growth of the reference strain was markedly inhibited by one dilution higher concentration  $\geq 160$  mg/mL of HICA (Figure 4c). Growth inhibition in the presence of HICA was dose-dependent for 2 reference strains and the clinical isolate of *C. acnes* tested at 48 hours.

## Discussion

The main finding of this study was that HICA showed broad-spectrum antibacterial activity against obligate anaerobic bacteria. All bacterial strains and isolates included were susceptible to HICA at a concentration  $\geq 160$  mg/mL. At

48 hours, the  $IC_{\geq 90}$  results were 1 to 3 dilutions lower for each isolate compared to the MBC at 72 hours. Growth of the reference strain and the clinical isolate of *P. gingivalis* in the negative control was optimal and the growth of the reference strain and the clinical isolate of the *P. micra* was the slightest in this study. HICA was bactericidal against species associated with periodontal disease. In addition, 1 reference strain and 2 clinical isolates of *C. acnes*, which constituted as comparators were killed in the highest test concentration of HICA.

At 72 hours, the MBC of HICA was  $\geq 80$  mg/mL for the reference strain and the clinical isolate of *P. micra*, for the reference strain of *P. gingivalis*, and for the clinical isolate of



**Figure 4.** Growth inhibition of gram-positive rods in 10 to 160 mg/mL of HICA. The activity of increasing concentrations of HICA in thioglycollate broth on 2 clinical isolates (a and b) and the reference strain (c) of *C. acnes* at pH 6.5 in anaerobiosis up to 72 hours is shown. Chlorhexidine digluconate 0.6 mg/mL was used as a positive control and thioglycollate broth as a negative control.

*T. forsythia*. However, the clinical isolate of *P. gingivalis* was susceptible for HICA at  $\geq 20$  mg/mL concentration. The MBC of HICA was  $\geq 40$  mg/mL for the reference strain and the clinical isolate of *F. nucleatum* and for the b serotype reference strain of *A. actinomycetemcomitans*. The reference strain of *A. actinomycetemcomitans* serotype a and the non-serotyped clinical isolate were the most sensitive to HICA, exhibiting an MBC in  $\geq 10$  mg/mL, which was even a 4-fold lower dilution than with the most tolerant reference strains and clinical isolates in this study. In general, gram-positive bacteria showed to be more resistant to HICA than gram-negative bacteria. This may be due to the differences between their cell wall structures. It is possible that HICA may permeate through the cell wall of gram-negative rods more easily than the cell wall with thick peptidoglycan layer of gram-positive bacteria under the test conditions used here.

In comparison to the results of earlier studies the MBC results for the reference strain and the clinical isolate of *F. nucleatum* were similar. The MBC for *F. nucleatum* was earlier shown to be 4.5 mg/mL in pH 5.2.<sup>24</sup> In the present study, they were probably killed in 9 times higher concentration because the growth conditions were more favorable at pH 6.5. The activity of HICA against other obligate anaerobic bacterial species included in this study were tested for the first time. The experimental design showed that the growth of all bacterial strains tested is inhibited and prevented by HICA. This is not necessarily enough evidence to prove its efficacy in clinical conditions. However, the results are promising because HICA, which

is well-tolerated and metabolized by human cells<sup>34,35</sup> might have a potential effect on the dysbiotic polymicrobial biofilm, virulence factors of the corner-stone bacterial species or host responses in periodontitis because of its broad-spectrum antimicrobial activity,<sup>10,24-27,37</sup> the inhibitory activity against microbial biofilms,<sup>28</sup> and proteolytic enzyme activities.<sup>29</sup>

In a clinical study, in which the effect of oral prophylaxis before root planning and scaling was monitored, it showed no improvements in the treatment outcomes.<sup>38</sup> If the living circumstances of dysbiotic microflora is worsened by mechanical treatment and could be aided by a safe antibacterial agent, commensal bacteria may have a better chance to recolonize previously anaerobic oral niches. This is a prerequisite for balancing oral cavity microbiome and healing of periodontitis. New and safe topical antimicrobial agents are needed for the prevention and treatment of superficial and dysbiotic infections to avoid potential adverse effects of presently used antiseptics<sup>33</sup> and antibiotics,<sup>8</sup> as well as the emergence of antibiotic resistance.<sup>39</sup> HICA may prove to be a useful agent in the treatment and prevention of topical infections such as moderately advanced and difficult-to-treat periodontal infections.

This study met the objectives of the study and answered the study question as to whether HICA is active against a range of gram-positive and gram-negative obligately anaerobic bacterial species associated with periodontitis. Clinical studies about the potential efficacy of HICA together with root planning and scaling in the treatment of periodontitis are needed in the future.

## Author Contributions

Conception or design of the work was done by MS, RR-R and TS. MS collected the data. Data analysis and interpretation were performed by MS, RR-R, SS and TS. All authors contributed to drafting and critical revision of the article. All authors approved the final version of the article to be published.

## ORCID iD

Marjut Sakko  <https://orcid.org/0000-0003-2278-3643>

## REFERENCES

- Hajishengallis G, Liang S, Payne MA, et al. Low-abundance biofilm species orchestrates inflammatory periodontal disease through the commensal microbiota and complement. *Cell Host Microbe*. 2011;10:497-506.
- Casadevall A, Pirofski LA. Virulence factors and their mechanisms of action: the view from a damage-response framework. *J Water Health*. 2009;7:S2-S18.
- Lang ML, Zhu L, Kreth J. Keeping the bad bacteria in check: interactions of the host immune system with oral cavity biofilms. *Endod Topics*. 2010;22:17-32.
- Hanes PJ, Purvis JP. Local anti-infective therapy: pharmacological agents. A systematic review. *Ann Periodontol*. 2003;8:79-98.
- Jepsen K, Jepsen S. Antibiotics/antimicrobials: systemic and local administration in the therapy of mild to moderately advanced periodontitis. *Periodontol 2000*. 2016;71:82-112.
- Rovai ES, Souto ML, Ganhito JA, Holzhausen M, Chambrone L, Pannuti CM. Efficacy of local antimicrobials in the non-surgical treatment of patients with periodontitis and diabetes: a systematic review. *J Periodontol*. 2016;87:1406-1417.
- Herrera D, Matesanz P, Martín C, Oud V, Feres M, Teughels W. Adjunctive effect of locally delivered antimicrobials in periodontitis therapy: a systematic review and meta-analysis. *J Clin Periodontol*. 2020;47:239-256.
- Teughels W, Feres M, Oud V, Martín C, Matesanz P, Herrera D. Adjunctive effect of systemic antimicrobials in periodontitis therapy: a systematic review and meta-analysis. *J Clin Periodontol*. 2020;47:257-281.
- Williamson DA, Carter GP, Howden BP. Current and emerging topical antibacterials and antiseptics: agents, action, and resistance patterns. *Clin Microbiol Rev*. 2017;30:827-860.
- Hietala PK, Westermarck HW, Jaarma M. Identification of antimicrobial alpha-hydroxyacids in *Lactobacillus plantarum*-fermented animal protein. *Nutr Metab*. 1979;23:227-234.
- Hummel W, Schütte H, Kula MR. D-2-hydroxyisocaproate dehydrogenase from *Lactobacillus casei*: a new enzyme suitable for stereospecific reduction of 2-ketocarboxylic acids. *Appl Microbiol Biotechnol*. 1985;21:7-15.
- Yamazaki Y, Maeda H. Enzymatic synthesis of optically pure (R)-(-)-mandelic acid and other 2-hydroxycarboxylic acids: screening for the enzyme, and its purification, characterization and use. *Agric Biol Chem*. 1986;50:2621-2631.
- Kallwass HKW. Potential of R-2-hydroxyisocaproate dehydrogenase from *Lactobacillus casei* for stereospecific reductions. *Enzyme Microb Technol*. 1992;14:28-35.
- Bernard N, Johnsen K, Ferain T, et al. NAD<sup>+</sup>-dependent D-2-hydroxyisocaproate dehydrogenase of *Lactobacillus delbrueckii* subsp. *bulgaricus*. *Eur J Biochem*. 1994;224:439-446.
- Niefind K, Hecht HJ, Schomburg D. Crystallization and preliminary characterization of crystals of D-2-hydroxyisocaproate dehydrogenase from *Lactobacillus casei*. *J Mol Biol*. 1994;240:400-402.
- Niefind K, Hecht HJ, Schomburg D. Crystal structure of L-2-hydroxyisocaproate dehydrogenase from *Lactobacillus confusus* at 2.2 Å resolution. An example of strong asymmetry between subunits. *J Mol Biol*. 1995;251:256-281.
- Dengler U, Niefind K, Kiess M, Schomburg D. Crystal structure of a ternary complex of D-2-hydroxyisocaproate dehydrogenase from *Lactobacillus casei*, NAD<sup>+</sup> and 2-oxoisocaproate at 1.9 Å resolution. *J Mol Biol*. 1997;267:640-660.
- Yvon M, Rijnen L. Cheese flavour formation by amino acid catabolism. *Int Dairy J*. 2001;11(4):185-201. doi:10.1016/s0958-6946(01)00049-8
- Broadbent JR, Gummalla S, Hughes JE, Johnson ME, Rankin SA, Drake MA. Overexpression of *Lactobacillus casei* D-hydroxyisocaproic acid dehydrogenase in cheddar cheese. *Appl Environ Microbiol*. 2004;70:4814-4820.
- Smit BA, Engels WJ, Wouters JT, Smit G. Diversity of L-leucine catabolism in various microorganisms involved in dairy fermentations, and identification of the rate-controlling step in the formation of the potent flavour component 3-methylbutanal. *Appl Microbiol Biotechnol*. 2004;64:396-402.
- Smit G, Smit BA, Engels WJ. Flavour formation by lactic acid bacteria and biochemical flavour profiling of cheese products. *FEMS Microbiol Rev*. 2005;29:591-610.
- Chambellon E, Rijnen L, Lorquet F, et al. The D-2-hydroxyacid dehydrogenase incorrectly annotated panE is the sole reduction system for branched-chain 2-keto acids in *Lactococcus lactis*. *J Bacteriol*. 2009;191:873-881.
- Park B, Hwang H, Chang JY, et al. Identification of 2-hydroxyisocaproic acid production in lactic acid bacteria and evaluation of microbial dynamics during kimchi ripening. *Sci Rep*. 2017;7:10904.
- Sakko M, Tjäderhane L, Sorsa T, et al. 2-hydroxyisocaproic acid (HICA): a new potential topical antibacterial agent. *Int J Antimicrob Agents*. 2012;39:539-540.
- Broberg A, Jacobsson K, Ström K, Schnürer J. Metabolite profiles of lactic acid bacteria in grass silage. *Appl Environ Microbiol*. 2007;73:5547-5552.
- Ndagano D, Lamoureux T, Dortu C, Vandermoten S, Thonart P. Antifungal activity of 2 lactic acid bacteria of the *Weissella* genus isolated from food. *J Food Sci*. 2011;76:M305-M311.
- Sakko M, Moore C, Novak-Frazer L, et al. 2-hydroxyisocaproic acid is fungicidal for *Candida* and *Aspergillus* species. *Mycoses*. 2014;57:214-221.
- Nieminen MT, Novak-Frazer L, Rautemaa W, et al. A novel antifungal is active against *Candida albicans* biofilms and inhibits mutagenic acetaldehyde production *in vitro*. *PLoS One*. 2014;9:e101859.
- Nieminen MT, Hernandez M, Novak-Frazer L, et al. DL-2-hydroxyisocaproic acid attenuates inflammatory responses in a murine *Candida albicans* biofilm model. *Clin Vaccine Immunol*. 2014;21:1240-1245.
- EFSA Panel on Food Contact Materials, Enzymes, Flavourings and Processing Aids (CEF). Scientific opinion on flavouring group evaluation 10, revision 3 (FGE.10Rev3): Aliphatic primary and secondary saturated and unsaturated alcohols, aldehydes, acetals, carboxylic acids and esters containing an additional oxygenated functional group and lactones from chemical groups 9, 13 and 30. *EFSA J*. 2012;10:2563.
- Ehling S, Reddy TM. Direct analysis of leucine and its metabolites β-hydroxy-β-methylbutyric acid, α-ketoisocaproic acid, and α-hydroxyisocaproic acid in human breast milk by liquid chromatography-mass spectrometry. *J Agric Food Chem*. 2015;63:7567-7573.
- Selis D, Pande Y, Smoczer C, et al. Cytotoxicity and genotoxicity of a new intracanal medicament, 2-hydroxyisocaproic acid: an *in vitro* study. *J Endod*. 2019;45:578-583.
- Verma UP, Gupta A, Yadav RK, Tiwari R, Sharma R, Balapure AK. Cytotoxicity of chlorhexidine and neem extract on cultured human gingival fibroblasts through fluorescence-activated cell sorting analysis: an *in-vitro* study. *Eur J Dent*. 2018;12:344-349.
- Hoffer LJ, Taveroff A, Robitaille L, Mamer OA, Reimer MLJ. α-Keto and α-Hydroxy branched-chain acid interrelationships in normal humans. *J Nutr*. 1993;123:1513-1521.
- Dallmann R, Viola AU, Tarokh L, Cajochen C, Brown SA. The human circadian metabolome. *Proc Natl Acad Sci U S A*. 2012;109:2625-2629.
- Clinical and Laboratory Standards Institute. *Methods for Antimicrobial Susceptibility Testing of Anaerobic Bacteria; Approved Standard*. 5th ed. Document M11-A5 CLSI; 2001.
- Pahalagedara ASNW, Flint S, Palmer J, Subbaraj A, Brightwell G, Gupta TB. Antimicrobial activity of soil *Clostridium* enriched conditioned media against *Bacillus mycoides*, *Bacillus cereus*, and *Pseudomonas aeruginosa*. *Front Microbiol*. 2020;11:608998.
- Jentsch HFR, Heusinger T, Weickert A, Eick S. Professional tooth cleaning prior to non-surgical periodontal therapy: a randomized clinical trial. *J Periodontol*. 2020;91:174-182.
- Davies J, Davies D. Origins and evolution of antibiotic resistance. *Microbiol Mol Biol Rev*. 2010;74:417-433.