

<https://helda.helsinki.fi>

ESHRE guideline : endometriosis

Endometriosis Guideline Core Grp

2022-03-04

Endometriosis Guideline Core Grp , ESHRE Endometriosis Guideline Grp , Becker , C M , Bokor , A , Heikinheimo , O & Vermeulen , N 2022 , ' ESHRE guideline : endometriosis ' , Human reproduction open , vol. 2022 , no. 2 , 009 . <https://doi.org/10.1093/hropen/hoac009>

<http://hdl.handle.net/10138/343040>

<https://doi.org/10.1093/hropen/hoac009>

cc_by_nc

publishedVersion






Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.


This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

ESHRE guideline: endometriosis[†]

The members of the Endometriosis Guideline Core Group, Christian M. Becker ^{1,*}, Attila Bokor², Oskari Heikinheimo ³, Andrew Horne ⁴, Femke Jansen⁵, Ludwig Kiesel⁶, Kathleen King⁷, Marina Kvaskoff ⁸, Annemiek Nap⁹, Katrine Petersen¹⁰, Ertan Saridogan^{11,12}, Carla Tomassetti^{13,14}, Nehalennia van Hanegem¹⁵, Nicolas Vulliemoz¹⁶, and Nathalie Vermeulen ¹⁷; on behalf of the ESHRE Endometriosis Guideline Group[‡]

¹Nuffield Department of Women's and Reproductive Health, Endometriosis CaRe Centre, University of Oxford, Oxford, UK ²Department of Obstetrics and Gynecology, Semmelweis University, Budapest, Hungary ³Department of Obstetrics & Gynecology, University of Helsinki and Helsinki University Hospital, Helsinki, Finland ⁴EXPPECT Centre for Endometriosis and Pelvic Pain, MRC Centre for Reproductive Health, University of Edinburgh, Edinburgh, UK ⁵EndoHome—Endometriosis Association Belgium, Belgium ⁶Department of Gynecology and Obstetrics, University Hospital Muenster, Muenster, Germany ⁷Individual Endometriosis Advocate, Ireland ⁸Paris-Saclay University, UVSQ, Univ. Paris-Sud, Inserm, Gustave Roussy, "Exposome and Heredity" Team, CESP, Villejuif, France ⁹Department of Gynaecology and Obstetrics, Radboudumc, Nijmegen, The Netherlands ¹⁰Pain Management Centre, UCLH, London, UK ¹¹Department of Obstetrics and Gynaecology, University College London Hospital, London, UK ¹²Elizabeth Garrett Anderson Institute for Women's Health, University College London, London, UK ¹³Department of Obstetrics and Gynaecology, Leuven University Fertility Center, University Hospitals Leuven, Leuven, Belgium ¹⁴Faculty of Medicine, Department of Development and Regeneration, LEERM (Lab of Endometrium, Endometriosis and Reproductive Medicine), KU Leuven, Leuven, Belgium ¹⁵Department of Reproductive Medicine and Gynecology, University Medical Center Utrecht, Utrecht, The Netherlands ¹⁶Department of Woman Mother Child, Fertility Medicine and Gynaecological Endocrinology, Lausanne University Hospital, Lausanne, Switzerland ¹⁷European Society of Human Reproduction and Embryology, Strombeek-Bever, Belgium

*Correspondence address. Nuffield Department of Women's and Reproductive Health, Endometriosis CaRe Centre, John Radcliffe Hospital, University of Oxford, Level 3, Women's Centre, Oxford OX3 9DU, UK. E-mail: christian.becker@wrh.ox.ac.uk; guidelines@eshre.eu  <https://orcid.org/0000-0002-9870-9581>

Submitted on February 10, 2022; editorial decision on February 25, 2022

STUDY QUESTION: How should endometriosis be diagnosed and managed based on the best available evidence from published literature?

SUMMARY ANSWER: The current guideline provides 109 recommendations on diagnosis, treatments for pain and infertility, management of disease recurrence, asymptomatic or extrapelvic disease, endometriosis in adolescents and postmenopausal women, prevention and the association with cancer.

WHAT IS KNOWN ALREADY: Endometriosis is a chronic condition with a plethora of presentations in terms of not only the occurrence of lesions, but also the presence of signs and symptoms. The most important symptoms include pain and infertility.

STUDY DESIGN, SIZE, DURATION: The guideline was developed according to the structured methodology for development of ESHRE guidelines. After formulation of key questions by a group of experts, literature searches and assessments were performed. Papers published up to 1 December 2020 and written in English were included in the literature review.

PARTICIPANTS/MATERIALS, SETTING, METHODS: Based on the collected evidence, recommendations were formulated and discussed within specialist subgroups and then presented to the core guideline development group (GDG) until consensus was reached. A stakeholder review was organized after finalization of the draft. The final version was approved by the GDG and the ESHRE Executive Committee.

MAIN RESULTS AND THE ROLE OF CHANCE: This guideline aims to help clinicians to apply best care for women with endometriosis. Although studies mostly focus on women of reproductive age, the guideline also addresses endometriosis in adolescents and postmenopausal women. The guideline outlines the diagnostic process for endometriosis, which challenges laparoscopy and histology as gold standard diagnostic tests. The options for treatment of endometriosis-associated pain symptoms include analgesics, medical treatments and surgery. Non-pharmacological treatments are also discussed. For management of endometriosis-associated infertility, surgical treatment and/or medically assisted reproduction are feasible. While most of the more recent studies confirm previous ESHRE recommendations, there are five

[†]ESHRE Pages content is not externally peer reviewed. The manuscript has been approved by the Executive Committee of ESHRE.

[‡]The members of the ESHRE Endometriosis Guideline Group are given in the Appendix.

© The Author(s) 2022. Published by Oxford University Press on behalf of European Society of Human Reproduction and Embryology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

topics in which significant changes to recommendations were required and changes in clinical practice are to be expected.

LIMITATIONS, REASONS FOR CAUTION: The guideline describes different management options but, based on existing evidence, no firm recommendations could be formulated on the most appropriate treatments. Also, for specific clinical issues, such as asymptomatic endometriosis or extrapelvic endometriosis, the evidence is too scarce to make evidence-based recommendations.

WIDER IMPLICATIONS OF THE FINDINGS: The guideline provides clinicians with clear advice on best practice in endometriosis care, based on the best evidence currently available. In addition, a list of research recommendations is provided to stimulate further studies in endometriosis.

STUDY FUNDING/COMPETING INTEREST(S): The guideline was developed and funded by ESHRE, covering expenses associated with the guideline meetings, with the literature searches and with the dissemination of the guideline. The guideline group members did not receive payments. C.M.B. reports grants from Bayer Healthcare and the European Commission; Participation on a Data Safety Monitoring Board or Advisory Board with ObsEva (Data Safety Monitoring Group) and Myovant (Scientific Advisory Group). A.B. reports grants from FEMaLE executive board member and European Commission Horizon 2020 grant; consulting fees from Ethicon Endo Surgery, Medtronic; honoraria for lectures from Ethicon; and support for meeting attendance from Gedeon Richter; A.H. reports grants from MRC, NIHR, CSO, Roche Diagnostics, Astra Zeneca, Ferring; Consulting fees from Roche Diagnostics, Nordic Pharma, Chugai and Benevolent AI Bio Limited all paid to the institution; a pending patent on Serum endometriosis biomarker; he is also Chair of TSC for STOP-OHSS and CERM trials. O.H. reports consulting fees and speaker's fees from Gedeon Richter and Bayer AG; support for attending meetings from Gedeon-Richter, and leadership roles at the Finnish Society for Obstetrics and Gynecology and the Nordic federation of the societies of obstetrics and gynecology. L.K. reports consulting fees from Gedeon Richter, AstraZeneca, Novartis, Dr KADE/Besins, Palleos Healthcare, Roche, Mithra; honoraria for lectures from Gedeon Richter, AstraZeneca, Novartis, Dr KADE/Besins, Palleos Healthcare, Roche, Mithra; support for attending meetings from Gedeon Richter, AstraZeneca, Novartis, Dr KADE/Besins, Palleos Healthcare, Roche, Mithra; he also has a leadership role in the German Society of Gynecological Endocrinology (DGGEF). M.K. reports grants from French Foundation for Medical Research (FRM), Australian Ministry of Health, Medical Research Future Fund and French National Cancer Institute; support for meeting attendance from European Society for Gynaecological Endoscopy (ESGE), European Congress on Endometriosis (EEC) and ESHRE; She is an advisory Board Member, FEMaLe Project (Finding Endometriosis Using Machine Learning), Scientific Committee Chair for the French Foundation for Research on Endometriosis and Scientific Committee Chair for the ComPaRe-Endometriosis cohort. A.N. reports grants from Merck SA and Ferring; speaker fees from Merck SA and Ferring; support for meeting attendance from Merck SA; Participation on a Data Safety Monitoring Board or Advisory Board with Nordic Pharma and Merck SA; she also is a board member of medical advisory board, Endometriosis Society, the Netherlands (patients advocacy group) and an executive board member of the World Endometriosis Society. E.S. reports grants from National Institute for Health Research UK, Rosetrees Trust, Barts and the London Charity; Royalties from De Gruyter (book editor); consulting fees from Hologic; speakers fees from Hologic, Johnson & Johnson, Medtronic, Intuitive, Olympus and Karl Storz; Participation in the Medicines for Women's Health Expert Advisory Group with Medicines and Healthcare Products Regulatory Agency (MHRA); he is also Ambassador for the World Endometriosis Society. C.T. reports grants from Merck SA; Consulting fees from Gedeon Richter, Nordic Pharma and Merck SA; speaker fees from Merck SA, all paid to the institution; and support for meeting attendance from Ferring, Gedeon Richter and Merck SA. The other authors have no conflicts of interest to declare.

DISCLAIMER: This guideline represents the views of ESHRE, which were achieved after careful consideration of the scientific evidence available at the time of preparation. In the absence of scientific evidence on certain aspects, a consensus between the relevant ESHRE stakeholders has been obtained.

Adherence to these clinical practice guidelines does not guarantee a successful or specific outcome, nor does it establish a standard of care. Clinical practice guidelines do not replace the need for application of clinical judgement to each individual presentation, nor variations based on locality and facility type.

ESHRE makes no warranty, express or implied, regarding the clinical practice guidelines and specifically excludes any warranties of merchantability and fitness for a particular use or purpose (Full disclaimer available at www.eshre.eu/guidelines).

Key words: endometriosis / guideline / fertility / pelvic pain / adolescent / surgery / ESHRE guideline

WHAT DOES THIS MEAN FOR PATIENTS?

Endometriosis is a chronic condition with a large impact not only on the patient's quality of life, but also on social contacts and work. Endometriosis is characterized mainly by symptoms of pain (often linked to the menstrual cycle) and infertility. Often, endometriosis is subdivided according to the type and location of the lesions into peritoneal endometriosis, deep endometriosis and endometrioma.

The current paper summarizes the ESHRE guideline on endometriosis, providing clinicians with evidence-based recommendations on the diagnosis and management of endometriosis-associated symptoms, including medical treatment, surgery and assisted reproduction. Information and recommendations are also provided on other topics related to endometriosis, such as prevention, pregnancy and cancer.

Introduction

Endometriosis is a disease characterized by the presence of endometrium-like epithelium and/or stroma outside the endometrium and myometrium, usually with an associated inflammatory process (International Working Group of AAGL, ESGE, ESHRE and WES *et al.*, 2021). The exact prevalence of endometriosis is unknown, but estimates range from 2% to 10% within the general female population and up to 50% in infertile women (Eskenazi and Warner, 1997; Meuleman *et al.*, 2009; Zondervan *et al.*, 2020).

The ESHRE Guideline for the Diagnosis and Treatment of Endometriosis (2005) and the ESHRE Guideline: Management of women with endometriosis (2013) have been a reference point for best clinical care in endometriosis for years (Kennedy *et al.*, 2005; Dunselman *et al.*, 2014). Based on continuous new research and developments, it was considered that the last recommendations formulated in 2013/2014 required a revision.

Materials and methods

The guideline was developed according to a well-documented methodology that is universal to ESHRE guidelines (Vermeulen *et al.*, 2019). The core guideline development group (GDG) was composed of past members of the guideline group from 2013 and additional experts selected from applicants to a call for experts. All other European experts applying to the call were included as subgroup members, assisting a core group member preparing the guideline on a certain topic. The GDG included two patient representatives, and five patient organizations were represented in the subgroups.

Forty-two key questions were formulated by the GDG, of which seven were answered as narrative questions, and 35 as PICO (Patient, Intervention, Comparison, Outcome) questions. For each PICO question, databases (PubMed/MEDLINE and the Cochrane library) were searched from inception to 1 December 2020, limited to studies written


in English. From the literature searches, studies were selected based on the PICO questions, assessed for quality and summarized in evidence tables. GDG subgroup meetings were organized, face-to-face and online, for presentation and discussion of the evidence and draft recommendations by the assigned core group member. Proposed recommendations by the subgroups were then discussed in core group meetings until a consensus was reached. Each recommendation was labelled as strong or weak and a grade was assigned based on the strength of the supporting evidence (High ⊕⊕⊕⊕, Moderate ⊕⊕⊕○, Low ⊕⊕○○ and Very low ⊕○○○). Good practice points (GPPs) based on clinical expertise were added where relevant to clarify the recommendations or to provide further practical advice. 'Research only' recommendations were also made, and those interventions should be applied only within the context of research, with appropriate precautions and ethical approval.

Strong recommendations should be used as a recommendation to be applied for most patients, while weak recommendations require discussion and shared decision-making (Fig. 1).

For the narrative questions, a similar literature search was conducted. Collected data were summarized in a narrative summary and conclusions were formulated. In case of insufficient data to provide recommendations in reply to a PICO question, a conclusion was also added. For clarity, these conclusions are labelled 'conclusion, not recommendation' in the current paper.

The guideline draft and an invitation to participate in the stakeholder review were published on the ESHRE website between 24 June and 15 August 2021. All comments were processed by the core group, either by adapting the content of the guideline and/or by replying to the reviewer. The review process was summarized in the review report, which is published on the ESHRE website (www.eshre.eu/Guidelines). Overall, 56.5% of the 253 comments resulted in an adaptation or correction in the guideline text.

This guideline will be considered for update 4 years after publication, with an intermediate assessment of the need for updating 2 years after publication.



Results

Key questions and recommendations

The scope of the ESHRE guideline on endometriosis is to provide guidance on the management of endometriosis; either diagnosed or strongly suspected. In line with endometriosis research, terminology and discussion, the guideline is focused on females and menstruation. The GDG recognizes that there are individuals living with endometriosis who are transgender, who do not menstruate, who do not have a uterus or who do not identify with the terms used in the literature. Throughout, the term 'women with endometriosis' is used, but this is not intended to isolate, exclude or diminish any individual's experience nor to discriminate against any group.

Diagnosis of endometriosis

The recommended diagnostic process for endometriosis is summarized in Fig. 2.

Can clinical symptoms predict the presence of endometriosis?

The GDG recommends that clinicians should consider the diagnosis of endometriosis in individuals presenting with the following cyclical and non-cyclical signs and symptoms:

dysmenorrhoea, deep dyspareunia, dysuria, dyschezia, painful rectal bleeding or haematuria, shoulder tip pain, catamenial pneumothorax, cyclical cough/haemoptysis/chest pain, cyclical scar swelling and pain, fatigue and infertility (Forman et al., 1993; Eskenazi et al., 2001; Ballard et al., 2008; Nnoaham et al., 2012).

GPP

Does the use of symptom diaries or questionnaires compared to traditional history taking lead to improved or earlier diagnosis of endometriosis?

As no recommendation could be made, the following conclusion was formulated. Although currently no evidence exists that a symptom diary/questionnaire/app reduces the time to diagnosis or leads to earlier diagnosis, the GDG considers their potential benefit in complementing the traditional history taking process as it aids in objectifying pain and empowering women to demonstrate their symptoms (conclusion, not recommendation).

Does clinical examination of symptomatic women reliably predict the presence of endometriosis?

Clinical examination, including vaginal examination where appropriate, should be considered to identify deep nodules or endometriomas in patients with suspected endometriosis, although the diagnostic accuracy is low (Ripps and Martin, 1992; Nezhad et al., 1994; Koninckx et al., 1996; Eskenazi et al., 2001; Chapron et al., 2002; Condous et al., 2007; Bazot et al., 2009; Khawaja et al., 2009; Hudelist et al., 2011; Paulson and Paulson, 2011).

In women with suspected endometriosis, further diagnostic steps, including imaging, should be considered even if the clinical examination is normal.

Strong recommendation
⊕○○○

Strong recommendation
⊕○○○

Are medical technologies reliable in diagnosing endometriosis and establishing the extent of the disease?

Clinicians should not use measurement of biomarkers in endometrial tissue, blood, menstrual or uterine fluids to diagnose endometriosis (Mol et al., 1998; May et al., 2010; May et al., 2011; Liu et al., 2015; Cosar et al., 2016; Gupta et al., 2016; Hirsch et al., 2016; Nisenblat et al., 2016a; Vanhie et al., 2019; Moustafa et al., 2020).

Strong recommendation
⊕⊕⊕○

Clinicians are recommended to use imaging (ultrasound (US) or MRI) in the diagnostic work-up for endometriosis, but they need to be aware that a negative finding does not exclude endometriosis, particularly superficial peritoneal disease (Bazot et al., 2009; Manganaro et al., 2012; Guerriero et al., 2014; Thomeer et al., 2014; Nisenblat et al., 2016b; Moura et al., 2019).

Strong recommendation
⊕⊕○○

In patients with negative imaging results or where empirical treatment was unsuccessful or inappropriate, the GDG recommends that clinicians consider offering laparoscopy for the diagnosis and treatment of suspected endometriosis.

GPP

The GDG recommends that laparoscopic identification of endometriotic lesions is confirmed by histology although negative histology does not entirely rule out the disease.

GPP

Does diagnostic laparoscopy compared to empirical medical treatment result in better symptom management in women suspected of endometriosis?

As there is no evidence of superiority of either approach (Chapron et al., 1998; Byrne et al., 2018; Bafort et al., 2020), the GDG concluded that both diagnostic laparoscopy and imaging combined with empirical treatment (hormonal contraceptives or progestogens) can be considered in women suspected of endometriosis. Pros and cons should be discussed with the patient (conclusion, not recommendation).

Is long-term monitoring of women with endometriosis beneficial in preventing adverse outcomes (recurrence, complications, malignancy)?

Follow-up and psychological support should be considered in women with confirmed endometriosis, particularly deep and ovarian endometriosis, although there is currently no evidence of benefit of regular long-term monitoring for early detection of recurrence, complications, or malignancy (Pittaway, 1990; Matalliotakis et al., 1994; Chen et al., 1998).

Weak recommendation
⊕○○○

The appropriate frequency and type of follow-up or monitoring is unknown and should be individualized based on previous and current treatments, and severity of the disease and symptoms.

GPP

Does early diagnosis of endometriosis versus late diagnosis lead to better quality of life?

Although no adequate studies exist to support the benefits of early versus late diagnosis, the GDG recommends that in symptomatic women,

DIAGNOSIS OF ENDOMETRIOSIS


SIGNS AND SYMPTOMS

Consider Endometriosis when the woman reports one or more of these symptoms

- | | |
|-------------------------|--|
| Dysmenorrhoea | Shoulder tip pain |
| Deep dyspareunia | Catamenial pneumothorax |
| Dysuria | Cyclical cough/haemoptysis /chest pain |
| Dyschezia | Cyclical scar swelling and pain |
| Painful Rectal bleeding | Fatigue |
| Haematuria | Infertility |

A symptom diary or app can be helpful in the history taking process

Explore a diagnosis of endometriosis



¹ EMPIRICAL TREATMENT = Combined hormonal contraceptives or Progestogens
² Be aware that negative histology does not rule out endometriosis



Figure 2. The recommended diagnostic process for endometriosis. DE, deep endometriosis; US, ultrasound.

attempts should be made to relieve symptoms, either by empirical treatment or after a diagnosis of endometriosis (*conclusion, not recommendation*).

Treatment of endometriosis-associated pain

The recommendations for treatment of pain symptoms linked to endometriosis are summarized in Fig. 3.

Are analgesics effective for symptomatic relief of painful symptoms associated with endometriosis?

Women may be offered non-steroidal anti-inflammatory drugs (NSAIDs) or other analgesics (either alone or in combination with other treatments) to reduce endometriosis-associated pain (Brown et al., 2017).

Weak
recommendation
⊕○○○

Are hormone therapies effective for painful symptoms associated with endometriosis?

It is recommended to offer women hormone treatment (combined hormonal contraceptives, progestogens, GnRH agonists or GnRH antagonists) as one of the options to reduce endometriosis-associated pain.

Strong
recommendation
⊕⊕⊕○

The GDG recommends that clinicians take a shared decision-making approach and take individual preferences, side effects, individual efficacy, costs and availability into consideration when choosing hormone treatments for endometriosis-associated pain.

GPP

It is recommended to prescribe women a combined hormonal contraceptive (oral, vaginal ring or transdermal) to reduce endometriosis-associated dyspareunia, dysmenorrhoea and non-menstrual pain (Brown et al., 2018; Jensen et al., 2018; Grandi et al., 2019).

Strong
recommendation
⊕⊕○○

Women suffering from endometriosis-associated dysmenorrhoea can be offered the continuous use of a combined hormonal contraceptive pill (Hee et al., 2013; Zorbas et al., 2015; Muzii et al., 2016b).

Weak
recommendation
⊕⊕○○

It is recommended to prescribe women progestogens to reduce endometriosis-associated pain (Momoeda et al., 2009; Brown et al., 2012; Petraglia et al., 2012; Andres et al., 2015; Dragoman and Gaffield, 2016).

Strong
recommendation
⊕⊕○○

The GDG recommends that clinicians take the different side effect profiles of progestogens into account when prescribing them.

GPP

It is recommended to prescribe women a levonorgestrel-releasing intrauterine (LNG-IUS) system or an etonogestrel-releasing subdermal implant to reduce endometriosis-associated pain (Lan et al., 2013; Margatho et al., 2020).

Strong
recommendation
⊕⊕⊕○

It is recommended to prescribe women GnRH agonists to reduce endometriosis-associated pain, although evidence is limited regarding dosage or duration of treatment (Brown et al., 2010; Tang et al., 2017).

Strong
recommendation
⊕⊕○○

The GDG recommends that GnRH agonists are prescribed as second-line (e.g. if hormonal contraceptives or progestogens have been ineffective) due to their side effect profile.

GPP

Clinicians should consider prescribing combined hormonal add-back therapy alongside GnRH agonist therapy to prevent bone loss and hypo-oestrogenic symptoms (Wu et al., 2014; Sauerbrun-Cutler and Alvero, 2019).

Strong
recommendation
⊕⊕⊕○

(continued)

It can be considered to prescribe women GnRH antagonists to reduce endometriosis-associated pain, although evidence is limited regarding dosage or duration of treatment (Taylor et al., 2017; Donnez et al., 2020; Osuga et al., 2021).

Weak
recommendation
⊕⊕⊕○

The GDG recommends that GnRH antagonists are prescribed as second-line (e.g. if hormonal contraceptives or progestogens have been ineffective) owing to their side effect profile.

GPP

In women with endometriosis-associated pain refractory to other medical or surgical treatment, it is recommended to prescribe aromatase inhibitors, as they reduce endometriosis-associated pain. Aromatase inhibitors may be prescribed in combination with oral contraceptives, progestogens, GnRH agonists or GnRH antagonists (Ferrero et al., 2011; Almassinokiani et al., 2014; Agarwal and Foster, 2015).

Strong
recommendation
⊕⊕○○

Is surgery effective for treatment of pain associated with endometriosis?

It is recommended to offer surgery as one of the options to reduce endometriosis-associated pain (Sutton et al., 1994; Franck et al., 2018; Arcoverde et al., 2019; Bafort et al., 2020).

Strong
recommendation
⊕⊕○○

When surgery is performed, clinicians may consider excision instead of ablation of endometriosis to reduce endometriosis-associated pain (Wright et al., 2005; Healey et al., 2014; Pundir et al., 2017).

Weak
recommendation
⊕⊕○○

It can be concluded that laparoscopic uterosacral nerve ablation (LUNA) is not beneficial as an additional procedure to conventional laparoscopic surgery for endometriosis, as it offers no additional benefit over surgery alone (Proctor et al., 2005). Presacral neurectomy (PSN) is beneficial for treatment of endometriosis-associated midline pain as an adjunct to conventional laparoscopic surgery, but it should be stressed that PSN requires a high degree of skill and is associated with an increased risk of adverse effects such as intraoperative bleeding, and postoperative constipation, urinary urgency and painless first stage of labour (Miller et al., 2020) (*conclusion, not recommendation*).

When performing surgery in women with ovarian endometrioma, clinicians should perform cystectomy instead of drainage and coagulation, as cystectomy reduces recurrence of endometrioma and endometriosis-associated pain (Hart et al., 2008; Candiani et al., 2020).

Strong
recommendation
⊕⊕○○

When performing surgery in women with ovarian endometrioma, clinicians can consider both cystectomy and CO₂ laser vaporization, as both techniques appear to have similar recurrence rates beyond the first year after surgery. Early post-surgical recurrence rates may be lower after cystectomy (Muzii et al., 2005, 2016a; Mossa et al., 2010; Porpora et al., 2010; Carmona et al., 2011; Shaltout et al., 2019).


Weak
recommendation
⊕○○○

When performing surgery for ovarian endometrioma, specific caution should be used to minimize ovarian damage (Busacca et al., 2006; Muzii et al., 2015; Muzii et al., 2016a; Shaltout et al., 2019; Younis et al., 2019).

Strong
recommendation
⊕○○○

(continued)

TREATMENTS FOR ENDOMETRIOSIS



NON-PHARMACOLOGICAL TREATMENTS FOR PAIN?

*Discuss non-medical strategies to address quality of life and well-being.
No recommendation can be made for a specific intervention*



Figure 3. Summary of the recommendations for treatment of pain symptoms linked to endometriosis. NSAID, non-steroidal anti-inflammatory.

Clinicians can consider performing surgical removal of deep endometriosis, as it may reduce endometriosis-associated pain and improves quality of life (Stepniowska et al., 2010; De Cicco et al., 2011; Meuleman et al., 2011; Byrne et al., 2018; Arcoverde et al., 2019; Bendifallah et al., 2020).

The GDG recommends that women with deep endometriosis are referred to a centre of expertise.

The GDG recommends that patients undergoing surgery, particularly for deep endometriosis, are informed of potential risks, benefits and long-term effect on quality of life.

Weak
recommendation
⊕⊕○○

GPP

GPP

Owing to the heterogeneity of patient populations, surgical approaches, preferences and techniques, the GDG decided not to make any conclusions or recommendations on the techniques to be applied for treatment of pain associated with deep endometriosis (*conclusion, not recommendation*).

Is there a subgroup of women with confirmed endometriosis who respond better to surgery than others?

Clinicians can consider hysterectomy (with or without removal of the ovaries) with removal of all visible endometriosis lesions, in those women who no longer wish to conceive and failed to respond to more conservative treatments. Women should be informed that hysterectomy will not necessarily cure the symptoms or the disease.

When a decision is made whether to remove the ovaries, the long-term consequences of early menopause and possible need for hormone replacement therapy should be considered.

The GDG recommends that when hysterectomy is performed, a total hysterectomy is preferred (Namnoum et al., 1995; Sandström et al., 2020; Shakiba et al., 2008).

Weak
recommendation
⊕⊕○○

GPP

GPP

There are currently no prognostic markers that can be used to select patients that would benefit from surgery. Such markers would need to be assessed prior to surgery and predict a clinically meaningful improvement of pain symptoms. In the absence of prognostic markers, no recommendation could be formulated (*conclusion, not recommendation*).

Are medical therapies effective as an adjunct to surgical therapy?

It is not recommended to prescribe preoperative hormone treatment to improve the immediate outcome of surgery for pain in women with endometriosis (Chen et al., 2020).

Women may be offered postoperative hormone treatment to improve the immediate outcome of surgery for pain in women with endometriosis if not desiring immediate pregnancy (Tanmahasamut et al., 2012; Chen et al., 2020).

Strong
recommendation
⊕⊕○○

Weak
recommendation
⊕⊕○○

Are surgical therapies more effective than medical therapies for women with endometriosis with pain symptoms?

The GDG recommends that clinicians take a shared decision-making approach and take individual preferences, side effects, individual efficacy, costs and availability into consideration when choosing between hormone treatments and surgical treatments for endometriosis-associated pain.

GPP

What non-medical management strategies are effective for symptoms associated with endometriosis (pain and quality of life)?

The GDG recommends that clinicians discuss non-medical strategies to address quality of life and psychological well-being of women managing symptoms of endometriosis. However, no recommendations can be made for any specific non-medical intervention (Chinese medicine, nutrition, electrotherapy, acupuncture, physiotherapy, exercise and psychological interventions) to reduce pain or improve quality of life measures in women with endometriosis, as the potential benefits and harms are unclear.

GPP

Treatment of endometriosis-associated infertility

The recommendations for treatment of endometriosis-associated infertility are summarized in Fig. 4.

Are hormone/medical therapies effective for treatment of endometriosis-associated infertility?

In infertile women with endometriosis, clinicians should not prescribe ovarian suppression treatment to improve fertility (Hughes et al., 2007).

Strong
recommendation
⊕⊕○○

Women seeking pregnancy should not be prescribed postoperative hormone suppression with the sole purpose to enhance future pregnancy rates (Chen et al., 2020).

Strong
recommendation
⊕⊕○○

Those women who cannot attempt to or decide not to conceive immediately after surgery may be offered hormone therapy as it does not negatively impact their fertility and improves the immediate outcome of surgery for pain (Chen et al., 2020).

Weak
recommendation
⊕⊕○○

In infertile women with endometriosis, clinicians should not prescribe pentoxifylline, other anti-inflammatory drugs or letrozole outside ovulation-induction to improve natural pregnancy rates (Alborzi et al., 2011; Lu et al., 2012).

Strong
recommendation
⊕○○○

TREATMENTS FOR ENDOMETRIOSIS




Figure 4. Summary of the recommendations on treatment of endometriosis-associated infertility. EFI, endometriosis fertility index; MAR, medically assisted reproduction.

In women with endometriosis, is surgery effective to increase the chance of natural pregnancy?

Operative laparoscopy could be offered as a treatment option for endometriosis-associated infertility in revised American Society for Reproductive Medicine (rASRM) stage I/II endometriosis as it improves the rate of ongoing pregnancy (Jin and Ruiz Beguerie, 2014; Bafort et al., 2020; Hodgson et al., 2020).

Weak recommendation
⊕⊕○○

Clinicians may consider operative laparoscopy for the treatment of endometrioma-associated infertility as it may increase their chance of natural pregnancy, although no data from comparative studies exist (Dan and Limin, 2013; Alborzi et al., 2019; Candiani et al., 2020).

Weak recommendation
⊕○○○

Although no compelling evidence exists that operative laparoscopy for deep endometriosis improves fertility, operative laparoscopy may represent a treatment option in symptomatic patients wishing to conceive (Meuleman et al., 2011; Vercellini et al., 2012; Iversen et al., 2017).

Weak recommendation
⊕○○○

The GDG recommends that the decision to perform surgery should be guided by the presence or absence of pain symptoms, patient age and preferences, history of previous surgery, presence of other infertility factors, ovarian reserve and the estimated endometriosis fertility index (EFI).

GPP

Which patients need treatment with assisted reproduction technology after surgery?

While no recommendation could be formulated, the GDG concluded that women should be counselled of their chances of becoming pregnant after surgery. To identify patients that may benefit from ART after surgery, the EFI should be used as it is validated, reproducible and cost-effective. The results of other fertility investigations, such as their partner's sperm analysis, should be taken into account (conclusion, not recommendation).

Is medically assisted reproduction effective for infertility associated with endometriosis?

In infertile women with rASRM stage I/II endometriosis, clinicians may perform IUI with ovarian stimulation, instead of expectant management or IUI alone, as it increases pregnancy rates (Nulsen et al., 1993; Tummon et al., 1997; Omland et al., 1998).

Weak recommendation
⊕○○○

Although the value of IUI in infertile women with rASRM stage III/IV endometriosis with tubal patency is uncertain, the use of IUI with ovarian stimulation could be considered (van der Houwen et al., 2014).

Weak recommendation
⊕○○○

ART can be performed for infertility associated with endometriosis, especially if tubal function is compromised, if there is male factor infertility, in case of low EFI and/or if other treatments have failed (Harb et al., 2013; Hamdan et al., 2015b; Senapati et al., 2016; Murta et al., 2018; Muteshi et al., 2018; Alshehre et al., 2021).

Weak recommendation
⊕⊕○○

A specific protocol for ART in women with endometriosis cannot be recommended. Both GnRH antagonist and agonist protocols can be offered based on patients' and physicians' preferences as no difference in pregnancy or live birth rate has been demonstrated (Pabuccu et al., 2007; Rodriguez-Purata et al., 2013; Bastu et al., 2014; Kolanska et al., 2017; Drakopoulos et al., 2018).

Weak recommendation
⊕○○○

Women with endometriosis can be reassured regarding the safety of ART since the recurrence rates are not increased compared to those women not undergoing ART (Benaglia et al., 2008; Somigliana et al., 2019).

Weak recommendation
⊕⊕⊕○

In women with endometrioma, clinicians may use antibiotic prophylaxis at the time of oocyte retrieval, although the risk of ovarian abscess formation following follicle aspiration is low.

GPP

Are medical therapies effective as an adjunct to medically assisted reproduction for endometriosis-associated infertility?

The extended administration of GnRH agonist prior to ART treatment to improve live birth rate in infertile women with endometriosis is not recommended, as the benefit is uncertain (Georgiou et al., 2019; Cao et al., 2020; Kaponis et al., 2020).

Strong recommendation
⊕○○○

There is insufficient evidence to recommend prolonged administration of the combined oral contraceptives (COC)/progestogens as a pre-treatment to ART to increase live birth rates (de Ziegler et al., 2010).

Weak recommendation
⊕○○○

Are surgical therapies effective as an adjunct prior to medically assisted reproduction for endometriosis-associated infertility?

Clinicians are not recommended to routinely perform surgery prior to ART to improve live birth rates in women with rASRM stage I/II endometriosis, as the potential benefits are unclear (Opoien et al., 2011; Hamdan et al., 2015b).

Strong recommendation
⊕⊕○○

Clinicians are not recommended to routinely perform surgery for ovarian endometrioma prior to ART to improve live birth rates, as the current evidence shows no benefit and surgery is likely to have a negative impact on ovarian reserve (Coccia et al., 2014; Hamdan et al., 2015a; Nickkho-Amiry et al., 2018; Şükür et al., 2021).

Strong recommendation
⊕⊕○○

Surgery for endometrioma prior to ART can be considered to improve endometriosis-associated pain or accessibility of follicles.

GPP

The decision to offer surgical excision of deep endometriosis lesions prior to ART should be guided mainly by pain symptoms and patient preference as its effectiveness on reproductive outcome is uncertain owing to lack of randomized studies (Bianchi et al., 2009; Soriano et al., 2016; Bendifallah et al., 2017; Breteau et al., 2020).

Strong recommendation
⊕○○○

What non-medical management strategies are effective for infertility associated with endometriosis?

Regarding non-medical strategies on infertility, there is no clear evidence that any non-medical interventions for women with endometriosis will be of benefit to increase the chance of pregnancy. No recommendation can be made to support any non-medical interventions (nutrition, Chinese medicine, electrotherapy, acupuncture, physiotherapy, exercise and psychological interventions) to increase fertility in women with endometriosis. The potential benefits and harms are unclear (conclusion, not recommendation).

(continued)

Is endometriosis an indication for fertility preservation (ovarian tissue/oocytes)?

In case of extensive ovarian endometriosis, clinicians should discuss the pros and cons of fertility preservation with women with endometriosis. The true benefit of fertility preservation in women with endometriosis remains unknown (Cobo *et al.*, 2020; Kim *et al.*, 2020).

Strong recommendation
⊕○○○

What is the impact of endometriosis on pregnancy and obstetric outcomes?

Patients should not be advised to become pregnant with the sole purpose of treating endometriosis, as pregnancy does not always lead to improvement of symptoms or reduction of disease progression (Leeners *et al.*, 2018).

Strong recommendation
⊕○○○

Endometriomas may change in appearance during pregnancy. In case of finding an atypical endometrioma during US in pregnancy, it is recommended to refer the patient to a centre with appropriate expertise (Leone Roberti Maggiore *et al.*, 2016).

Strong recommendation
⊕○○○

Complications related directly to pre-existing endometriosis lesions are rare, but probably under-reported. Such complications may be related to their decidualization, adhesion formation/stretching and endometriosis-related chronic inflammation. Although rare, they may represent life-threatening situations that may require surgical management (Leone Roberti Maggiore *et al.*, 2016; Leone Roberti Maggiore *et al.*, 2017; Lier *et al.*, 2017; Glavind *et al.*, 2018).

Clinicians should be aware that there may be an increased risk of first trimester miscarriage and ectopic pregnancy in women with endometriosis (Leone Roberti Maggiore *et al.*, 2016; Santulli *et al.*, 2016; Kohl Schwartz *et al.*, 2017; Saraswat *et al.*, 2017; Yong *et al.*, 2020).

Strong recommendation
⊕⊕○○

Clinicians should be aware of endometriosis-associated complications in pregnancy, although these are rare. As these findings are based on low/moderate quality studies, these results should be interpreted with caution and currently do not warrant increased antenatal monitoring or dissuade women from becoming pregnant (Leone Roberti Maggiore *et al.*, 2016; Lalani *et al.*, 2018; Perez-Lopez *et al.*, 2018; Horton *et al.*, 2019).

Strong recommendation
⊕⊕○○

The recommendations and information on endometriosis and pregnancy are summarized in Fig. 5.

Endometriosis recurrence

Is there a role for secondary prevention of recurrence of disease and painful symptoms in patients treated for endometriosis?

When surgery is indicated in women with an endometrioma, clinicians should perform ovarian cystectomy, instead of drainage and electrocoagulation, for the secondary prevention of endometriosis-associated dysmenorrhoea, dyspareunia and non-menstrual pelvic pain. However, the risk of reduced ovarian reserve should be taken into account.

Strong recommendation
⊕⊕○○

Clinicians should consider prescribing the postoperative use of a LNG-IUS system (52 mg) or a combined hormonal contraceptive for at least 18–24 months for the secondary prevention of endometriosis-associated dysmenorrhoea (Seracchioli *et al.*, 2009; Lee *et al.*, 2018; Song *et al.*, 2018; Chen *et al.*, 2020; Zakhari *et al.*, 2021).

Strong recommendation
⊕⊕○○

After surgical management of ovarian endometrioma in women not immediately seeking conception, clinicians are recommended to offer long-term hormone treatment (e.g. combined hormonal contraceptives) for the secondary prevention of endometrioma and endometriosis-associated related symptom recurrence.

Strong recommendation
⊕○○○

For the prevention of recurrence of deep endometriosis and associated symptoms, long-term administration of postoperative hormone treatment can be considered.

Weak recommendation
⊕○○○

Clinicians can perform ART in women with deep endometriosis, as it does not seem to increase endometriosis recurrence per se (Somigliana *et al.*, 2019).

Weak recommendation
⊕⊕⊕○

How should patients with reoccurring endometriosis or recurring symptoms be managed? Is repetitive surgery effective for symptoms associated with endometriosis?

Any hormone treatment or surgery can be offered to treat recurring pain symptoms in women with endometriosis (Candiani *et al.*, 1991; Hornstein *et al.*, 1997; Vercellini *et al.*, 2002; Razzi *et al.*, 2007; Muzii *et al.*, 2015; Abdou *et al.*, 2018; Koshiba *et al.*, 2018; Lee *et al.*, 2018).

Weak recommendation
⊕○○○

Endometriosis and adolescence

Which diagnostic procedures should be applied in adolescents with possible endometriosis?

In adolescents, clinicians should take a careful history to identify possible risk factors for endometriosis, such as a positive family history, obstructive genital malformations, early menarche or short menstrual cycle (Geysenbergh *et al.*, 2017).

Strong recommendation
⊕○○○

Clinicians may consider endometriosis in young women presenting with (cyclical) absenteeism from school, or with use of oral contraceptives for treatment of dysmenorrhoea (Chapron *et al.*, 2011).

Weak recommendation
⊕○○○

In adolescents, clinicians should take a careful history and consider the following symptoms as suggestive of the presence of endometriosis:

chronic or acyclical pelvic pain, particularly combined with nausea, dysmenorrhoea, dyschezia, dysuria, dyspareunia;
cyclical pelvic pain (Greene *et al.*, 2009; Treloar *et al.*, 2010; Vicino *et al.*, 2010; Yang *et al.*, 2012; DiVasta *et al.*, 2018).

Strong recommendation
⊕○○○

In the absence of evidence for adolescents specifically, the recommendations for clinical examination in adults can be applied.

(continued)

ENDOMETRIOSIS AND PREGNANCY




Figure 5. Summary of the recommendations and information on endometriosis and pregnancy.

ENDOMETRIOSIS AND CANCER

Absolute risk of developing cancer in a woman's lifetime




Figure 6. Infographic on the absolute risk of developing cancer in a woman's lifetime.

The GDG recommends that before performing vaginal examination and/or rectal examination in adolescents, the acceptability should be discussed with the adolescent and her caregiver, taking into consideration the patient's age and cultural background.	GPP
Transvaginal US is recommended to be used in adolescents in whom it is appropriate, as it is effective in diagnosing ovarian endometriosis. If a transvaginal scan is not appropriate, MRI, transabdominal, transperineal or transrectal scan may be considered (Yang et al., 2012; Brosens et al., 2013; Martire et al., 2020).	Strong recommendation ⊕⊕○○
Serum biomarkers (e.g. CA-125) are not recommended for diagnosing or ruling out endometriosis in adolescents (Seckin et al., 2018; Sasamoto et al., 2020).	Strong recommendation ⊕⊕○○
In adolescents with suspected endometriosis where imaging is negative and medical treatments (with NSAIDs and/or hormonal contraceptives) have not been successful, diagnostic laparoscopy may be considered (Vicino et al., 2010; Shah and Missmer, 2011; Yang et al., 2012; Brosens et al., 2013).	Weak recommendation ⊕○○○

Should diagnosis of endometriosis in adolescents be confirmed by histology?

If a laparoscopy is performed, clinicians should consider taking biopsies to confirm the diagnosis histologically, although negative histology does not entirely rule out the disease (Janssen et al., 2013).	Strong recommendation ⊕⊕○○
---	-------------------------------

What is the best treatment for adolescents with (suspected) endometriosis?

In adolescents with severe dysmenorrhoea and/or endometriosis-associated pain, clinicians should prescribe hormonal contraceptives or progestogens (systemically or via LNG-IUS) as first-line hormonal hormone therapy because they may be effective and safe. However, it is important to note that some progestogens may decrease bone mineral density (Davis et al., 2005; Yoost et al., 2013; Ebert et al., 2017).	Strong recommendation ⊕○○○
The GDG recommends clinicians consider NSAIDs as treatment for endometriosis-associated pain in adolescents with (suspected) endometriosis, especially if first-line hormone treatment is not an option.	GPP
In adolescents with laparoscopically confirmed endometriosis and associated pain in whom hormonal contraceptives or progestogen therapy failed, clinicians may consider prescribing GnRH agonists for up to 1 year, as they are effective and safe when combined with add-back therapy (DiVasta et al., 2015; Gallagher et al., 2017; Gallagher et al., 2018).	Weak recommendation ⊕⊕○○
The GDG recommends that in young women and adolescents, if GnRH agonist treatment is considered, it should be used only after careful consideration and discussion of potential side effects and potential long-term health risks with a practitioner in a secondary or tertiary care setting.	GPP

(continued)

In adolescents with endometriosis, clinicians may consider surgical removal of endometriosis lesions to manage endometriosis-related symptoms. However, symptom recurrence rates may be considerable, especially when surgery is not followed by hormone treatment (Roman, 2010; Tandoi et al., 2011; Yeung et al., 2011; Lee et al., 2017).	Weak recommendation ⊕○○○
The GDG recommends that if surgical treatment is indicated in adolescents with endometriosis, it should be performed laparoscopically by an experienced surgeon, and, if possible, complete laparoscopic removal of all present endometriosis should be performed.	GPP
In adolescents with endometriosis, clinicians should consider postoperative hormone therapy, as this may suppress recurrence of symptoms (Doyle et al., 2009; Seo et al., 2017).	Strong recommendation ⊕○○○

Is endometriosis in adolescents an indication for fertility preservation (ovarian tissue/oocytes)?

The GDG recommends that adolescents with endometriosis are informed of the potential detrimental effect of ovarian endometriosis and surgery on ovarian reserve and future fertility.	GPP
Fertility preservation options exist and the GDG recommends that adolescents are informed about them, although the true benefit, safety and indications in adolescents with endometriosis remain unknown.	GPP

Endometriosis and menopause

Is endometriosis still active during menopause and, if so, how should the symptoms be treated?

The GDG concluded that clinicians should be aware that endometriosis can still be active/symptomatic after menopause (conclusion, not recommendation).

Is surgical/medical treatment effective and safe in women with a history of endometriosis?

Clinicians may consider surgical treatment for postmenopausal women presenting with signs of endometriosis and/or pain to enable histological confirmation of the diagnosis of endometriosis (Redwine, 1994; Clayton et al., 1999; Morotti et al., 2012; Sun et al., 2013; Behera et al., 2006).	Weak recommendation ⊕○○○
The GDG recommends that clinicians acknowledge the uncertainty towards the risk of malignancy in postmenopausal women. If a pelvic mass is detected, the work-up and treatment should be performed according to national oncology guidelines.	GPP
For postmenopausal women with endometriosis-associated pain, clinicians may consider aromatase inhibitors as a treatment option especially if surgery is not feasible (Polyzos et al., 2011; Pavone and Bulun, 2012).	Weak recommendation ⊕○○○

Is hormone treatment effective and safe for relief of menopausal symptoms in women with a history of endometriosis?

Clinicians may consider combined menopausal hormone therapy for the treatment of postmenopausal symptoms in women (both after natural and surgical menopause) with a history of endometriosis (Matorras *et al.*, 2002; Gemmell *et al.*, 2017).

Weak recommendation
⊕⊕○○

Clinicians should avoid prescribing oestrogen-only regimens for the treatment of vasomotor symptoms in postmenopausal women with a history of endometriosis, as these regimens may be associated with a higher risk of malignant transformation (Gemmell *et al.*, 2017).

Strong recommendation
⊕⊕○○

The GDG recommends that clinicians continue to treat women with a history of endometriosis after surgical menopause with combined oestrogen–progestogen at least up to the age of natural menopause.

GPP

Are women with endometriosis at higher risk of experiencing menopause-related major health concerns?

Clinicians should be aware that women with endometriosis who have undergone an early bilateral salpingo-oophorectomy as part of their treatment have an increased risk of diminished bone density, dementia and cardiovascular disease. It is also important to note that women with endometriosis have an increased risk of cardiovascular disease, irrespective of whether they have had an early surgical menopause (*conclusion, not recommendation*).

Extrapelvic endometriosis

How reliable is imaging for diagnosing extrapelvic endometriosis?

Clinicians should be aware of symptoms of extrapelvic endometriosis, such as cyclical shoulder pain, cyclical spontaneous pneumothorax, cyclical cough or nodules which enlarge during menses.

GPP

It is advisable to discuss diagnosis and management of extrapelvic endometriosis in a multidisciplinary team in a centre with sufficient expertise.

GPP

Does treatment for extrapelvic endometriosis relieve symptoms?

For abdominal extrapelvic endometriosis, surgical removal is the preferred treatment, when possible, to relieve symptoms. Hormone treatment may also be an option when surgery is not possible or acceptable (Horton *et al.*, 2008; Zhu *et al.*, 2017; Andres *et al.*, 2020; Hirata *et al.*, 2020).

Weak recommendation
⊕○○○

For thoracic endometriosis, hormone treatment can be offered. If surgery is indicated, it should be performed in a multidisciplinary manner involving a thoracic surgeon and/or other relevant specialists (Joseph and Sahn, 1996; Ceccaroni *et al.*, 2013; Nezhat *et al.*, 2014; Gil and Tulandi, 2020; Andres *et al.*, 2020; Viguera Smith *et al.*, 2021; Ciriaco *et al.*, 2022).

Weak recommendation
⊕○○○

Asymptomatic endometriosis

Is treatment beneficial for incidental finding of asymptomatic endometriosis?

The GDG recommends that clinicians should inform and counsel women about any incidental finding of endometriosis.

GPP

Clinicians should not routinely perform surgical excision/ablation for an incidental finding of asymptomatic endometriosis at the time of surgery (Moen and Stokstad, 2002).

Strong recommendation
⊕○○○

Clinicians should not prescribe medical treatment in women with incidental finding of endometriosis.

Strong recommendation
⊕○○○

Is long term monitoring of women with asymptomatic endometriosis beneficial in preventing adverse outcomes?

Routine US monitoring of asymptomatic endometriosis can be considered (Maouris, 1991; Alcázar *et al.*, 2005; Pearce *et al.*, 2012; Serati *et al.*, 2013).

Weak recommendation
⊕○○○

Primary prevention of endometriosis

Is there a role for primary prevention of endometriosis?

Although there is no direct evidence of benefit in preventing endometriosis in the future, women can be advised of aiming for a healthy lifestyle and diet, with reduced alcohol intake and regular physical activity (Hansen and Knudsen, 2013; Parazzini *et al.*, 2013a,b; Bravi *et al.*, 2014; Ricci *et al.*, 2016; Harris *et al.*, 2018; Nodler *et al.*, 2020; Qiu *et al.*, 2020).

Weak recommendation
⊕⊕○○

The usefulness of hormonal contraceptives for the primary prevention of endometriosis is uncertain (Vercellini *et al.*, 2011).

Weak recommendation
⊕⊕○○

Genetic testing in women with suspected or confirmed endometriosis should only be performed within a research setting.

RESEARCH-ONLY

Endometriosis and cancer

Are patients with endometriosis at increased risk of cancer?

Clinicians should inform women with endometriosis requesting information on their risk of developing cancer that endometriosis is not associated with a significantly higher risk of cancer overall (Fig. 6). Although endometriosis is associated with a higher risk of ovarian, breast and thyroid cancers in particular, the increase in absolute risk compared with women in the general population is low (Kvaskoff *et al.*, 2021).

Strong recommendation
⊕⊕○○

What information could clinicians provide to women with endometriosis regarding their risk of developing cancer?

The GDG recommends that clinicians reassure women with endometriosis with regards to their cancer risk and address their concern to reduce their risk by recommending general cancer prevention measures (avoiding smoking, maintaining a healthy weight, exercising regularly, having a balanced diet with high intakes of fruits and vegetables and low intakes of alcohol, and using sun protection).

GPP

Are somatic mutations in deep endometriosis of patients without cancer predictive for ovarian cancer development and/or progression?

Based on the limited literature and controversial findings, there is little evidence that somatic mutations in patients with deep endometriosis may be predictive of development and/or progression of ovarian cancer (*conclusion, not recommendation*).

Does the use of hormone treatments increase the risk of cancer?

Clinicians should reassure women with endometriosis about the risk of malignancy associated with the use of hormonal contraceptives (Smith et al., 2003; Zucchetto et al., 2009; Gierisch et al., 2013; Havrilesky et al., 2013; Braganza et al., 2014; Berlanda et al., 2016; Wentzensen et al., 2016; Butt et al., 2018; Michels et al., 2018).

Strong
recommendation
⊕⊕○○

Should women with endometriosis be monitored for detection of malignancy?

In women with endometriosis, clinicians should not systematically perform cancer screening beyond the existing population-based cancer screening guidelines (Kvaskoff et al., 2021).

Strong
recommendation
⊕⊕○○

Clinicians can consider cancer screening according to local guidelines in individual patients that have additional risk factors, e.g. strong family history, specific germline mutations.

GPP

Does surgery for endometriosis change the future risk of cancer?

Clinicians should be aware that there is epidemiological data, mostly on ovarian endometriosis, showing that complete excision of visible endometriosis may reduce the risk of ovarian cancer. The potential benefits should be weighed against the risks of surgery (morbidity, pain and ovarian reserve) (Rossing et al., 2008; Melin et al., 2013; Haraguchi et al., 2016).

Strong
recommendation
⊕⊕○○

Discussion

This paper provides an overview of recommendations for diagnosis of endometriosis and treatment of associated symptoms during different stages of life. In addition, guidance is provided on the possible connection with development of cancer, and with regards to prevention. Overall, 109 recommendations have been formulated, 79 supported by research data and 30 GPPs based primarily on clinical expertise. The guidelines are based on the best available evidence or, where data of sufficient quality were absent, on recommendations by the GDG (GPPs).

The current guideline and recommendations are an update of the ESHRE endometriosis guidelines published in 2013 and 2005 (Kennedy et al., 2005; Dunselman et al., 2014). The key questions and topics covered in the guideline of 2013 were updated based on data published between 2013 and 2021, where available, and in accordance with changes in clinical practice. The latter applied, for example, to the oral use of danazol and anti-progestogens as a medical treatment and to LUNA, PSN and anti-adhesion agents as surgical interventions. These interventions are still discussed in the guideline, but no longer discussed in recommendations for clinical practice.

While most of the more recent studies confirm previous ESHRE recommendations, there are five topics in which significant changes in clinical practice are to be expected. The first change, primarily based on clinical practice rather than published data, is the evolution in the diagnostic process. While previously a laparoscopy was regarded as the diagnostic gold standard, it is now only recommended in patients with negative imaging results and/or where empirical treatment was unsuccessful or inappropriate. Secondly, studies on GnRH antagonist treatments support their use as an additional (second-line) treatment option. Thirdly, recent data indicate that postoperative medical treatment may be beneficial towards pain management and support a recommendation to offer it to women not desiring immediate pregnancy. Fourthly, the extended administration of GnRH agonist prior to ART treatment to improve live birth rate in infertile women with endometriosis (ultralong protocol) is no longer recommended because of unclear benefits. Finally, the EFI was added as a step in the treatment as it can support decision-making for the most appropriate option to achieve pregnancy after surgery.

In addition to the topics discussed in the previous guideline, the current guideline addresses highly important previous gaps in clinical management, with an additional chapter on adolescent endometriosis, information on pregnancy and fertility preservation, and extended information on endometriosis in menopause, as well as data on the link between endometriosis and cancer.

Despite our best efforts to provide clear guidance on the management of endometriosis using all available evidence, there is still an urgent need for more research both to achieve more clarity on the most appropriate diagnostic and treatment options, and to answer very basic questions as to the natural course of the disease. This guideline provides 30 recommendations for research written to inspire researchers and hopefully also facilitate funding for endometriosis studies (Supplementary Data).

In summary, the 2022 ESHRE Guideline: Endometriosis is a comprehensive update of the existing evidence and should assist healthcare professionals in their decision making and patients in their understanding of the management suggestions. Active involvement and input by patient representatives at all stages was central to the success of this endeavour. As such, the guideline was created by medical professionals, patient representatives and specialists in epidemiology and guideline methodology. The detailed guideline document and a patient-friendly version can be accessed via the ESHRE website (<https://www.eshre.eu/Guideline/Endometriosis>).

Supplementary data

Supplementary data are available at *Human Reproduction Open* online.

Data availability

The full guideline and supporting data (literature report, evidence tables) are available on www.eshre.eu/guidelines.

Acknowledgements

The authors would like to thank the experts that participated in the stakeholder review for their constructive and helpful comments.

Authors' roles

The guideline core group was chaired by C.M.B. N.V. provided methodological support. All other authors contributed equally to writing the guideline. All authors have revised and approved the final version. The collaborators supported the guideline core group for the individual chapters.

Funding

The guideline was developed and funded by ESHRE, covering expenses associated with the guideline meetings, with the literature searches and with the dissemination of the guideline. The guideline group members did not receive payment.

Conflict of interest

C.M.B. reports grants from Bayer Healthcare and the European Commission; Participation on a Data Safety Monitoring Board or Advisory Board with ObsEva (Data Safety Monitoring Group) and Myovant (Scientific Advisory Group). A.B. reports grants from FEMaLE

executive board member and European Commission Horizon 2020 grant; consulting fees from Ethicon Endo Surgery, Medtronic; honoraria for lectures from Ethicon; and support for meeting attendance from Gedeon Richter; A.H. reports grants from MRC, NIHR, CSO, Roche Diagnostics, Astra Zeneca, Ferring; Consulting fees from Roche Diagnostics, Nordic Pharma, Chugai and Benevolent AI Bio Limited all paid to the institution; a pending patent on Serum endometriosis biomarker; he is also Chair of TSC for STOP-OHSS and CERM trials. O.H. reports consulting fees and speaker's fees from Gedeon Richter and Bayer AG; support for attending meetings from Gedeon-Richter, and leadership roles at the Finnish Society for Obstetrics and Gynecology and the Nordic federation of the societies of obstetrics and gynecology. L.K. reports consulting fees from Gedeon Richter, AstraZeneca, Novartis, Dr KADE/Besins, Palleos Healthcare, Roche, Mithra; honoraria for lectures from Gedeon Richter, AstraZeneca, Novartis, Dr KADE/Besins, Palleos Healthcare, Roche, Mithra; support for attending meetings from Gedeon Richter, AstraZeneca, Novartis, Dr KADE/Besins, Palleos Healthcare, Roche, Mithra; he also has a leadership role in the German Society of Gynecological Endocrinology (DGGEF). M.K. reports grants from French Foundation for Medical Research (FRM), Australian Ministry of Health, Medical Research Future Fund and French National Cancer Institute; support for meeting attendance from European Society for Gynaecological Endoscopy (ESGE), European Congress on Endometriosis (EEC) and ESHRE; She is an advisory Board Member, FEMaLe Project (Finding Endometriosis Using Machine Learning), Scientific Committee Chair for the French Foundation for Research on Endometriosis and Scientific Committee Chair for the ComPaRe-Endometriosis cohort. A.N. reports grants from Merck SA and Ferring; speaker fees from Merck SA and Ferring; support for meeting attendance from Merck SA; Participation on a Data Safety Monitoring Board or Advisory Board with Nordic Pharma and Merck SA; she also is a board member of medical advisory board, Endometriosis Society, the Netherlands (patients advocacy group) and an executive board member World Endometriosis Society. E.S. reports grants from National Institute for Health Research UK, Rosetrees Trust, Barts and the London Charity; Royalties from De Gruyter (book editor); consulting fees from Hologic; speakers fees from Hologic, Johnson & Johnson, Medtronic, Intuitive, Olympus and Karl Storz; Participation in the Medicines for Women's Health Expert Advisory Group with Medicines and Healthcare Products Regulatory Agency (MHRA); he is also Ambassador for the World Endometriosis Society. C.T. reports grants from Merck SA; Consulting fees from Gedeon Richter, Nordic Pharma and Merck SA; speaker fees from Merck SA, all paid to the institution; and support for meeting attendance from Ferring, Gedeon Richter, Merck SA. The other authors have no conflicts of interest to declare.

Appendix

Members of the ESHRE Endometriosis Guideline Group

Signe Altmäe	Department of Biochemistry and Molecular Biology, Faculty of Sciences, University of Granda, Spain; Div. Obstetrics and Gynaecology, CLINTEC, Karolinska Institutet, Sweden.
Baris Ata	Koc University School of Medicine, Turkey
Elizabeth Ball	The Royal London Hospital, Bartshealth NHS Trust and Queen Mary University of London, London, UK; City University London, London, UK
Fabio Barra	Academic Unit of Obstetrics and Gynecology, IRCCS Ospedale Policlinico San Martino, Genoa, Italy; Department of Neurosciences, Rehabilitation, Ophthalmology, Genetics, Maternal and Child Health (DiNOGMI), University of Genoa, Genoa, Italy.
Ercan Bastu	Department of Obstetrics and Gynecology, University of Massachusetts Chan Medical School, USA
Alexandra Bianco-Anil	EndoFrance, French patients' Association, France
Ulla Breth Knudsen	Aarhus University, Aarhus, Denmark
Réka Brubel	Semmelweis University, Faculty of Medicine, Budapest, Hungary
Julia Cambitzi	Pain Management Centre, University College London Hospitals (UCLH), London, UK
Astrid Cantineau	University of Groningen, University Medical Center Groningen, Groningen, The Netherlands
Ying Cheong	University of Southampton, Complete Fertility Southampton, Southampton, UK
Angelos Daniilidis	2nd University Department of Obstetrics & Gynecology, Hippokratia General Hospital, Aristotle University of Thessaloniki, Greece
Bianca De Bie	Endometriose Stichting, The Netherlands
Caterina Exacoustos	Department of Surgical Sciences, Obstetric and Gynecological Unit, University of Rome 'Tor Vergata', Rome, Italy
Simone Ferrero	Academic Unit of Obstetrics and Gynecology, IRCCS Ospedale Policlinico San Martino, Genoa, Italy; Department of Neurosciences, Rehabilitation, Ophthalmology, Genetics, Maternal and Child Health (DiNOGMI), University of Genoa, Genoa, Italy
Tarek Gelbaya	University Hospitals of Leicester, Leicester, UK
Josepha Goetz-Collinet	EndoFrance, French patients' Association, France
Gernot Hudelist	Hospital St. John of God Vienna, Vienna, Austria
Munawar Hussain	Southend University Hospital, UK
Tereza Indrielle-Kelly	Burton and Derby hospitals NHS Trust, Burton on Trent, UK
Shaheen Khazali	Centre for Endometriosis and Minimally Invasive Gynaecology (CEMIG) at The HCA Lister Hospital, Chelsea, London, UK
Sujata Lalit Kumar	Department of Women's and Children's Health, Karolinska Institutet, Stockholm, Sweden; Stockholm IVF, Stockholm, Sweden
Umberto Leone Roberti Maggiore	Department of Gynecologic Oncology, IRCCS National Cancer Institute, Milan, Italy
Jacques WM Maas	Maastricht University Medical Centre, Department of Obstetrics and Gynecology and GROW—School for Oncology and Developmental Biology, Maastricht, the Netherlands.
Helen McLaughlin	Endometriosis advocate, London; Endometriosis UK
José Metello	CIRMA, Hospital Garcia de Orta, Almada; GINEMED, MaloClinics, Lisboa, Portugal
Velja Mijatovic	Academic Endometriosis Center Amsterdam UMC, Amsterdam, The Netherlands
Yasaman Miremadi	The Austrian Society of Sterility, Fertility and Endocrinology, Austria; TFP kinderwunschlinik Wien, Austria
Charles Muteshi	University of Oxford, Oxford, UK
Michelle Nisolle	University of Liege/CHR Citadelle, Liege, Belgium
Engin Oral	Department of Obstetrics and Gynecology, Bezmialem Vakif University Medical Faculty, Istanbul, Turkey
George Pados	Aristotle University of Thessaloniki, 1st Dept. OB-GYN, 'Papageorgiou' General Hospital, Thessaloniki and Centre for Endoscopic Surgery 'DIAVALKANIKO' hospital, Thessaloniki, Greece
Dana Parades	Endometriosis Association, Finland
Nicola Pluchino	Division of Gynecology and Obstetrics, University Hospital of Geneva, Geneva, Switzerland
Prasanna Raj Supramaniam	Endometriosis CaRe Centre Oxford, Nuffield Department of Women's and Reproductive Health, University of Oxford, UK; Oxford University Hospitals NHS Foundation Trust, Oxford, UK
Maren Schick	Institute of Medical Psychology, Center for Psychosocial Medicine, University Hospital Heidelberg, Heidelberg, Germany

(continued)

Beata Seeber

Department of Gynecologic Endocrinology and Reproductive Medicine, Medical University of Innsbruck, Innsbruck, Austria

Renato Seracchioli

Division of Gynecology and Human Reproduction Physiopathology, IRCCS Azienda Ospedaliero-Universitaria di Bologna, S. Orsola Hospital, University of Bologna, Bologna, Italy

Antonio Simone Laganà

Department of Obstetrics and Gynecology, 'Filippo Del Ponte' Hospital, University of Insubria, Varese, Italy

Andreas Stavroulis

Endometriosis and Fertility Center, Cyprus; American Medical Center, Nicosia, Cyprus

Linda Tebache

University of Liege/CHR Citadelle, Liege, Belgium

Gürkan Uncu

Uludag University, Bursa, Turkey

Uschi Van den Broeck

Leuven University Fertility Center (LUFC), University Hospitals Leuven, Leuven, Belgium

Arno van Peperstraten

University Medical Center Utrecht, The Netherlands

Attila Vereczkey

Versys Clinics Human Reproduction Institute, Budapest, Hungary

Albert Wolthuis

University Hospitals Leuven, Leuven, Belgium

Pinar Yalçın Bahat

Istanbul Sağlık Bilimleri University Kanuni Sultan Süleyman Training and Research Hospital, Istanbul, Turkey

Chadi Yazbeck

Cherest Fertility Center—Reprogyne Institute, Paris, France; Foch University Hospital, Dept Obstetrics Gynecology and Reproductive Medicine, Suresnes, France

References

- Abdou AM, Ammar IMM, Alnemr AAA, Abdelrhman AA. Dienogest versus leuprolide acetate for recurrent pelvic pain following laparoscopic treatment of endometriosis. *J Obstet Gynaecol India* 2018; **68**:306–313.
- Agarwal SK, Foster WG. Reduction in endometrioma size with three months of aromatase inhibition and progestin add-back. *Biomed Res Int* 2015; **2015**:878517.
- Alborzi S, Hamed B, Omidvar A, Dehbashi S, Alborzi S, Alborzi M. A comparison of the effect of short-term aromatase inhibitor (letrozole) and GnRH agonist (triptorelin) versus case control on pregnancy rate and symptom and sign recurrence after laparoscopic treatment of endometriosis. *Arch Gynecol Obstet* 2011; **284**: 105–110.
- Alborzi S, Zahiri Sorouri Z, Askari E, Poordast T, Chamanara K. The success of various endometrioma treatments in infertility: a systematic review and meta-analysis of prospective studies. *Reprod Med Biol* 2019; **18**:312–322.
- Alcázar JL, Castillo G, Jurado M, García GL. Is expectant management of sonographically benign adnexal cysts an option in selected asymptomatic premenopausal women? *Hum Reprod* 2005; **20**: 3231–3234.
- Almassinokiani F, Almasi A, Akbari P, Saberifard M. Effect of Letrozole on endometriosis-related pelvic pain. *Med J Islam Repub Iran* 2014; **28**:107.
- Alshehre SM, Narice BF, Fenwick MA, Metwally M. The impact of endometrioma on in vitro fertilisation/intra-cytoplasmic injection IVF/ICSI reproductive outcomes: a systematic review and meta-analysis. *Arch Gynecol Obstet* 2021; **303**:3–16.
- Andres MDP, Lopes LA, Baracat EC, Podgaec S. Dienogest in the treatment of endometriosis: systematic review. *Arch Gynecol Obstet* 2015; **292**:523–529.
- Andres MP, Arcoverde FVL, Souza CCC, Fernandes LFC, Abrão MS, Kho RM. Extrapelvic endometriosis: a systematic review. *J Minim Invasive Gynecol* 2020; **27**:373–389.
- Arcoverde FVL, Andres MP, Borrelli GM, Barbosa PA, Abrão MS, Kho RM. Surgery for endometriosis improves major domains of quality of life: a systematic review and meta-analysis. *J Minim Invasive Gynecol* 2019; **26**:266–278.
- Bafort C, Beebejaun Y, Tomassetti C, Bosteels J, Duffy JMN; Cochrane Gynaecology and Fertility Group. Laparoscopic surgery for endometriosis. *Cochrane Database Syst Rev* 2020; **10**: CD011031.
- Ballard KD, Seaman HE, de Vries CS, Wright JT. Can symptomatology help in the diagnosis of endometriosis? Findings from a national case-control study—part I. *BJOG* 2008; **115**:1382–1391.
- Bastu E, Yasa C, Dural O, Mutlu MF, Celik C, Ugurlucan FG, Buyru F. Comparison of ovulation induction protocols after endometrioma resection. *JLS* 2014; **18**:e2014.00128.
- Bazot M, Lafont C, Rouzier R, Roseau G, Thomassin-Naggara I, Darai E. Diagnostic accuracy of physical examination, transvaginal sonography, rectal endoscopic sonography, and magnetic resonance imaging to diagnose deep infiltrating endometriosis. *Fertil Steril* 2009; **92**:1825–1833.
- Behera M, Vilos GA, Hollett-Caines J, Abu-Rafea B, Ahmad R. Laparoscopic findings, histopathologic evaluation, and clinical outcomes in women with chronic pelvic pain after hysterectomy and bilateral salpingo-oophorectomy. *J Minim Invasive Gynecol* 2006; **13**: 431–435.
- Benaglia L, Somigliana E, Iemello R, Colpi E, Nicolosi AE, Ragni G. Endometrioma and oocyte retrieval-induced pelvic abscess: a clinical concern or an exceptional complication? *Fertil Steril* 2008; **89**: 1263–1266.
- Bendifallah S, Roman H, Mathieu d'Argent E, Touleimat S, Cohen J, Darai E, Ballester M. Colorectal endometriosis-associated infertility: should surgery precede ART? *Fertil Steril* 2017; **108**:525–531.e524.
- Bendifallah S, Vesale E, Darai E, Thomassin-Naggara I, Bazot M, Tuech JJ, Abo C, Roman H. Recurrence after surgery for colorectal endometriosis: a systematic review and meta-analysis. *J Minim Invasive Gynecol* 2020; **27**:441–451.e442.
- Berlanda N, Somigliana E, Vigano P, Vercellini P. Safety of medical treatments for endometriosis. *Expert Opin Drug Saf* 2016; **15**: 21–30.
- Bianchi PH, Pereira RM, Zanatta A, Alegretti JR, Motta EL, Serafini PC. Extensive excision of deep infiltrative endometriosis before

- in vitro fertilization significantly improves pregnancy rates. *J Minim Invasive Gynecol* 2009;**16**:174–180.
- Braganza MZ, de González AB, Schonfeld SJ, Wentzensen N, Brenner AV, Kitahara CM. Benign breast and gynecologic conditions, reproductive and hormonal factors, and risk of thyroid cancer. *Cancer Prev Res (Phila)*. 2014;**7**:418–425.
- Bravi F, Parazzini F, Cipriani S, Chiaffarino F, Ricci E, Chiantera V, Vigano P, La Vecchia C. Tobacco smoking and risk of endometriosis: a systematic review and meta-analysis. *BMJ Open* 2014;**4**:e006325.
- Breteau P, Chanavaz-Lacheray I, Rubod C, Turck M, Sanguin S, Pop I, Resch B, Roman H. Pregnancy rates after surgical treatment of deep infiltrating endometriosis in infertile patients with at least 2 previous in vitro fertilization or intracytoplasmic sperm injection failures. *J Minim Invasive Gynecol* 2020;**27**:1148–1157.
- Brosens I, Gordts S, Benagiano G. Endometriosis in adolescents is a hidden, progressive and severe disease that deserves attention, not just compassion. *Hum Reprod* 2013;**28**:2026–2031.
- Brown J, Crawford TJ, Allen C, Hopewell S, Prentice A. Nonsteroidal anti-inflammatory drugs for pain in women with endometriosis. *Cochrane Database Syst Rev* 2017;**1**:CD004753.
- Brown J, Crawford TJ, Datta S, Prentice A. Oral contraceptives for pain associated with endometriosis. *Cochrane Database Syst Rev* 2018;**5**:CD001019.
- Brown J, Kives S, Akhtar M; Cochrane Gynaecology and Fertility Group. Progestagens and anti-progestagens for pain associated with endometriosis. *Cochrane Database Syst Rev* 2012;**2012**:CD002122.
- Brown J, Pan A, Hart RJ. Gonadotrophin-releasing hormone analogues for pain associated with endometriosis. *Cochrane Database Syst Rev* 2010;**2010**:CD008475.
- Busacca M, Riparini J, Somigliana E, Oggioni G, Izzo S, Vignali M, Candiani M. Postsurgical ovarian failure after laparoscopic excision of bilateral endometriomas. *Am J Obstet Gynecol* 2006;**195**:421–425.
- Butt SA, Lidegaard Ø, Skovlund C, Hannaford PC, Iversen L, Fielding S, Mørch LS. Hormonal contraceptive use and risk of pancreatic cancer – a cohort study among premenopausal women. *PLoS One* 2018;**13**:e0206358.
- Byrne D, Curnow T, Smith P, Cutner A, Saridogan E, Clark TJ; BSGE Endometriosis Centres. Laparoscopic excision of deep rectovaginal endometriosis in BSGE endometriosis centres: a multicentre prospective cohort study. *BMJ Open* 2018;**8**:e018924.
- Candiani GB, Fedele L, Vercellini P, Bianchi S, Di Nola G. Repetitive conservative surgery for recurrence of endometriosis. *Obstet Gynecol* 1991;**77**:421–424.
- Candiani M, Ottolina J, Schimberni M, Tandoi I, Bartiromo L, Ferrari S. Recurrence rate after “One-Step” CO(2) fiber laser vaporization versus cystectomy for ovarian endometrioma: a 3-year follow-up study. *J Minim Invasive Gynecol* 2020;**27**:901–908.
- Cao X, Chang H-Y, Xu J-Y, Zheng Y, Xiang Y-G, Xiao B, Geng X-J, Ni L-L, Chu X-Y, Tao S-B et al. The effectiveness of different down-regulating protocols on in vitro fertilization-embryo transfer in endometriosis: a meta-analysis. *Reprod Biol Endocrinol* 2020;**18**:16.
- Carmona F, Martínez-Zamora MA, Rabanal A, Martínez-Román S, Balasch J. Ovarian cystectomy versus laser vaporization in the treatment of ovarian endometriomas: a randomized clinical trial with a five-year follow-up. *Fertil Steril* 2011;**96**:251–254.
- Ceccaroni M, Roviglione G, Giampaolino P, Clarizia R, Bruni F, Ruffo G, Patrelli TS, De Placido G, Minelli L. Laparoscopic surgical treatment of diaphragmatic endometriosis: a 7-year single-institution retrospective review. *Surg Endosc* 2013;**27**:625–632.
- Chapron C, Dubuisson JB, Pansini V, Vieira M, Fauconnier A, Barakat H, Dousset B. Routine clinical examination is not sufficient for diagnosing and locating deeply infiltrating endometriosis. *J Am Assoc Gynecol Laparosc* 2002;**9**:115–119.
- Chapron C, Lafay-Pillet MC, Monceau E, Borghese B, Ngo C, Souza C, de Ziegler D. Questioning patients about their adolescent history can identify markers associated with deep infiltrating endometriosis. *Fertil Steril* 2011;**95**:877–881.
- Chapron C, Querleu D, Bruhat MA, Madelenat P, Fernandez H, Pierre F, Dubuisson JB. Surgical complications of diagnostic and operative gynaecological laparoscopy: a series of 29,966 cases. *Hum Reprod* 1998;**13**:867–872.
- Chen FP, Soong YK, Lee N, Lo SK. The use of serum CA-125 as a marker for endometriosis in patients with dysmenorrhea for monitoring therapy and for recurrence of endometriosis. *Acta Obstet Gynecol Scand* 1998;**77**:665–670.
- Chen I, Veth VB, Choudhry AJ, Murji A, Zakhari A, Black AY, Agarpao C, Maas JW. Pre- and postsurgical medical therapy for endometriosis surgery. *Cochrane Database Syst Rev* 2020;**11**:CD003678.
- Ciriaco P, Muriana P, Lembo R, Carretta A, Negri G. Treatment of thoracic endometriosis syndrome: a meta-analysis and review. *Ann Thorac Surg* 2022;**113**:324–336.
- Clayton RD, Hawe JA, Love JC, Wilkinson N, Garry R. Recurrent pain after hysterectomy and bilateral salpingo-oophorectomy for endometriosis: evaluation of laparoscopic excision of residual endometriosis. *Br J Obstet Gynaecol* 1999;**106**:740–744.
- Cobo A, Giles J, Paoletti S, Pellicer A, Remohí J, García-Velasco JA. Oocyte vitrification for fertility preservation in women with endometriosis: an observational study. *Fertil Steril* 2020;**113**:836–844.
- Coccia ME, Rizzello F, Barone S, Pinelli S, Rapalini E, Parri C, Caracciolo D, Papageorgiou S, Cima G, Gandini L. Is there a critical endometrioma size associated with reduced ovarian responsiveness in assisted reproduction techniques? *Reprod Biomed Online* 2014;**29**:259–266.
- Condous G, Van Calster B, Van Huffel S, Lam A. What is the value of preoperative bimanual pelvic examination in women undergoing laparoscopic total hysterectomy? *J Minim Invasive Gynecol* 2007;**14**:334–338.
- Cosar E, Mamillapalli R, Ersoy GS, Cho S, Seifer B, Taylor HS. Serum microRNAs as diagnostic markers of endometriosis: a comprehensive array-based analysis. *Fertil Steril* 2016;**106**:402–409.
- Dan H, Limin F. Laparoscopic ovarian cystectomy versus fenestration/coagulation or laser vaporization for the treatment of endometriomas: a meta-analysis of randomized controlled trials. *Gynecol Obstet Invest* 2013;**76**:75–82.
- Davis AR, Westhoff C, O’Connell K, Gallagher N. Oral contraceptives for dysmenorrhea in adolescent girls: a randomized trial. *Obstet Gynecol* 2005;**106**:97–104.
- De Cicco C, Corona R, Schonman R, Mailova K, Ussia A, Koninckx P. Bowel resection for deep endometriosis: a systematic review. *BJOG* 2011;**118**:285–291.

- de Ziegler D, Gayet V, Aubriot FX, Fauque P, Streuli I, Wolf JP, de Mouzon J, Chapron C. Use of oral contraceptives in women with endometriosis before assisted reproduction treatment improves outcomes. *Fertil Steril* 2010;**94**:2796–2799.
- DiVasta AD, Feldman HA, Sadler Gallagher J, Stokes NA, Laufer MR, Hornstein MD, Gordon CM. Hormonal add-back therapy for females treated with gonadotropin-releasing hormone agonist for endometriosis: a randomized controlled trial. *Obstet Gynecol* 2015;**126**:617–627.
- DiVasta AD, Vitonis AF, Laufer MR, Missmer SA. Spectrum of symptoms in women diagnosed with endometriosis during adolescence vs adulthood. *Am J Obstet Gynecol* 2018;**218**:324.e321–324.e311.
- Donnez J, Taylor HS, Taylor RN, Akin MD, Tatarchuk TF, Wilk K, Gotteland JP, Lecomte V, Bestel E. Treatment of endometriosis-associated pain with linzagolix, an oral gonadotropin-releasing hormone-antagonist: a randomized clinical trial. *Fertil Steril* 2020;**114**:44–55.
- Doyle JO, Missmer SA, Laufer MR. The effect of combined surgical-medical intervention on the progression of endometriosis in an adolescent and young adult population. *J Pediatr Adolesc Gynecol* 2009;**22**:257–263.
- Dragoman MV, Gaffield ME. The safety of subcutaneously administered depot medroxyprogesterone acetate (104mg/0.65mL): a systematic review. *Contraception* 2016;**94**:202–215.
- Drakopoulos P, Rosetti J, Pluchino N, Blockeel C, Santos-Ribeiro S, de Brucker M, Drakakis P, Camus M, Tournaye H, Polyzos NP. Does the type of GnRH analogue used, affect live birth rates in women with endometriosis undergoing IVF/ICSI treatment, according to the rAFS stage? *Gynecol Endocrinol* 2018;**34**:884–889.
- Dunselman GAJ, Vermeulen N, Becker C, Calhaz-Jorge C, D'Hooghe T, De Bie B, Heikinheimo O, Horne AW, Kiesel L, Nap A et al.; European Society of Human Reproduction and Embryology. ESHRE guideline: management of women with endometriosis. *Hum Reprod* 2014;**29**:400–412.
- Ebert AD, Dong L, Merz M, Kirsch B, Francuski M, Bottcher B, Roman H, Suvitie P, Hlavackova O, Gude K et al. Dienogest 2 mg daily in the treatment of adolescents with clinically suspected endometriosis: the ViSanne study to Assess safety in ADolescents. *J Pediatr Adolesc Gynecol* 2017;**30**:560–567.
- Eskenazi B, Warner M, Bonsignore L, Olive D, Samuels S, Vercellini P. Validation study of nonsurgical diagnosis of endometriosis. *Fertil Steril* 2001;**76**:929–935.
- Eskenazi B, Warner ML. Epidemiology of endometriosis. *Obstet Gynecol Clin North Am* 1997;**24**:235–258.
- Ferrero S, Gillott DJ, Venturini PL, Remorgida V. Use of aromatase inhibitors to treat endometriosis-related pain symptoms: a systematic review. *Reprod Biol Endocrinol* 2011;**9**:89.
- Forman RG, Robinson JN, Mehta Z, Barlow DH. Patient history as a simple predictor of pelvic pathology in subfertile women. *Hum Reprod* 1993;**8**:53–55.
- Franck C, Poulsen MH, Karampas G, Giraldi A, Rudnicki M. Questionnaire-based evaluation of sexual life after laparoscopic surgery for endometriosis: a systematic review of prospective studies. *Acta Obstet Gynecol Scand* 2018;**97**:1091–1104.
- Gallagher JS, Feldman HA, Stokes NA, Laufer MR, Hornstein MD, Gordon CM, DiVasta AD. The effects of gonadotropin-releasing hormone agonist combined with add-back therapy on quality of life for adolescents with endometriosis: a randomized controlled trial. *J Pediatr Adolesc Gynecol* 2017;**30**:215–222.
- Gallagher JS, Missmer SA, Hornstein MD, Laufer MR, Gordon CM, DiVasta AD. Long-term effects of gonadotropin-releasing hormone agonists and add-back in adolescent endometriosis. *J Pediatr Adolesc Gynecol* 2018;**31**:376–381.
- Gemmell LC, Webster KE, Kirtley S, Vincent K, Zondervan KT, Becker CM. The management of menopause in women with a history of endometriosis: a systematic review. *Hum Reprod Update* 2017;**23**:481–500.
- Georgiou EX, Melo P, Baker PE, Sallam HN, Arici A, Garcia-Velasco JA, Abou-Setta AM, Becker C, Granne IE. Long-term GnRH agonist therapy before in vitro fertilisation (IVF) for improving fertility outcomes in women with endometriosis. *Cochrane Database Syst Rev* 2019;**2019**:CD013240.
- Geysenbergh B, Dancet EAF, D'Hooghe T. Detecting endometriosis in adolescents: why not start from self-report screening questionnaires for adult women? *Gynecol Obstet Invest* 2017;**82**:322–328.
- Gierisch JM, Coeytaux RR, Urrutia RP, Havrilesky LJ, Moorman PG, Lowery WJ, Dinan M, McBroom AJ, Hasselblad V, Sanders GD et al. Oral contraceptive use and risk of breast, cervical, colorectal, and endometrial cancers: a systematic review. *Cancer Epidemiol Biomarkers Prev* 2013;**22**:1931–1943.
- Gil Y, Tulandi T. Diagnosis and treatment of catamenial pneumothorax: a systematic review. *J Minim Invasive Gynecol* 2020;**27**:48–53.
- Glavind MT, Mollgaard MV, Iversen ML, Arendt LH, Forman A. Obstetrical outcome in women with endometriosis including spontaneous hemoperitoneum and bowel perforation: a systematic review. *Best Pract Res Clin Obstet Gynaecol* 2018;**51**:41–52.
- Grandi G, Barra F, Ferrero S, Sileo FG, Bertucci E, Napolitano A, Facchinetti F. Hormonal contraception in women with endometriosis: a systematic review. *Eur J Contracept Reprod Health Care* 2019;**24**:61–70.
- Greene R, Stratton P, Cleary SD, Ballweg ML, Sinai N. Diagnostic experience among 4,334 women reporting surgically diagnosed endometriosis. *Fertil Steril* 2009;**91**:32–39.
- Guerriero S, Saba L, Ajossa S, Peddes C, Angiolucci M, Perniciano M, Melis GB, Alcazar JL. Three-dimensional ultrasonography in the diagnosis of deep endometriosis. *Hum Reprod* 2014;**29**:1189–1198.
- Gupta D, Hull ML, Fraser I, Miller L, Bossuyt PM, Johnson N, Nisenblat V. Endometrial biomarkers for the non-invasive diagnosis of endometriosis. *Cochrane Database Syst Rev* 2016;**4**:CD012165.
- Hamdan M, Dunselman G, Li TC, Cheong Y. The impact of endometrioma on IVF/ICSI outcomes: a systematic review and meta-analysis. *Hum Reprod Update* 2015a;**21**:809–825.
- Hamdan M, Omar SZ, Dunselman G, Cheong Y. Influence of endometriosis on assisted reproductive technology outcomes: a systematic review and meta-analysis. *Obstet Gynecol* 2015b;**125**:79–88.
- Hansen SO, Knudsen UB. Endometriosis, dysmenorrhoea and diet. *Eur J Obstet Gynecol Reprod Biol* 2013;**169**:162–171.
- Haraguchi H, Koga K, Takamura M, Makabe T, Sue F, Miyashita M, Urata Y, Izumi G, Harada M, Hirata T et al. Development of ovarian cancer after excision of endometrioma. *Fertil Steril* 2016;**106**:1432–1437.e1432.
- Harb HM, Gallos ID, Chu J, Harb M, Coomarasamy A. The effect of endometriosis on in vitro fertilisation outcome: a systematic review and meta-analysis. *BJOG* 2013;**120**:1308–1320.

- Harris HR, Eke AC, Chavarro JE, Missmer SA. Fruit and vegetable consumption and risk of endometriosis. *Hum Reprod* 2018;**33**: 715–727.
- Hart RJ, Hickey M, Maouris P, Buckett W; Cochrane Gynaecology and Fertility Group. Excisional surgery versus ablative surgery for ovarian endometriomata. *Cochrane Database Syst Rev* 2008: CD004992.
- Havrilesky LJ, Moorman PG, Lowery WJ, Gierisch JM, Coeytaux RR, Urrutia RP, Dinan M, McBroom AJ, Hasselblad V, Sanders GD et al. Oral contraceptive pills as primary prevention for ovarian cancer: a systematic review and meta-analysis. *Obstet Gynecol* 2013;**122**:139–147.
- Healey M, Cheng C, Kaur H. To excise or ablate endometriosis? A prospective randomized double-blinded trial after 5-year follow-up. *J Minim Invasive Gynecol* 2014;**21**:999–1004.
- Hee L, Kettner LO, Vejtorp M. Continuous use of oral contraceptives: an overview of effects and side-effects. *Acta Obstet Gynecol Scand* 2013;**92**:125–136.
- Hirata T, Koga K, Kitade M, Fukuda S, Neriishi K, Taniguchi F, Honda R, Takazawa N, Tanaka T, Kurihara M et al. A national survey of umbilical endometriosis in Japan. *J Minim Invasive Gynecol* 2020;**27**:80–87.
- Hirsch M, Duffy J, Davis CJ, Nieves Plana M, Khan KS; on behalf of the International Collaboration to Harmonise Outcomes and Measures for Endometriosis. Diagnostic accuracy of cancer antigen 125 for endometriosis: a systematic review and meta-analysis. *BJOG* 2016;**123**:1761–1768.
- Hodgson RM, Lee HL, Wang R, Mol BW, Johnson N. Interventions for endometriosis-related infertility: a systematic review and network meta-analysis. *Fertil Steril* 2020;**113**:374–382.e372.
- Hornstein MD, Yuzpe AA, Burry K, Buttram VC Jr, Heinrichs LR, Soderstrom RM, Steinberger E, Lin JS. Retreatment with nafarelin for recurrent endometriosis symptoms: efficacy, safety, and bone mineral density. *Fertil Steril* 1997;**67**:1013–1018.
- Horton J, Sterrenburg M, Lane S, Maheshwari A, Li TC, Cheong Y. Reproductive, obstetric, and perinatal outcomes of women with adenomyosis and endometriosis: a systematic review and meta-analysis. *Hum Reprod Update* 2019;**25**:592–632.
- Horton JD, Dezee KJ, Ahnfeldt EP, Wagner M. Abdominal wall endometriosis: a surgeon's perspective and review of 445 cases. *Am J Surg* 2008;**196**:207–212.
- Hudelist G, Ballard K, English J, Wright J, Banerjee S, Mastoroudes H, Thomas A, Singer CF, Keckstein J. Transvaginal sonography vs. clinical examination in the preoperative diagnosis of deep infiltrating endometriosis. *Ultrasound Obstet Gynecol* 2011;**37**:480–487.
- Hughes E, Brown J, Collins JJ, Farquhar C, Fedorkow DM, Vanderkerchove P; Cochrane Gynaecology and Fertility Group. Ovulation suppression for endometriosis for women with subfertility. *Cochrane Database Syst Rev* 2007;**2007**:CD000155.
- International Working Group of AAGL, ESGE, ESHRE and WES; Tomassetti C, Johnson NP, Petrozza J, Abrao MS, Einarsson JI, Horne AW, Lee TTM, Missmer S, Vermeulen N, Zondervan KT et al. An international terminology for endometriosis, 2021. *Hum Reprod Open* 2021;**2021**:hoab029. hoab029.
- Iversen ML, Seyer-Hansen M, Forman A. Does surgery for deep infiltrating bowel endometriosis improve fertility? A systematic review. *Acta Obstet Gynecol Scand* 2017;**96**:688–693.
- Janssen EB, Rijkers AC, Hoppenbrouwers K, Meuleman C, D'Hooghe TM. Prevalence of endometriosis diagnosed by laparoscopy in adolescents with dysmenorrhea or chronic pelvic pain: a systematic review. *Hum Reprod Update* 2013;**19**:570–582.
- Jensen JT, Schlaff W, Gordon K. Use of combined hormonal contraceptives for the treatment of endometriosis-related pain: a systematic review of the evidence. *Fertil Steril* 2018;**110**:137–152.e131.
- Jin X, Ruiz Beguerie J. Laparoscopic surgery for subfertility related to endometriosis: a meta-analysis. *Taiwan J Obstet Gynecol* 2014;**53**: 303–308.
- Joseph J, Sahn SA. Thoracic endometriosis syndrome: new observations from an analysis of 110 cases. *Am J Med* 1996;**100**:164–170.
- Kaponis A, Chatzopoulos G, Paschopoulos M, Georgiou I, Paraskevaidis V, Zikopoulos K, Tsiveriotis K, Taniguchi F, Adonakis G, Harada T. Ultralong administration of gonadotropin-releasing hormone agonists before in vitro fertilization improves fertilization rate but not clinical pregnancy rate in women with mild endometriosis: a prospective, randomized, controlled trial. *Fertil Steril* 2020;**113**:828–835.
- Kennedy S, Bergqvist A, Chapron C, D'Hooghe T, Dunselman G, Greb R, Hummelshoj L, Prentice A, Saridogan E; ESHRE Special Interest Group for Endometriosis and Endometrium Guideline Development Group. ESHRE guideline for the diagnosis and treatment of endometriosis. *Hum Reprod* 2005;**20**:2698–2704.
- Khawaja UB, Khawaja AA, Gowani SA, Shoukat S, Ejaz S, Ali FN, Rizvi J, Nawaz FH. Frequency of endometriosis among infertile women and association of clinical signs and symptoms with the laparoscopic staging of endometriosis. *J Pak Med Assoc* 2009;**59**: 30–34.
- Kim SJ, Kim SK, Lee JR, Suh CS, Kim SH. Oocyte cryopreservation for fertility preservation in women with ovarian endometriosis. *Reprod Biomed Online* 2020;**40**:827–834.
- Kohl Schwartz AS, Wolfler MM, Mitter V, Rauchfuss M, Haeberlin F, Eberhard M, von Orelli S, Imthurn B, Imesch P, Fink D et al. Endometriosis, especially mild disease: a risk factor for miscarriages. *Fertil Steril* 2017;**108**:806–814.e2.
- Kolanska K, Cohen J, Bendifallah S, Selleret L, Antoine JM, Chabbert-Buffet N, Darai E, d'Argent EM. Pregnancy outcomes after controlled ovarian hyperstimulation in women with endometriosis-associated infertility: GnRH-agonist versus GnRH-antagonist. *J Gynecol Obstet Hum Reprod* 2017;**46**:681–686.
- Koninckx PR, Meuleman C, Oosterlynck D, Cornillie FJ. Diagnosis of deep endometriosis by clinical examination during menstruation and plasma CA-125 concentration. *Fertil Steril* 1996;**65**:280–287.
- Koshiba A, Mori T, Okimura H, Akiyama K, Kataoka H, Takaoka O, Ito F, Matsushima H, Kusuki I, Kitawaki J. Dienogest therapy during the early stages of recurrence of endometrioma might be an alternative therapeutic option to avoid repeat surgeries. *J Obstet Gynaecol Res* 2018;**44**:1970–1976.
- Kvaskoff M, Mahamat-Saleh Y, Farland LV, Shigesu N, Terry KL, Harris HR, Roman H, Becker CM, As-Sanie S, Zondervan KT et al. Endometriosis and cancer: a systematic review and meta-analysis. *Hum Reprod Update* 2021;**27**:393–420.
- Lalani S, Choudhry AJ, Firth B, Bacal V, Walker M, Wen SW, Singh S, Amath A, Hodge M, Chen I. Endometriosis and adverse maternal, fetal and neonatal outcomes, a systematic review and meta-analysis. *Hum Reprod* 2018;**33**:1854–1865.

- Lan S, Ling L, Jianhong Z, Xijing J, Lihui W. Analysis of the levonorgestrel-releasing intrauterine system in women with endometriosis. *J Int Med Res* 2013;**41**:548–558.
- Lee KH, Jung YW, Song SY, Kang BH, Yang JB, Ko YB, Lee M, Han HY, Yoo HJ. Comparison of the efficacy of dienogest and levonorgestrel-releasing intrauterine system after laparoscopic surgery for endometriosis. *J Obstet Gynaecol Res* 2018;**44**:1779–1786.
- Lee SY, Kim ML, Seong SJ, Bae JW, Cho YJ. Recurrence of ovarian endometrioma in adolescents after conservative, laparoscopic cyst enucleation. *J Pediatr Adolesc Gynecol* 2017;**30**:228–233.
- Leeners B, Damaso F, Ochsenbein-Kolble N, Farquhar C. The effect of pregnancy on endometriosis-facts or fiction? *Hum Reprod Update* 2018;**24**:290–299.
- Leone Roberti Maggiore U, Ferrero S, Mangili G, Bergamini A, Inversetti A, Giorgione V, Vigano P, Candiani M. A systematic review on endometriosis during pregnancy: diagnosis, misdiagnosis, complications and outcomes. *Hum Reprod Update* 2016;**22**:70–103.
- Leone Roberti Maggiore U, Inversetti A, Schimberni M, Vigano P, Giorgione V, Candiani M. Obstetrical complications of endometriosis, particularly deep endometriosis. *Fertil Steril* 2017;**108**:895–912.
- Lier MCI, Malik RF, Ket JCF, Lambalk CB, Brosens IA, Mijatovic V. Spontaneous hemoperitoneum in pregnancy (SHiP) and endometriosis – a systematic review of the recent literature. *Eur J Obstet Gynecol Reprod Biol* 2017;**219**:57–65.
- Liu E, Nisenblat V, Farquhar C, Fraser I, Bossuyt PM, Johnson N, Hull ML. Urinary biomarkers for the non-invasive diagnosis of endometriosis. *Cochrane Database Syst Rev* 2015;**2015**:CD012019.
- Lu D, Song H, Li Y, Clarke J, Shi G. Pentoxifylline for endometriosis. *Cochrane Database Syst Rev* 2012;**1**:CD007677.
- Manganaro L, Fierro F, Tomei A, Irimia D, Lodise P, Sergi ME, Vinci V, Sollazzo P, Porpora MG, Delfini R et al. Feasibility of 3.0T pelvic MR imaging in the evaluation of endometriosis. *Eur J Radiol* 2012;**81**:1381–1387.
- Maouris P. Asymptomatic mild endometriosis in infertile women: the case for expectant management. *Obstet Gynecol Surv* 1991;**46**:548–551.
- Margatho D, Carvalho NM, Bahamondes L. Endometriosis-associated pain scores and biomarkers in users of the etonogestrel-releasing subdermal implant or the 52-mg levonorgestrel-releasing intrauterine system for up to 24 months. *Eur J Contracept Reprod Health Care* 2020;**25**:133–140.
- Martire FG, Lazzeri L, Conway F, Siciliano T, Pietropolli A, Piccione E, Solima E, Centini G, Zupi E, Exacoustos C. Adolescence and endometriosis: symptoms, ultrasound signs and early diagnosis. *Fertil Steril* 2020;**114**:1049–1057.
- Matalliotakis I, Makrigiannakis A, Karkavitsas N, Psaroudakis E, Froudarakis G, Koumantakis E. Use of CA-125 in the diagnosis and management of endometriosis: influence of treatment with danazol. *Int J Fertil Menopausal Stud* 1994;**39**:100–104.
- Matorras R, Elorriaga MA, Pijoan JI, Ramon O, Rodriguez-Escudero FJ. Recurrence of endometriosis in women with bilateral adnexectomy (with or without total hysterectomy) who received hormone replacement therapy. *Fertil Steril* 2002;**77**:303–308.
- May KE, Conduit-Hulbert SA, Villar J, Kirtley S, Kennedy SH, Becker CM. Peripheral biomarkers of endometriosis: a systematic review. *Hum Reprod Update* 2010;**16**:651–674.
- May KE, Villar J, Kirtley S, Kennedy SH, Becker CM. Endometrial alterations in endometriosis: a systematic review of putative biomarkers. *Hum Reprod Update* 2011;**17**:637–653.
- Melin AS, Lundholm C, Malki N, Swahn ML, Sparen P, Bergqvist A. Hormonal and surgical treatments for endometriosis and risk of epithelial ovarian cancer. *Acta Obstet Gynecol Scand* 2013;**92**:546–554.
- Meuleman C, Tomassetti C, D'Hoore A, Van Cleynenbreugel B, Penninckx F, Vergote I, D'Hooghe T. Surgical treatment of deeply infiltrating endometriosis with colorectal involvement. *Hum Reprod Update* 2011;**17**:311–326.
- Meuleman C, Vandenabeele B, Fieuws S, Spiessens C, Timmerman D, D'Hooghe T. High prevalence of endometriosis in infertile women with normal ovulation and normospermic partners. *Fertil Steril* 2009;**92**:68–74.
- Michels KA, Pfeiffer RM, Brinton LA, Trabert B. Modification of the associations between duration of oral contraceptive use and ovarian, endometrial, breast, and colorectal cancers. *JAMA Oncol* 2018;**4**:516–521.
- Miller LE, Bhattacharyya R, Miller VM. Clinical utility of presacral neurectomy as an adjunct to conservative endometriosis surgery: systematic review and meta-analysis of controlled studies. *Sci Rep* 2020;**10**:6901.
- Moen MH, Stokstad T. A long-term follow-up study of women with asymptomatic endometriosis diagnosed incidentally at sterilization. *Fertil Steril* 2002;**78**:773–776.
- Mol BW, Bayram N, Lijmer JG, Wiegerinck MA, Bongers MY, van der Veen F, Bossuyt PM. The performance of CA-125 measurement in the detection of endometriosis: a meta-analysis. *Fertil Steril* 1998;**70**:1101–1108.
- Momoeda M, Harada T, Terakawa N, Aso T, Fukunaga M, Hagino H, Taketani Y. Long-term use of dienogest for the treatment of endometriosis. *J Obstet Gynaecol Res* 2009;**35**:1069–1076.
- Morotti M, Remorgida V, Venturini PL, Ferrero S. Endometriosis in menopause: a single institution experience. *Arch Gynecol Obstet* 2012;**286**:1571–1575.
- Mossa B, Ebano V, Tucci S, Rega C, Dolce E, Frega A, Marziani R. Laparoscopic surgery for the management of ovarian endometriomas. *Med Sci Monit* 2010;**16**:MT45–50.
- Moura APC, Ribeiro HSAA, Bernardo WM, Simões R, Torres US, D'Ippolito G, Bazot M, Ribeiro PAAG. Accuracy of transvaginal sonography versus magnetic resonance imaging in the diagnosis of rectosigmoid endometriosis: systematic review and meta-analysis. *PLoS One* 2019;**14**:e0214842.
- Moustafa S, Burn M, Mamillapalli R, Nematian S, Flores V, Taylor HS. Accurate diagnosis of endometriosis using serum microRNAs. *Am J Obstet Gynecol* 2020;**223**:e551–e557.
- Murta M, Machado RC, Zegers-Hochschild F, Checa MA, Sampaio M, Geber S. Endometriosis does not affect live birth rates of patients submitted to assisted reproduction techniques: analysis of the Latin American Network Registry database from 1995 to 2011. *J Assist Reprod Genet* 2018;**35**:1395–1399.
- Muteshi CM, Ohuma EO, Child T, Becker CM. The effect of endometriosis on live birth rate and other reproductive outcomes in ART cycles: a cohort study. *Hum Reprod Open* 2018;**2018**:hoy016.
- Muzii L, Achilli C, Bergamini V, Candiani M, Garavaglia E, Lazzeri L, Lecce F, Maiorana A, Maneschi F, Marana R et al. Comparison

- between the stripping technique and the combined excisional/ablative technique for the treatment of bilateral ovarian endometriomas: a multicentre RCT. *Hum Reprod* 2016a;**31**:339–344.
- Muzii L, Achilli C, Lecce F, Bianchi A, Franceschetti S, Marchetti C, Perniola G, Panici PB. Second surgery for recurrent endometriomas is more harmful to healthy ovarian tissue and ovarian reserve than first surgery. *Fertil Steril* 2015;**103**:738–743.
- Muzii L, Bellati F, Palaia I, Plotti F, Mancini N, Zullo MA, Angioli R, Panici PB. Laparoscopic stripping of endometriomas: a randomized trial on different surgical techniques. Part I: clinical results. *Hum Reprod* 2005;**20**:1981–1986.
- Muzii L, Di Tucci C, Achilli C, Di Donato V, Musella A, Palaia I, Panici PB. Continuous versus cyclic oral contraceptives after laparoscopic excision of ovarian endometriomas: a systematic review and metaanalysis. *Am J Obstet Gynecol* 2016b;**214**:203–211.
- Namnoum AB, Hickman TN, Goodman SB, Gehlbach DL, Rock JA. Incidence of symptom recurrence after hysterectomy for endometriosis. *Fertil Steril* 1995;**64**:898–902.
- Nezhat C, Main J, Paka C, Nezhat A, Beygui RE. Multidisciplinary treatment for thoracic and abdominopelvic endometriosis. *JSL* 2014;**18**:e2014.00312.
- Nezhat C, Santolaya J, Nezhat FR, Nezhat C. Comparison of transvaginal sonography and bimanual pelvic examination in patients with laparoscopically confirmed endometriosis. *J Am Assoc Gynecol Laparosc* 1994;**1**:127–130.
- Nickkho-Amiry M, Savant R, Majumder K, Edi-O'sagie E, Akhtar M. The effect of surgical management of endometrioma on the IVF/ICSI outcomes when compared with no treatment? A systematic review and meta-analysis. *Arch Gynecol Obstet* 2018;**297**:1043–1057.
- Nisenblat V, Bossuyt PM, Shaikh R, Farquhar C, Jordan V, Scheffers CS, Mol BW, Johnson N, Hull ML. Blood biomarkers for the non-invasive diagnosis of endometriosis. *Cochrane Database Syst Rev* 2016a:CD012179.
- Nisenblat V, Bossuyt PM, Farquhar C, Johnson N, Hull ML; Cochrane Gynaecology and Fertility Group. Imaging modalities for the non-invasive diagnosis of endometriosis. *Cochrane Database Syst Rev* 2016b;**2**:CD009591.
- Nnoaham KE, Hummelshoj L, Kennedy SH, Jenkinson C, Zondervan KT; World Endometriosis Research Foundation Women's Health Symptom Survey Consortium. Developing symptom-based predictive models of endometriosis as a clinical screening tool: results from a multicenter study. *Fertil Steril* 2012;**98**:692–701.e695.
- Nodler JL, Harris HR, Chavarro JE, Frazier AL, Missmer SA. Dairy consumption during adolescence and endometriosis risk. *Am J Obstet Gynecol* 2020;**222**:257.e1–257.e16.
- Nulsen JC, Walsh S, Dumez S, Metzger DA. A randomized and longitudinal study of human menopausal gonadotropin with intrauterine insemination in the treatment of infertility. *Obstet Gynecol* 1993;**82**:780–786.
- Omland AK, Tanbo T, Dale PO, Abyholm T. Artificial insemination by husband in unexplained infertility compared with infertility associated with peritoneal endometriosis. *Hum Reprod* 1998;**13**:2602–2605.
- Opoien HK, Fedorcsak P, Byholm T, Tanbo T. Complete surgical removal of minimal and mild endometriosis improves outcome of subsequent IVF/ICSI treatment. *Reprod Biomed Online* 2011;**23**:389–395.
- Osuga Y, Seki Y, Tanimoto M, Kusumoto T, Kudou K, Terakawa N. Relugolix, an oral gonadotropin-releasing hormone receptor antagonist, reduces endometriosis-associated pain in a dose-response manner: a randomized, double-blind, placebo-controlled study. *Fertil Steril* 2021;**115**:397–405.
- Pabuccu R, Onalan G, Kaya C. GnRH agonist and antagonist protocols for stage I-II endometriosis and endometrioma in in vitro fertilization/intracytoplasmic sperm injection cycles. *Fertil Steril* 2007;**88**:832–839.
- Parazzini F, Cipriani S, Bravi F, Pelucchi C, Chiaffarino F, Ricci E, Vigano P. A metaanalysis on alcohol consumption and risk of endometriosis. *Am J Obstet Gynecol* 2013a;**209**:106.e1–e101–110.
- Parazzini F, Vigano P, Candiani M, Fedele L. Diet and endometriosis risk: a literature review. *Reprod Biomed Online* 2013b;**26**:323–336.
- Paulson JD, Paulson JN. Anterior vaginal wall tenderness (AVWT) as a physical symptom in chronic pelvic pain. *JSL* 2011;**15**:6–9.
- Pavone ME, Bulun SE. Aromatase inhibitors for the treatment of endometriosis. *Fertil Steril* 2012;**98**:1370–1379.
- Pearce CL, Templeman C, Rossing MA, Lee A, Near AM, Webb PM, Nagle CM, Doherty JA, Cushing-Haugen KL, Wicklund KG et al. Association between endometriosis and risk of histological subtypes of ovarian cancer: a pooled analysis of case-control studies. *Lancet Oncol* 2012;**13**:385–394.
- Perez-Lopez FR, Villagrasa-Boli P, Munoz-Olarte M, Morera-Grau A, Cruz-Andres P, Hernandez AV. Association between endometriosis and preterm birth in women with spontaneous conception or using assisted reproductive technology: a systematic review and meta-analysis of cohort studies. *Reprod Sci* 2018;**25**:311–319.
- Petraglia F, Hornung D, Seitz C, Faustmann T, Gerlinger C, Luisi S, Lazzeri L, Strowitzki T. Reduced pelvic pain in women with endometriosis: efficacy of long-term dienogest treatment. *Arch Gynecol Obstet* 2012;**285**:167–173.
- Pittaway DE. The use of serial CA 125 concentrations to monitor endometriosis in infertile women. *Am J Obstet Gynecol* 1990;**163**:1032–1035; discussion 1035–1037.
- Polyzos NP, Fatemi HM, Zavos A, Grimbizis G, Kyrou D, Velasco JG, Devroey P, Tarlatzis B, Papanikolaou EG. Aromatase inhibitors in post-menopausal endometriosis. *Reprod Biol Endocrinol* 2011;**9**:90.
- Porpora MG, Pallante D, Ferro A, Crisafi B, Bellati F, Benedetti Panici P. Pain and ovarian endometrioma recurrence after laparoscopic treatment of endometriosis: a long-term prospective study. *Fertil Steril* 2010;**93**:716–721.
- Proctor M, Latthe P, Farquhar C, Khan K, Johnson N; Cochrane Gynaecology and Fertility Group; Surgical interruption of pelvic nerve pathways for primary and secondary dysmenorrhoea. *Cochrane Database Syst Rev* 2005;
- Pundir J, Omanwa K, Kovoor E, Pundir V, Lancaster G, Barton-Smith P. Laparoscopic excision versus ablation for endometriosis-associated pain: an updated systematic review and meta-analysis. *J Minim Invasive Gynecol* 2017;**24**:747–756.
- Qiu Y, Yuan S, Wang H. Vitamin D status in endometriosis: a systematic review and meta-analysis. *Arch Gynecol Obstet* 2020;**302**:141–152.

- Razzi S, Luisi S, Ferretti C, Calonaci F, Gabbanini M, Mazzini M, Petraglia F. Use of a progestogen only preparation containing desogestrel in the treatment of recurrent pelvic pain after conservative surgery for endometriosis. *Eur J Obstet Gynecol Reprod Biol* 2007; **135**:188–190.
- Redwine DB. Endometriosis persisting after castration: clinical characteristics and results of surgical management. *Obstet Gynecol* 1994; **83**:405–413.
- Ricci E, Vigano P, Cipriani S, Chiaffarino F, Bianchi S, Rebonato G, Parazzini F. Physical activity and endometriosis risk in women with infertility or pain: systematic review and meta-analysis. *Medicine (Baltimore)* 2016; **95**:e4957.
- Ripps BA, Martin DC. Correlation of focal pelvic tenderness with implant dimension and stage of endometriosis. *J Reprod Med* 1992; **37**:620–624.
- Rodriguez-Purata J, Coroleu B, Tur R, Carrasco B, Rodriguez I, Barri PN. Endometriosis and IVF: are agonists really better? Analysis of 1180 cycles with the propensity score matching. *Gynecol Endocrinol* 2013; **29**:859–862.
- Roman JD. Adolescent endometriosis in the Waikato region of New Zealand—a comparative cohort study with a mean follow-up time of 2.6 years. *Aust N Z J Obstet Gynaecol* 2010; **50**:179–183.
- Rossing MA, Cushing-Haugen KL, Wicklund KG, Doherty JA, Weiss NS. Risk of epithelial ovarian cancer in relation to benign ovarian conditions and ovarian surgery. *Cancer Causes Control* 2008; **19**:1357–1364.
- Sandström A, Bixo M, Johansson M, Bäckström T, Turkmen S. Effect of hysterectomy on pain in women with endometriosis: a population-based registry study. *BJOG* 2020; **127**:1628–1635.
- Santulli P, Marcellin L, Menard S, Thubert T, Khoshnood B, Gayet V, Goffinet F, Ancel PY, Chapron C. Increased rate of spontaneous miscarriages in endometriosis-affected women. *Hum Reprod* 2016; **31**:1014–1023.
- Saraswat L, Ayansina DT, Cooper KG, Bhattacharya S, Miligkos D, Horne AW, Bhattacharya S. Pregnancy outcomes in women with endometriosis: a national record linkage study. *BJOG* 2017; **124**:444–452.
- Sasamoto N, DePari M, Vitonis AF, Laufer MR, Missmer SA, Shafir AL, Terry KL. Evaluation of CA125 in relation to pain symptoms among adolescents and young adult women with and without surgically-confirmed endometriosis. *PLoS One* 2020; **15**:e0238043.
- Sauerbrun-Cutler MT, Alvero R. Short- and long-term impact of gonadotropin-releasing hormone analogue treatment on bone loss and fracture. *Fertil Steril* 2019; **112**:799–803.
- Seckin B, Ates MC, Kirbas A, Yesilyurt H. Usefulness of hematological parameters for differential diagnosis of endometriomas in adolescents/young adults and older women. *Int J Adolesc Med Health* 2018; **33**. <https://doi.org/10.1515/ijamh-2018-0078>.
- Senapati S, Sammel MD, Morse C, Barnhart KT. Impact of endometriosis on in vitro fertilization outcomes: an evaluation of the Society for Assisted Reproductive Technologies Database. *Fertil Steril* 2016; **106**:164–171.e161.
- Seo JW, Lee DY, Yoon BK, Choi D. The efficacy of postoperative cyclic oral contraceptives after gonadotropin-releasing hormone agonist therapy to prevent endometrioma recurrence in adolescents. *J Pediatr Adolesc Gynecol* 2017; **30**:223–227.
- Seracchioli R, Mabrouk M, Manuzzi L, Vicenzi C, Frasca C, Elmakky A, Venturoli S. Post-operative use of oral contraceptive pills for prevention of anatomical relapse or symptom-recurrence after conservative surgery for endometriosis. *Hum Reprod* 2009; **24**:2729–2735.
- Serati M, Cattoni E, Braga A, Uccella S, Cromi A, Ghezzi F. Deep endometriosis and bladder and detrusor functions in women without urinary symptoms: a pilot study through an unexplored world. *Fertil Steril* 2013; **100**:1332–1336.
- Shah DK, Missmer SA. Scientific investigation of endometriosis among adolescents. *J Pediatr Adolesc Gynecol* 2011; **24**:S18–S19.
- Shakiba K, Bena JF, McGill KM, Minger J, Falcone T. Surgical treatment of endometriosis: a 7-year follow-up on the requirement for further surgery. *Obstet Gynecol* 2008; **111**:1285–1292.
- Shaltout MF, Elsheikhah A, Maged AM, Elsherbini MM, Zaki SS, Dahab S, Elkomy RO. A randomized controlled trial of a new technique for laparoscopic management of ovarian endometriosis preventing recurrence and keeping ovarian reserve. *J Ovarian Res* 2019; **12**:66.
- Smith JS, Green J, Berrington de Gonzalez A, Appleby P, Peto J, Plummer M, Franceschi S, Beral V. Cervical cancer and use of hormonal contraceptives: a systematic review. *Lancet* 2003; **361**:1159–1167.
- Somigliana E, Vigano P, Benaglia L, Busnelli A, Paffoni A, Vercellini P. Ovarian stimulation and endometriosis progression or recurrence: a systematic review. *Reprod Biomed Online* 2019; **38**:185–194.
- Song SY, Park M, Lee GW, Lee KH, Chang HK, Kwak SM, Yoo HJ. Efficacy of levonorgestrel releasing intrauterine system as a postoperative maintenance therapy of endometriosis: a meta-analysis. *Eur J Obstet Gynecol Reprod Biol* 2018; **231**:85–92.
- Soriano D, Adler I, Bouaziz J, Zolti M, Eisenberg VH, Goldenberg M, Seidman DS, Elizur SE. Fertility outcome of laparoscopic treatment in patients with severe endometriosis and repeated in vitro fertilization failures. *Fertil Steril* 2016; **106**:1264–1269.
- Stepniewska A, Pomini P, Guerriero M, Scioscia M, Ruffo G, Minelli L. Colorectal endometriosis: benefits of long-term follow-up in patients who underwent laparoscopic surgery. *Fertil Steril* 2010; **93**:2444–2446.
- Şükür YE, Özmen B, Yakıştıran B, Atabekoğlu CS, Berker B, Aytaç R, Sönmezer M. Endometrioma surgery is associated with increased risk of subsequent assisted reproductive technology cycle cancellation; a retrospective cohort study. *J Obstet Gynaecol* 2021; **41**:259–262.
- Sun PR, Leng JH, Jia SZ, Lang JH. Postmenopausal endometriosis: a retrospective analysis of 69 patients during a 20-year period. *Chin Med J (Engl)* 2013; **126**:4588–4589.
- Sutton CJ, Ewen SP, Whitelaw N, Haines P. Prospective, randomized, double-blind, controlled trial of laser laparoscopy in the treatment of pelvic pain associated with minimal, mild, and moderate endometriosis. *Fertil Steril* 1994; **62**:696–700.
- Tandoi I, Somigliana E, Riparini J, Ronzoni S, Vigano P, Candiani M. High rate of endometriosis recurrence in young women. *J Pediatr Adolesc Gynecol* 2011; **24**:376–379.
- Tang H, Wu R, Li X, Zhou Y, Liu Z, Wang C, Chen Y, Zhang F. Curative effect of 1.88-mg and 3.75-mg gonadotrophin-releasing hormone agonist on stage III-IV endometriosis: randomized controlled study. *J Obstet Gynaecol Res* 2017; **43**:1550–1554.

- Tanmahasamut P, Rattanachaiyanont M, Angsuwathana S, Techatrakaisak K, Indhavivadhana S, Leerasiri P. Postoperative levonorgestrel-releasing intrauterine system for pelvic endometriosis-related pain: a randomized controlled trial. *Obstet Gynecol* 2012;**119**:519–526.
- Taylor HS, Giudice LC, Lessey BA, Abrao MS, Kotarski J, Archer DF, Diamond MP, Surrey E, Johnson NP, Watts NB et al. Treatment of endometriosis-associated pain with elagolix, an oral GnRH antagonist. *N Engl J Med* 2017;**377**:28–40.
- Thomeer MG, Steensma AB, van Santbrink EJ, Willemsen FE, Wielopolski PA, Hunink MG, Spronk S, Laven JS, Krestin GP. Can magnetic resonance imaging at 3.0-Tesla reliably detect patients with endometriosis? Initial results. *J Obstet Gynaecol Res* 2014;**40**:1051–1058.
- Treloar SA, Bell TA, Nagle CM, Purdie DM, Green AC. Early menstrual characteristics associated with subsequent diagnosis of endometriosis. *Am J Obstet Gynecol* 2010;**202**:534–e531–536.
- Tummon IS, Asher LJ, Martin JS, Tulandi T. Randomized controlled trial of superovulation and insemination for infertility associated with minimal or mild endometriosis. *Fertil Steril* 1997;**68**:8–12.
- van der Houwen LE, Schreurs AM, Schats R, Heymans MW, Lambalk CB, Hompes PG, Mijatovic V. Efficacy and safety of intrauterine insemination in patients with moderate-to-severe endometriosis. *Reprod Biomed Online* 2014;**28**:590–598.
- Vanhie A, O D, Peterse D, Beckers A, Cuellar A, Fassbender A, Meuleman C, Mestdagh P, D'Hooghe T. Plasma miRNAs as biomarkers for endometriosis. *Hum Reprod* 2019;**34**:1650–1660.
- Vercellini P, Barbara G, Buggio L, Frattaruolo MP, Somigliana E, Fedele L. Effect of patient selection on estimate of reproductive success after surgery for rectovaginal endometriosis: literature review. *Reprod Biomed Online* 2012;**24**:389–395.
- Vercellini P, De Giorgi O, Mosconi P, Stellato G, Vicentini S, Crosignani PG. Cyproterone acetate versus a continuous monophasic oral contraceptive in the treatment of recurrent pelvic pain after conservative surgery for symptomatic endometriosis. *Fertil Steril* 2002;**77**:52–61.
- Vercellini P, Eskenazi B, Consonni D, Somigliana E, Parazzini F, Abbiati A, Fedele L. Oral contraceptives and risk of endometriosis: a systematic review and meta-analysis. *Hum Reprod Update* 2011;**17**:159–170.
- Vermeulen N, Le Clef N, Mcheik S, D'Angelo A, Tilleman K, Veleva Z, Nelen W. Manual for ESHRE guideline development. 2019. <https://www.eshre.eu/Guidelines-and-Legal/Guidelines/Guideline-development-process> (14 March 2022, date last accessed).
- Vicino M, Parazzini F, Cipriani S, Frontino G. Endometriosis in young women: the experience of GISE. *J Pediatr Adolesc Gynecol* 2010;**23**:223–225.
- Vigueras Smith A, Cabrera R, Kondo W, Ferreira H. Diaphragmatic endometriosis minimally invasive treatment: a feasible and effective approach. *J Obstet Gynaecol* 2021;**41**:176–186.
- Wentzensen N, Poole EM, Trabert B, White E, Arslan AA, Patel AV, Setiawan VW, Visvanathan K, Weiderpass E, Adami HO et al. Ovarian cancer risk factors by histologic subtype: an analysis from the ovarian cancer cohort consortium. *J Clin Oncol* 2016;**34**:2888–2898.
- Wright J, Lotfallah H, Jones K, Lovell D. A randomized trial of excision versus ablation for mild endometriosis. *Fertil Steril* 2005;**83**:1830–1836.
- Wu D, Hu M, Hong L, Hong S, Ding W, Min J, Fang G, Guo W. Clinical efficacy of add-back therapy in treatment of endometriosis: a meta-analysis. *Arch Gynecol Obstet* 2014;**290**:513–523.
- Yang Y, Wang Y, Yang J, Wang S, Lang J. Adolescent endometriosis in China: a retrospective analysis of 63 cases. *J Pediatr Adolesc Gynecol* 2012;**25**:295–299.
- Yeung P Jr, Sinervo K, Winer W, Albee RB Jr. Complete laparoscopic excision of endometriosis in teenagers: is postoperative hormonal suppression necessary? *Fertil Steril* 2011;**95**:1909–1912, 1912.e1901.
- Yong PJ, Matwani S, Brace C, Quaiattini A, Bedaiwy MA, Albert A, Allaire C. Endometriosis and ectopic pregnancy: a meta-analysis. *J Minim Invasive Gynecol* 2020;**27**:352–361.e352.
- Yoost J, LaJoie AS, Hertweck P, Loveless M. Use of the levonorgestrel intrauterine system in adolescents with endometriosis. *J Pediatr Adolesc Gynecol* 2013;**26**:120–124.
- Younis JS, Shapso N, Fleming R, Ben-Shlomo I, Izhaki I. Impact of unilateral versus bilateral ovarian endometriotic cystectomy on ovarian reserve: a systematic review and meta-analysis. *Hum Reprod Update* 2019;**25**:375–391.
- Zakhari A, Delpero E, McKeown S, Tomlinson G, Bougie O, Murji A. Endometriosis recurrence following post-operative hormonal suppression: a systematic review and meta-analysis. *Hum Reprod Update* 2021;**27**:96–107.
- Zhu X, Chen L, Deng X, Xiao S, Ye M, Xue M. A comparison between high-intensity focused ultrasound and surgical treatment for the management of abdominal wall endometriosis. *BJOG* 2017;**124** Suppl 3:53–58.
- Zondervan KT, Becker CM, Missmer SA. Endometriosis. *N Engl J Med* 2020;**382**:1244–1256.
- Zorbas KA, Economopoulos KP, Vlahos NF. Continuous versus cyclic oral contraceptives for the treatment of endometriosis: a systematic review. *Arch Gynecol Obstet* 2015;**292**:37–43.
- Zucchetto A, Serraino D, Polesel J, Negri E, De Paoli A, Dal Maso L, Montella M, La Vecchia C, Franceschi S, Talamini R. Hormone-related factors and gynecological conditions in relation to endometrial cancer risk. *Eur J Cancer Prev* 2009;**18**:316–321.