

<https://helda.helsinki.fi>

Dietary fat intake and quality in long-term care residents in two cohorts assessed 10 years apart

Jyväkorpi , Satu K.

2022-04-12

Jyväkorpi , S K , Suominen , M H , Strandberg , T E , Salminen , K , Niskanen , R T , Roitto , H-M , Saarela , R K T & Pitkälä , K H 2022 , ' Dietary fat intake and quality in long-term care residents in two cohorts assessed 10 years apart ' , BMC Nutrition , vol. 8 , no. 1 , 31 . <https://doi.org/10.1186/s40795-022-00524-9>

<http://hdl.handle.net/10138/343064>

<https://doi.org/10.1186/s40795-022-00524-9>

cc_by

publishedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

RESEARCH

Open Access



Dietary fat intake and quality in long-term care residents in two cohorts assessed 10 years apart

K. Jyväkorpi Satu^{1*}, H. Suominen Merja¹, E. Strandberg Timo^{1,2}, Karoliina Salminen¹, T. Niskanen Riikka¹, Hanna-Maria Raitto^{1,3}, K. T. Saarela Riitta⁴ and H. Pitkälä Kaisu¹

Abstract

Purpose: To describe and compare detailed dietary fat intake, fat quality and associative factors between two measuring points 10 years apart of residents living in long-term care facilities, and to reflect how fat composition and fat quality corresponds to current nutrition recommendations.

Methods: In 2007 long-term care residents ($n=374$) of 25 assisted-living facilities and nursing homes and in 2017–18 long-term care residents ($n=486$) of 17 respective facilities in Helsinki metropolitan area were recruited for this study. Information on the residents' heights, demographic information and use of calcium and vitamin D supplementation were retrieved from medical records. Residents' clinical assessment included Clinical Dementia Rating (CDR), the Mini Nutritional Assessment (MNA) and questionnaire related to nutrition care. Participants' energy and fat intake were determined from 1–2-day food diaries kept by the ward nurses, and fat quality indicators calculated.

Results: Age, gender distribution, MNA score or body mass index did not differ between the two cohorts. Residents' cognitive status, subjective health and mobility were poorer in 2017 compared to 2007. Total fat and saturated fatty acid (SFA) intakes were higher and fat quality indicators lower in the 2017 cohort residents than in the 2007 cohort residents. Sugar intake, male gender, eating independently, eating larger amounts and not having dry mouth predicted higher SFA intake in the 2017 cohort.

Conclusions: The fat quality in long-term care residents in our study worsened in spite of official recommendations between the two measurement points.

Keywords: Fat quality, Fat composition, Saturated fatty acids, Monounsaturated fatty acids, Polyunsaturated fatty acids, Long-term care

Background

Fat is an important macronutrient in the human diet. It is a major source of energy and it facilitates the absorption of fat-soluble vitamins, carotenoids and other phytochemicals [1]. It is also a source for essential fatty acids

(EFA) present particularly in vegetable oils, nuts, seeds and fish, which are necessary for development, normal function of nervous and cardiovascular systems and general health [2]. EFAs cannot be synthesized by humans and must be provided by the diet [3]. Fat also contributes to the acceptability and texture of foods, and improves taste by enhancing flavor and aroma.

Institutionalized older people often have cognitive and physical impairments along with many chronic illnesses [4]. Malnutrition is thus commonly encountered

*Correspondence: satujyvakorpi@gery.fi

¹ Department of General Practice and Primary Health Care, and Helsinki University Central Hospital, Unit of Primary Health Care, University of Helsinki, Tukholmankatu 8, 00014 Helsingin yliopisto Helsinki, Finland
Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

in long-term care residents and it weakens quality of life and increases morbidity and mortality [5–7]. Frailty, poor cognitive status, polypharmacy, gastro-intestinal symptoms, poor dentition, and swallowing difficulties often lead to inadequate energy and nutrient intakes, weight loss and malnutrition in long-term care residents [8–11]. It is a common practice in long-term care facilities to use more fat to prepare foods in order to increase acceptability and energy content of served foods [12]. Thus, fat has an important role in the diets of long-term care residents.

Recently in Finland, a special nutrition recommendation for older people was published, emphasizing along with sufficient protein intake, diet quality and other issues, the importance of fat quality also in the frailest older adults in long-term care [13]. In the recommendation -- similarly to many other national authorities' and expert groups' recommendations -- good fat quality is defined as keeping saturated fatty acid (SFA) intake lower than 10% of energy (E%) and replacing SFAs with monounsaturated fatty acids (MUFA) and polyunsaturated fatty acids (PUFA) [14–18]. Good fat quality has been previously shown to have an important role in cardiovascular and cognitive health, and may thus be important also for the health of the frailest older adults [14, 19, 20].

The purpose of this study was to describe and compare dietary fat intake, its detailed composition and quality and associative factors in two cohorts of residents living in long-term care facilities and in the same geographical area, measured 10 years apart, and to investigate how the fat composition and quality corresponds to the current nutrition recommendations.

Methods

In 2007, a sample of long-term care residents from 25 assisted-living facilities and nursing homes in Helsinki metropolitan area were recruited for the original nutrition study. In 2017/18 we conducted a follow-up study and recruited volunteer residents from a sample of 17 assisted-living facilities and nursing homes in Helsinki. The long-term care facilities were randomly selected within voluntary institutions. Six of the original facilities were included in the 2017/18 sample. In all the institutions both in those included in 2007 and in 2017/18, registered nurses were in charge of the wards and constant 24/7 assistance was available.

The inclusion criteria for this present study were: age \geq 65 years, living permanently in long-term care facility, sufficient information available on demographic factors and nutritional care, a filled 1--2-day food diary. Trained nurses collected the data both 2007 and 2017/18. The participants' weights were measured. Their heights were obtained from the medical records,

and body mass index (BMI) was calculated as weight divided by height squared (kg/m^2). Information on the residents' demographic information and use of calcium and vitamin D supplementation were retrieved from medical records. The cognitive status of the residents in both cohorts (2007, 2017/18) was measured using Clinical Dementia Rating (CDR) [21]. Mobility was assessed with one item in MNA questionnaire and categorized as: bed or chair bound, able to get out of bed/chair, but does not go out or goes out. Similarly, self-rated health was also assessed by one item in MNA and categorized as: considers oneself healthy or quite healthy, considers oneself sick or very sick, not able to answer. Nutritional status was assessed, using the Mini Nutritional Assessment (MNA) long version [22]. In addition, several questions associated with nutritional care were asked. These included: 1) amounts of eaten foods (very little or little, normal or a lot); 2) whether resident eats snack (yes, no); 3) texture of food (liquid or puree, soft, normal); 4) use of oral nutritional supplements (yes, no); 5) frequency of weight monitoring (2–6 times a year, >6 times a year); 6) chewing problems (yes, no); 7) dry mouth (yes, no); 8) pain in the mouth (yes, no); 9) dysphagia (yes, no); and 10) need of help with eating (yes, no).

Participants' energy and detailed fat intakes were determined from 1--2-day food diaries kept by the ward nurses. Prior to the data collection, the nurses participated in comprehensive training sessions on how to fill food diaries for the residents organized by the study's investigators (MS) in 2007 and (MS, SKJ) in 2017. In 2007, the food diaries were analyzed using Nutrica dietary software (version 3.11, Kela, Turku, Finland), and in 2017, using Aivo Diet dietary software (version 2.2.0.0, Aivo Oy, Turku, Finland) both containing the Fineli Food Composition database including foods and recipes for the typical Finnish mixed dishes that are customarily served in long-term care. The instruction was to record all the foods and beverages consumed by the resident. The nurses estimated portion sizes, using household measures. For prepacked products, the exact brand and product name were required. Dietary data including energy, total fat intake and fat composition of diet (including SFA, MUFA, and PUFA) were analyzed from food diaries. Using these data, we calculated fat quality indicators (MUFA:SFA- and PUFA:SFA- ratios), and percentage of energy from SFAs, MUFAs and PUFAs in the diets of the residents. In 2017, we also have data of more detailed fat composition, such as amounts of n-3 fatty acids, n-6 fatty acids, their ratios, or amounts of trans fatty acids. Due to limitations of the earlier dietary analyzing tool, these data were not available in 2007.

Statistical analysis

The data from the two cohorts were combined and descriptive statistics presented. The differences between the baseline characteristics, energy and detailed fat intakes and fat quality of groups were analyzed between the two cohorts (2007, 2017), using the Chi² -test or Fisher's exact test for categorical variables and t-test or Mann Whitney U-test for continuous variables. SFA intake from 2017/18 cohort was divided into quartiles and nutrition related factors classified to those quartiles accordingly. The SFA intake quartiles of 2017/18 cohort were as follows; Q₁<23.5263 g; Q₂=23.5263 g--29.7475 g; Q₃=29.7476 g–37.7125 g; Q₄>37.7125 g. Differences between nutritional factors classified into SFA quartiles were analyzed using Cochran-Armitage test for trend. Analysis of covariance (ANCOVA) test was used to investigate independent associations with SFA intake. Univariate general linear model was used to explore these associations. We explored associations of 2017 and of 2007 cohort separately. For the 2017 cohort: model 1 includes intercept, age, sex and sugar intake, and model 2 additionally includes need of help with eating (yes vs. no) and excludes age as a covariate. For the 2007 cohort: model 1 included the same characteristics that in 2017 cohort, in the model 2 additionally need of help with eating, and in the model 3 additionally total MNA score excluding need of help with eating and sex as covariates.

Selection of covariates

Age was selected as a covariate since older age is associated with more risk of malnutrition. Gender was selected as covariate, because females and males might differ in dietary preferences even in long- term care. Higher degree of dependency is a known risk factor for malnutrition. Thus needing help with eating was selected as a covariate [23] and total MNA score as a covariate, since it is a validated instrument to identify malnutrition [22]. Dietary sugar intake was selected as a covariate for the model because along with fat, increment of sugar is common in long- term care to increase acceptability and energy content of the diet and thereby prevent weight loss [24]. Accordingly, sugar in various forms (sugary drinks and snacks; adding sugar to coffee, tea, porridge, and gruel, etc.) is frequently served. The statistical tests were performed using IBM SPSS statistical program (version 26, Chicago, US).

Ethics

The ethics committee approvals were obtained for both 2007 and 2017/18 by the ethics committee of the Department of Medicine at Helsinki University Hospital and City of Helsinki (ethical approval number:

HUS/2042/2016). Informed written consent was asked from all participants or in cases of moderate to severe dementia (MMSE <20 points), from their closest proxies.

Results

In total, 860 volunteer residents, which included 374 residents from 2007 cohort and 486 residents from 2017/18 cohort, were included in the study. Gender distribution, BMI nor MNA scores did not differ between the two cohorts (Table 1). Participants' cognitive state measured using CDR, mobility and subjective health were poorer in the 2017/18 cohort compared to the 2007 cohort. Use of calcium supplements was more frequent in 2007 (47.2%) compared to 2017/18 (35.1%), $p < 0.001$, whereas vitamin D supplementation was more frequent in 2017/18 (82.5%) compared to 2007 (54.7%), $p < 0.001$.

Energy intake in females was lower in 2017/18 (1584 kcal) compared to 2007 (1653 kcal), whereas no differences were observed in energy intake in male residents between the two cohorts (Table 2). Total fat and SFA intakes were significantly higher in males and females in the 2017/18 cohort than in 2007, respectively, whereas the total amounts of MUFA or PUFA did not differ between the two cohorts in female or male residents.

Fat quality indicators PUFA:SFA- ratio and MUFA:SFA- ratio differed significantly between the two cohorts. The females and males in the 2017/18 cohort had lower PUFA:SFA- ratios than observed in the 2007 cohort. Similarly, MUFA:SFA- ratios in both females and males were lower in the participants in the 2017/18 cohort compared to the 2007 cohort.

Mean PUFA n-3 intake was 3 g and n-6 intake 5 g in 2017/18 in participating residents, and ratio between n-3:n-6 was 0.6. We did not have this data from the 2007 cohort. Percentage of energy from total fat (Fat E%) and SFA were significantly higher in the 2017/18 cohort than 2007 cohort, 35 E% and 31 E%, and 17 E% and 13 E%, respectively.

The SFA intake quartiles were not associated with texture of the food, eating snacks, using oral nutritional supplements, more frequent weight monitoring, chewing problems, dysphagia, whereas they were linearly associated with the amount of eaten foods, and inversely associated with having dry mouth (Table 3). Those who had dry mouth consumed less energy than those who did not have dry mouth (1483 kcal vs. 1657 kcal). In addition, eating independently was linearly associated with higher SFA intake. Figure 1 illustrates how the SFA intakes are in line with nutrition recommendation.

In the ANCOVA model 1, dietary sugar intake and male gender were associated with higher SFA intake in the 2017 cohort, whereas age was not associated with SFA intake (adjusted $R^2=0.236$). In the model 2, dietary

Table 1 Baseline characteristics of two long-term care cohorts

Characteristics	Cohort of 2007; n = 374	Cohort of 2017/2018; n = 486	CI 95%	p-value ¹
Females, %	82.4	79.4		0.30
Age, years (SD)	83.3 (7.4)	82.4 (7.6)	-0.507, 1.505	0.33
MNA, total score (SD)	20.2 (3.5)	20.3 (3.4)	-0.588, 0.373	0.15
Nutritional status (MNA), %				
Good nutritional status, > 23	17.4	16.2		0.82
Risk of malnutrition 23.5–17	68.2	68.0		
Malnourished < 17	14.4	15.8		
BMI, kg/m ² (SD)	25.3 (4.8)	26.0 (5.0)	-1.339, 0.025	0.74
CDR, total score (SD) classification, %	1.8 (1.0)	2.6 (0.7)	-0.838, -0.610	< 0.001
0.5–1 Mild dementia	25.6	8.2		< 0.001
2 Moderate	39.5	26.5		
3 Severe	35.0	65.3		
Diabetes mellitus, %	16.2	16.6		0.45
Coronary heart disease, %	26.4	17.4		0.001
Coronary thrombosis, %	11.6	3.2		< 0.001
Stroke or TIA	24.1	22.9		0.38
Dementia	78.6	80.0		0.33
Subjective health, %				
considers oneself healthy or quite healthy	74.7	48.4		< 0.001
considers oneself sick or very sick	25.4	12.3		
Not able to answer	0	39.3		
Mobility, %				
bed or chair bound	13.6	47.1		< 0.001
able to get out of bed/chair but does not go out	45.5	26.4		
goes out	40.9	26.6		
Use of calcium supplementation, %	47.2	35.1		< 0.001
Use of vitamin D supplementation, %	54.7	82.5		< 0.001

SD Standard deviation, CI Confidence Interval, MNA Mini Nutritional Assessment, BMI Body mass index, kg kilogram, m meter; CDR Clinical Dementia Rating

¹ Statistical significance for p-value was set to < 0.05

sugar intake, eating independently and male gender were associated with higher SFA intake, adjusted ($R^2=0.244$). In the 2007 cohort, in model 1 higher SFA intake was associated with sugar intake and age, whereas gender was not associated with SFA intake, adjusted $R^2=0.061$ (Table 4). In the model 2 SFA intake was associated with sugar intake, age, whereas gender or eating independently were not associated with SFA intake (adjusted $R^2=0.065$). In the model 3 SFA intake was associated with sugar intake, age and total MNA score (adjusted $R^2=0.068$).

Discussion

We observed that dietary fat quality in the long-term care facilities was poor in 2007 but even worse in 2017/18, in spite of the official nutrition recommendations. This worsening was due to significant increase of SFA intake and consequently dietary fat quality indicators worsened in older long-term care residents. Higher SFA intake

of the 2017 cohort was predicted by sugar intake, male gender, eating independently, eating higher amounts of foods, and not having dry mouth.

Very few studies have described fat composition profile of long-term care residents. In a study by Rodríguez-Rejón et al. [25] fat composition of Spanish long-term care facilities was reported, but quality indicators were not calculated. In their study, SFA intake was about half of that in 2017/18 cohort in our study. In a Canadian study, SFA intake evaluated from long-term care facilities' menus was about the same as in the 2007 cohort, but MUFA and PUFA intakes were considerably higher than in either of the cohorts in our study [26].

Recommendations on fat quality are uniform across various national authorities and expert groups [14–18]. Generally, all authorities encourage reducing SFA intake to less than 10 E% or even reducing it even more and replacing it with MUFA and PUFA [15]. In our study, fat quality in the long-term care residents was already

Table 2 Energy, detailed fat intake and fat quality indicators between long-term cohorts of long term care residents in 2007 and 2017

Energy and fat intakes, fat quality and vitamins D and E intakes, mean (SD)	Cohort of 2007 n=374	Cohort of 2017/2018 n=486	CI 95%	p-value ¹	Nutrition Recommendation ^a
Energy, kcal (SD)	1691 (443)	1630 (397)	4.29, 118.71	0.04	
females	1653 (409)	1584 (393)	9.16, 129.13	0.02	
males	1870 (545)	1809 (363)	-90.89, 212.08	0.43	
Total fat, g (SD)	59 (21)	64 (20)	-7.80, -2.34	< 0.001	
females	57 (20)	61 (20)	-7.14, -1.19	0.006	
males	65 (22)	73 (18)	-13.75, -1.33	0.018	
SFA, g (SD)	24 (10)	31 (11)	-7.98, -5.20	< 0.001	
females	24 (10)	30 (10)	-7.40, -4.37	< 0.001	
males	26 (10)	35 (11)	-12.33, -5.79	< 0.001	
TRANS FA, g (SD)		1.3 (0.4)			
females	N/A	1.2 (0.4)			
males		1.4 (0.4)			
MUFA, g (SD)	18 (7)	19 (6)	-1.54, 0.19	0.13	
females	18 (7)	18 (6)	-1.50, 0.40	0.26	
males	21 (7)	21 (5)	-2.85, 1.27	0.45	
PUFA, g (SD)	7 (3)	7 (3)	-0.21, 0.62	0.33	
females	7 (3)	7 (2)	-0.28, 0.59	0.48	
males	9 (4)	8 (2)	-0.45, 1.75	0.25	
N-3		3 (1)			
N-6	N/A	5 (2)			
N-6:N-3		1.7 (1)			
PUFA/SFA (SD)	0.32 (0.17))	0.24 (0.14)	0.05, 0.10	< 0.001	
females	0.31 (0.17)	0.24 (0.09)	0.05, 0.10	< 0.001	
males	0.34 (0.17)	0.26 (0.26)	0.12, 0.16	0.022	
MUFA/SFA (SD)	0.78 (0.23)	0.63 (0.16)	0.12, 0.18	< 0.001	
females	0.78 (0.24)	0.63 (0.14)	0.12, 0.17	< 0.001	
males	0.81 (0.21)	0.64 (0.21)	0.11, 0.24	< 0.001	
FAT E%	31 E%	35 E%	-0.05, -0.03	< 0.001	25–40 E%
SFA E%	13 E%	17 E%	-0.05, -0.04	< 0.001	< 10 E%
MUFA E%	9.7 E%	10.4 E%	-0.07, -0.04	< 0.001	10–20 E%
PUFA E%	3.8 E%	4.3 E%	-0.01, -0.04	< 0.001	5–10 E%

SD Standard deviation, CI Confidence Interval, SFA Saturated fatty acids, MUFA Monounsaturated fatty acids, PUFA Polyunsaturated fatty acids, E% Percentage of total energy

¹ Statistical significance for p-value was set to < 0.05

^a Nordic Nutrition Recommendation 2014

poor in 2007, but even worse in 2017: SFA intake of the residents was on average 17 E% compared to the 13 E% in 2007. Moreover, in both of the cohorts, intake of PUFAs was lower than the recommended 5–10 E%. Malnutrition and its risk in the residents was high (> 80%) in both measuring points, which is common in long-term care facilities [5–7]. In order to avoid residents' weight loss it is a common practice to increase energy content of the served meals by adding fat to the meals during preparation [12]. Traditionally this has meant adding butter or cream to various foods and using whole milk products instead of low fat or fat free products. In the newly


published recommendation in Finland targeted especially for older people, fat quality is seen as an important issue, and the use of vegetable oils and soft margarines are encouraged instead of butter and cream [13]. This clearly has not happened in the long-term care settings. Although the data in both cohorts was gathered prior to the publication of the new recommendation, the general recommendation for diet quality has been published in 2014 [15].

Good fat quality may slow some aspects of age-related decline in health. High SFA intakes elevate liver fat and serum cholesterol, whereas increase of MUFA and PUFA

Table 3 Percentage of nutrition related issues according to SFA quartiles in the long-term care cohort of 2017

SFA Quartiles Nutritional problems	SFA Q ₁ <i>n</i> =119	SFA Q ₂ <i>n</i> =122	SFA Q ₃ <i>n</i> =121	SFA Q ₄ <i>n</i> =119	p-value ¹
Amount of eaten foods, %					<i>p</i> <0.001
very little or little	30	21	17	13	
normal	64	75	75	77	
a lot or very much	6	4	8	11	
Eats snacks, %					<i>p</i> =0.76
yes	81	84	81	84	
Texture of food, %					<i>p</i> =0.96
liquid, puree or soft	36	34	34	36	
normal	65	67	67	64	
Uses oral nutritional supplements, %					<i>p</i> =0.22
yes	22	16	20	14	
Weight monitoring, %					<i>p</i> =0.63
twice to six times a year	20	19	29	21	
>6 times a year	79	81	71	79	
Chewing problems, %					<i>p</i> =0.28
yes	29	30	24	24	
Dry mouth, %					<i>p</i> =0.018
yes	18	17	12	8	
Pain in the mouth, %					<i>p</i> =0.43
yes	10	6	5	8	
Dysphagia, %					<i>p</i> =0.90
yes	19	14	16	18	
Needs help eating, %					<i>p</i> <0.001
yes	83	75	71	63	

SFA Saturated fatty acids, Q Quartiles

¹ Statistical significance for p-value was set to <0.05**Fig. 1** Saturated fatty acid (SFA) intake recommendation, mean SFA intake in 2007 and 2017/18 cohorts showed as percentage of energy (E%). E% = percentage of energy

seems to be beneficial for modulation of liver fat and lipid metabolism [27–30]. Moreover, high MUFA and PUFA diets may improve insulin sensitivity, [31] reduce type 2

diabetes risk [32] and improve cardiovascular outcomes [33]. Healthy brain is also very much dependent on good cardiovascular health [33]. Fatty acids take part in multiple functions in the body and interact with other dietary components as well as microbiome and thus dietary fat composition may be either pro- or anti-inflammatory [33, 34]. Cardiovascular disease, Alzheimer's disease and frailty have all been associated with increased chronic inflammation [35–37]. It has been suggested that n-6:n-3 ratio is also important for cardiovascular health [38], although the optimal n-6:n-3 ratio for human health remains under debate [39]. In our study, the n-6:n-3 ratio was reasonable good, but the problem with fat quality had more to do with low PUFA intakes in general. Thus, good fat quality and sufficient intake of n-3 fatty acids are important also for the oldest-old individuals.

Although the residents of the two cohorts were of similar age and did not differ in MNA score or BMI, the residents in the 2017/18 cohort had worse physical and mental health than the residents in the 2007 cohort. This reflects stricter national guidelines for admission

Table 4 Univariate general linear model of associative factors of saturated fat intake

2017 Cohort	B	95% confidence interval		P-value
		Lower Bound	Upper Bound	
Model 1				
Intercept	16.68	6.64	26.73	0.001
Age	0.26	-0.08	0.14	0.64
Sex (females vs. males)	-4.87	-6.94	-2.81	< 0.001
Sugar intake	0.13	0.11	0.15	< 0.001
Adjusted R ²	0.236			
Model 2				
Intercept	16.86	6.08	27.63	0.002
Sex (females vs. males)	-4.86	-7.11	-2.61	< 0.001
Sugar intake	0.13	0.11	0.16	< 0.001
Need of help with eating (no vs. yes)	-3.17	-5.23	1.11	0.003
Adjusted R ²	0.242			
Model 3				
Intercept	12.35	-0.53	25.23	0.060
Sex (females vs. males)	-4.77	-7.07	-2.47	< 0.001
Sugar intake	0.13	0.10	0.16	< 0.001
Total MNA score	0.20	-0.08	0.48	0.164
Adjusted R ²	0.229			
2007 Cohort				
Model 1	B	95% confidence interval		P-value
		Lower Bound	Upper Bound	
Intercept	33.88	22.45	45.29	< 0.001
Age	-0.15	-0.29	-0.02	0.029
Sex (females vs. males)	-1.69	-4.34	0.97	0.21
Sugar intake	0.063	0.03	0.09	< 0.001
Adjusted R ²	0.052			
Model 2				
Intercept	31.05	19.61	42.49	< 0.001
Age	-0.14	-0.28	-0.09	0.036
Sex (females vs. males)	-1.63	-4.25	0.99	0.22
Sugar intake	0.07	0.04	0.10	< 0.001
Need of help with eating (yes vs. no)	3.01	1.02	5.00	.003
Adjusted R ²	0.073			
Model 3				
Intercept	25.41	11.82	39.00	< 0.001
Age	-0.14	-0.28	-0.01	0.038
Total MNA score	0.31	0.02	0.59	0.035
Sugar intake	0.07	0.03	0.10	< 0.001
Adjusted R ²	0.060			

to long-term care facilities [40]. In Finland due to public policy institutionalized care for older people have been reduced to the minimum and people are expected to live in their homes as long as possible [38]. Thus, only those who have very severe dementia, mobility disability, or other severe health complications due to multiple chronic diseases, are offered a place in a nursing home

or assisted-living facility type of long-term care. This can also be seen in the participants of the 2017/2018 cohort, who had very poor cognition measured with CDR (Table 1) compared to the participants in the 2007 cohort.

However, factors related to nutritional care such as intensive nutritional care, use of oral nutrient

supplements, dysphagia, chewing problems, or other nutrition related issues that might be associated with poor nutrition, were not associated with higher SFA intake in this study. Of specific nutritional issues, only dry mouth was inversely associated with SFA intake. This reflects higher food consumption, as those with dry mouth consumed considerably less energy than those who did not. Similarly, residents eating independently and residents reported consuming larger amounts of foods had higher SFA intake than those needing help with eating or those who only reported eating very little. Thus, all the nutritional care related associations were associated with amount of eaten foods.

It is quite interesting, that despite of official recommendations, the fat quality has worsened. This might be because during the past 10 years, specific education on how to identify and treat malnutrition in long-term care has been provided for nurses working in these facilities. This seems to have led to an increment of SFAs to the served foods in order to avoid weight loss of the residents. As sugar intake was one of the strongest predictors of higher SFA intake, it is likely that sugar is also added to the diet in order to make the offered foods more palatable for the residents at risk of weight loss.

The strengths of our study include its large sample of long-term care residents in both 2007 and 2017/18. To best of our knowledge this is the first study that specifically explores detailed fat composition and quality and how it has changed in recent years in the diets of these people. Trained nurses or nutritionists performed all the measurements in both 2007 and 2017/18 cohorts and all the questionnaires and measurements were validated. Moreover, demographic information, and nutritional supplements were retrieved from medical records, which increase the reliability of our results. However, our study also has many limitations. Food diaries may be subject to error. However, since trained nurses filled in the diaries for the residents, subjective under or over reporting is unlikely. More problematically, when only the 1–2 day food diaries are assessed, they may differ from the person's average food intake over a longer period of time. However, although the individual food intake may vary on a daily basis, our results are relevant at the group level [41]. Moreover, practices that favor using SFA sources such as whole milk products, spreads with high SFA content, butter and cream are not likely to vary considerably from day to day in long-term care facilities. One limitation has to do with the food diary data analyzing tool used in 2007 (Nutrica). Data obtained in the later cohort 2017/18 was more detailed than in 2007, and included also the amounts of trans fatty acids, n-3 and n-6 fatty acids. Therefore, very detailed comparison of fat composition between the two cohorts was not possible. Change

in food diary analyzing tools between the two cohorts should not affect the reliability of the dietary intakes, since both programs are based on the same Finnish food database (Fineli) and are validated tools. The residents in the 2017/18 cohort had poorer health, cognition and mobility, which make them more vulnerable to malnutrition compared to the residents of 2007. However, the residents also had many similarities, and did not differ significantly in respect to nutrition between the cohorts. A further limitation in our study is that the mean time of stay in long-term care is only 2 years; therefore it was not possible to follow the same residents over time. However, although changes in dietary fat intake may have been due to numerous confounding factors e.g. dentition, BMI, health status or other dietary factors, in our study higher SFA intakes were only related to the amount of eaten foods.

The 2017/18 sample was randomly selected within voluntary facilities. The latter sample included six long-term care facilities originally included in the 2007 sample. Although, not all the same facilities were compared, all the long-term care facilities are operated by the city of Helsinki and all facilities should follow the same nutritional guidelines. Moreover, they have same resources, same kind of care and residents are alike spending their last years of their lives there. The participation of the facilities in this study was voluntary and the investigators could not influence the participation. Although fewer facilities participated in the cohort on 2017/2018, in fact higher number of residents took part of the study compared to the 2007 study. The results were obtained from long-term care facilities in the Helsinki metropolitan area with residents mostly of Caucasian origin. The data may thus not be applicable to other ethnic groups. Finally, due to observational nature of our study, no causal relationships can be drawn from these results.

Conclusions

We observed that fat quality has worsened during the last decade among the long-term care residents in spite of the official nutrition recommendations. Nurses and food service personal should be educated about the nutrition recommendations and taught how to increase energy content of foods by using good instead of low quality fats in the diets of long-term residents. This could contribute to better resident health and well-being.

Abbreviations

BMI: Body mass index; CDR: Clinical dementia rating; EFA: Essential fatty acids; FA: Fatty acid; MNA: Mini Nutritional Assessment; MUFA: Monounsaturated fatty acids; PUFA: Polyunsaturated fatty acids; SFA: Saturated fatty acids; Q: Quartile.

Acknowledgements

Open access funded by Helsinki University Library.

Authors' contributions

SKJ conceptualized and designed the article, carried out the statistical analysis and drafted the first version of the manuscript. SKJ and MHS trained the nurses to keep food diaries for the residents. KS, RTN and HMR took part of the measurements and examination of the residents. KHP supervised and obtained funding the study, and all the authors wrote the paper and accepted the final version.

Funding

This work was supported by Päivikki and Sakari Sohlberg foundation (SKJ) and VTR-funding of the Helsinki University Hospital (EVO) (KHP).

Availability of data and materials

The dataset analysed during the current study are not publicly available due to privacy regulations of City of Helsinki. Data inquiries should be addressed to professor Kaisu Pitkälä, email: kaisu.pitkala@helsinki.fi

Declarations

Ethics approval and consent to participate

The study was conducted according to the guidelines of the Declaration of Helsinki, and approved by the Ethics Committee of the Department of Medicine at Helsinki University Hospital and City of Helsinki (HUS/2042/2016). Informed, written consent was obtained from all participants or in cases of moderate to severe dementia (MMSE < 20 points), from their closest proxies such as authorized legal representatives, guardians or caretakers.

Consent for publication

NA.

Competing interests

The authors report no conflict of interest. The sponsors had no role in the design, execution, interpretation, or writing of the study.

Author details

¹Department of General Practice and Primary Health Care, and Helsinki University Central Hospital, Unit of Primary Health Care, University of Helsinki, Tukholmankatu 8, 00014 Helsingin yliopisto Helsinki, Finland. ²Center for Life Course Health Research, University of Oulu, Oulu, Finland. ³City of Helsinki, Department of Social Services and Health Care, Geriatric Clinic, Helsinki Hospital, Helsinki, Finland. ⁴City of Helsinki, Department of Social Services and Health Care, Oral Health Care, Helsinki, Finland.

Received: 30 September 2021 Accepted: 6 April 2022

Published online: 12 April 2022

References

- Borel P, Desmarchelier C. Bioavailability of fat-soluble vitamins and phytochemicals in humans: effects of genetic variation. *Annu Rev Nutr*. 2018;38:69–96. <https://doi.org/10.1146/annurev-nutr-082117-051628>.
- Crawford MA, Bazinet RP, Sinclair AJ. Fat intake and CNS functioning: ageing and disease. *Ann Nutr Metab*. 2009;55:202–28. <https://doi.org/10.1159/000229003>.
- Uauy R. Dietary fat quality for optimal health and well-being: overview of recommendations. *Ann Nutr Metab*. 2009;54:2–7. <https://doi.org/10.1159/000220821>.
- Boscart V, Crutchlow LE, Sheibani Taucar L, et al. Chronic disease management models in nursing homes: a scoping review. *BMJ Open*. 2020;10:e032316. <https://doi.org/10.1136/bmjopen-2019-032316>.
- Suominen M, Muurinen S, Routasalo P, et al. Malnutrition and associated factors among aged residents in all nursing homes in Helsinki. *Eur J Clin Nutr*. 2005;59:578–83. <https://doi.org/10.1038/sj.ejcn.1602111>.
- Soini H, Routasalo P, Lagström H. Characteristics of the Mini-nutritional assessment in elderly home-care patients. *Eur J Clin Nutr*. 2004;58:64–70. <https://doi.org/10.1038/sj.ejcn.1601748>.
- Vikstedt T, Suominen MH, Joki A, Muurinen S, Soini H, Pitkälä KH. Nutritional status, energy, protein, and micronutrient intake of older service house residents. *J Am Med Dir Assoc*. 2011;12:302–7. <https://doi.org/10.1016/j.jamda.2010.12.098>.
- Thiyagalingam S, Kulinski AE, Thorsteinsdottir B, Shindelar KL, Takahashi PY. Dysphagia in older adults. *Mayo Clin Proc*. 2021;96:488–97. <https://doi.org/10.1016/j.mayocp.2020.08.001>.
- Dominguez Castro P, Reynolds C, Bizzaro MG, et al. Characteristics and determinants of high volume dispensing in long-term oral nutritional supplement users in primary care: a secondary analysis. *BJGP Open*. 2021. <https://doi.org/10.3399/BJGPO.2020.0131>.
- Blaum CS, Fries BE, Fiatarone MA. Factors associated with low body mass index and weight loss in nursing home residents. *J Gerontol A Biol Sci Med Sci*. 1995;50:M162–8. <https://doi.org/10.1093/gerona/50a.3.m162>.
- Fried LP, Tangen CM, Walston J, et al. Frailty in older adults: evidence for a phenotype. *J Geront Ser A*. 2001;56:M146–57.
- Odlund Olin A, Armyr I, Soop M, et al. (2003) energy-dense meals improve energy intake in elderly residents in a nursing home. *Clin Nutr*. 2003;22:125–31. <https://doi.org/10.1054/cn.2002.0610>.
- Valtion ravitsemusneuvottelukunta ja Terveyden ja hyvinvoinnin laitos (2020). Vireytä seniorivuosiin – ikääntyneiden ruokasuoitusketut 2020. Terveyden ja hyvinvoinnin laitos. PunaMusta Oy Helsinki. https://www.julkari.fi/bitstream/handle/10024/139415/THL_OHJ_4_2020_Vireytt%C3%A4%20seniorivuosiin_verkko.pdf?sequence=4&isAllowed=y. (in Finnish). Accessed 4 April 2021.
- Krauss RM, Kris-Etherton PM. Public health guidelines should recommend reducing saturated fat. Consumption as much as possible: debate consensus. *Am J Clin Nutr*. 2020;112:25–6. <https://doi.org/10.1093/ajcn/nqaa134>.
- Nordic Nutrition Recommendations. Integrating nutrition and physical activity. Copenhagen: Nordic Council Ministers; 2014. <https://doi.org/10.6027/Nord2014-002>. Accessed 1 April 2021
- AFSSA (Agence Française de Sécurité Sanitaire des Aliments), 2009. Avis de l'Agence française de sécurité sanitaire des aliments sur l'estimation des apports en acides gras trans de la population française. Request 2007-SA-220.
- US Department of Health and Human Services; US Department of Agriculture. 2015–2020 Dietary guidelines for Americans. 8th ed. Washington (DC): US Government Printing Office; 2015. https://health.gov/dietaryguidelines/2015/resources/2015-2020_Dietary_Guidelines.pdf. Accessed 2 April 2021
- Arnett DK, Blumenthal RS, Albert MA, et al. ACC/AHA guideline on the primary prevention of cardiovascular disease: a report of the American College of Cardiology/American Heart Association task force on clinical practice guidelines. *Circulation*. 2019;140:e596–646. <https://doi.org/10.1161/JACC.2019.03.009> Accessed 5 April 2021.
- Jyväkorpi SK, Urtamo A, Strandberg TE. Dietary fat composition and frailty in oldest-old men. *J Am Geriatr Soc*. 2020;68:1346–8. <https://doi.org/10.1111/jgs.16402>.
- Sandoval-Hnsausti H, Pérez-Tasigchana RF, López-García E, et al. Macro-nutrients intake and incident frailty in older adults: a prospective cohort study. *J Gerontol A Biol Sci Med Sci*. 2016;71:1329–34. <https://doi.org/10.1093/gerona/glw033>.
- Hughes CP, Berg L, Danziger WL, Coben LA, Martin RL. A new clinical scale for the staging of dementia. *Br J Psychiatry*. 1982;140:566–72.
- Vellas B, Guigoz Y, Baumgartner M, Garry PJ, Laque S, Albareda JL. Relationships between nutritional markers and the mini-nutritional assessment in 155 older persons. *J Am Geriatr Soc*. 2000;48:1300–9.
- Volkert D, Beck AM, Cederholm T, et al. Management of malnutrition in older patients-current approaches, evidence and open questions. *J Clin Med*. 2019;8:974. <https://doi.org/10.3390/jcm8070974>.
- Jyväkorpi SK, Pitkälä KH, Puranen TM, et al. High intake of nonmilk extrinsic sugars is associated with protein and micronutrient dilution in home-dwelling and institutionalized older people. *J Am Med Dir Assoc*. 2017;18:301–5. <https://doi.org/10.1016/j.jamda.2016.09.023>.
- Rodríguez-Rejón AJ, Ruiz-López MD, Artacho R. Dietary intake and associated factors in long-term care homes in Southeast Spain. *Nutrients*. 2019;11:266. <https://doi.org/10.3390/nu11020266>.
- Viveky N, Billinsky J, Thorpe L, et al. Challenges in planning long-term care menus that meet dietary recommendations. *Can J Diet Pract Res*. 2013;74:84–7. <https://doi.org/10.3148/74.2.2013.84>.

27. Mensink RP. Effects of saturated fatty acids on serum lipids and lipoproteins: a systematic review and regression analysis. Geneva: World Health Organization; 2016.
28. Björmo H, Iggman D, Kullberg J, et al. Effects of n-6 PUFAs compared with SFAs on liver fat, lipoproteins, and inflammation in abdominal obesity: a randomized controlled trial. *Am J Clin Nutr.* 2012;95:1003–12. <https://doi.org/10.3945/ajcn.111.030114>.
29. Rosqvist F, Iggman D, Kullberg J, et al. Overfeeding polyunsaturated and saturated fat causes distinct effects on liver and visceral fat accumulation in humans. *Diabetes.* 2014;63:2356–68. <https://doi.org/10.2337/db13-1622>.
30. Yki-Järvinen H. Nutritional modulation of non-alcoholic fatty liver disease and insulin resistance. *Nutrients.* 2015;7:9127–38. <https://doi.org/10.3390/nu7115454>.
31. Imamura F, Micha R, Wu JH, et al (2016) Effects of saturated fat, polyunsaturated fat, monounsaturated fat, and carbohydrate on glucose-insulin homeostasis: a systematic review and meta-analysis of randomised controlled feeding trials. *Plos Med.* <https://doi.org/10.1371/journal.pmed.1002087>.
32. Salas-Salvadó J, Bulló M, Estruch R, et al. Prevention of diabetes with Mediterranean diets: a subgroup analysis of a randomized trial. *Ann Intern Med.* 2014;160:1–10. <https://doi.org/10.7326/M13-1725>.
33. Guasch-Ferré M, Babio N, Martínez-González MA, et al. Dietary fat intake and risk of cardiovascular disease and all-cause mortality in a population at high risk of cardiovascular disease. *Am J Clin Nutr.* 2015;102:1563–73. <https://doi.org/10.3945/ajcn.115.116046>.
34. Crichton GE, Elias MF, Davey A, Alkerwi A. Cardiovascular health and cognitive function: the Maine-Syracuse longitudinal study. *Plos One.* 2014;9:e89317. <https://doi.org/10.1371/journal.pone.0089317>.
35. Fritsche KL. The science of fatty acids and inflammation. *Adv Nutr.* 2015;6:293S–301S. <https://doi.org/10.3945/an.114.006940>.
36. Soysal P, Stubbs B, Lucato P, et al. Inflammation and frailty in the elderly: a systematic review and meta-analysis. *Ageing Res Rev.* 2016;31:1–8. <https://doi.org/10.1016/j.arr.2016.08.006>.
37. Ferrucci L, Fabbri E. Inflammaging: chronic inflammation in ageing, cardiovascular disease, and frailty. *Nat Rev Cardiol.* 2018;15:505–22. <https://doi.org/10.1038/s41569-018-0064-2>.
38. Wijendran V, Hayes KC. Dietary n-6 and n-3 fatty acid balance and cardiovascular health. *Annu Rev Nutr.* 2004;24:597–615. <https://doi.org/10.1146/annurev.nutr.24.012003.132106> PMID: 15189133.
39. Deckelbaum RJ. N-6 and n-3 fatty acids and atherosclerosis, ratios or amounts? *Atherosclerosis.* *Arterioscler Thromb Vasc Biol.* 2010;30:2325–6. <https://doi.org/10.1161/ATVBAHA.110>.
40. THL. Kotihoito 2020. <https://thl.fi/fi/web/ikaantyminen/muuttuvat-vanhaspalvelut/kotihoito>. Accessed 6 Feb 2021.
41. Basiotis PP, Welsh SO, Cronin FJ, Kelsay JL, Mertz W. Number of days of food intake records required to estimate individual and group nutrient intakes with defined confidence. *J Nutr.* 1987;117:1638–41. <https://doi.org/10.1093/jn/117.9.1638>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

