



<https://helda.helsinki.fi>

Standardised data collection in prehospital critical care : a comparison of medical problem categories and discharge diagnoses

Tommila, Miretta

2022-04-12

Tommila , M , Pappinen , J , Raatiniemi , L , Saviluoto , A , Toivonen , T , Björkman , J & Nurmi , J 2022 , ' Standardised data collection in prehospital critical care : a comparison of medical problem categories and discharge diagnoses ' , Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine , vol. 30 , no. 1 , 26 . <https://doi.org/10.1186/s13049-022-01013-5>

<http://hdl.handle.net/10138/343086>

<https://doi.org/10.1186/s13049-022-01013-5>

cc_by
publishedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

ORIGINAL RESEARCH

Open Access



Standardised data collection in prehospital critical care: a comparison of medical problem categories and discharge diagnoses

Miretta Tommila¹, Jukka Pappinen², Lasse Raatinen^{3,4}, Anssi Saviluoto^{2,5}, Tuukka Toivonen², Johannes Björkman^{2,6} and Jouni Nurmi^{2,6*}

Abstract

Background: Prehospital medical problem reporting is essential in the management of helicopter emergency medical services (HEMS) operations. The consensus-based template for reporting and documenting in physician-staffed prehospital services exists and the classification of medical problems presented in the template is widely used in research and quality improvement. However, validation of the reported prehospital medical problem is lacking. This study aimed to describe the in-hospital diagnoses, patient characteristics and medical interventions in different categories of medical problems.

Methods: This retrospective, observational registry study examined the 10 most common in-hospital International Statistical Classification of Disease (ICD-10) diagnoses within different prehospital medical problem categories, defined by the HEMS physician/paramedic immediately after the mission was completed. Data were gathered from a national HEMS quality registry and a national hospital discharge registry. Patient characteristics and medical interventions related to different medical problem categories are also described.

Results: A total of 33,844 patients were included in the analyses. All the medical problem categories included a broad spectrum of ICD-10 diagnoses (the number of diagnosis classes per medical problem category ranged from 73 to 403). The most frequent diagnoses were mainly consistent with the reported medical problems. Overlapping of ICD-10 diagnoses was mostly seen in two medical problem categories: stroke and acute neurology excluding stroke. Additionally, typical patient characteristics and disturbances in vital signs were related to adequate medical problem categories.

Conclusions: Medical problems reported by HEMS personnel have adequate correspondence to hospital discharge diagnoses. However, the classification of cerebrovascular accidents remains challenging.

Keywords: Air ambulances, Emergency medical services, Critical care, Medical problem reporting, Prehospital, Documenting

Introduction

Documenting and reporting are important parts of prehospital care. Conscientious documenting is a legal necessity as a part of a patient's treatment, but it is also an important way to acquire data for clinical quality improvement and benchmarking and for research purposes. Certainly, medical problem categorisation is a

*Correspondence: jouni.nurmi@hus.fi

⁶ Emergency Medicine and Services, Helsinki University Hospital and University of Helsinki, FinnHEMS 10, Vesikuja 9, 01530 Vantaa, Finland
Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

prerequisite in the management of helicopter emergency medical services (HEMS) operations. Designed to standardise prehospital documenting and reporting policies, a consensus-based template for physician-staffed prehospital services was first published in 2011 [1]. The feasibility of this template has been demonstrated, and it has been recently updated [2, 3].

Despite the existing international template, there are still variations in the ways patients' medical problems are reported in different HEMS systems. In Nordic HEMS, the reporting rests on the principles of the international template, although each system has its own modifications [4–6]. In comparison to other European services, there is even more divergence in reporting of medical problems [7, 8]. The categories in the consensus template are widely used in research reports, for example in studies regarding airway management [9] and physiological disturbances [10] in prehospital critical care. However, the differences in reporting make it challenging to compare these services. Moreover, the lack of data on exact medical conditions in each category makes it challenging to generalise the findings.

Currently, validation of the reported prehospital medical problem is lacking. The parity between medical problem categorisation and in-hospital diagnoses is unknown. Moreover, it is not known if the categorisations made by providers in the field are accurate. In the current study, we aimed to describe the in-hospital diagnosis classes and patient characteristics of the patients classified by HEMS personnel into different medical problem categories, according to the consensus template of reporting physician-provided prehospital care [1]. Furthermore, we introduce patient characteristics, medical interventions and 30-day mortality associated with each medical problem category in the Finnish HEMS.

Methods

Study design

We performed a retrospective, observational registry study describing the 10 most common International Statistical Classification of Diseases (ICD-10) diagnoses within medical problem categories as defined in the consensus-based template for physician-staffed prehospital services [1]. Additionally, we examined typical patient characteristics and medical interventions related to different medical problem categories. Ethical permission was granted by the Ethical Board of the University of Helsinki (HUS/3115/2019 §194). Study permission was granted by all the participating hospital districts. The reporting of the study adheres to Strengthening the Reporting of Observational studies in Epidemiology (STROBE) guidelines [11].

Setting

The Finnish HEMS system consists of five physician-staffed units and one paramedic-staffed unit. The paramedic-staffed unit serves the vast and sparsely populated Lapland. The units are primarily dispatched to emergencies by the Emergency Response Centres, with slight regional differences in the dispatch criteria. The dispatch criteria, mission profile and the national HEMS have recently been described elsewhere [6].

Participants

All patients encountered by HEMS units between 1 January 2012 and 31 December 2018 were included in the study. The timeframe depicts the foundation of the database and the latest practical point. Due to the study's retrospective design, no power calculation was performed and all available data were included.

Data sources

FinnHEMS database (FHDB)

The FHDB is a national HEMS quality registry containing details on all HEMS missions in Finland. After each mission, the HEMS crew enters the mission details into the database. There have been no changes related to reporting of medical problem categorisation or other investigated parameters during the study period. FHDB and data collection have recently been described in detail in previous papers [6, 12].

The national hospital discharge register (HILMO)

The HILMO database is maintained by the Finnish Institute for Health and Welfare. All organisations providing inpatient care are required by law to enter data on all hospital admissions and discharge. Upon discharge, up to five main diagnoses—following the ICD-10 nomenclature—are then entered into the HILMO register. The entered data also include additional information, such as provided treatment (Nordic Classification of Surgical Procedures [NCSP]) and length-of-stay. The completeness and accuracy of the HILMO register have been evaluated to be appropriate, and the validity of the register has been stated to be suitable for health services research [13]. For our study, all the main diagnoses from consecutive hospital admissions between 1 January 2012 and 31 December 2018 were acquired from the HILMO database.

Finnish digital and population data service agency

Information on living status or date of death was acquired for all the encountered patients from the

Finnish Digital and Population Data Service Agency until the end of October 2019.

Variables

In the FHDB, each patient is assigned one of 10 possible mission medical problems, according to the HEMS provider's assessment on the true medical condition after the mission is completed. The medical problem categories, as defined in the consensus-based template for physician-staffed prehospital services, are: cardiac arrest, trauma, breathing difficulties, chest pain, stroke, acute neurology excluding stroke, psychiatry including intoxication, obstetrics and childbirth, infection, and other [1].

Outcome was defined as the main diagnosis classes at hospital discharge, extracted from the HILMO register. Since the discharge diagnoses might not be arranged in order of importance, all the main diagnoses were included. The diagnoses are reported as the ICD-10 main categories, i.e. without numbers after the decimal separator (for example, I21 corresponding to acute myocardial infarction, including subheadings, e.g. I21.1 for acute transmural myocardial infarction of anterior wall).

In addition to the hospital discharge diagnoses, we gathered typical patient characteristics categorised by each medical problem as follows: sex, age, first vital parameter values recorded by HEMS personnel (oxygen saturation, heart rate, systolic blood pressure, and


Glasgow Coma Score [GCS]). We calculated the proportion of hypoxic and hypotensive patients per every medical problem category. The definition of hypoxia was an oxygen saturation below 90%; hypotension was defined as systolic blood pressure under 90 mmHg [14]. We also determined the proportion of patients treated with advanced airway management techniques used or with vasoactive medication given within each category. Finally, we assessed 30-day mortality in different medical problem categories.

Statistical methods

Quantitative variables are presented as means with standard deviation (\pm SD) or median with the interquartile range (25th–75th percentile), depending on the distribution. Categorical variables are presented as n (%). Missing data were not included in the analysis; however, the percentages of missing data are reported as n (%). We used IBM SPSS Statistics 25 (IBM Corporation, Armonk, NY, USA) for all statistical analyses.

Results

A total of 33,844 patients were recorded in the FHDB as encountered by HEMS personnel within the study timeframe. All these patients were included in the study (Fig. 1). Due to the possibility of several ICD-10 diagnoses for each individual patient, the total number of



ICD-10 diagnoses was 36,483. The number of diagnoses under each medical problem category ranged from 73 to 403, and the proportion of the diagnoses not covered by the 10 most frequent diagnoses varied 16–67% between the categories.

The 10 most frequent diagnoses under each specific medical problem category are described in Table 1. There was a vast heterogeneity of ICD-10 diagnoses in every medical problem category, but the diagnoses were mainly consistent with the reported medical problem. The greatest variation was noticed under the medical problem category of stroke, in which that category was mixed with the acute neurology excluding stroke category. Among the more typical diagnoses in the medical problem category of cardiac arrest, non-traumatic subarachnoid haemorrhage was ranked fifth. Furthermore, intracranial injury represented 25% of the ICD-10 diagnoses categorised under trauma.

The characteristics of the patients in different categories are presented in Table 2. The patients in the trauma, psychiatric including intoxication, and obstetrics and childbirth categories were, on average, younger than the patients in the other medical problem categories. The lowest values of GCS were found in the cardiac arrest, stroke, acute neurology excluding stroke and psychiatric including intoxication categories.

The prevalence of hypoxia and hypotension as well as interventions performed in different patient categories are presented in Table 3. Hypoxia was most common in the breathing difficulties and infection categories. Hypotension was most frequently noticed in the infection category and the other diagnoses category.

Vasoactive medication was most often used for the patients in the cardiac arrest, breathing difficulties, chest pain, stroke, and infection categories. Advanced airway was most common in the categories which also had the lowest GCS values. Significant variation in 30-day mortality was observed between the groups.

Patients' medical problem and whether vasoactive medication was given were available for all patients, whereas sex and age was missing for 383 (1.1%) and 28 (0.01%) patients respectively. Out of the vital signs at the time of encounter systolic blood pressure could not be measured for 885 (2.6%) and was missing or not measured for 8492 (25%) patients. Heart rate could not be measured for 537 (1.6%) patients and was missing for 7711 (23%) patients, oxygen saturation could not be measured for 5184 (15%) and was missing for 4236 (13%) patients. GCS was missing for 2076 (6.1%) patients. Whether an advanced airway was used was missing for 160 (0.5%), 30-day mortality was unavailable for 2079 (6.1%) and no discharge diagnoses were available for 7630 (23%) patients.

Discussion

The main finding of this study is that the medical problems reported by HEMS personnel were comparable to the hospital discharge diagnoses, excluding the medical problem reported as stroke. Furthermore, we observed noticeable heterogeneity and a broad spectrum of the diagnoses in the patients treated by HEMS personnel.

To the best of our knowledge, this is the first study that has investigated the hospital diagnoses of the patients treated by HEMS personnel and classified according to the consensus-based reporting template. The strength of the study is its use of large, nationwide registries as a data source. A significant number of HEMS missions was included in the analysis, and the overall mission reporting quality has been reported to be at an advisable level [6, 12]. The data collection in the HEMS database is mainly based on the international guidelines [1], although with slight modifications [6].

However, our study has several limitations. As a registry study, the major limitation is its retrospective nature. A comparison of the databases is somewhat flawed: data entered into HILMO have shown excellent validity for single disease groups but lacking in, for example, polytrauma patients [15]. Moreover, according to our hypothesis, the discharge diagnosis is assumed to be correct; however, that is not always the case and not all patients receive a definitive diagnosis. Furthermore, the accuracy of the prehospital classification was not evaluated in an objective and systematic way as there is no golden standard for prehospital category for each discharge diagnosis. Thus, the conclusion of the study is subjective and needs to be interpreted according to the results obtained. The inexact nature of medical science and inter-operator differences in clinical judgement must be considered and accepted when interpreting the results of medical studies.

In our study, many of the patients diagnosed with stroke had a discharge diagnosis more suitable for the medical problem of acute neurology excluding stroke. In fact, the definition of a stroke also broadly includes intracerebral haemorrhage and subarachnoid haemorrhage in addition to an ischemic cerebral event [16]. Our finding reflects the difficulty of diagnosing different intracranial disorders and neurological emergencies solely based on clinical symptoms [17, 18]. Moreover, patients presenting classical stroke symptoms without gross impairment of consciousness are mainly treated by paramedics without HEMS assistance [19]. Hence, the stroke situations that have demanded HEMS involvement have probably been more complex.

The medical problem reported as trauma revealed that the proportion of traumatic brain injuries (TBI) is remarkably high: approximately one in four trauma

Table 1 Ten most frequent ICD-10 top level diagnoses categorized by medical problem. Presented as patient category (number of separate top-level diagnoses) or n (%)

| Diagnoses | N | % | Diagnoses | N | % |
|---|----------|----------|--|----------|----------|
| Cardiac arrest (238 diagnosis blocks in total) | | | Trauma (368) | | |
| Cardiac arrest | 1904 | 44.1 | Intracranial injury | 2455 | 24.6 |
| Myocardial infarction | 583 | 13.5 | Fracture of rib(s), sternum and thoracic spine | 655 | 6.6 |
| III-defined and unknown cause of mortality | 324 | 7.5 | Fracture of skull and facial bones | 480 | 4.8 |
| Other cardiac arrhythmias | 172 | 4.0 | Fracture of lumbar spine and pelvis | 461 | 4.6 |
| Nontraumatic subarachnoid hemorrhage | 98 | 2.3 | Injury of other and unspecified intrathoracic organs | 380 | 3.8 |
| Chronic ischemic heart disease | 95 | 2.2 | Open wound of head | 375 | 3.8 |
| Other disorders of brain | 53 | 1.2 | Fracture of lower leg, including ankle | 284 | 2.8 |
| Pulmonary embolism | 46 | 1.1 | Fracture of femur | 270 | 2.7 |
| Heart failure | 39 | 0.9 | Fracture of neck | 247 | 2.5 |
| Pneumonia, unspecified organism | 37 | 0.9 | Injury of intra-abdominal organs | 224 | 2.2 |
| Other diagnoses | 963 | 22.3 | Other diagnoses | 4144 | 41.5 |
| Breathing difficulties (273) | | | Chest pain (123) | | |
| Heart failure | 403 | 14.5 | Myocardial infarction | 749 | 42.2 |
| Pneumonia, organism unspecified | 347 | 12.5 | Pain in throat and chest | 243 | 13.7 |
| Other chronic obstructive pulmonary disease | 212 | 7.6 | Angina pectoris | 173 | 9.7 |
| Abnormalities of breathing | 156 | 5.6 | Chronic ischemic heart disease | 93 | 5.2 |
| Acute myocardial infarction | 133 | 4.8 | Atrial fibrillation and flutter | 75 | 4.2 |
| Other respiratory disorders | 92 | 3.3 | Paroxysmal tachycardia | 46 | 2.6 |
| Bacterial pneumonia, not elsewhere classified | 75 | 2.7 | Heart failure | 40 | 2.3 |
| Adverse effects, not elsewhere classified | 65 | 2.3 | Syncope and collapse | 23 | 1.3 |
| Asthma | 56 | 2.0 | Dissection of aorta | 20 | 1.1 |
| Foreign body in respiratory tract | 54 | 1.9 | Abdominal and pelvic pain | 18 | 1.0 |
| Other diagnoses | 1188 | 42.7 | Other diagnoses | 295 | 16.6 |
| Stroke (191) | | | Other neurological (332) | | |
| Cerebral infarction | 724 | 22.0 | Convulsions, not elsewhere classified | 994 | 17.9 |
| Nontraumatic intracerebral hemorrhage | 538 | 16.3 | Epilepsy and recurrent seizures | 994 | 17.9 |
| Nontraumatic subarachnoid hemorrhage | 492 | 14.9 | Status epilepticus | 363 | 6.5 |
| Epilepsy and recurrent seizures | 153 | 4.6 | Nontraumatic intracerebral hemorrhage | 289 | 5.2 |
| Intracranial injury | 143 | 4.3 | Mental and behavioural disorders due to use of alcohol | 245 | 4.4 |
| Convulsions, not elsewhere classified | 142 | 4.3 | Personality disorder | 237 | 4.3 |
| Transient cerebral ischemic attacks and related syndromes | 109 | 3.3 | Syncope and collapse | 210 | 3.8 |
| Somnolence, stupor and coma | 77 | 2.3 | Cerebral infarction | 204 | 3.7 |
| Syncope and collapse | 71 | 2.2 | Somnolence, stupor and coma | 181 | 3.3 |
| Status epilepticus | 53 | 1.6 | Intracranial injury | 175 | 3.2 |
| Other diagnoses | 796 | 31.8 | Other diagnoses | 1663 | 29.9 |
| Psychiatric including intoxication (220) | | | Gynecologic or obstetric (73) | | |
| Poisoning or adverse effect of medicament | 1348 | 37.8 | Full-term uncomplicated delivery | 368 | 47.4 |
| Mental and behav. disorders due to alcohol | 556 | 15.6 | Maternal postpartum care and examination | 46 | 5.9 |
| Depression | 137 | 3.8 | Supervision of normal pregnancy | 45 | 5.8 |
| Poisoning due to narcotics or hallucinogens | 122 | 3.4 | False labor | 43 | 5.5 |
| Toxic effect of alcohol | 99 | 2.8 | Single delivery by caesarean section | 34 | 4.4 |
| Somnolence, stupor and coma | 81 | 2.3 | Abdominal and pelvic pain | 22 | 2.8 |
| Recurrent major depression | 53 | 1.5 | Obstetrical hemorrhage | 17 | 2.2 |
| Opioid related disorders | 49 | 1.4 | Single delivery by forceps and vacuum extractor | 14 | 1.8 |
| Other psychoactive substance related disorders | 49 | 1.4 | Placenta previa | 14 | 1.8 |
| Convulsions, not elsewhere classified | 46 | 1.3 | Premature rupture of membranes | 13 | 1.7 |
| Other diagnoses | 1030 | 28.9 | Other diagnoses | 160 | 20.6 |

Table 1 (continued)

| Diagnoses | N | % | Diagnoses | N | % |
|---|----------|----------|--|----------|----------|
| Infection (146) | | | Oher (403) | | |
| Other sepsis | 84 | 14.6 | Syncope and collapse | 205 | 5.3 |
| Pneumonia, unspecified organism | 71 | 12.3 | Paroxysmal tachycardia | 162 | 4.2 |
| Convulsions, not elsewhere classified | 50 | 8.7 | Dissection of aorta | 131 | 3.4 |
| Bacterial infection of unspecified site | 35 | 6.1 | Hypothermia | 127 | 3.3 |
| Bacterial pneumonia, not elsewhere classified | 15 | 2.6 | Other diseases of digestive system | 124 | 3.2 |
| Infectious gastroenteritis and colitis, unspecified | 14 | 2.4 | Poisoning or adverse effect of medicament | 122 | 3.2 |
| Erysipelas | 11 | 1.9 | Mental and behav. disorders due to alcohol | 118 | 3.1 |
| Streptococcal sepsis | 10 | 1.7 | Adverse effects, not elsewhere classified | 115 | 3.0 |
| Acute tubulo-interstitial nephritis | 9 | 1.6 | Atrial fibrillation and flutter | 92 | 2.4 |
| Abdominal and pelvic pain | 9 | 1.6 | Abdominal and pelvic pain | 90 | 2.3 |
| Other diagnoses | 268 | 46.5 | Other diagnoses | 2577 | 66.7 |

Table 2 Patient characteristics categorized by medical problem. Presented as N, proportion (%) or median (25th–75th percentile)

| Medical problem | N | Sex; male (%) | Age (years) | Oxygen saturation (%) | Heart rate (1/min) | Systolic blood pressure (mmHg) | GCS |
|--------------------------------------|----------|----------------------|--------------------|------------------------------|---------------------------|---------------------------------------|----------------------|
| Cardiac arrest | 6900 | 71.8 | 68 | 58–78 | * | * | 3 3–3 |
| Trauma | 8897 | 71.8 | 40 | 23–59 | 98 95–99 | 90 80–105 | 130 115–150 15 12–15 |
| Breathing difficulties | 2077 | 53.9 | 69 | 52–80 | 90 80–96 | 108 90–126 | 140 118–169 15 13–15 |
| Chest pain | 1380 | 68.0 | 68 | 60–78 | 97 95–98 | 80 65–100 | 140 112–162 15 15–15 |
| Stroke | 2295 | 50.5 | 72 | 61–80 | 97 94–98 | 83 70–100 | 157 133–185 8 4–13 |
| Acute neurology (excluding stroke) | 4366 | 57.3 | 58 | 32–72 | 97 94–98 | 98 80–119 | 134 113–158 9 5–14 |
| Psychiatric (including intoxication) | 3318 | 55.1 | 37 | 25–50 | 97 94–99 | 90 75–105 | 120 105–137 8 3–13 |
| Gynecologic or obstetric | 689 | 1.9 | 30 | 26–35 | 98 97–99 | 90 80–100 | 125 115–139 15 15–15 |
| Infection | 422 | 56.6 | 60 | 19–75 | 95 90–98 | 110 90–130 | 110 90–130 14 9–15 |
| Other | 3500 | 63.0 | 62 | 42–76 | 97 94–99 | 90 70–112 | 120 96–143 15 15–15 |

All values are first values recorded by HEMS

GCS: Glasgow coma score

HEMS: Helicopter emergency medical services

*Initial values are not reported as marked proportion of the patients are encountered with cardiac arrest and unmeasurable vital signs

Table 3 Performed medical interventions, 30-day mortality and proportions of patients suffering from hypoxia or hypotension when encountered by HEMS. Reported as percentage (%)

| Medical problem | Hypoxic (%) | Advanced airway (%) | Hypotensive (%) | Vasoactive medication (%) | 30-day mortality (%) |
|--------------------------------------|--------------------|----------------------------|------------------------|----------------------------------|-----------------------------|
| Cardiac arrest | * | 49 | * | 59 | 79 |
| Trauma | 6 | 17 | 6 | 11 | 11 |
| Breathing difficulties | 47 | 11 | 7 | 23 | 18 |
| Chest pain | 8 | 3 | 12 | 34 | 8 |
| Stroke | 9 | 38 | 3 | 24 | 38 |
| Acute neurology (excluding stroke) | 11 | 25 | 6 | 14 | 13 |
| Psychiatric (including intoxication) | 11 | 25 | 11 | 12 | 2 |
| Gynecologic or obstetric | 0.8 | 1 | 3 | 0.4 | 0.2 |
| Infection | 20 | 13 | 24 | 26 | 18 |
| Other | 9 | 11 | 19 | 18 | 14 |

*Initial values cannot be measured

patients treated by HEMS personnel suffered from a TBI. The high proportion can be explained by the fact that these patients are included in the dispatch criteria for HEMS and they have been shown to benefit from prehospital critical care [20–22].

As expected, there were no dramatic findings related to typical patient characteristics in the different medical problem categories. Sudden illness or trauma can affect people of various ages, but patients suffering trauma or psychiatric or obstetric problems seem to be younger than patients with acute cardiovascular or cerebrovascular events. Disturbances in certain vital organ functions were associated with the adequate medical problem category; for example, hypoxia was most common during respiratory distress and infections, hypotension was most common during infections and in the miscellaneous category (containing for example severe arrhythmias) and declined consciousness was common during cardiac arrest, cerebrovascular events and intoxication. Evidently, advanced airway management was most frequently associated with medical problems causing declined consciousness, but vasoactive medication was used more often not only in categories linked to hypotension, but also in patients with chest pain, stroke and breathing difficulties. However, in this study only initial values of the vital parameters are investigated, so it seems likely that patients who got vasoactive medication also presented more or less unstable hemodynamics at some stage of prehospital treatment.

The highest 30-day mortality was seen most often among cardiac arrest patients, as expected. Patients categorised under stroke had the second highest fatality. This might be explained by the fact that HEMS mainly involves in the most demanding stroke cases. Indeed, increased mortality also among patients suffering respiratory distress or infections probably reflects the severity of the typical diagnoses classified under these categories, such as serious cardiovascular events, or critical infections like sepsis or pneumonia. Furthermore, fairly notable mortality, 14%, was found in category “other”, and nearly one fifth of these patients needed also vasoactive medication. Based on these findings, this quite large group of patients seems to be severely ill, and it would be important to be able to classify these patients more thoroughly.

In addition to that, the substantial number of diagnosis groups in each patient category raises the question as to the extent to which this crude classification is appropriate in the prehospital critical care research. Adjusting outcome analyses based on this classification may not be sufficient as the groups are highly heterogeneous in terms of pathophysiology. An attempt to address the effectiveness of HEMS and the prehospital critical care provided

by the physicians has been made by conventional clinical studies, including cohort and case-control designs [23]. However, the intervention (prehospital critical care provided by the HEMS physician) includes many different procedures, logistic approaches, tactics, complicated clinical pathways and human factors; thus, the intervention is complex instead of simple (pharmaceutical). This complexity is increased by the significant heterogeneity of the patients being treated, as demonstrated by the current study. Thus, it is unlike to adequately summarise the evidence of the effectiveness, in general, of using conventional clinical study designs and review methods. Instead, the potential role of HEMS in treating different patient groups could be better understood using the application theories and frameworks of complex health care interventions [24].

Despite the lack of totally uniform reporting policies, our results are likely to be generalisable to other similar HEMS organisations, but for comparison to different services, for instances ambulances, more research is needed. In the future, a uniform, standardised reporting system would alleviate the comparison between different HEMS systems, both from a clinical/operational viewpoint and a scientific viewpoint.

Based on our study, it is especially important to improve the accuracy in the medical problem category of stroke. In the recently updated documenting and reporting template, this area is now divided into smaller units: stroke, acute neurology excluding stroke, reduced level of consciousness and poisoning/intoxication. This division might improve the accuracy of the classification of unconscious patients, although prehospital diagnostics discerning between stroke and other neurological emergencies is still challenging. Finally, in addition to our findings about the medical problems categories, the significant prevalence of subarachnoid haemorrhage among cardiac arrest patients must be kept in mind, and this diagnosis should be examined more closely.

Conclusion

The medical problems categorised by HEMS personnel can be seen as valid. The greatest challenges remain in the classification of stroke and other neurological emergencies. Patients treated by HEMS personnel are characterised by significant heterogeneity and the crude classification based on the reporting template should be used cautiously in scientific research.

Abbreviations

ALS: Advanced life support; FHDB: FinnHEMS database; GCS: Glasgow Coma Scale; HEMS: Helicopter emergency medical services; HILO: National hospital discharge register; ICD-10: International statistical classification of diseases; NCSP: Nordic classification of surgical procedures; OHCA: Out-of-hospital

cardiac arrest; SD: Standard deviation; STROBE: Strengthening the reporting of observational studies in epidemiology; TBI: Traumatic brain injury.

Acknowledgements

Not applicable.

Author contributions

All the authors contributed to the conception and design of the study. JP, AS and JN analysed the data. MT, TT and AS drafted the manuscript. All the authors contributed to the data interpretation and critical revision of the manuscript. All the authors have read and approved the final manuscript.

Funding

This work was supported by FinnHEMS Research and Development Unit and Helsinki University Hospital (state funding, VTR TYH2019243).

Availability of data and materials

The dataset used during the current study is available from the corresponding author upon reasonable request.

Declarations

Ethics approval and consent to participate

We requested and were granted permission to conduct this study from each participating hospital district (Oulu University Hospital 200/2019 2.7.2019, Helsinki University Hospital HUS/280/2019 9.7.2019, Turku University Hospital J30/19 4.8.2019, Hospital District of Lapland 32/2019 22.8.2019, Kuopio University Hospital RPL 102/2019 22.8.2019, Tampere University Hospital RTL-R19580 2.9.2019). Ethical permission, although not necessary for observational studies, was granted by the Ethical Board of the University of Helsinki (HUS/3115/2019 §194 27.11.2019). The Finnish Institute for Health and Welfare granted permission to the HILMO data (Dnro THL/2231/5.05.00/2019 21.2.2020). The study was purely observational and did not affect patient treatment in any way. Thus, patient consent was not required according to Finnish legislation, nor was it sought.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Perioperative Services, Intensive Care Medicine and Pain Management, Turku University Hospital and University of Turku, Turku, Finland.
²FinnHEMS Ltd, HEMS Operations, Vantaa, Finland.
³Centre for Prehospital Emergency Care, Oulu, Finland.
⁴Research Group of Surgery, Anaesthesiology and Intensive Care, Division of Anesthesiology Oulu University Hospital, Medical Research Centre, University of Oulu, Oulu, Finland.
⁵University of Eastern Finland, Kuopio, Finland.
⁶Emergency Medicine and Services, Helsinki University Hospital and University of Helsinki, FinnHEMS 10, Vesikuja 9, 01530 Vantaa, Finland.

Received: 13 December 2021 Accepted: 31 March 2022

Published online: 12 April 2022

References

- Krüger AJ, Lockey D, Kurola J, Bartolomeo SD, Castrén M, Mikkelsen S, et al. A consensus-based template for documenting and reporting in physician-staffed pre-hospital services. *Scand J Trauma Resusc Emerg Med*. 2011;19:71–71.
- Tønsager K, Krüger AJ, Ringdal KG, Rehn M, Hossfeld B, Breitenmoser I, et al. Template for documenting and reporting data in physician-staffed pre-hospital services: a consensus-based update. *Scand J Trauma Resusc Emerg Med*. 2020;28:25.
- Lossius HM, Krüger AJ, Ringdal KG, Sollid SJ, Lockey DJ. Developing templates for uniform data documentation and reporting in critical care using a modified nominal group technique. *Scand J Trauma Resusc Emerg Med*. 2013;21:80–80.
- Kornhall D, Näslund R, Klingberg C, Schiborr R, Gellerfors M. The mission characteristics of a newly implemented rural helicopter emergency medical service. *BMC Emerg Med*. 2018;18:28.
- Alstrup K, Petersen JAK, Barfod C, Knudsen L, Rognås L, Møller TP. The Danish helicopter emergency medical service database: high quality data with great potential. *Scand J Trauma Resusc Emerg Med*. 2019;27:38.
- Saviluoto A, Björkman J, Olkinuora A, Virkkunen I, Kirves H, Setälä P, et al. The first seven years of nationally organized helicopter emergency medical services in Finland—the data from quality registry. *Scand J Trauma Resusc Emerg Med*. 2020;28:46.
- Koivulahti O, Tommila M, Häavisto E. The accuracy of preliminary diagnoses made by paramedics—a cross-sectional comparative study. *Scand J Trauma Resusc Emerg Med*. 2020;28:70.
- Rzońca P, Gałązkowski R, Wójcik-Fatla A, Panasiuk L, Gotlib J. Missions of the helicopter emergency medical service in rural and urban areas in Poland—a comparative retrospective analysis. *Ann Agric Environ Med*. 2019;26:355–60.
- Gellerfors M, Fevang E, Bäckman A, Krüger A, Mikkelsen S, Nurmi J, et al. Pre-hospital advanced airway management by anaesthetist and nurse anaesthetist critical care teams: a prospective observational study of 2028 pre-hospital tracheal intubations. *Brit J Anaesth*. 2018;120:1103–9.
- Björkman J, Raatinen I, Setälä P, Nurmi J. Shock index as a predictor for short term mortality in helicopter emergency medical services: a registry study. *Acta Anaesth Scand*. 2020;65:816–23.
- von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandebroucke JP, et al. The strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *Int J Surg*. 2014;12:1495–9.
- Heino A, Iirola T, Raatinen I, Nurmi J, Olkinuora A, Laukanen-Nevala P, et al. The reliability and accuracy of operational system data in a nationwide helicopter emergency medical services mission database. *BMC Emerg Med*. 2019;19:53.
- Sund R. Quality of the finnish hospital discharge register: a systematic review. *Scand J Public Health*. 2012;40:505–15.
- Badjatia N, Carney N, Crocco TJ, Fallat ME, Hennes HMA, Jagoda AS, et al. Guidelines for prehospital management of traumatic brain injury 2nd edition. *Prehosp Emerg Care*. 2009;12:1–52.
- Heinänen M, Brinck T, Handolin L, Mattila VM, Söderlund T. Accuracy and coverage of diagnosis and procedural coding of severely injured patients in the finnish hospital discharge register: comparison to patient files and the helsinki trauma registry. *Scand J Surg*. 2016;106:269–77.
- Sacco RL, Kasner SE, Broderick JP, Caplan LR, Connors JJB, Culebras A, et al. An updated definition of stroke for the 21st century: a statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke J Cereb Circ*. 2013;44:2064–89.
- ter Avest E, Taylor S, Wilson M, Lyon RL. Prehospital clinical signs are a poor predictor of raised intracranial pressure following traumatic brain injury. *Emerg Med J*. 2021;38:21–26.
- Lutz M, Möckel M, Lindner T, Ploner CJ, Braun M, Schmidt WU. The accuracy of initial diagnoses in coma: an observational study in 835 patients with non-traumatic disorder of consciousness. *Scand J Trauma Resusc Emerg Med*. 2021;29:15.
- Saviluoto A, Harve-Rytälä H, Lääperi M, Kirves H, Jäntti H, Nurmi J. A potential method of identifying stroke and other intracranial lesions in a prehospital setting. *Scand J Trauma Resusc Emerg Med*. 2020;28:39.
- Pakkanen T, Nurmi J, Huhtala H, Silfvast T. Prehospital on-scene anaesthetist treating severe traumatic brain injury patients is associated with lower mortality and better neurological outcome. *Scand J Trauma Resusc Emerg Med*. 2019;27:9.
- Andruszkow H, Schweigkofler U, Lefering R, Frey M, Horst K, Pfeifer R, et al. Impact of helicopter emergency medical service in traumatized patients: Which patient benefits most? *PLOS ONE*. 2016;11:e0146897.
- Taylor C, Jan S, Curtis K, Tzannes A, Li Q, Palmer C, et al. The cost-effectiveness of physician staffed Helicopter Emergency Medical Service (HEMS) transport to a major trauma centre in NSW, Australia. *Inj*. 2012;43:1843–9.

23. Risgaard B, Draegert C, Baekgaard JS, Steinmetz J, Rasmussen LS. Impact of physician-staffed helicopters on pre-hospital patient outcomes: a systematic review. *Acta Anaesth Scand.* 2020;64:691–704.
24. Kelly MP, Noyes J, Kane RL, Chang C, Uhl S, Robinson KA, et al. AHRQ series on complex intervention systematic reviews—paper 2: defining complexity, formulating scope, and questions. *J Clin Epidemiol.* 2017;90:11–8.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

