

<https://helda.helsinki.fi>

---

## Environmental triggers of Parkinson's disease-Implications of the Braak and dual-hit hypotheses

Chen, Honglei

2022-02

---

Chen , H , Wang , K , Scheperjans , F & Killinger , B 2022 , ' Environmental triggers of Parkinson's disease-Implications of the Braak and dual-hit hypotheses ' , *Neurobiology of Disease* , vol. 163 , 105601 . <https://doi.org/10.1016/j.nbd.2021.105601>

---

<http://hdl.handle.net/10138/343243>

<https://doi.org/10.1016/j.nbd.2021.105601>

---

cc\_by\_nc\_nd

publishedVersion


---

*Downloaded from Helda, University of Helsinki institutional repository.*

*This is an electronic reprint of the original article.*

*This reprint may differ from the original in pagination and typographic detail.*

*Please cite the original version.*



# Environmental triggers of Parkinson's disease – Implications of the Braak and dual-hit hypotheses

Honglei Chen<sup>a,\*</sup>, Keran Wang<sup>a</sup>, Filip Scheperjans<sup>b</sup>, Bryan Killinger<sup>c</sup>

<sup>a</sup> Epidemiology and Biostatistics, College of Human Medicine, Michigan State University, East Lansing, MI 48824, USA

<sup>b</sup> Department of Neurology, Helsinki University Hospital, and Clinicum, University of Helsinki, Haartmaninkatu 4, 00290 Helsinki, Finland

<sup>c</sup> Graduate College, Rush University Medical Center, Chicago, IL 60612, USA

## ARTICLE INFO

### Keywords:

Parkinson's disease  
Environmental risk factors  
Olfactory impairment  
Microbiota  
Gut-brain axis

## ABSTRACT

Idiopathic Parkinson's disease (PD) may take decades to develop, during which many risk or protective factors may come into play to initiate the pathogenesis or modify its progression to clinical PD. The lack of understanding of this prodromal phase of PD and the factors involved has been a major hurdle in the study of PD etiology and preventive strategies. Although still controversial, the Braak and dual-hit hypotheses that PD may start peripherally in the olfactory structures and/or the gut provides a theoretical platform to identify the triggers and modifiers of PD prodromal development and progression. This is particularly true for the search of environmental causes of PD as the olfactory structures and gut are the major human mucosal interfaces with the environment. In this review, we lay out our personal views about how the Braak and dual-hit hypotheses may help us search for the environmental triggers and modifiers for PD, summarize available experimental and epidemiological evidence, and discuss research gaps and strategies.

## 1. Introduction

Idiopathic Parkinson's disease (PD) is the second most prevalent neurodegenerative disease, affecting 1–2% of the older adults. Multiple recent reports show that PD emerges as the fastest growing neurological disease in terms of both prevalence and death (Darweesh et al., 2018; GBD Parkinson's Disease Collaborators, 2018), which cannot be solely explained by population aging. There is preliminary albeit inconsistent evidence that the incidence of PD might have been increasing over the past decades (Savica et al., 2017). Despite this increasing burden of PD, research on the environmental triggers and modifiers for PD development has been stagnant. While a portion of the PD cases have a clear genetic component, most late-onset PD are likely the results of environmental contributions and their interactions with genetic susceptibility. The search for environmental causes of PD has been hampered by the current lack of understanding of the disease's natural history. In particular, by the time of PD diagnosis with bothering cardinal motor signs, the disease might have undergone decades of “insidious” prodromal development, during which many factors may come into play to initiate pathology or modify its progression. While some experimental models (e.g., Thy1-aSyn mouse) recapitulates key pathogenesises (e.g.,

alpha-synuclein accumulation and progressive changes in dopamine release and striatal content) and phenotypical features of prodromal PD (Chesselet et al., 2012), animal models and human studies are yet to fully unveil these complex, decades-long processes. To some extent, the lack of understanding of this “black box” of PD prodromal development has led to the “blind men and an elephant” dilemma in the research of PD etiology. Consequently, even the most robust epidemiological findings on non-genetic risk factors for PD (e.g., smoking or coffee drinking) are subject to alternative explanations such as reverse causation or unmeasured confounding (Ritz and Rhodes, 2010). On the other hand, the prolonged prodromal development of PD may also present an unprecedented opportunity to understand PD etiology and disease prevention. For example, by identifying factors that contribute to various stages of PD prodromal development, we may be able to eventually reveal the sequence of events in PD development and their causes, fundamentally improving understanding of PD etiology and developing preventive strategies.

The scientific premises of such research endeavors however critically depend on a better understanding of the stages of PD prodromal development and the identification of readily assessable staging markers. To this end, the Braak hypothesis (Braak et al., 2006) presents

\* Corresponding author at: Department of Epidemiology and Biostatistics, College of Human Medicine, Michigan State University, 909 Wilson Rd, East Lansing, MI 48824, USA.

E-mail address: [chenho19@msu.edu](mailto:chenho19@msu.edu) (H. Chen).

<https://doi.org/10.1016/j.nbd.2021.105601>

Received 8 September 2021; Received in revised form 5 December 2021; Accepted 22 December 2021

Available online 23 December 2021

0969-9961/© 2021 The Authors.

Published by Elsevier Inc.

This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

the exact hypothetical framework that dissects the steps of PD prodromal development, which posits that the PD pathogenesis may first initiate in the olfactory structures and the gut enteric nerves, years if not decades, before spreading to the substantia nigra where dopaminergic neuron death occurs and PD motor dysfunction arises. Although the hypothesis itself is still somewhat controversial as clinical PD is heterogeneous and contradictory pathological findings exist, it raises the intriguing possibility that, at least in some PD patients, the pathogenesis may start in the peripheral system, and some of the long-observed nonmotor symptoms in PD are likely integral part of the disease's natural history and prodromal development. In support, accumulating evidence shows that olfactory impairment (Chen et al., 2017; Ross et al., 2008), REM sleep behavior disorder (RBD) (Postuma et al., 2009; Postuma et al., 2015), and constipation (Abbott et al., 2001; Gao et al., 2011; Savica et al., 2009) strongly predict future risk of PD, and they may have developed years or decades before PD clinical diagnosis. Most symptoms, with the exception of polysomnography-confirmed RBD, lack clinical specificity to PD. However, research on these prodromal symptoms, coupled with studies on other PD biomarkers (e.g., neuroimaging, blood, saliva, or skin-biopsy based), has opened the window to characterize at-risk populations for PD, define PD prodromal phenotype, and identify potential causes and preventive measures to slow down or stop its progression to overt clinical PD.


More importantly, these advances may enable us to entertain the fundamental question of PD etiology and prevention - when, where, and how does PD pathogenesis start, develop and progress? Fig. 1 presents a hypothetical scenario (Chen, 2018; Chen and Ritz, 2018). At birth or young age, the person may have some of the genetic variants or nongenetic events (e.g., head injury) that put them at a higher risk of PD. With occupational uses of certain pesticides or organic solvents, PD pathogenesis may begin to develop at mid-adulthood. This process may be insidious initially, but over time as the pathogenesis progresses, non-specific symptoms (e.g., poor olfaction, constipation, sleep problems) or even the specific RBD symptoms may develop. Occupational hazards continue, and additional environmental factors may come into play (e.g., stopped smoking, or another head injury). Eventually, the more notable cardinal motor signs develop, leading to a PD diagnosis later in life. It is important to note that this process is slowly progressive and

may take decades, and its clinical presentations and sequential orders are likely heterogeneous. While there are still substantial challenges to adequately define prodromal PD, by assessing these early nonmotor and motor symptoms as intermediate phenotypes, we may be able to bring new insights into this “black-box” of PD prodromal development by identifying factors that initiate PD pathogenesis, lead to its prodromal phenotypes, or modify progression to clinical PD.


The dual-hit hypothesis further speculates that environmental toxins such as pesticides, air pollutants, metals or virus, may enter the body via the nasal cavity or the gut, initiate PD pathogenesis, gain access to the brain via the olfactory pathway or the vagus nerve, and eventually lead to clinical PD (Fig. 2) (Hawkes et al., 2007; Hawkes et al., 2009). Notably, the nasal cavity and gut are the two anatomic sites where the human mucosal surfaces directly interact with the environment, where inflammation commonly occurs, and where paths to the brain are well established. Although the PD dual-hit hypothesis itself is yet to be systematically examined, if proven true, it will have profound implications for PD prevention and treatment as these environmental factors are modifiable. In recent reviews (Chen, 2018; Chen and Ritz, 2018), we have speculated how this prodromal framework may help advance the understanding of environmental causes for PD, major research directions, and challenges. In this review, we focus on summarizing available epidemiological and experimental evidence for this PD dual-hit hypothesis. The potential roles of the gut-brain axis and gut microbiota in PD etiology have stimulated substantial research interests and have been the topic of extensive recent reviews (Elfil et al., 2020; Killinger and Labrie, 2019; Lionnet et al., 2018; Scheperjans et al., 2018; Sharabi et al., 2021; Travagli et al., 2020), we hereby first discussed in detail the olfactory pathway and its environmental relevance in the context of PD etiological research.

## 2. The olfactory pathway

In the Braak staging paradigm, the olfactory bulb is one of the sites where PD pathogenesis may start. While poor olfaction has long been recognized as a prodromal symptom of PD (Ross et al., 2008), it is difficult to estimate its timeline leading to PD clinical onset. Nonetheless, population-based cohort studies have shown that poor olfaction



**Fig. 1.** A Systematic Life-long Approach to Study Environmental Triggers and Modifiers of PD. Late-onset sporadic PD takes decades to develop and has a prolonged prodromal stage. By assessing PD prodromal symptoms as intermediate phenotypes, we may be able to bring new insights into the “black-box” of PD etiology by identifying factors that trigger PD pathogenesis, lead to its prodromal phenotypes, or modify its phenotypical conversion to clinical PD at various stages of disease development. Modified from *J Parkinsons Dis.* Chen & Ritz, “The Search for Environmental Causes of Parkinson’s Disease: Moving Forward” 8 (2018) S9–S17. Copyright (2018), with permission from IOS Press. The original publication is available at IOS Press through <https://doi.org/10.3233/JPD-181493>.



**Fig. 2.** The Environmental Dual-Hits that May Trigger PD Pathogenesis. Environmental exposures may enter the body via the nose or the mouth, which may initiate PD pathogenesis in the olfactory structures and/or the gut enteric nerves in susceptible individuals via mechanisms such as inflammation or microbiota dysbiosis; over years, the pathology may progress to the central olfactory structures and/or the lower brain stem, and eventually lead to dopaminergic neuron loss in the substantia nigra. [Created with [BioRender.com](https://www.biorender.com)] Part of the figure was modified from Ambrosini et al. *Frontiers in Aging Neuroscience* (2019), Volume 11, Article 130 doi: <https://doi.org/10.3389/fnagi.2019.00130>

strongly predicts PD risk with up to 10 years of follow-up (Chen et al., 2017; Mahlknecht et al., 2018). Further, in a reconstructed timeline based on data from 154 idiopathic RBD patients, poor olfaction appeared to be the first symptom of clinical synucleinopathy, starting >20 years before phenotypical conversion to PD or Lewy body dementia (Fereshtehnejad et al., 2019). In comparison, constipation began to develop ~10 years before the phenotypical conversion and motor signs ~5 years before.

The causes for poor olfaction in older adults also remain elusive. Except for age, male sex, and a few acquired causes (Doty, 2009), the roles of environmental exposures in age-related olfactory impairment are largely unknown (Doty, 2015). We hereby critically evaluate the rationale and empirical evidence for key environmental exposures in relation to poor olfaction and their potential relevance to PD, focusing on pesticides, air pollutants, cigarette smoking, heavy metals and

organic solvents (Table 1). These airborne environmental exposures have been studied in relation to PD risk and, at least theoretically, they can gain access to the brain via the nasal and olfactory pathway bypassing the blood-brain barrier (Maher et al., 2016; Prediger et al., 2011; Sasajima et al., 2015).

### 2.1. Epidemiological evidence

Overwhelming epidemiological and experimental evidence supports roles of pesticides in PD development, albeit data on specific pesticides, detrimental exposure levels, windows of exposure, and routes of entry are still limited. One can, however, reasonably hypothesize that pesticides may trigger PD pathogenesis through the nasal olfactory pathway. When airborne, pesticides may damage the olfactory epithelium, impairing nerve function, inducing local acute or chronic inflammation,

**Table 1**  
Summary of epidemiological evidence <sup>a</sup> on selected potential airborne environmental “risk factors” for PD in relation to olfaction.

	Study design	Study setting and participants	Exposure assessment	Outcome assessment	Primary findings	Comments
Pesticides	Cross-sectional, case-control, & longitudinal studies	Mostly occupational setting (e.g., farmers or farm workers vs. others); a few population-based studies; mostly working age population or older; primarily men	Self-reported pesticide exposures; farmer as a surrogate for pesticide exposure; biospecimen concentration (e.g., urinary sample)	Sniffin-Stick odor identification and threshold tests; self-reported olfactory impairment	Mostly positive associations (OR: 1.02–1.49), but data are not entirely consistent, especially on which olfaction domain was impaired.	Exposure assessment errors; self-reported olfaction not accurate; no data on whether the impairment was transitory, progressive or permanent (i.e., age related decline)
Air pollutants	Cross-sectional, & case-control studies	Population-based studies; wide age range (i.e., young, middle, & old age); both men and women	Ecological (e.g., residents of Mexico city vs. another city); criteria air pollutants (e.g., NO <sub>2</sub> or PM <sub>2.5</sub> )	Sniffin-Stick odor identification and threshold tests; clinical anosmia based on ICD codes	Mostly positive associations (OR: 1.20–1.73), but some null association findings	No longitudinal studies; unsure if the impairment was transitory or age-related; possible effect modification by age
Cigarette smoking	Cross-sectional, & longitudinal studies	Population-based studies; PD patients vs. others; wide age range, including young, middle, and old age, both men and women	Self-reported smoking history (e.g., never, past, current), pack-years, years of smoking, smoking cessation	Sniffin-Stick odor threshold and odor identification; UPSIT; MODSIT; SDOIT; self-reported OI; B-SIT; SOIT; Connecticut Chemosensory Clinical Research Center test; other non-commercial tests (e.g., identification of series dilution of common odors)	Mostly positive associations (OR: 1.36–2.43) especially for current smoking, but not entirely consistent. In PD patients, smoking was not positively associated with poor olfaction	Unsure if the impairment was transitory or age-related; analyses limited to PD may lead to a biased association of smoking with poor olfaction
<sup>4</sup> Heavy metals	Cross-sectional studies	Predominantly occupational settings (e.g., welders vs. others); a few population-based studies; mostly working age population or older, occasionally children	Surrogate (e.g., welder); biospecimen concentrations (blood, urine, hair, fingernail, bone); most studied metals include manganese, cadmium, chromium and lead	Olfactory acuity by olfactometer; Sniffin-Stick odor threshold and odor identification; UPSIT; self-reported olfaction other non-commercial tests (e.g., identification of series dilution of common odors)	Manganese: inconsistent - positive, negative or null associations; cadmium, & lead: inconsistent - positive or null associations; Chromium: positive associations, but data are limited	Small sample sizes, cross-sectional, and occupational setting; unsure if the impairment was transitory (due to occupational hazards) or age-related.
Organic solvents	Cross-sectional studies	Occupational settings (e.g., workers in paint manufacturing facilities, printing workers, or medical laboratory employee vs others); working age; men and women	Surrogate (e.g., job-title based exposure assessment)	Self-reported olfaction; Sniffin-Stick odor threshold and odor identification; UPSIT	Mostly positive associations, but some show null associations. There is a potential effect modification by smoking	Small sample sizes, cross-sectional, and occupational setting; unsure if the impairment was transitory (due to occupational hazards) or age-related.

Abbreviations: OR: odds ratio; UPSIT: University of Pennsylvania Smell Identification Test; MODSIT: Modular Smell Identification Test; SDOIT: The San Diego Odor Identification Test; B-SIT: Brief Smell Identification Test; SOIT: Scandinavian Odor Identification Test.

<sup>a</sup> See references and citations in the text due to space limit.

and disrupting the xenobiotic function, immune protection, and microbiome of the olfactory mucosa (Doty, 2015). Pesticides may find their way to the brain via olfactory structures, and in susceptible individuals, may initiate or exacerbate the abnormal aggregation of  $\alpha$ -synuclein (Kumar et al., 2016; Maturana et al., 2015; Rey et al., 2016; Rey et al., 2018; Rokad et al., 2017). The nasal mucosa, at the interface between the body and the environment, once damaged, may have diminished mechanical and/or immune protection and thus impart a higher susceptibility to future virus invasions or environmental neurotoxins that lead to PD. Alternatively, pesticides may also enter the body via the digestive tract and initiate synucleinopathy in the gut, which may later spread to the brain (Braak et al., 2006). Over time, these mechanisms may individually or synergistically contribute to age-related poor olfaction and neurodegeneration.

Despite such plausible mechanisms, there are few empirical studies of pesticides and olfaction in humans. A case-report clearly documented progressive and irreversible loss of the sense of smell in an Italian physician following acute indoor exposure to high levels of a pyrethrin-based insecticide (Gobba and Abbacchini, 2012). Several other studies (Ahman et al., 2001; Holmstrom et al., 2008; Quandt et al., 2017; Quandt et al., 2016) reported that farmers or farm workers were more likely to report poor olfaction or fail a smell test than the comparison group; however, these studies are mostly cross-sectional and did not directly examine pesticide and poor olfaction. We examined both occupational pesticide uses and high pesticide exposure events (HPEE, e.g., major accidental spills) in relation to self-reported poor olfaction ~20 years later among farmers in the Agricultural Health Study (Shrestha et al., 2019; Shrestha et al., 2021). In the analysis of occupation data from 20,409 farmers, we found modest associations with self-reported poor olfaction for 20 out of the 50 specific pesticides (odds ratios between 1.11 and 1.33) (Shrestha et al., 2021). In another analysis of 11,232 farmers (Shrestha et al., 2019), a history of HPEE was associated with a 49% higher odds of reporting poor olfaction ~20 years later, and statistically significant associations were found for two organochlorine insecticides (DDT and lindane) and four herbicides (alachlor, metolachlor, 2,4-D, and pendimethalin). These preliminary data, once confirmed, may suggest that these pesticides contribute to the early stages of PD development. To our knowledge, no epidemiological study has examined whether pesticides modify the phenotypical conversion from poor olfaction to clinical PD.

Many of the same biological rationales for linking pesticides to poor olfaction and/or PD can be easily extended to air pollutants. For example, the fine particulate matter (PM<sub>2.5</sub>) may induce  $\alpha$ -synuclein pathology in the olfactory bulb via mechanisms of neuroinflammation and oxidative stress (Block et al., 2012), and alternatively, they may also gain access to the human body via the gastrointestinal tract, for example, by nasal dripping or saliva swallowing as the dual-hit hypothesis has suggested (Hawkes et al., 2009). Unlike pesticides, the epidemiological evidence on major air pollutants and PD risk has been inconsistent (Jo et al., 2021; Liu et al., 2016; Palacios et al., 2017; Ritz et al., 2016; Toro et al., 2019; Willis et al., 2010). For example, in the most recent comprehensive analysis of the Korean National Health Insurance data, the only statistical difference in PD risk was found when comparing the highest vs. lowest exposure categories of nitrogen dioxide (NO<sub>2</sub>), whereas no associations were found of PD with other major criteria pollutants (i.e., PM<sub>2.5</sub>, PM<sub>10</sub>, ozone, or carbon monoxide) (Jo et al., 2021).

Data on air pollutants and poor olfaction are also limited. Earlier studies are mostly ecological in design and reported worse olfaction in residents of highly polluted cities such as Mexico City as compared to their counterparts in surrounding areas of cleaner air (Calderon-Garciduenas et al., 2010; Guarneros et al., 2009; Hudson et al., 2006; Sorokowska et al., 2015; Sorokowska et al., 2013). The association of ambient air pollutants with olfaction was also examined in two large cross-sectional studies (Adams et al., 2016; Ajmani et al., 2016) and a case-control study (Zhang et al., 2021). In a US population, positive

associations with poor olfaction were found for both exposures to PM<sub>2.5</sub> (Ajmani et al., 2016) and NO<sub>2</sub> (Adams et al., 2016); however, the association with PM<sub>2.5</sub> was limited to younger participants ages 57–64 and participants from the Northeast regions. In a recent case-control study, PM<sub>2.5</sub> exposure was associated with the odds of having anosmia (Zhang et al., 2021). Therefore, the evidence on air pollution and olfactory impairment is preliminary but suggestive.

Unlike pesticides or air pollutants, cigarette smoking was inversely associated with the risk of PD (Chen et al., 2010). This relationship is inarguably the best-established epidemiological observation for PD, albeit its causality is still debatable. If this association is indeed causal that smoking reduces the risk of developing PD, one would logically expect smoking may protect against poor olfaction which is a prodromal marker of PD. The overall evidence to date however suggests the opposite. Most available studies are cross-sectional in design and found current smokers had a higher odds of having poor olfaction, as summarized in a 2017 meta-analysis (Ajmani et al., 2017). This observation is further supported by two recent large cohort analyses, both of which found current smokers are about 2–3 times more likely to develop poor olfaction (Palmquist et al., 2020; Schubert et al., 2021). Further, several recent studies specifically investigated this association in PD patients and found that PD smokers had better olfaction than PD nonsmokers (Lucassen et al., 2014; Sharer et al., 2015). This type of analysis however can be misleading in the causal inference of smoking and olfaction in prodromal PD due to potential collider bias (Hernan et al., 2004). As such, the evidence is not consistent with the possibility that smoking protects against hyposmia. Interestingly, a few studies also examined the association of smoking with RBD, the other most prominent prodromal symptom of PD. In these studies, cigarette smoking is associated modestly with a higher risk of RBD (Postuma et al., 2012; Shrestha et al., 2018) but not with its phenotypical conversion to PD or other types of clinical synucleinopathy (Postuma et al., 2015). Therefore, data to date do not reconcile with the strong inverse association of smoking with PD. While they do not necessarily rule out a causal relationship that cigarette smoking reduces PD risk for several reasons (Chen, 2018), they offer no support that smoking is protective in the early stage of PD development.

Finally, the Braak and dual-hit hypotheses also provide a platform to systematically examine the roles of several other environmental exposures such as heavy metals (e.g., manganese) (Mezzaroba et al., 2019) and organic solvents (e.g. trichloroethylene) (Lock et al., 2013) in relation to olfactory impairment in the context of PD prodromal development. Available studies were often conducted in occupational settings, participants were often young, and it was unclear whether the olfaction deficit was transitory or permanent. Data are not entirely consistent. Several studies reported that occupational exposures to cadmium, lead, and chromium were associated with poor olfaction (Adams and Crabtree, 1961; Casjens et al., 2018; Grashow et al., 2015; Kitamura et al., 2003; Liu et al., 1985; Mascagni et al., 2003; Potts, 1965; Sulkowski et al., 2000). In contrast, a couple of studies reported that occupational exposure to manganese or higher blood manganese level was associated with better olfaction performance (Antunes et al., 2007; Casjens et al., 2017; Lucchini et al., 1997), whereas children from areas with high manganese exposures performed worse on olfactory identification than their peers from low exposure areas (Guarneros et al., 2020). Studies on solvent exposures were predominantly published before 2010, and most found workers with occupational exposures to solvents or related chemicals were more likely to report poor olfaction or perform worse on the olfaction identification or threshold tests than their counterparts without such exposures (Ahlström et al., 1986; Baelum et al., 1982; Bolla et al., 1995; Holmström et al., 1995; Lee et al., 2018; Lucchini et al., 1997; Sandmark et al., 1989; Schwartz et al., 1989; Schwartz et al., 1990; Yu et al., 2004; Zibrowski and Robertson, 2006). Therefore, the overall literature appears to support associations of occupational exposures to certain heavy metals and solvents with poor olfaction, but the relevance of these associations to PD development is yet to be assessed.

## 2.2. Experimental evidence

Compared to epidemiological data, experimental studies have more directly examined the Braak and dual-hit hypotheses in the context of PD research, as summarized below. It is important to note that although there is experimental evidence for self-propagating alpha-synuclein assemblies spread from the gut or the olfactory bulb to the brain, most evidence is based on studies of alpha-synuclein preformed fibril (PFFs) injections (Challis et al., 2020; Kim et al., 2019; Van Den Berge et al., 2021), and thus the factors that may trigger the de novo formation of “spread-competent” alpha-synuclein assemblies remain unknown. Furthermore, it is still unclear if this spreading process is driven by a prion-like or cell-autonomous mechanism. Therefore, the environmental “triggers” considered here should be viewed in the context of the experimental setting and the observed phenotype, which often lacks the hallmark pathological alpha-synuclein inclusions. Nevertheless, animal studies have provided preliminary evidence that the alpha-synuclein pathology can spread from the olfactory bulb throughout the brain to secondary sites, and to some extent, mimic the distribution of synucleinopathy in human diseases such as PD (Rey et al., 2013; Rey et al., 2016).

Experimental studies showed that several pesticides, when administered through the nasal cavity, can damage dopamine-containing neurons in the brain and recapitulate some of the PD phenotypes (Sasajima et al., 2015; Sasajima et al., 2017; Toyoda et al., 2020; Voronkov et al., 2017). Rotenone is a pesticide that inhibits the complex I of the mitochondrial electron transport chain, which leads to the inhibition of ATP synthesis, excessive production of oxidative radicals, and the death of dopaminergic neurons. Intranasally administered rotenone to mice impairs mitral cell activity in the olfactory bulb, possibly due to the denervation of inhibitory dopaminergic inputs (Sasajima et al., 2017). Mitral cells and their projections are common Lewy pathology containing structure in PD olfactory bulb (Sengoku et al., 2008). It is possible that mitral cell dysfunction following rotenone exposure results in Lewy pathology which then could project to several brain nuclei involved in PD (Cersosimo, 2018). Chronically administered intranasal rotenone to mice may damage the nigrostriatal dopamine system, specifically axons of the dorsal striatum (Ferrante et al., 1997), although data are not consistent across studies (Rojo et al., 2007). Furthermore, structures resembling Lewy pathology have been observed in the substantia nigra of mice with chronic exposure to rotenone (Betarbet et al., 2000). Intranasally administered pesticide paraquat accumulates in structures throughout the mouse brain and results in impairment of olfactory discrimination task (Anderson et al., 2021). However, like rotenone, the PD phenotype produced by paraquat inhalation in rodents is mild (Rojo et al., 2007), and therefore it is yet to demonstrate if nasal exposure to these pesticides alone is sufficient to initiate PD pathogenesis. MPTP has a chemical structure similar to paraquat, and intranasal exposure closely recapitulates many PD phenotypes, including olfactory dysfunction, motor abnormalities, and dopaminergic neuron loss in the olfactory bulb and substantia nigra (Prediger et al., 2010; Prediger et al., 2006; Prediger et al., 2009; Rojo et al., 2006).

Metals have long been implicated in PD pathogenesis. Various metals may promote alpha-synuclein aggregation (Jinsmaa et al., 2014; Paik et al., 1999; Wright et al., 2009), but intranasal metal exposure has not been extensively investigated in a PD model. Nevertheless, the olfactory bulb absorbs metal fumes (Sunderman, 2001), and some (e.g., manganese, nickel, and zinc) may be further transported throughout the brain via secondary olfactory neurons (Gianutsos et al., 1997; Sunderman, 2001). Recent evidence further suggests that manganese may promote the spread of alpha-synuclein oligomers via exosomes (Harischandra et al., 2019). Furthermore, mice exposed to intranasal vanadium, a metal in welding fumes that has been associated with PD (Park et al., 2005; Racette et al., 2001), showed a loss of tyrosine hydroxylase positive neurons of the olfactory bulb glomerular layer (Ngwa et al., 2014). However, the Lewy pathology was not assessed in the study (Ngwa et al.,

2014). As to organic solvents, although systemically administered trichloroethylene may kill midbrain dopamine neurons (Guehl et al., 1999; Keane et al., 2019) and induce alpha-synuclein accumulation (Liu et al., 2010), to our knowledge, its potential adverse effects via inhalation has not been directly investigated.

In summary, despite the high biological plausibility that environmental toxicants may trigger PD pathogenesis through the olfactory pathway, relevant research is still in its infancy. Preliminary data from animal models suggest that the intranasal administration of certain pesticides or heavy metals may lead to the initiation and spreading of abnormal alpha-synuclein and aggregation that, to some extent, recapitulate the human Lewy pathology. But conclusive systematic research is yet to be conducted. Epidemiological evidence on environmental risk factors for olfactory impairment is preliminary. To our knowledge, none were conducted in the context of PD research, and thus far has provided limited clues to environmental triggers and accelerators of PD pathogenesis in humans.

## 3. The gut-to-brain pathway

The Braak hypothesis also posits that synucleinopathy may initiate in the enteric nerves of the gut and later spread along the vagus nerve into the brain (Hawkes et al., 2009). In fact, the presence of Lewy pathology in the enteric nervous system of PD patients was reported decades earlier (Wakabayashi et al., 1988). Further, constipation has been robustly identified as an early prodromal symptom of PD (Abbott et al., 2001; Gao et al., 2011; Savica et al., 2009), albeit the strength of the association was modest as compared with that for poor olfaction or RBD. The pivotal roles of the gut-brain axis to PD pathogenesis have been the topic of extensive recent reviews (Elfil et al., 2020; Killinger and Labrie, 2019; Lionnet et al., 2018; Scheperjans et al., 2018; Sharabi et al., 2021; Travagli et al., 2020). We hereby briefly summarized the key epidemiological and experimental evidence on this topic and experimental findings on the potential roles of environmental factors.

### 3.1. Epidemiological evidence

Besides the well-established observation of constipation in prodromal PD, multiple lines of human empirical evidence support a critical role of the gut-brain axis in PD pathogenesis. First, in a systematic histopathology study of PD patients ( $n = 17$ ) and other synucleinopathies, Beach et al. reported phosphorylated alpha-synuclein throughout the gastrointestinal system (Beach et al., 2010). Interestingly, they did not find isolated synucleinopathy in the gut, challenging the gut-to-brain spreading in prodromal PD. In their recent publication with more PD patients ( $n = 53$ ), 81% had synucleinopathy in the stomach and 89% in the vagus nerve adjacent to the neck carotid artery, as compared with no positivity in control subjects, supporting the involvement of the gastrointestinal tract and vagus nerve in PD pathogenesis (Beach et al., 2021). Second, multiple register-based epidemiologic analyses reported a modest positive association of inflammatory bowel diseases and PD with a meta-analyzed risk ratio of 1.24 (95% confidence interval: 1.15–1.34) (Zhu et al., 2022). Interestingly, IBD patients with anti-tumor necrosis factor treatment had a 78% lower risk than those without the therapy (Peter et al., 2018). These findings are consistent with the hypothesis that chronic intestinal inflammation or a “leaky gut” may contribute to PD pathogenesis. Third, epidemiological evidence suggests that *Helicobacter pylori* infection is not only associated with clinical PD and its severity (Dardiotis et al., 2018), it may also play a role in developing PD. In a retrospective cohort analysis of insurance data from Taiwan, *H. pylori* infection was associated with a > 2 fold higher PD risk (Huang et al., 2018). Fourth, several studies examined surgical removal of the appendix or tonsil, the two prominent mucosa-associated lymphatic tissues in the gastrointestinal tract, in relation to PD risk. Findings are not consistent (Marras et al., 2016; Palacios et al., 2018; Tysnes et al., 2015), but analyses of the Swedish patient registers

showed a lower PD risk in people with appendectomy (Killinger et al., 2018; Liu et al., 2020). Finally, two independent studies, using data from the Swedish and Danish patient registers, reported that people with truncal vagotomy had about half of the PD risk compared to people without (Liu et al., 2017a; Svensson et al., 2015), supporting the possibility of the gut-to-brain spreading pathway via the vagus nerve. These data, taken together, lend supports to the roles of gut inflammation and gut-brain axis in the development of PD. Notably, much of the evidence was from analyses of administrative databases. While this type of data has clear strengths (e.g., large samples and comprehensive coverage), they were not collected for research purposes, have limited data on potential confounders, and identify cases from diagnostic codes (Mues et al., 2017). Therefore, caution is needed in analyzing administrative data and interpreting their results.

### 3.2. Experimental evidence

Animal experimental studies provided more direct support to this spreading hypothesis, demonstrating that peripheral  $\alpha$ -synuclein aggregates can be actively transported to the brain via the vagal nerve and can initiate a PD-like phenotype. Following injection, PFFs rapidly disseminate to neural circuits (Okuzumi et al., 2018) based on connectivity and possibly  $\alpha$ -synuclein expression patterns (Henderson et al., 2019). It appears that they are actively trafficked via axonal transport (Freundt et al., 2012; Holmqvist et al., 2014), possibly by interacting/binding with mobile membranous organelles and vesicles. PFFs injected into the gut travel along the vagal nerve to the dorsal motor nucleus (Ulusoy et al., 2017; Ulusoy et al., 2013), but other routes are also possible (Borghammer, 2021). Bidirectional spread between the central and peripheral nervous systems is also possible. A recent comprehensive analysis in rodents found gut-to-brain spread, caudal to rostral progressive  $\alpha$ -synuclein pathology, motor/cognitive impairments, gastrointestinal dysfunction, and dopaminergic neuron loss in the midbrain (Kim et al., 2019). Convincingly, both in wide-type mice with a vagotomy and SNCA knockout mice, the phenotype following PFFs injections into the gut was abolished (Kim et al., 2019). In summary, the spread of synucleinopathy from peripheral sites to the brain is largely supported by experimental evidence. However, PFFs injected into the gut do not invariably spread to the CNS and progressively throughout the brain (Manfredsson et al., 2018), and several critical factors including dose, site of injection, and the PFF strain/species are important. To date, the evidence of “gut only”  $\alpha$ -synuclein pathology in humans has yet to be demonstrated (Beach et al., 2010) which is required for the gut origin hypothesis of PD. However the lack of evidence could be due to the relatively low sampling of gut tissues per volume, akin to searching for a needle in a haystack.

### 3.3. Potential roles of environmental factors

One key implication of the gut-to-brain transmission of PD pathogenesis is the potential environmental contributions. Research on the roles of environmental toxicants and the gut-to-brain synucleinopathy spreading has been largely limited to pesticides in experimental studies. Orally administered rotenone causes apparent progressive pathology in the intermedialateral nucleus of the spinal cord and substantia nigra (Pan-Montojo et al., 2010), which can be prevented by vagotomy (Pan-Montojo et al., 2012). However, the pathology was determined by total  $\alpha$ -synuclein protein staining, which has a low specificity for the disease process. A separate study determined that TLR4 knockout mice are somewhat protected from the inflammatory and neurotoxic effects of oral rotenone treatment (Perez-Pardo et al., 2019), suggesting a critical role of inflammation in this process. Indeed, experimental studies have found that both acute and chronic gut inflammation may modulate the expression and possibly aggregation of  $\alpha$ -synuclein in rodents (Kishimoto et al., 2019; Prigent et al., 2019), but it is yet to be demonstrated that gut inflammation can lead to prion-like synucleinopathy. Besides

rotenone, paraquat was also tested in animal models. Orally administered paraquat to mice expressing A53T  $\alpha$ -synuclein had accelerated PSER129 pathology formation throughout the small intestine (Naudet et al., 2017). Although pesticides might promote abnormal  $\alpha$ -synuclein aggregates in the gut, other compounds might exacerbate their toxic effects. For example, paraquat produces a mild and often poorly reproducible PD-like phenotype in mice, but small doses of paraquat taken with dietary lectins produce a phenotype with several key features of PD, including the peripheral PSER129 inclusion (Anselmi et al., 2018). Authors of this study hypothesize that lectins may promote the absorption of the orally administered paraquat and thus enhance its toxic effects (Anselmi et al., 2018).

## 4. Microbiota

In recent years, the potential role of the microbiota in PD pathogenesis and progression has gained much attention (Boertien et al., 2019; Elfil et al., 2020) as it modulates the immune system and the absorption and metabolism of nutrients, medications, and environmental toxins. Certain aspects of microbiota composition are already determined in earliest childhood, e.g., by mode of delivery, breastfeeding, and antibiotic exposure. During vaginal delivery, the first colonization of the offspring occurs through the maternal microbiota, potentially affecting the inheritance of certain vulnerabilities independently of genotype (Sandoval-Motta et al., 2017). Throughout the lifespan, the microbiota colonizing the outer and inner surfaces of our body continues to be at the interface between the environment and the human organism and potentially influences their interactions. Further, converging evidence suggests that the gut microbiota influences the development, function, and aging of the brain (Heijtz et al., 2021), although the fundamental mechanisms by which the microbiota influences PD pathogenesis remain poorly understood.

### 4.1. Epidemiological evidence

Many studies have linked the gut microbiota to PD and to its motor and nonmotor symptoms. Despite some variabilities in the results and adjustments for medications, comorbidities and diet, study findings suggest a shift to a potentially proinflammatory microbiome composition in PD patients (Romano et al., 2021) with increased markers of gut inflammation and permeability (Aho et al., 2021; Schwiertz et al., 2018). Other potential pathways by which microbiota may influence PD development and progression include bacterial metabolites as well as potential cross seeding between bacterial amyloid proteins and  $\alpha$ -synuclein (Aho et al., 2021; Chen et al., 2016; Cirstea et al., 2020). The most consistent metabolic alteration so far has been reduced fecal levels of bacterially produced short-chain fatty acids (SCFAs), in particular butyrate, in PD patients (Aho et al., 2021; Unger et al., 2016). SCFAs can have a broad influence on the human host and brain through impact on (neuro)inflammation, epigenetic regulation, and other mechanisms (Silva et al., 2020).

The few longitudinal studies on microbiota in PD suggest links between gut microbiota composition and disease progression in established PD patients (Aho et al., 2019). Studies about microbiota contributions in the prodromal phase are scarce. To this end, one study found substantial overlaps in differential microbiome abundant taxa between PD patients and idiopathic RBD patients as compared with healthy controls, suggesting microbiota changes before PD clinical diagnosis (Heintz-Buschart et al., 2018). However, another study found that microbiota changes were more pronounced in established PD as compared to iRBD (Nishiwaki et al., 2020), suggesting that changes may evolve as disease progresses. A recent large study examined microbiota associations with known PD prodromal and risk markers and found differential microbiota abundance linked to constipation, possible RBD, smoking, and subthreshold parkinsonism (Heinzel et al., 2021), but not to olfactory impairment or other markers of prodromal PD.



Few human studies have examined nasal and oral microbiota in PD. Reliable sampling of the microbiota and DNA amplification from the nasal cavity has turned out to be challenging due to the low bacterial biomass encountered in nasal swabs. Therefore, there is a considerable risk of laboratory contaminants to influence the results when many amplification cycles are needed. Two recent studies (Pereira et al., 2017; Li et al., 2021) failed to find significant differential compositions of nasal microbiota between PD patients and controls based on swab sampling. The third study (Heintz-Buschart et al., 2018) used a nasal wash sampling technique and found no convincing differences of nasal microbiota across PD patients, iRBD patients and healthy controls. With respect to the oral microbiota, several studies reported significant differences between PD patients and controls. For example, the above referenced two studies (Pereira et al., 2017; Li et al., 2021) reported low abundance of *Neisseria* and *Neisseriaceae* and increased abundance in *Lactobacillaceae* /*Lactobacillales* in PD oral swab samples. A recent study (Rozas et al., 2021) further reported increased oral soft tissue abundance of *Lactobacillus* together with an increase of *Tannerella forsythia*, *Prevotella intermedia*, and opportunistic pathogens. Also oral hard tissue samples showed increased of opportunistic oral pathogens in PD. Factors that significantly influenced soft tissue beta-diversity and microbiota composition included dysphagia, drooling, and salivary pH. Taken together, while studies to date failed to find convincing links of nasal microbiota to PD, changes in oral microbiota have been suggested.

#### 4.2. Experimental evidence

The role of gut microbiota in PD has also been investigated in experimental studies. When raised germ-free or when the microbiota was eradicated by antibiotics, alpha-synuclein overexpressing mice showed little synucleinopathy, neuroinflammation, and motor symptoms (Sampson et al., 2016), suggesting gut microbiota are required for PD pathogenesis. Conversely, introducing PD-patient microbiota to these mice resulted in PD pathology and motor impairment (Sampson et al., 2016), which may be related to SCFAs produced by the microbiota. On the other hand, several studies suggest a beneficial or even neuroprotective role of SCFAs such as butyrate in PD experimental models (Liu et al., 2017b; Paiva et al., 2017).

#### 4.3. Potential roles of environmental factors

It is also important to consider that the gut microbiota acts at the interface of the human organism with the environment, and thus may play a role in how environmental influences impact disease initiation and progression. One such influence is the regulation of gut wall permeability (Ambrosini et al., 2019). Indeed, increased gut wall permeability or "leaky gut" has been described in PD (Forsyth et al., 2011), although this finding has not always been consistent and may vary between gut segments. Such increased permeability may facilitate access of potentially harmful environmental factors to the gut tissue and systemic circulation as well as immune activation.

Gut microbiota may influence how pesticides impact PD pathogenesis. Experimental evidence suggests that oral application of rotenone affects gut permeability only if a gut microbiota is present, but not in germ-free mice (Bhattarai et al., 2021). Rotenone induces microbiota changes, and these may have proinflammatory effects that are dependent on TLR4 signaling (Perez-Pardo et al., 2019; Perez-Pardo et al., 2018). Smoking and coffee drinking, two factors associated with lower PD risk in observational studies, may influence microbiota composition (González et al., 2020; Huang and Shi, 2019), raising the intriguing possibility that their roles in PD may be modulated or mediated by the gut microbiota. Further, there is evidence that children living in rural areas are more likely to have a microbiota that is linked to a lower risk of autoimmune diseases (Kirjavainen et al., 2019), the potential relevance of this finding to PD is yet to be investigated. Finally, a recent study (Mertsalmi et al., 2020) reported a higher PD risk in people with past

exposure to multiple courses of specific antibiotics treatments, in particular macrolides and lincosamides, antianaerobic, and broad-spectrum antibiotics, along with antifungal medications. Since these antibiotics have the highest impact on gut microbiota, it is likely that this finding was due to altered microbiota following the use of antibiotics (Mertsalmi et al., 2020). Interestingly, these associations were the strongest for exposures 10–15 years before PD diagnosis, consistent with the concept of a prolonged prodromal phase of PD.

Taken together, recent research not only supports a critical role of the gut microbiota in PD pathogenesis but also raises the possibility that they may modulate or mediate the effects of environmental risk factors for PD.

## 5. Research gaps and future directions

In the past several decades, research into the environmental causes of PD has significantly lagged behind genetic research, owing largely to the lack of understanding of the disease's natural history and its prolonged and complex prodromal development, as well as the lack of tools to adequately assess complex environmental exposures over an extended life period. The Braak and dual-hit hypotheses, albeit still controversial, have provided a timely theoretical framework to peek into the complexity of PD prodromal development (Braak et al., 2006; Hawkes et al., 2009). These hypotheses, coupled with empirical advances in understanding PD prodromal symptoms and markers, have opened an unprecedented window to define the environmental triggers and modifiers at various stages of PD development and to query novel etiological hypotheses of PD that have not been speculated before. These activities have been best illustrated by the preliminary inquiries of the etiological relevance of the gut-brain axis and gut microbiota in PD, as detailed above.

While promising, these lines of research are still in their infancy, with substantial research gaps and challenges ahead. First, the Braak hypothesis itself is still, to some extent, controversial (Lionnet et al., 2018). For example, neuropathological studies are yet to document standalone synucleinopathy in the gut without brain involvement (Beach et al., 2010), and the body-first vs. brain-first etiology of PD has emerged as an important topic of investigation (Borghammer et al., 2021). Researchers must be fully mindful of the etiological, pathological, and clinical complexity of PD development and presentations (Berg et al., 2021). Second, the potential roles of Lewy pathology and its spreading in PD pathogenesis and symptomatic presentation remain speculative. Although preliminary animal experimental research has provided provocative evidence supporting for the spreading of alpha-synuclein pathology induced by PFFs from olfactory and intestinal structures to the brain, but several aspects of the PFF model fall short. For example, the spreading in many instances does not seem progressive but instead occurs discretely among interconnected neurons, but over time the spread stops. This brings into question whether this spread is really a disease driving mechanism in PD prodromal and clinical progression (Killinger and Kordower, 2019). Further, PFF seeded Lewy pathology does not closely resemble bona fide Lewy pathology seen in the human PD brain. Many Lewy bodies in the human brain have exquisite halo structure of radiating filaments intermingled with damaged organelles, hinting at cellular regulated processes. These fine structural characteristics do not seem to be recapitulated with PFFs, or are not typically experimentally determined. Third, despite contributions from animal models of PD, it is challenging to have experimental models that recapitulate the scope of PD pathological and clinical endpoints in humans, including dopaminergic neuron loss, Lewy pathology, motor and nonmotor impairment as well as their sequential development. Fourth, roles of environmental toxins remain to be investigated in PD experimental models, at levels relevant to human health, via routes that mimic human exposure, and during various stages of prodromal development. This is particularly relevant to PD research as the disease is chronic and progressive, and the dual-hit hypothesis specifically demands experimental studies via

inhalation and gastrointestinal routes. Finally, humans are exposed to mixtures of chemicals, physical agents, microbial, and other environmental exposures throughout our lifetime, and new exposures emerge, yet PD research has been almost exclusively focused upon its known or suspected environmental causes (e.g., pesticides, metals, and air pollutants). Future research should capitalize on the new technology, such as large-scale screening to search for novel environmental chemicals for PD pathogenesis.

In human studies, it is also critical to target PD prodromal development, for example, by identifying triggers of PD pathogenesis and modifiers for its prodromal progression and phenotypical conversion (Chen and Ritz, 2018). Conducting such research is, however, very challenging. Although the Braak hypothesis provides a staging scheme of prodromal development, there are substantial knowledge gaps in identifying reliable markers and characterizing their sequential orders (if any) and phenotypical heterogeneity. Prodromal symptoms identified so far, except for PSG-confirmed RBD, lack specificity to PD (Chen et al., 2015). Research into the risk factors for RBD and factors that modify its phenotypical conversion is informative (Postuma et al., 2015; Postuma et al., 2012), but only about a quarter of PD patients have prodromal RBD (Chen et al., 2015). The International Movement Disorder Society recently updated its probability-based research criteria for prodromal PD, based on PD risk factors, prodromal nonmotor symptoms, subtle motor signs, neuroimaging, and other markers (Heinzel et al., 2019). It remains to see if whether and how this concept may lead to the identification and characterization of environmental contributions to prodromal PD.

Unlike the human genome, we do not have readily available tools to assess our environmental exposures, which are complex mixtures and consistently change throughout our lifetime (Patel et al., 2017). Further, it is yet to be studied whether the potential effects of exposures may be differential by exposure windows (e.g., prenatal, childhood, early, mid vs. late adulthood) (Gao et al., 2015) and stages of disease development (e.g., initiation vs. phenotypical conversion) (Postuma et al., 2015; Postuma et al., 2012). Finally, such effects are likely complex, cumulative, and interactive with other environmental and genetic factors, making them extremely difficult to assess (Chen and Ritz, 2018). Such complexity demands coordinated efforts from multiple large prospective cohorts with longitudinal and comprehensive assessments of environmental exposures, genetic, and other molecular signatures (e.g., epigenomic, metabolomic, microbiomic, and proteomic). Such studies should also collect and bank biospecimen (e.g., blood, feces, saliva, nasal mucosa) for future research needs (Marras et al., 2019). Principal investigators of these studies should work collaboratively to assess currently available resources, develop strategies to harmonize data and pool resources for PD prodromal research, and identify future research opportunities and strategies (Marras et al., 2019). This has been exemplified by the ever-growing international PD genetic research consortia (Aligning Science Across Parkinson's (ASAP) Initiative, 2021). Pooling cohorts for non-genetic research will be much more challenging and risky (Smith-Warner et al., 2006), but its success will be extremely rewarding by identifying modifiable causes of PD to prevent or delay the disease's clinical onset (Marras et al., 2019).

In conclusion, PD prodromal research represents an unprecedented opportunity for identifying the environmental triggers and effect modifiers of PD (Chen and Ritz, 2018). Such research, however, will be very challenging, calling for multidisciplinary efforts across biologists, epidemiologists, and clinical scientists to identify priorities, strategize approaches, and implement research activities (De Miranda et al., 2021). Further research should also take a life-long approach and leverage emerging tools to assess exposures and their biological effects to understand how PD starts and develops and how environmental factors contribute to these processes.

## Declaration of competing interest

Dr. Scheperjans is founder and CEO of NeuroInnovation Oy and NeuroBiome Ltd., is a member of the scientific advisory board and has received consulting fees and stock options from Axial Biotherapeutics. He has patents issued (FI127671B & US10139408B2) and pending (US16/186,663 & EP3149205) that are assigned to NeuroBiome Ltd. Other authors did not report and financial conflict of interest.

## Acknowledgements

This work is supported in part by the National Institute of Environmental Health Sciences (R01ES029227), the Office of the Assistant Secretary of Defense for Health Affairs through the Parkinson's Research Program (Award No. W81XWH-17-1-0536), and the Parkinson's Foundation (Grant No. PF-IMP-1825). Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the National Institutes of Health and the US Department of Defense. Dr. Scheperjans is supported by the Academy of Finland (295724, 310835, 342758), the Hospital District of Helsinki and Uusimaa (UAK1014004, UAK1014005, UAK1024MTM, TYH2018224, TYH2020335), the Olvi Foundation, the Konung Gustaf V:s och Drottning Victorias Frimurarestiftelse, and the Stockmann Foundation.

## References

- Abbott, R.D., et al., 2001. Frequency of bowel movements and the future risk of Parkinson's disease. *Neurology*. 57, 456–462.
- Adams, R.G., Crabtree, N., 1961. Anosmia in alkaline battery workers. *Br. J. Ind. Med.* 18, 216–221.
- Adams, D.R., et al., 2016. Nitrogen dioxide pollution exposure is associated with olfactory dysfunction in older U.S. adults. *Int. Forum Allerg. Rhinol.* 6, 1245–1252.
- Ahlström, R., et al., 1986. Impaired odor perception in tank cleaners. *Scand. J. Work Environ. Health* 12, 574–581.
- Ahman, M., et al., 2001. Nasal symptoms and pathophysiology in farmers. *Int. Arch. Occup. Environ. Health* 74, 279–284.
- Aho, V.T.E., et al., 2019. Gut microbiota in Parkinson's disease: temporal stability and relations to disease progression. *EBioMedicine*. 44, 691–707.
- Aho, V.T.E., et al., 2021. Relationships of gut microbiota, short-chain fatty acids, inflammation, and the gut barrier in Parkinson's disease. *Mol. Neurodegener.* 16, 6.
- Ajmani, G.S., et al., 2016. Fine particulate matter exposure and olfactory dysfunction among urban-dwelling older US adults. *Environ. Res.* 151, 797–803.
- Ajmani, G.S., et al., 2017. Smoking and olfactory dysfunction: a systematic literature review and meta-analysis. *Laryngoscope*. 127, 1753–1761.
- Aligning Science Across Parkinson's (ASAP) Initiative, 2021. Global Parkinson's Genetics Program. <https://www.michaeljfox.org/grant/global-parkinsons-genetics-program-gp2>.
- Amrosini, Y.M., et al., 2019. The gut-brain axis in neurodegenerative diseases and relevance of the canine model: a review. *Front. Aging Neurosci.* 11.
- Anderson, T., et al., 2021. Paraquat inhalation, a translationally relevant route of exposure: disposition to the brain and male-specific olfactory impairment in mice. *Toxicol. Sci.* 180, 175–185.
- Anselmi, L., et al., 2018. Ingestion of subthreshold doses of environmental toxins induces ascending Parkinsonism in the rat. *NPJ Parkinson's Dis.* 4, 30.
- Antunes, M.B., et al., 2007. San Francisco/Oakland Bay bridge welder study: olfactory function. *Neurology* 69, 1278–1284.
- Baelum, J., et al., 1982. Acute and subacute symptoms among workers in the printing industry. *Br. J. Ind. Med.* 39, 70–75.
- Beach, T.G., et al., 2010. Multi-organ distribution of phosphorylated alpha-synuclein histopathology in subjects with Lewy body disorders. *Acta Neuropathol.* 119, 689–702.
- Beach, T.G., et al., 2021. Vagus nerve and stomach synucleinopathy in Parkinson's disease, incidental Lewy body disease, and Normal elderly subjects: evidence against the "body-first" hypothesis. *J. Parkinsons Dis.* 11, 1833–1843.
- Berg, D., et al., 2021. Prodromal Parkinson disease subtypes - key to understanding heterogeneity. *Nat. Rev. Neurol.* 17, 349–361.
- Betarbet, R., et al., 2000. Chronic Systemic Pesticide Exposure Reproduces Features of Parkinson's Disease, vol. 3, pp. 1301–1306.
- Bhattarai, Y., et al., 2021. Role of gut microbiota in regulating gastrointestinal dysfunction and motor symptoms in a mouse model of Parkinson's disease. *Gut Microbes* 13, 1866974.
- Block, M.L., et al., 2012. The outdoor air pollution and brain health workshop. *Neurotoxicology*. 33, 972–984.
- Boertien, J.M., et al., 2019. Increasing comparability and utility of gut microbiome studies in Parkinson's disease: a systematic review. *J. Parkinsons Dis.* 9, S297–S312.
- Bolla, K.I., et al., 1995. Comparison of neurobehavioral function in workers exposed to a mixture of organic and inorganic lead and in workers exposed to solvents. *Am. J. Ind. Med.* 27, 231–246.

- Borghammer, P., 2021. The  $\alpha$ -Synuclein origin and connectome model (SOC model) of Parkinson's disease: explaining motor asymmetry, non-motor phenotypes, and cognitive decline. *J. Parkinsons Dis.* 11, 455–474.
- Borghammer, P., et al., 2021. Neuropathological evidence of body-first vs. brain-first Lewy body disease. *Neurobiol. Dis.* 161, 105557 <https://doi.org/10.1016/j.nbd.2021.105557>. Epub 2021 Nov 8.
- Braak, H., et al., 2006. Stanley Fahn lecture 2005: the staging procedure for the inclusion body pathology associated with sporadic Parkinson's disease reconsidered. *Mov. Disord.* 21, 2042–2051.
- Calderon-Garciduenas, L., et al., 2010. Urban air pollution: influences on olfactory function and pathology in exposed children and young adults. *Exp. Toxicol. Pathol.* 62, 91–102.
- Casjens, S., et al., 2017. Associations between former exposure to manganese and olfaction in an elderly population: results from the Heinz Nixdorf Recall Study. *Neurotoxicology.* 58, 58–65.
- Casjens, S., et al., 2018. Associations between blood lead, olfaction and fine-motor skills in elderly men: results from the Heinz Nixdorf Recall Study. *Neurotoxicology.* 68, 66–72.
- Cersosimo, M.G., 2018. Propagation of alpha-synuclein pathology from the olfactory bulb: possible role in the pathogenesis of dementia with Lewy bodies. *Cell Tissue Res.* 373, 233–243.
- Challis, C., et al., 2020. Gut-seeded alpha-synuclein fibrils promote gut dysfunction and brain pathology specifically in aged mice. *Nat. Neurosci.* 23, 327–336.
- Chen, H., 2018. The changing landscape of Parkinson epidemiologic research. *J. Parkinsons Dis.* 8, 1–12.
- Chen, H., Ritz, B., 2018. The search for environmental causes of Parkinson's disease: moving forward. *J. Parkinsons Dis.* 8, S9–S17.
- Chen, H., et al., 2010. Smoking duration, intensity, and risk of Parkinson disease. *Neurology* 74, 878–884.
- Chen, H., et al., 2015. Meta-analyses on prevalence of selected Parkinson's nonmotor symptoms before and after diagnosis. *Transl. Neurodegener.* 4, 1.
- Chen, S.G., et al., 2016. Exposure to the functional bacterial amyloid protein Curli enhances alpha-synuclein aggregation in aged Fischer 344 rats and *Caenorhabditis elegans*. *Sci. Rep.* 6, 34477.
- Chen, H., et al., 2017. Olfaction and incident Parkinson disease in US white and black older adults. *Neurology* 89, 1441–1447.
- Chesselet, M.F., et al., 2012. A progressive mouse model of Parkinson's disease: the Thy1-aSyn ("Line 61") mice. *Neurotherapeutics* 9, 297–314.
- Cirstea, M.S., et al., 2020. Microbiota composition and metabolism are associated with gut function in Parkinson's disease. *Mov. Disord.* 35, 1208–1217.
- Dardiotti, E., et al., 2018. H. Pylori and Parkinson's disease: meta-analyses including clinical severity. *Clin. Neurol. Neurosurg.* 175, 16–24.
- Darweesh, S.K.L., et al., 2018. Parkinson matters. *J. Parkinsons Dis.* 8, 495–498.
- De Miranda, B.R., et al., 2021. Preventing Parkinson's disease: an environmental agenda. *J. Parkinsons Dis.* <https://doi.org/10.3233/JPD-212922>. Online ahead of print.
- Doty, R.L., 2009. The olfactory system and its disorders. *Semin. Neurol.* 29, 74–81.
- Doty, R.L., 2015. Neurotoxic exposure and impairment of the chemical senses of taste and smell. *Handb. Clin. Neurol.* 131, 299–324.
- Elfil, M., et al., 2020. Implications of the gut microbiome in Parkinson's disease. *Mov. Disord.* 35, 921–933.
- Fereshtehnejad, S.M., et al., 2019. Evolution of prodromal Parkinson's disease and dementia with Lewy bodies: a prospective study. *Brain* 142, 2051–2067.
- Ferrante, R.J., et al., 1997. Systemic administration of rotenone produces selective damage in the striatum and globus pallidus, but not in the substantia nigra. *Brain Res.* 753, 157–162.
- Forsyth, C.B., et al., 2011. Increased intestinal permeability correlates with sigmoid mucosa alpha-synuclein staining and endotoxin exposure markers in early Parkinson's disease. *PLoS One* 6, e28032.
- Freundt, E.C., et al., 2012. Neuron-to-neuron transmission of  $\alpha$ -synuclein fibrils through axonal transport. *Ann. Neurol.* 72, 517–524.
- Gao, X., et al., 2011. A prospective study of bowel movement frequency and risk of Parkinson's disease. *Am. J. Epidemiol.* 174, 546–551.
- Gao, J., et al., 2015. Head injury, potential interaction with genes, and risk for Parkinson's disease. *Parkinsonism Relat. Disord.* 21, 292–296.
- GBD Parkinson's Disease Collaborators, 2018. Global, regional, and national burden of Parkinson's disease, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet Neurol.* 17, 939–953.
- Gianutos, G., et al., 1997. Accumulation of manganese in rat brain following intranasal administration. *Fundam. Appl. Toxicol.* 37, 102–105.
- Gobba, F., Abbacchini, C., 2012. Anosmia after exposure to a pyrethrin-based insecticide: a case report. *Int. J. Occup. Med. Environ. Health* 25, 506–512.
- González, S., et al., 2020. Long-term coffee consumption is associated with fecal microbial composition in humans. *Nutrients* 12, 1287.
- Grashow, R., et al., 2015. Cumulative lead exposure is associated with reduced olfactory recognition performance in elderly men: the normative aging study. *Neurotoxicology* 49, 158–164.
- Guarneros, M., et al., 2009. Mexico City air pollution adversely affects olfactory function and intranasal trigeminal sensitivity. *Chem. Senses* 34, 819–826.
- Guarneros, M., et al., 2020. Metal-containing particulate matter and associated reduced olfactory identification ability in children from an area of high atmospheric exposure in Mexico City. *Chem. Senses* 45, 45–58.
- Guehl, D., et al., 1999. Trichloroethylene and parkinsonism: a human and experimental observation. *Eur. J. Neurol.* 6, 609–611.
- Harischandra, D.S., et al., 2019. Manganese promotes the aggregation and prion-like cell-to-cell exosomal transmission of  $\alpha$ -synuclein. *Sci. Signal.* 12, eaau4543.
- Hawkes, C.H., et al., 2007. Parkinson's disease: a dual-hit hypothesis. *Neuropathol. Appl. Neurobiol.* 33, 599–614.
- Hawkes, C.H., et al., 2009. Parkinson's disease: the dual hit theory revisited. *Ann. N. Y. Acad. Sci.* 1170, 615–622.
- Heijtz, R.D., et al., 2021. Young microbiota rejuvenates the aging brain. *Nat. Aging* 1, 625–627.
- Heintz-Buschart, A., et al., 2018. The nasal and gut microbiome in Parkinson's disease and idiopathic rapid eye movement sleep behavior disorder. *Mov. Disord.* 33, 88–98.
- Heinzel, S., et al., 2019. Update of the MDS research criteria for prodromal Parkinson's disease. *Mov. Disord.* 34, 1464–1470.
- Heinzel, S., et al., 2021. Gut microbiome signatures of risk and prodromal markers of Parkinson disease. *Ann. Neurol.* 90 (3), E1–E12. <https://doi.org/10.1002/ana.26128>. Epub 2021 Jul 16.
- Henderson, M.X., et al., 2019. Spread of  $\alpha$ -synuclein pathology through the brain connectome is modulated by selective vulnerability and predicted by network analysis. *Nat. Neurosci.* 22, 1248–1257.
- Hernan, M.A., et al., 2004. A structural approach to selection bias. *Epidemiology* 15, 615–625.
- Holmqvist, S., et al., 2014. Direct evidence of Parkinson pathology spread from the gastrointestinal tract to the brain in rats. *Acta Neuropathol.* 128, 805–820.
- Holmström, M., et al., 1995. Upper airway symptoms and function in wood surface coating industry workers. *Am. J. Ind. Med.* 28, 207–220.
- Holmstrom, M., et al., 2008. Nasal complaints and signs of disease in farmers—a methodological study. *Acta Otolaryngol.* 128, 193–200.
- Huang, C., Shi, G., 2019. Smoking and microbiome in oral, airway, gut and some systemic diseases. *J. Transl. Med.* 17, 225.
- Huang, H.K., et al., 2018. Helicobacter pylori infection is associated with an increased risk of Parkinson's disease: a population-based retrospective cohort study. *Parkinsonism Relat. Disord.* 47, 26–31.
- Hudson, R., et al., 2006. Effect of air pollution on olfactory function in residents of Mexico City. *Chem. Senses* 31, 79–85.
- Jinsmaa, Y., et al., 2014. Divalent metal ions enhance DOPAL-induced oligomerization of alpha-synuclein. *Neurosci. Lett.* 569, 27–32.
- Jo, S., et al., 2021. Association of NO2 and other air pollution exposures with the risk of Parkinson disease. *JAMA Neurol.* 78, 800–808.
- Keane, P.C., et al., 2019. Trichloroethylene and its metabolite TaClo lead to degeneration of substantia nigra dopaminergic neurons: effects in wild type and human A30P mutant  $\alpha$ -synuclein mice. *Neurosci. Lett.* 711, 134437.
- Killinger, B.A., Kordower, J.H., 2019. Spreading of alpha-synuclein - relevant or epiphenomenon? *J. Neurochem.* 150, 605–611.
- Killinger, B., Labrie, V., 2019. The appendix in Parkinson's disease: from vestigial remnant to vital organ? *J. Parkinsons Dis.* 9, S345–S358.
- Killinger, B.A., et al., 2018. The vermiform appendix impacts the risk of developing Parkinson's disease. *Sci. Transl. Med.* 10.
- Kim, S., et al., 2019. Transneuronal propagation of pathologic  $\alpha$ -synuclein from the gut to the brain models Parkinson's disease. *Neuron* 103, 627–641.e7.
- Kirjavainen, P.V., et al., 2019. Farm-like indoor microbiota in non-farm homes protects children from asthma development. *Nat. Med.* 25, 1089–1095.
- Kishimoto, Y., et al., 2019. Chronic mild gut inflammation accelerates brain neuropathology and motor dysfunction in alpha-Synuclein mutant mice. *NeuroMolecular Med.* 21, 239–249.
- Kitamura, F., et al., 2003. Increase of olfactory threshold in plating factory workers exposed to chromium in Korea. *Ind. Health* 41, 279–285.
- Kumar, A., et al., 2016. Role of cytochrome c in alpha-synuclein radical formation: implications of alpha-synuclein in neuronal death in Maneb- and paraquat-induced model of Parkinson's disease. *Mol. Neurodegener.* 11, 70.
- Lee, S.-J., et al., 2018. Risk of olfactory dysfunction of the workers in the automobile repair, printing, shoemaking and plating industries in Korea: a cross-sectional study. *BMJ Open* 8, e022678.
- Li, Z., et al., 2021. Oral, nasal, and gut microbiota in Parkinson's disease. *Neuroscience* 480, 65–78.
- Lionnet, A., et al., 2018. Does Parkinson's disease start in the gut? *Acta Neuropathol.* 135, 1–12.
- Liu, Y.Z., et al., 1985. Effects of cadmium on cadmium smelter workers. *Scand. J. Work Environ. Health* 11 (Suppl. 4), 29–32.
- Liu, M., et al., 2010. Trichloroethylene induces dopaminergic neurodegeneration in fisher 344 rats. *J. Neurochem.* 112, 773–783.
- Liu, R., et al., 2016. Ambient air pollution exposures and risk of Parkinson disease. *Environ. Health Perspect.* 124, 1759–1765.
- Liu, B., et al., 2017a. Vagotomy and Parkinson disease: a Swedish register-based matched-cohort study. *Neurology* 88, 1996–2002.
- Liu, J., et al., 2017b. Sodium butyrate exerts protective effect against Parkinson's disease in mice via stimulation of glucagon like peptide-1. *J. Neurol. Sci.* 381, 176–181.
- Liu, B., et al., 2020. Appendectomy, tonsillectomy and Parkinson's disease risk: a Swedish register-based study. *Front. Neurol.* 11, 510.
- Lock, E.A., et al., 2013. Solvents and Parkinson disease: a systematic review of toxicological and epidemiological evidence. *Toxicol. Appl. Pharmacol.* 266, 345–355.
- Lucassen, E.B., et al., 2014. History of smoking and olfaction in Parkinson's disease. *Mov. Disord.* 29, 1069–1074.
- Lucchini, R., et al., 1997. Motor function, olfactory threshold, and hematological indices in manganese-exposed ferroalloy workers. *Environ. Res.* 73, 175–180.
- Maher, B.A., et al., 2016. Magnetite pollution nanoparticles in the human brain. *Proc. Natl. Acad. Sci. U. S. A.* 113, 10797–10801.
- Mahlknecht, P., et al., 2018. Reader response: olfaction and incident Parkinson disease in US white and black older adults. *Neurology* 90, 940.

- Manfredsson, F.P., et al., 2018. Induction of alpha-synuclein pathology in the enteric nervous system of the rat and non-human primate results in gastrointestinal dysmotility and transient CNS pathology. *Neurobiol. Dis.* 112, 106–118.
- Marras, C., et al., 2016. Appendectomy in mid and later life and risk of Parkinson's disease: a population-based study. *Mov. Disord.* 31, 1243–1247.
- Marras, C., et al., 2019. Environment, lifestyle, and Parkinson's disease: implications for prevention in the next decade. *Mov. Disord.* 34, 801–811.
- Mascagni, P., et al., 2003. Olfactory function in workers exposed to moderate airborne cadmium levels. *Neurotoxicology* 24, 717–724.
- Maturana, M.G., et al., 2015. Unveiling the role of the pesticides paraquat and rotenone on alpha-synuclein fibrillation in vitro. *Neurotoxicology* 46, 35–43.
- Mertsalmi, T.H., et al., 2020. Antibiotic exposure and risk of Parkinson's disease in Finland: a nationwide case-control study. *Mov. Disord.* 35, 431–442.
- Mezzaroba, L., et al., 2019. The role of zinc, copper, manganese and iron in neurodegenerative diseases. *Neurotoxicology* 74, 230–241.
- Mues, K.E., et al., 2017. Use of the Medicare database in epidemiologic and health services research: a valuable source of real-world evidence on the older and disabled populations in the US. *Clin. Epidemiol.* 9, 267–277.
- Naudet, N., et al., 2017. Oral exposure to Paraquat triggers earlier expression of phosphorylated  $\alpha$ -synuclein in the enteric nervous system of A53T mutant human  $\alpha$ -synuclein transgenic mice. *J. Neurochem. Exp. Neurol.* 76, 1046–1057.
- Ngwa, H.A., et al., 2014. Vanadium exposure induces olfactory dysfunction in an animal model of metal neurotoxicity. *Neurotoxicology* 43, 73–81.
- Nishiwaki, H., et al., 2020. Short-chain fatty acid-producing gut microbiota is decreased in Parkinson's disease but not in rapid-eye-movement sleep behavior disorder. *mSystems* 5.
- Okuzumi, A., et al., 2018. Rapid dissemination of alpha-synuclein seeds through neural circuits in an in-vivo prion-like seeding experiment. *Acta Neuropathol. Commun.* 6, 96.
- Paik, S.R., et al., 1999. Copper(II)-induced self-oligomerization of alpha-synuclein. *Biochem. J.* 340 (Pt 3), 821–828.
- Paiva, L., et al., 2017. Sodium butyrate rescues dopaminergic cells from alpha-synuclein-induced transcriptional deregulation and DNA damage. *Hum. Mol. Genet.* 26, 2231–2246.
- Palacios, N., et al., 2017. Air pollution and risk of Parkinson's disease in a large prospective study of men. *Environ. Health Perspect.* 125, 087011.
- Palacios, N., et al., 2018. Appendectomy and risk of Parkinson's disease in two large prospective cohorts of men and women. *Mov. Disord.* 33, 1492–1496.
- Palmquist, E., et al., 2020. A prospective study on risk factors for olfactory dysfunction in aging. *J. Gerontol. A Biol. Sci. Med. Sci.* 75, 603–610.
- Pan-Montojo, F., et al., 2010. Progression of Parkinson's disease pathology is reproduced by intragastric administration of rotenone in mice. *PLoS One* 5, e8762.
- Pan-Montojo, F., et al., 2012. Environmental toxins trigger PD-like progression via increased alpha-synuclein release from enteric neurons in mice. *Sci. Rep.* 2. <https://doi.org/10.1038/srep00898> (Epub 2012 Nov 30).
- Park, R.M., et al., 2005. Potential occupational risks for neurodegenerative diseases. *Am. J. Ind. Med.* 48, 63–77.
- Patel, C.J., et al., 2017. Opportunities and challenges for environmental exposure assessment in population-based studies. *Cancer Epidemiol. Biomark. Prev.* 26, 1370–1380.
- Pereira, P.A.B., et al., 2017. Oral and nasal microbiota in Parkinson's disease. *Parkinsonism Relat. Disord.* 38, 61–67.
- Perez-Pardo, P., et al., 2018. Gut bacterial composition in a mouse model of Parkinson's disease. *Benefic. Microbes* 9, 799–814.
- Perez-Pardo, P., et al., 2019. Role of TLR4 in the gut-brain axis in Parkinson's disease: a translational study from men to mice. *Gut.* 68, 829–843.
- Peter, I., et al., 2018. Anti-tumor necrosis factor therapy and incidence of Parkinson disease among patients with inflammatory bowel disease. *JAMA Neurol.* 75, 939–946.
- Postuma, R.B., et al., 2009. Quantifying the risk of neurodegenerative disease in idiopathic REM sleep behavior disorder. *Neurology* 72, 1296–1300.
- Postuma, R.B., et al., 2012. Environmental risk factors for REM sleep behavior disorder: a multicenter case-control study. *Neurology* 79, 428–434.
- Postuma, R.B., et al., 2015. Risk factors for neurodegeneration in idiopathic rapid eye movement sleep behavior disorder: a multicenter study. *Ann. Neurol.* 77, 830–839.
- Potts, C.L., 1965. Cadmium proteinuria—the health of battery workers exposed to cadmium oxide dust. *Ann. Occup. Hyg.* 8, 55–61.
- Prediger, R.D.S., et al., 2006. The risk is in the air: intranasal administration of MPTP to rats reproducing clinical features of Parkinson's disease. *Exp. Neurol.* 202, 391–403.
- Prediger, R.D.S., et al., 2009. Risk is in the air: an intranasal MPTP (1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine) rat model of Parkinson's disease. *Ann. N. Y. Acad. Sci.* 1170, 629–636.
- Prediger, R.D.S., et al., 2010. Single intranasal administration of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine in C57BL/6 mice models early preclinical phase of Parkinson's disease. *Neurotox. Res.* 17, 114–129.
- Prediger, R.D., et al., 2011. Intranasal administration of neurotoxins in animals: support for the olfactory vector hypothesis of Parkinson's disease. *Neurotox. Res.* 21, 90–116.
- Prigent, A., et al., 2019. Acute inflammation down-regulates alpha-synuclein expression in enteric neurons. *J. Neurochem.* 148, 746–760.
- Quandt, S.A., et al., 2016. Olfactory function in Latino farmworkers: subclinical neurological effects of pesticide exposure in a vulnerable population. *J. Occup. Environ. Med.* 58, 248–253.
- Quandt, S.A., et al., 2017. Olfactory function in Latino farmworkers over 2 years: longitudinal exploration of subclinical neurological effects of pesticide exposure. *J. Occup. Environ. Med.* 59, 1148–1152.
- Racette, B.A., et al., 2001. Welding-related parkinsonism: clinical features, treatment, and pathophysiology. *Neurology* 56, 8–13.
- Rey, N.L., et al., 2013. Transfer of human  $\alpha$ -synuclein from the olfactory bulb to interconnected brain regions in mice. *Acta Neuropathol.* 126, 555–573.
- Rey, N.L., et al., 2016. Widespread transneuronal propagation of alpha-synucleinopathy triggered in olfactory bulb mimics prodromal Parkinson's disease. *J. Exp. Med.* 213, 1759–1778.
- Rey, N.L., et al., 2018. The olfactory bulb as the entry site for prion-like propagation in neurodegenerative diseases. *Neurobiol. Dis.* 109, 226–248.
- Ritz, B., Rhodes, S.L., 2010. After half a century of research on smoking and PD, where do we go now? *Neurology* 74, 870–871.
- Ritz, B., et al., 2016. Traffic-related air pollution and Parkinson's disease in Denmark: a case-control study. *Environ. Health Perspect.* 124, 351–356.
- Rojo, A.I., et al., 2006. Persistent penetration of MPTP through the nasal route induces Parkinson's disease in mice. *Eur. J. Neurosci.* 24, 1874–1884.
- Rojo, A.I., et al., 2007. Chronic inhalation of rotenone or paraquat does not induce Parkinson's disease symptoms in mice or rats. *Exp. Neurol.* 208, 120–126.
- Rokad, D., et al., 2017. Role of neurotoxins and traumatic brain injury in alpha-synuclein protein misfolding and aggregation. *Brain Res. Bull.* 133, 60–70.
- Romano, S., et al., 2021. Meta-analysis of the Parkinson's disease gut microbiome suggests alterations linked to intestinal inflammation. *NPJ Parkinsons Dis.* 7, 27.
- Ross, G.W., et al., 2008. Association of olfactory dysfunction with risk for future Parkinson's disease. *Ann. Neurol.* 63, 167–173.
- Rozas, N.S., et al., 2021. Oral factors that impact the oral microbiota in Parkinson's disease. *Microorganisms* 9.
- Sampson, T.R., et al., 2016. Gut microbiota regulate motor deficits and Neuroinflammation in a model of Parkinson's disease. *Cell* 167, 1469–1480.e12.
- Sandmark, B., et al., 1989. Olfactory function in painters exposed to organic solvents. *Scand. J. Work Environ. Health* 15, 60–63.
- Sandoval-Motta, S., et al., 2017. Evolving ecosystems: inheritance and selection in the light of the microbiome. *Arch. Med. Res.* 48, 780–789.
- Sasajima, H., et al., 2015. Intranasal administration of rotenone in mice attenuated olfactory functions through the lesion of dopaminergic neurons in the olfactory bulb. *Neurotoxicology* 51, 106–115.
- Sasajima, H., et al., 2017. Intranasal administration of rotenone to mice induces dopaminergic neurite degeneration of dopaminergic neurons in the substantia nigra. *Biol. Pharm. Bull.* 40, 108–112.
- Savica, R., et al., 2009. Medical records documentation of constipation preceding Parkinson disease: a case-control study. *Neurology* 73, 1752–1758.
- Savica, R., et al., 2017. Time trends in the incidence of Parkinson's disease: a 30-year study. *JAMA Neurol.* 32, 227–234.
- Scheperjans, F., et al., 2018. The gut and Parkinson's disease: hype or hope? *J. Parkinsons Dis.* 8, S31–S39.
- Schubert, C.R., et al., 2021. Exposure to cadmium, lead, and tobacco smoke and the 10-year cumulative incidence of olfactory impairment: the beaver dam offspring study. *JAMA Otolaryngol. Head Neck Surg.* 147, 510–517.
- Schwartz, B.S., et al., 1989. Olfactory function in chemical workers exposed to acrylate and methacrylate vapors. *Am. J. Public Health* 79, 613–618.
- Schwartz, B.S., et al., 1990. Solvent-associated decrements in olfactory function in paint manufacturing workers. *Am. J. Ind. Med.* 18, 697–706.
- Schwartz, A., et al., 2018. Fecal markers of intestinal inflammation and intestinal permeability are elevated in Parkinson's disease. *Parkinsonism Relat. Disord.* 50, 104–107.
- Sengoku, R., et al., 2008. Incidence and extent of Lewy body-related alpha-synucleinopathy in aging human olfactory bulb. *J. Neurochem. Exp. Neurol.* 67, 1072–1083.
- Sharabi, Y., et al., 2021. Parkinson's disease outside the brain: targeting the autonomic nervous system. *Lancet Neurol.* 20, 868–876.
- Sharer, J.D., et al., 2015. Olfactory dysfunction in Parkinson's disease: positive effect of cigarette smoking. *Mov. Disord.* 30, 859–862.
- Shrestha, S., et al., 2018. Factors associated with dream enacting behaviors among US farmers. *Parkinsonism Relat. Disord.* 57, 9–15.
- Shrestha, S., et al., 2019. High pesticide exposure events and olfactory impairment among U.S. Farm. *Environ. Health Perspect.* 127, 17005.
- Shrestha, S., et al., 2021. Occupational pesticide use and self-reported olfactory impairment in US farmers. *Occup. Environ. Med.* 78, 179–191.
- Silva, Y.P., et al., 2020. The role of short-chain fatty acids from gut microbiota in gut-brain communication. *Front. Endocrinol. (Lausanne)* 11, 25.
- Smith-Warner, S.A., et al., 2006. Methods for pooling results of epidemiologic studies: the pooling project of prospective studies of diet and cancer. *Am. J. Epidemiol.* 163, 1053–1064.
- Sorokowska, A., et al., 2013. Olfaction and environment: 'Tsimane' of Bolivian rainforest have lower threshold of odor detection than industrialized German people. *PLoS One* 8, e69203.
- Sorokowska, A., et al., 2015. Determinants of human olfactory performance: a cross-cultural study. *Sci. Total Environ.* 506–507, 196–200.
- Sulkowski, W.J., et al., 2000. Smell impairment in workers occupationally exposed to cadmium. *Acta Otolaryngol.* 120, 316–318.
- Sunderman, F.W., 2001. Nasal toxicity, carcinogenicity, and olfactory uptake of metals. *Ann. Clin. Lab. Sci.* 31, 3–24.
- Svensson, E., et al., 2015. Vagotomy and subsequent risk of Parkinson's disease. *Ann. Neurol.* 78, 522–529.
- Toro, R., et al., 2019. Parkinson's disease and long-term exposure to outdoor air pollution: a matched case-control study in the Netherlands. *Environ. Int.* 129, 28–34.

- Toyoda, H., et al., 2020. Intranasal administration of rotenone reduces GABAergic inhibition in the mouse insular cortex leading to impairment of LTD and conditioned taste aversion memory. *Int. J. Mol. Sci.* 22, E259.
- Travagli, R.A., et al., 2020. Parkinson disease and the gut: new insights into pathogenesis and clinical relevance. *Nat. Rev. Gastroenterol. Hepatol.* 17, 673–685.
- Tysnes, O.B., et al., 2015. Does vagotomy reduce the risk of Parkinson's disease? *Ann. Neurol.* 78, 1011–1012.
- Ulusoy, A., et al., 2013. Caudo-rostral brain spreading of  $\alpha$ -synuclein through vagal connections. *EMBO Mol. Med.* 5, 1119–1127.
- Ulusoy, A., et al., 2017. Brain-to-stomach transfer of  $\alpha$ -synuclein via vagal preganglionic projections. *Acta Neuropathol.* 133, 381–393.
- Unger, M.M., et al., 2016. Short chain fatty acids and gut microbiota differ between patients with Parkinson's disease and age-matched controls. *Parkinsonism Relat. Disord.* 32, 66–72.
- Van Den Berge, N., et al., 2021. Ageing promotes pathological alpha-synuclein propagation and autonomic dysfunction in wild-type rats. *Brain* 144, 1853–1868.
- Voronkov, D.N., et al., 2017. Immunomorphological changes in the olfactory bulbs of rats after intranasal administration of rotenone. *Bull. Exp. Biol. Med.* 164, 203–206.
- Wakabayashi, K., et al., 1988. Parkinson's disease: the presence of Lewy bodies in Auerbach's and Meissner's plexuses. *Acta Neuropathol.* 76, 217–221.
- Willis, A.W., et al., 2010. Metal emissions and urban incident Parkinson disease: a community health study of Medicare beneficiaries by using geographic information systems. *Am. J. Epidemiol.* 172, 1357–1363.
- Wright, J.A., et al., 2009. Unique copper-induced oligomers mediate alpha-synuclein toxicity. *FASEB J.* 23, 2384–2393.
- Yu, I.T.-S., et al., 2004. Occupational exposure to mixtures of organic solvents increases the risk of neurological symptoms among printing workers in Hong Kong. *J. Occup. Environ. Med.* 46, 323–330.
- Zhang, Z., et al., 2021. Exposure to particulate matter air pollution and anosmia. *JAMA Netw. Open* 4, e2111606.
- Zhu, Y., et al., 2022. Association between inflammatory bowel diseases and Parkinson's disease: systematic review and meta-analysis. *Neural Regen. Res.* 17, 344–353.
- Zibrowski, E.M., Robertson, J.M., 2006. Olfactory sensitivity in medical laboratory workers occupationally exposed to organic solvent mixtures. *Occup. Med. (Oxford, England)* 56, 51–54.