

<https://helda.helsinki.fi>

Taxonomic bias in amphibian research : Are researchers responding to conservation need?

da Silva, Arthur F.

2020-08

da Silva , A F , Malhado , A C M , Correia , R A , Ladle , R J , Vital , M V C & Mott , T 2020 , ' Taxonomic bias in amphibian research : Are researchers responding to conservation need? ' , Journal for Nature Conservation , vol. 56 , 125829 . <https://doi.org/10.1016/j.jnc.2020.125829>

<http://hdl.handle.net/10138/343271>

<https://doi.org/10.1016/j.jnc.2020.125829>

cc_by_nc_nd

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

1 **Taxonomic bias in amphibian research:**
2 **Are researchers responding to conservation need?**

3

4 Authors: Arthur F. da Silva ^{a*}, Ana C. M. Malhado ^a, Ricardo A. Correia ^{a, b, c}, Richard
5 J. Ladle ^a, Marcos V. C. Vital ^a, Tamí Mott ^a

6

7 Affiliations:

8 a. Instituto de Ciências Biológicas e da Saúde, Universidade Federal de Alagoas,
9 Campus A. C. Simões, Avenida Lourival Melo Mota, Tabuleiro dos Martins, Maceió,
10 Alagoas, Brasil.

11 b. DBIO & CESAM - Centre for Environmental and Marine Studies, Universidade de
12 Aveiro, Campus Universitário de Santiago, Aveiro, Portugal.

13 c. Current address: Helsinki Lab of Interdisciplinary Conservation Science (HELICS),
14 Department of Geosciences and Geography, University of Helsinki, Finland.

15

16 *Corresponding author:

17 Arthur Filipe da Silva

18 Email: arthurfilipe.biologia@gmail.com

19 Phone number: +55 82 99687-7762.

20

21

22 Abstract:

23 Amphibians are very diverse, widely distributed, and the most endangered class of
24 vertebrates. As with other taxa, effective conservation of amphibians needs to be
25 supported by detailed scientific knowledge. However, species rich and broadly
26 distributed taxa are typically characterized by high variability in research effort. Our
27 objective was therefore to understand which factors (ecological and cultural) have led
28 some amphibian species to be more researched than others. We used two proxies of
29 research effort: i) the total number of articles on Web of Science (WoS) that mention
30 the scientific name (or synonyms) of each species, and; ii) the number of conservation
31 science articles on WoS that mention the scientific name (or synonyms) of each species.
32 These measures were used as dependent variables in zero hurdle regression models with
33 the aim of identifying the most important factors driving species-level knowledge
34 production. Well researched species (generally, and for conservation) tend to have a
35 longer history of scientific research, come from countries with high scientific capacity,
36 have large body size, and to be present in man-made habitats. Endangered species tend
37 to be less researched, generally and for conservation, possibly because they are often
38 more difficult to study: many endangered amphibians are restricted to small, fragmented
39 and remote habitats in countries with low scientific capacity. We conclude with a
40 discussion of how taxonomic biases in research effort on amphibians can be addressed
41 given the limited funds available for conservation research.

42 Keywords: Biodiversity, Conservation, Research effort, Scientific knowledge,
43 Bibliometrics.

44 **1. Introduction**

45 Amphibians are among the most endangered vertebrate groups (Ceballos et al., 2015;
46 Ripple et al., 2019). Several factors have been identified as responsible for amphibian
47 population die-offs across the world - including pollution, introduction of exotic species
48 and the infectious pathogens such as chytrids, ranaviruses, Perkinsea and trematodes -
49 with habitat loss identified as the most high-profile threat (Wake and Vredenburg, 2008;
50 Mann et al., 2009; Berger et al., 2016; DiRenzo and Grant, 2019; Scheele et al., 2019).
51 The way an amphibian responds to threats is linked to its biology, ecology and
52 evolution (Lips, 2016) and scientific knowledge about a species is therefore essential to
53 formulate effective conservation actions (Arlettaz et al., 2010; Canessa et al., 2019;
54 Lewis et al., 2019).

55 Despite the importance of scientific knowledge for conservation, many
56 amphibian species are very poorly known (Scheele et al., 2019). Indeed, the research
57 effort expended on different species is extremely patchy, with a few well studied species
58 and many species that are almost unknown to science (Clark and May, 2002; Murray et
59 al., 2015; Fleming and Bateman, 2016). The reasons for this patchiness are complex,
60 and may include geographical variation in the allocation of financial resources for
61 research, spatial and temporal variation in research capacity, and the intrinsic
62 characteristics of a species that makes it an ‘appropriate’ research target (Clark and
63 May, 2002). In this context, we hypothesise that species that are already well-known
64 scientifically (both generally and by a given individual or research group), of cultural
65 importance (e.g. threatened, invasive, economically important), and/or have traits that
66 make them convenient to study (e.g. large, conspicuous and diurnal) will be subject to
67 higher levels of research effort.

68 Here, we test the above hypothesis by: (i) quantifying research effort (both
69 general and conservation-related) for all extant amphibian species based on bibliometric
70 analysis, and; (ii) statistically identifying the main factors responsible for the observed
71 biases in the scientific knowledge production. In other words, we seek to understand
72 why some amphibian species are more researched than others and assess whether
73 conservation researchers are adequately responding to perceived conservation need.

74

75 **2. Methods**

76 *2.1. Global list of amphibian species*

77 We collected a list the names of all known extant amphibian species from the online
78 platform *Amphibian Species of the World*
79 (www.research.amnh.org/vz/herpetology/amphibia/). Name data was retrieved using the
80 *defrostR* package, within the R statistical environment, in February 2018. Our final
81 dataset included 7,668 species, distributed over three Orders (Anura: 6,752 species,
82 Caudata: 711 species, and Gymnophiona: 205 species). In addition to the currently
83 accepted scientific name for each amphibian species, we also retrieved all known
84 synonyms.

85

86 2.2. *Quantification of Scientific Knowledge Production*

87 Based on the assumption that more intensively studied species will be the subject of a
88 greater number of publications, we calculated as metric of research effort for every
89 amphibian species on our list the number of conservation-themed articles indexed in
90 WoS platform (www.webofknowledge.com) that mention its scientific name (or any of
91 its synonym) in the title, abstract or keywords. This metric was calculated by filtering
92 the search results to include on articles that appear in Journals in WoS's "Biodiversity
93 and Conservation" thematic area. We perform this filtering in order to rescue works that
94 have relevant implications for conservation.

95 Each amphibian species in our database was the subject of a unique search using
96 currently accepted scientific name of the species and any synonyms (e.g. "*Hylodes*
97 *gryllus*" OR "*Rana dorsalis*"). Including synonyms is an important strategy to
98 maximize data capture and to reduce biases caused by species that have undergone one
99 or more taxonomic revisions (Guala, 2016; Correia et al., 2018). Searches were
100 manually conducted between March 2018 and May 2018, and considered documents
101 registered between 1945 and 2018. We used the WoS' general search engine, that
102 consults all databases indexed to WoS.

103 Our metric of research effort is conservative in that it does not count all
104 potentially relevant articles. First, it excludes articles that only mention the common
105 name of a species in the title, abstract or keywords. Nevertheless, we considered that the
106 slight loss of data from excluding common names was outweighed by the reduced
107 biases and increased replicability of using scientific names and synonyms (Correia et
108 al., 2018). Second, it excludes articles where information on some species appears in the
109 main text of an article, but not in the title, abstract or keywords. Our metric therefore

110 only captures articles where the species was the focus or a major element of the
111 research, since this will typically result in a mention in the title, abstract or keywords.

112

113 2.3. Explanatory Variables

114 We considered a range of biocultural traits (explanatory variables) that may influence a
115 scientists' decision to study a particular amphibian species. While some of the factors
116 potentially affecting this decision (e.g. research funding) cannot be easily assessed for
117 the majority of species, many factors are quantifiable for most species. Specifically, the
118 following variables were considered for analysis:

- 119 (i) *Threat status*: researchers may be influenced by conservation need, with
120 more research effort being directed to highly threatened species. This
121 association should be most apparent for conservation research
122 production. Conversely, most threatened species have small populations
123 and restricted distributions, so may be less practical to study. The threat
124 status of each species was retrieved from the IUCN Red List
125 (www.iucnredlist.org). We excluded species that were classified as 'DD'
126 (*Data Deficient*), 'EX' (*Extinct*) and 'EW' (*Extinct in the Wild*), since, by
127 definition, for these species biological information is lacking or cannot
128 be studied. We placed the remaining species into three categories: 'LC'
129 (*Least Concern*), 'NT' (*Near Threatened*), and *Threatened*, which
130 included 'VU' (*Vulnerable*), 'EN' (*Endangered*) and 'CR' (*Critically*
131 *Endangered*);
- 132 (ii) *Research history* (based in the year of the first publication for each
133 species in the platform): we theorized that, due to the iterative nature of
134 scientific research, species that were the subject of previous research
135 would be more likely to be the target of future research;
- 136 (iii) *Scientific capacity*: based on the % contribution of range countries to
137 global publications in the environmental sciences (1996-2017), using
138 data from Scimago (www.scimago.com). We reasoned that species
139 present in countries with higher environmental science capacity would be
140 more likely to be studied, and for those studies to be published;
- 141 (iv) *Presence in anthropic environments*: we obtained from the IUCN Red
142 List website information on amphibian species that occur in man-made
143 habitats, both aquatic and terrestrial. Our prediction was that species

144 occurring in anthropogenic areas would be more researched because they
145 can often be found close to research centers;

146 (v) *Body size*: there is a large body of literature that suggests that larger
147 species of vertebrates generate more public interest (e.g. Frynta et al.,
148 2013; Correia et al., 2016; Roll et al., 2016), and may have more intrinsic
149 appeal to researchers. Larger species may also be easier to locate and
150 sample in the field, and may be more attractive for leveraging
151 conservation funding. We retrieved amphibians' body size information
152 (in millimeter) from AmphiBIO database (Oliveira et al., 2017).

153 After removing extinct/data deficient species and those with missing data points,
154 our final dataset used in the model contained 3,468 species.

155

156 2.4. Data Analysis

157 Because many species were not associated with even a single record in the Web of
158 Science, our response variables contained many zeros. To account for this fact, we used
159 a zero-inflated hurdle model. This model has two components: a hurdle component, that
160 takes into consideration the zero counts, and a truncated count component for positive
161 counts. To perform this analysis, we used the *pscl* R package. The variable 'research
162 history' was, necessarily, not included in the zero hurdle models. Since several
163 explanatory variables in our study may influence scientific research for certain
164 amphibian species, a single model will not be able to provide an accurate representation
165 of the current scenario. Therefore, we used a multi-model inference approach to
166 calculate the effect of each explanatory variable on scientific research (Burnham and
167 Anderson, 2004; Burnham et al., 2011). We evaluated all possible model combinations
168 taking into consideration the list of explanatory variables considered in this study, and
169 identified the set of most adequate models according to AIC corrected for small sample
170 size (AICc). We then carried out a model averaging process where using all models
171 which had a delta AIC of less than 5 in relation to the best model (i.e. that with the
172 lowest AICc score). All continuous explanatory variables were standardized by
173 subtracting the variable mean to each value and dividing it the variable standard
174 deviation before inclusion in the models. This approach allows a direct comparison of
175 the estimated effects of each variable on research effort (Schielzeth, 2010).

176

177 3. Results

178 From 3,468 amphibian species of our dataset, 334 species (310 anurans, 18 salamanders
179 and 6 caecilians) were not associated with any articles retrieved from WoS, from 1945
180 to February 2018. A total of 3,134 amphibian species and 209,098 articles were
181 retrieved. For 2,720 anuran species, 177,510 articles were registered. For 361
182 salamander species, 30,802 articles and for 53 caecilians, 786 records were obtained. In
183 a general scale, regarding to the distribution of number of articles, only 24 species had
184 more than 1,000 articles registered in the platform. Of these species, 9 had above 5,000
185 articles, and 5 above 10,000. Among the species that had less than 1,000 records, 13 had
186 between 500 and 950 articles, 42 between 200 and 490, and 70 had between 100 and
187 190. Thus, most of the species studied (95.7%) had below 100 WoS records (Figure 1).
188 The 10 most studied species were all classified as Least Concern the IUCN. African
189 clawed frog (*Xenopus laevis*) had the highest number of articles, with 46,021 documents
190 (Figure S1a). Among the 10 most studied endangered species for all areas, axolotl
191 (*Ambystoma mexicanum*) was the most studied, with 2,228 articles (Figure S1b). The
192 Iberian ribbed newt (*Pleurodeles waltl*) had 1,515 articles, and was the most researched
193 among species classified as Near Threatened.

194 Filtering searches for “Biodiversity and Conservation” thematic area, we
195 retrieved 18,824 articles about 2,214 species. We recovered a total of 1,926 anuran
196 species (14,873 articles), 264 salamander species (3,893 articles), and 24 caecilian
197 species (58 articles). The 10 most studied species were again all classified as Least
198 Concern. The common toad (*Bufo bufo*) was the most studied species, with 1,395
199 articles (Figure 2a). Among threatened species, the mountain yellow-legged frog (*Rana*
200 *muscosa*) had the greatest number (90) of conservation articles (Figure 2b). Of the
201 species classified as Near Threatened according to IUCN criteria, the hellbender
202 (*Cryptobranchus alleganiensis*) was the most studied (86 articles).

203 Our models revealed a very consistent pattern of associations between
204 biocultural traits and research effort (Figure 3). As predicted, larger amphibian species
205 that occur in countries with higher scientific capacity were more frequently the subjects
206 of research. Research volume also was significantly associated with species with a
207 longer history of research. Perhaps surprisingly, more threatened species were less
208 likely to be the subject of articles in conservation orientated journals.

209

210 **4. Discussion**

211 Most amphibians are not well studied: more than 95% of the amphibian species in our
212 database were associated with between zero and 100 articles. Threatened species were
213 more likely to be associated with no articles or a low volume of articles (Figure 3).
214 These results do not support the argument that the global extinction risk of a species is
215 an important driver of scientific research effort (Zhang et al., 2015; Jarić et al., 2019),
216 regarding amphibians. Thus, in general terms we can tentatively conclude that
217 conservation need is often outweighed by other, perhaps more practical, factors when
218 researchers are deciding which species would be the most appropriate subject of a
219 particular scientific study. One of these practical concerns could be the local
220 conservation need, e.g. a nationally threatened species that is not threatened at the
221 global scale considered by the IUCN. Another important concern are factors that might
222 increase or decrease the resources (financial and human) needed to successfully conduct
223 a field or lab-based research project. For example, easy access to a conveniently located
224 and abundant wild population will considerably reduce the resources needed for field-
225 based studies. Similarly, species that have characteristics that make them easier to
226 collect and observe (e.g. large body size, diurnal behaviour patterns) may have reduced
227 resource requirements. Conversely, species that are conservation priorities will tend to
228 be associated with increasing resource requirements for research since extinction risk is
229 a reflection of population decline and fragmentation, range reduction, and rarity
230 (Hartley and Kunin, 2003). Many endangered species are also endemic, and are
231 restricted to remote, poorly accessible regions (Howard et al., 2015; Xing et al., 2016)
232 that are unlikely to have good research infrastructure.

233 Resource requirements for scientific studies increase enormously when
234 researchers need to travel internationally, meaning that most field-based studies are
235 conducted in the researcher's country of residence. This explains why amphibian
236 species that are resident in countries with high capacity in environmental science are
237 more researched, both generally and also for conservation. Indeed, European and North
238 American anurans included some of the most studied species with conservation-related
239 focus. Financial resource restrictions on amphibian research may be particularly severe:
240 recent research suggests that amphibians, even if threatened, receive less investment for
241 conservation (Davies et al., 2018).

242 'Researchability' is also predicted to vary with how much scientific knowledge
243 already exists about a species (Engemann et al., 2015; dos Santos et al., 2020), since
244 science is an iterative process that constantly builds on the results of previous studies.

245 This is reflected in the positive association between years since first publication and
246 research volume. Researchers working on a poorly known endangered species may
247 therefore require a much greater research effort to generate data of sufficient interest
248 and novelty for an international journal. If such publications are a significant factor in
249 career advancement, this may lead to risk-averseness among conservation researchers
250 (Wilson et al., 2006) and their students. Indeed, Tim Caro recently observed a growing
251 tendency of graduate students studying animal behaviour to work on common species
252 that are considered, in some way, to be similar to a species of conservation concern
253 (Caro, 2017). Caro attributes this trend to the fact that rare species are “difficult to
254 locate and result in small sample sizes” (Caro, 2017) - presumably leading to studies
255 that are difficult to publish. In summary, our results broadly support the notion that
256 there may often be conflict between what needs to be studied (for conservation) and the
257 career aspirations of researchers.

258 Although endangered amphibian species in general have notably fewer articles
259 than non-endangered species, there are some interesting exceptions. The axolotl
260 (*Ambystoma mexicanum*), for example, is currently declining due to anthropic activities
261 (Ayala et al., 2018) but is well represented in the scientific literature. This is due to the
262 fact that the axolotl is commonly used as a model organism for development science
263 because of its high regenerative capacity (McCusker et al., 2016; Nowoshilow et al.,
264 2018). Moreover, some well-studied non-threatened species on our list may soon
265 become threatened. This may be the case for both the common toad (*Bufo bufo*) and
266 common frog (*Rana temporaria*). These species presented 1,395 and 831 articles related
267 to conservation, respectively, and 13,025 and 10,693 articles for all thematic areas.
268 Although widely distributed, classified as Least Concern and with stable trends in
269 IUCN, common toad populations have been suffering local declines due to pollution,
270 agricultural activities and road mortalities (Dmowski et al., 2015; Guillot et al., 2016;
271 Salazar et al., 2016; Kaczmarek et al., 2016). In addition, this species is victim to
272 *Bufoviridis* *herpesvirus 1*, a severe dermatitis which has caused mortality of these
273 organisms in Switzerland (Origgi et al., 2018). Likewise, the common frog, though
274 relatively abundant in Europe, is susceptible to *Ranavirus* and *Batrachochytrium*
275 *dendrobatidis*, that have already been implicated in the extinction of several amphibian
276 species (Bayley et al., 2013; Price et al., 2015). As pointed out by Petrovan and Schmidt
277 (2016), common toads and common frogs have suffered considerable declines in the
278 United Kingdom and Switzerland, even though they are widespread species. These

279 authors highlight the need for more research into common amphibian abundance trends
280 rather than focusing only on the most endangered species, as the decline of common
281 species can drastically affect ecosystem functions. This fact may reflect the reason why
282 our research has presented a larger number of articles for these and others widespread
283 and non-threatened species according the IUCN, thus perhaps demonstrating an interest
284 of researchers in a threat level locally experienced by the species.

285 The American bullfrog (*Lithobates catesbeianus*) was the second most
286 researched species, possibly reflecting its commercial importance as a food species for
287 human consumption and its use as a biological control agent (Dias et al., 2009;
288 Mendoza et al., 2012). This species is also invasive, having been introduced into many
289 regions around the globe (Silva et al., 2009; Mikula, 2015). Similarly, the Japanese
290 wrinkled frog (*Glandirana rugosa*), which was also highly targeted by researchers, was
291 introduced on the Hawaii Island as a biological control of pests, presenting an impact on
292 the local fauna, specially to endemic organisms (Kleeck and Holland, 2018). Something
293 similar happened with the cane toad (*Rhinella marina*), a highly invasive species,
294 causing many native organisms to decline (Griffiths and McKay, 2007; Tingley and
295 Shine, 2011; Ward-Fear et al., 2016). These cases demonstrate that even though these
296 species are not considered threatened, studying them can contribute positively to
297 conservation.

298 Iberian ribbed newt (*Pleurodeles waltl*), despite having shown a low number of
299 works (34 articles), was among the most researched species in the ‘Near Threatened’
300 category. This species endemic at Iberian Peninsula and Morocco (Beukema et al.,
301 2013) presented a significant decline highly due to the habitat loss, invasive species and
302 mortality on the roads (Montori et al., 2002). These aspects can make *P. waltl* attractive
303 for conservation research, although the fact that it is an endemic and declining species
304 can make it less accessible. The mountain yellow-legged frog (*Rana muscosa*) was the
305 most studied threatened species, although it was only associated with 90 documents in
306 our database. In comparison, the common toad (*Bufo bufo*), which was the most studied
307 species for conservation production, had 1,395 articles in that area. The (relatively) high
308 conservation output for *R. muscosa* can be explained by its presence in a high scientific
309 capacity country (the USA), even though it is physically small and is restricted to the
310 state of California. Its populations have declined rapidly in recent decades due to a
311 combination of predation by introduced fish species, exposure to pesticides and
312 chytridiomycosis infection (Rachowicz and Briggs, 2007; Sparling et al., 2015; Poorten

313 et al., 2017). Despite these factors that have led *R. muscosa* to the threatened level,
314 research on this species combined with practical conservation actions has favoured its
315 population increase. One example of such actions is the removal of introduced non-
316 native fish species, which has enable the recovery of anuran populations of this and
317 other species (Knapp et al., 2016; Poorten et al., 2017). Furthermore, scientific research
318 on these organisms can yield valuable results in several aspects. Studying their
319 abundance, for example, has allowed to detect changes in the abundance of species that
320 are affected by several life stages of these frogs, such as aquatic macroinvertebrates. In
321 addition, because it occurs in widely protected habitats, i.e., unaltered by development,
322 *R. muscosa* becomes ideal as a model of study on amphibian decline due to causes that
323 are not related to habitat loss. *Rana muscosa* was the first anuran species found to host
324 anti-Bd bacteria on the skin, thus contributing to the control of Bd (*Batrachochytrium*
325 *dendrobatidis*) outbreaks in persistent populations, and encouraging research into this
326 innate immunity mechanism in other anuran species (Reinke et al., 2019).

327 In general, our findings were able to present an overview of the current scenario
328 of the research effort directions for amphibians. However, from our discoveries it is also
329 possible to identify others taxa which have ecological and/or evolutionary traits similar
330 to amphibians and that may present resembling patterns of research effort. Similarly, it
331 is also possible to investigate whether the research effort for these others taxa would
332 follow different patterns from those of amphibians, and how this would relate to their
333 threat levels and conservation efforts. In addition, Davies et al. (2018) pointed out that
334 public interest in endangered species of birds and mammals has motivated
335 conservationist investments. On the other hand, threatened species of amphibians,
336 reptiles and fishes, which are comparatively less known to the public, receive smaller
337 conservation investments (Davies et al., 2018). Therefore, identifying potentially
338 emblematic amphibian species from our outcomes, and promoting them in conservation
339 programs can contribute to the preservation of both the amphibian community and other
340 biological groups.

341

342 **5. Conclusions**

343 As a taxonomic group, Amphibians are among the most threatened vertebrates on Earth
344 due to the impact of man-made climate change, habitat loss and fragmentation,
345 pollution and emerging diseases (Sodhi et al., 2008; Collins et al., 2009). Conserving
346 the world's amphibian species in the face of these threats requires: i) robust scientific

347 knowledge, and; ii) organizations and individuals with the capacity to use this
348 knowledge to mount effective conservation interventions: so-called ‘evidence-based
349 conservation’ (Sutherland et al., 2004). Our study demonstrates that one of the barriers
350 to evidence-based conservation of amphibians is the lack of knowledge about many
351 species, especially those identified as being at risk of extinction. However, although
352 scientific knowledge is essential, by itself it is not a sufficient measure for a species to
353 be conserved. In this context, an adequate communication between research and public
354 actions is highly necessary for efficient conservation strategies may be perform
355 (Arlettaz et al., 2010; Canessa et al., 2019; Grant et al., 2019; Lewis et al., 2019). For
356 this to occur, it is essential that public initiatives consider the generating causes of the
357 decline of species, as climate change, which are responsible for several losses of
358 amphibians (Winter et al., 2016). Nevertheless, our analysis also suggests some possible
359 strategies to reduce the biases in research effort. Firstly, dedicated research funding
360 streams targeted at endangered species may be effective at counter-balancing the
361 advantages of working on more abundant species. In addition, as indicated by Winter et
362 al. (2016), scientists should also focus on those under-represented species. In this
363 context, the EDGE of Existence Programme, of Zoological Society of London, which
364 aims to awareness and raise funds to conserve unique and threatened species, is an
365 important example of an initiative that can motivate research on such species. Secondly,
366 there is enormous scope for increasing international collaboration for research on
367 endangered amphibians, with the aim of reducing the negative impact of low
368 environmental science capacity in some developing countries. Finally, editors and
369 reviewers for conservation journals could adopt a more critical attitude to studies that
370 use abundant species as proxies for ecologically similar endangered species,
371 foregrounding the value of research on rarely studied amphibians where the
372 conservation need is the greatest.

373

374 **Acknowledgements**

375 AFS would like to thank the Coordenação de Aperfeiçoamento de Pessoal de Nível
376 Superior (CAPES) for a Master’s scholarship. AM, RL and TM are funded by the
377 Brazilian National Council for Scientific and Technological Development CNPq
378 fellowships – (#309980/2018-6 - AM; #309879/2019-1 - RL; 309904/2015-3 and
379 312291/2018-3 - TM). RAC is currently funded by a post-doctoral grant from the
380 Helsinki Institute for Sustainability Science (HELSUS) and the University of Helsinki.

381 We would also like to thank the board of examiners of Master's program Adrian Garda,
382 Tereza Thomé, Guilherme Ferreira and Robson Santos, for their valuable suggestions.
383 We also acknowledge Janisson Santos, from the 21st Century Conservation Lab
384 (Federal University of Alagoas), for suggestions with data collection. Finally, we also
385 thank two anonymous reviewers for their constructive comments.

386

387 **References**

388 Arlettaz, R., Schaub, M., Fournier, J., Reichlin, T.S., Sierro, A., Watson, J.E.M.,
389 Braunisch, V., 2010. From Publications to Public Actions: When Conservation
390 Biologists Bridge the Gap between Research and Implementation. *BioScience*, 60, 835-
391 842. <https://dx.doi.org/10.1525/bio.2010.60.10.10>.

392

393 Ayala, C., Ramos, A.G., Merlo, Á., Zambrano, L., 2018. Microhabitat selection of
394 axolotls, *Ambystoma mexicanum*, in artificial and natural aquatic systems.
395 *Hydrobiologia*, 828, 11-20. <https://doi.org/10.1007/s10750-018-3792-8>.

396

397 Bayley, A.E., Hill, B.J., Feist, S.W., 2013. Susceptibility of the European common frog
398 *Rana temporaria* to a panel of ranavirus isolates from fish and amphibian hosts.
399 *Diseases of Aquatic Organisms*, 103, 171-183. <https://doi.org/10.3354/dao02574>.

400

401 Berger, L., Roberts, A.A., Voyles, J., Longcore, J.E., Murray, K.A., Skerratt, L.F.,
402 2016. History and recent progress on chytridiomycosis in amphibians. *Fungal Ecology*,
403 19, 89-99. <http://dx.doi.org/10.1016/j.funeco.2015.09.007>.

404

405 Beukema, W., Pous, P., Donaire-Barroso, D., Bogaertes, S., Garcia-Porta, J., Escoriza,
406 D., Arribas, O.J., Mouden, E.H., Carranza, S., 2013. Review of the systematics,
407 distribution, biogeography and natural history of Moroccan amphibians. *Zootaxa*, 3661,
408 001-060. <http://dx.doi.org/10.11646/zootaxa.3661.1.1>.

409

410 Burnham, K.P., Anderson, D.R., 2004. Multimodel inference: understanding AIC and
411 BIC in model selection. *Sociological Methods & Research*, 33, 261-304.
412 <https://doi.org/10.1177/0049124104268644>.

413

- 414 Burnham, K.P., Anderson, D.R., Huyvaert, K.P., 2011. AIC model selection and
415 multimodel inference in behavioral ecology: some background, observations, and
416 comparisons. *Behavioral Ecology and Sociobiology*, 65, 23-35.
417 <https://doi.org/10.1007/s00265-010-1029-6>.
418
- 419 Canessa, S., Spitzen-van der Sluijs, A., Martel, A., Pasmans, F., 2019. Mitigation of
420 amphibian disease requires a stronger connection between research and management.
421 *Biological Conservation*, 236, 236-242. <http://dx.doi.org/10.1016/j.biocon.2019.05.030>.
422
- 423 Caro, T., 2007. Behavior and conservation: a bridge too far? *Trends in Ecology &*
424 *Evolution*, 22, 394-400. <https://doi.org/10.1016/j.tree.2007.06.003>.
425
- 426 Ceballos, G., Ehrlich, P.R., Barnosky, A.D., García, A., Pringle, R.M., Palmer, T.M.,
427 2015. Accelerated modern human-induced species losses: Entering the sixth mass
428 extinction. *Science Advances*, 1, e1400253. <https://doi.org/10.1126/sciadv.1400253>.
429
- 430 Clark, J.A., May, R.M., 2002. Taxonomic bias in conservation research. *Science*, 297,
431 191-192. <https://doi.org/10.1126/science.297.5579.191b>.
432
- 433 Collins, J.P., Crump, M.L., Lovejoy III, T.E., 2009. *Extinction in our times: global*
434 *amphibian decline*, Oxford University Press, New York.
435
- 436 Correia, R.A., Jepson, P., Malhado, A.C.M., Ladle, R.J., 2016. Familiarity breeds
437 content: assessing bird species popularity with culturomics. *PeerJ*, 4, e1728.
438 <https://doi.org/10.7717/peerj.1728>.
439
- 440 Correia, R.A., Jarić, I., Jepson, P., Malhado, A.C., Alves, J.A., Ladle, R.J., 2018.
441 Nomenclature instability in species culturomic assessments: Why synonyms matter.
442 *Ecological Indicators*, 90, 74-78. <https://doi.org/10.1016/j.ecolind.2018.02.059>.
443
- 444 Davies, T., Cowley, A., Bennie, J., Leyshon, C., Inger, R., Carter, H., Robinson, B.,
445 Duffy, J., Casalegno, S., Lambert, G., Gaston, K., 2018. Popular interest in vertebrates
446 does not reflect extinction risk and is associated with bias in conservation investment.
447 *PLOS ONE*, 13, e0203694. <https://doi.org/10.1371/journal.pone.0203694>.

448

449 Dias, D.D.C., De Stéfani, M.V., Ferreira, C.M., França, F.M., Ranzani-Paiva, M.J.T.,
450 Santos, A.A., 2009. Haematologic and immunologic parameters of bullfrogs, *Lithobates*
451 *catesbeianus*, fed probiotics. *Aquaculture Research*, 41, 1064-1071. 10.1111/j.1365-
452 2109.2009.02390.x.

453

454 DiRenzo, G.V., Grant, E.H.C., 2019. Overview of emerging amphibian pathogens and
455 modeling advances for conservation-related decisions. *Biological Conservation*, 236,
456 474-483. <https://doi.org/10.1016/j.biocon.2019.05.034>.

457

458 Dmowski, K., Rossa, M., Kowalska, J., Krasnodebska-Ostrega, B., 2015. Thallium in
459 spawn, juveniles, and adult common toads (*Bufo bufo*) living in the vicinity of a zinc-
460 mining complex, Poland. *Environmental Monitoring and Assessment*, 187, 4141.
461 10.1007/s10661-014-4141-7.

462

463 dos Santos, J.W., Correia, R.A., Malhado, A.C.M., Campos-Silva, J.V., Teles, D.,
464 Jepson, P., Ladle, R.J., *in press*. Drivers of taxonomic bias in conservation research: A
465 global analysis of terrestrial mammals. *Animal Conservation*.
466 <https://doi.org/10.1111/acv.12586>.

467

468 Engemann, K., Enquist, B.J., Sandel, B., Boyle, B., Jørgensen, P.M., Morueta-Holme,
469 N., et al., 2015. Limited sampling hampers “big data” estimation of species richness in a
470 tropical biodiversity hotspot. *Ecology and Evolution*, 5, 807-20.
471 <https://doi.org/10.1002/ece3.1405>.

472

473 Fleming, P.A., Bateman, P.W., 2016. The good, the bad, and the ugly: which Australian
474 terrestrial mammal species attract most research? *Mammal Review*, 46, 241-254.
475 <https://dx.doi.org/10.1111/mam.12066>.

476

477 Frost, D.R., 2018. *Amphibian Species of the World: An Online Reference*. Version 6.0
478 (February 2018). Electronic Database accessible at
479 <http://research.amnh.org/herpetology/amphibia/index.html>. American Museum of
480 Natural History, New York, USA.

481

- 482 Frynta, D., Šimková, O., Lišková, S., Landová, E., 2013. Mammalian Collection on
483 Noah's Ark: The Effects of Beauty, Brain and Body Size. PLOS ONE, 8, e63110.
484 <https://doi.org/10.1371/journal.pone.0063110>.
485
- 486 Guala, G.F., 2016. The importance of species name synonyms in literature searches.
487 PLOS ONE, 11, e0162648. <https://doi.org/10.1371/journal.pone.0162648>.
488
- 489 Guillot, H., Boissinot, A., Angelier, F., Lourdais, O., Bonnet, X., Brischoux, F., 2016.
490 Landscape influences the morphology of male common toads (*Bufo bufo*). Agriculture,
491 Ecosystems & Environment, 233, 106-110.
492 <http://dx.doi.org/10.1016/j.agee.2016.08.032>.
493
- 494 Grant, E.H.C., Muths, E., Schmidt, B.R., Petrovan, S.O., 2019. Amphibian conservation
495 in the Anthropocene. Biological Conservation, 236, 543-547.
496 <https://doi.org/10.1016/j.biocon.2019.03.003>.
497
- 498 Griffiths, A.D., McKay, J.L., 2007. Cane toads reduce the abundance and site
499 occupancy of Merten's water monitor (*Varanus mertensi*). Wildlife Research, 34, 609-
500 615. <https://doi.org/10.1071/WR07024>.
501
- 502 Hartley, S., Kunin, W.E., 2003. Scale dependency of rarity, extinction risk, and
503 conservation priority. Conservation Biology, 17, 1559-1570.
504 <https://doi.org/10.1111/j.1523-1739.2003.00015.x>.
505
- 506 Howard, J.K., Klausmeyer, K.R., Fesenmyer, K.A., Furnish, J., Gardali, T., Grantham,
507 T., et al., 2015. Patterns of Freshwater Species Richness, Endemism, and Vulnerability
508 in California. PLOS ONE, 10, e0130710. <https://doi.org/10.1371/journal.pone.0130710>.
509
- 510 Jarić, I., Correia, R.A., Roberts, D.L., Gessner, J., Meinard, Y., Courchamp, F., 2019.
511 On the overlap between scientific and societal taxonomic attentions - Insights for
512 conservation. Science of the Total Environment, 648, 772-778.
513 <https://doi.org/10.1016/j.scitotenv.2018.08.198>.
514

- 515 Kaczmarek, M., Kolenda, K., Rozenblut-Kościsty, B., Sośnicka, W., 2016. Phalangeal
516 bone anomalies in the European common toad *Bufo bufo* from polluted environments.
517 Environmental Science and Pollution Research, 23, 21940-21946. 10.1007/s11356-016-
518 7297-6.
- 519
- 520 Kleeck, M.J.V., Holland, B.S., 2018. Gut Check: Predatory Ecology of the Invasive
521 Wrinkled Frog (*Glandirana rugosa*) in Hawai'i. Pacific Science, 72, 199-208.
522 <https://doi.org/10.2984/72.2.2>.
- 523
- 524 Knapp, R.A., Fellers, G.M., Kleeman, P.M., Miller, D.A.W., Vredenburg, V.T.,
525 Rosenblum, E.B., Briggs, C.J., 2016. Large-scale recovery of an endangered amphibian
526 despite ongoing exposure to multiple stressors. Proceedings of the National Academy of
527 Sciences of the United States of America, 113, 11889-11894.
528 <https://doi.org/10.1073/pnas.1600983113>.
- 529
- 530 Lewis, C.H., Richards-Zawacki, C.L., Ibáñez, R., Luedtke, J., Voyles, J., Houser, P.,
531 Gratwicke, B., 2019. Conserving Panamanian harlequin frogs by integrating captive-
532 breeding and research programs. Biological Conservation, 236, 180-187.
533 <https://doi.org/10.1016/j.biocon.2019.05.029>.
- 534
- 535 Lips, K.R., 2016. Overview of chytrid emergence and impacts on amphibians.
536 Philosophical Transactions of the Royal Society B: Biological Sciences, 371, 20150465.
537 <http://dx.doi.org/10.1098/rstb.2015.0465>.
- 538
- 539 Mann, R.M., Hyne, R.V., Choung, C.B., Wilson, S.P., 2009. Amphibians and
540 agricultural chemicals: Review of the risks in a complex environment. Environmental
541 Pollution, 157, 2903-2927. <https://doi.org/10.1016/j.envpol.2009.05.015>.
- 542
- 543 McCusker, C.D., Diaz-Castillo, C., Sosnik, J., Phan, A., Gardiner, D.M., 2016.
544 Histological image data of limb skeletal tissue from larval and adult *Ambystoma*
545 *mexicanum*. Data in Brief. 8, 1206-1208. <https://doi.org/10.1016/j.dib.2016.07.028>.
- 546

- 547 Mendoza, G.M., Pasteris, S.E., Ale, C.E., Otero, M.C., Bühler, M.I., Nader-Macías,
548 M.E.F., 2012. Cultivable microbiota of *Lithobates catesbeianus* and advances in the
549 selection of lactic acid bacteria as biological control agents in raniculture. Research in
550 Veterinary Science, 93, 1160-1167. <https://doi.org/10.1016/j.rvsc.2012.05.007>.
- 551
- 552 Mikula, P., 2015. Fish and amphibians as bat predators. European Journal of Ecology,
553 1, 71-80. <https://doi.org/10.1515/eje-2015-0010>.
- 554
- 555 Montori, A., Llórente, G.A., Santos, X., Carretero, M.A., 2002. *Pleurodeles waltl*
556 Michahelles, 1830. Gallipato. In J. M. Pleguezuelos, R. Márquez, & M. Lizana (Eds),
557 Atlas y Libro Rojo de los Anfibios y Reptiles de España (pp. 51-53). Madrid: Dirección
558 General de Conservación de la Naturaleza-Asociación Herpetológica Española.
- 559
- 560 Murray, H.J., Green, E.J., Williams, D.R., Burfield, I.J., Brooke, M.D.L., 2015. Is
561 research effort associated with the conservation status of European bird species?
562 Endangered Species Research, 27, 193-206. <https://doi.org/10.3354/esr00656>.
- 563
- 564 Nowoshilow, S., Schloissnig, S., Fei, J., Dahl, A., Pang, A.W.C., Pippel, M., et al.,
565 2018. The axolotl genome and the evolution of key tissue formation regulators. Nature.
566 554, 50-65. <https://doi.org/10.1038/nature25458>.
- 567
- 568 Oliveira, B.F., São-Pedro, V.A., Santos-Barrera, G., Penone, C., Costa, G.C., 2017.
569 AmphiBIO, a global database for amphibian ecological traits. Scientific Data, 4,
570 170123. <https://doi.org/10.1038/sdata.2017.123>.
- 571
- 572 Origgi, F.C., Schmidt, B.R., Lohmann, P., Otten, P., Meier, R.K., Pisano, S.R., et al.,
573 2018. *Bufo* *herpesvirus 1* (BfHV1) associated dermatitis and mortality in free
574 ranging common toads (*Bufo bufo*) in Switzerland. Scientific Reports, 8, 14737.
575 <https://doi.org/10.1038/s41598-018-32841-0>.
- 576
- 577 Petrovan, S.O., Schmidt, B.R., 2016. Volunteer Conservation Action Data Reveals
578 Large-Scale and Long-Term Negative Population Trends of a Widespread Amphibian,
579 the Common Toad (*Bufo bufo*). PLOS ONE, 11, e0161943.
580 <https://doi.org/10.1371/journal.pone.0161943>.

581

582 Poorten, T.J., Knapp, R.A., Rosenblum, E.B., 2017. Population genetic structure of the
583 endangered Sierra Nevada yellow-legged frog (*Rana sierrae*) in Yosemite National Park
584 based on multi-locus nuclear data from swab samples. *Conservation Genetics*, 18, 731-
585 744. <https://doi.org/10.1007/s10592-016-0923-5>.

586

587 Price, S.J., Garner, T.W., Balloux, F., Ruis, C., Paszkiewicz, K.H., Moore, K., Griffiths,
588 A.G., 2015. A de novo assembly of the common frog (*Rana temporaria*) transcriptome
589 and comparison of transcription following exposure to *Ranavirus* and *Batrachochytrium*
590 *dendrobatidis*. *PLOS ONE*, 10, e0130500.
591 <https://doi.org/10.1371/journal.pone.0130500>.

592

593 Rachowicz, L.J., Briggs, C.J., 2007. Quantifying the disease transmission function:
594 effects of density on *Batrachochytrium dendrobatidis* transmission in the mountain
595 yellow-legged frog *Rana muscosa*. *Journal of Animal Ecology*, 76, 711-721.
596 <https://doi.org/10.1111/j.1365-2656.2007.01256.x>.

597

598 R Core Team, 2020. R: A language and environment for statistical computing. R
599 Foundation for Statistical Computing, Vienna, Austria (<https://www.R-project.org/>).

600

601 Reinke, B.A., Miller, D.A.W., Janzen, F.J. 2019. What Have Long-Term Field Studies
602 Taught Us About Population Dynamics? *Annual Review of Ecology, Evolution, and*
603 *Systematics*, 50, 1-18. <https://doi.org/10.1146/annurev-ecolsys-110218-024717>.

604

605 Ripple, W.J., Wolf, C., Newsome, T.M., Betts, M.G., Ceballos, G., Courchamp, F.,
606 Hayward, M.W., Valkenburgh, B.V., Wallach, A.D., Worm, B., 2019. Are we eating
607 the world's megafauna to extinction? *Conservation Letters*, 12, e12627.
608 [10.1111/conl.12627](https://doi.org/10.1111/conl.12627).

609

610 Roll, U., Mittermeier, J.C., Diaz, G.I., Novosolov, M., Feldman, A., Itescu, Y., et al.,
611 2016. Using Wikipedia page views to explore the cultural importance of global reptiles.
612 *Biological Conservation*, 204, 42-50. <https://doi.org/10.1016/j.biocon.2016.03.037>.

613

- 614 Salazar, R.D., Montgomery, R.A., Thresher, S.E., Macdonald, D.W., 2016. Mapping the
615 Relative Probability of Common Toad Occurrence in Terrestrial Lowland Farm Habitat
616 in the United Kingdom. PLOS ONE, 11, e0148269. 10.1371/journal.pone.0148269.
617
- 618 Scheele, B.C., Pasmans, F., Skerratt, L.F., Berger, L., Martel, A., Beukema, W., et al.,
619 2019. Amphibian fungal panzootic causes catastrophic and ongoing loss of biodiversity.
620 Science, 363, 1459-1463. 10.1126/science.aav0379.
621
- 622 Schielzeth, H., 2010. Simple means to improve the interpretability of regression
623 coefficients. Methods in Ecology and Evolution, 1, 103-113.
624 <https://doi.org/10.1111/j.2041-210X.2010.00012.x>.
625
- 626 Silva, E.T.D., Reis, E.P.D., Feio, R.N., Filho, O.P.R., 2009. Diet of the invasive frog
627 *Lithobates catesbeianus* (Shaw, 1802) (Anura: Ranidae) in Viçosa, Minas Gerais State,
628 Brazil. South American Journal of Herpetology, 4, 286-294.
629 <https://doi.org/10.2994/057.004.0312>.
630
- 631 Sodhi, N.S., Bickford, D., Diesmos, A.C., Lee, T.M., Koh, L.P., Brook, B.W.,
632 Sekercioglu, C.H., Bradshaw, C.J., 2008. Measuring the meltdown: drivers of global
633 amphibian extinction and decline. PLOS ONE, 3, e1636.
634 <https://doi.org/10.1371/journal.pone.0001636>.
635
- 636 Sparling, D.W., Bickham, J., Cowman, D., Fellers, G.M., Lacher, T., Matson, C.W.,
637 McConnell, L., 2015. In situ effects of pesticides on amphibians in the Sierra Nevada.
638 Ecotoxicology, 24, 262-278. <https://doi.org/10.1007/s10646-014-1375-7>.
639
- 640 Sutherland, W.J., Pullin, A.S., Dolman, P.M., Knight, T.M., 2004. The need for
641 evidence-based conservation. Trends in Ecology & Evolution, 19, 305-308.
642 <https://doi.org/10.1016/j.tree.2004.03.018>.
643
- 644 Tingley, R., Shine, R., 2011. Desiccation Risk Drives the Spatial Ecology of an
645 Invasive Anuran (*Rhinella marina*) in the Australian Semi-Desert. PLOS ONE, 6,
646 e25979. <https://doi.org/10.1371/journal.pone.0025979>.
647

648 Wake, D.B., Vredenburg, V.T., 2008. Are we in the midst of the sixth mass extinction?
649 A view from the world of amphibians. *Proceedings of the National Academy of*
650 *Sciences*, 105, 11466-11473. [10.1073/pnas.0801921105](https://doi.org/10.1073/pnas.0801921105).

651

652 Ward-Fear, G., Greenlees, M.J., Shine, R., 2016. Toads on Lava: Spatial Ecology and
653 Habitat Use of Invasive Cane Toads (*Rhinella marina*) in Hawai'i. *PLOS ONE*, 11,
654 e0151700. <https://doi.org/10.1371/journal.pone.0151700>.

655

656 Wilson, K.A., McBride, M.F., Bode, M., Possingham, H.P., 2006. Prioritizing global
657 conservation efforts. *Nature*, 440, 337-340. <https://doi.org/10.1038/nature04366>

658

659 Winter, M., Fiedler, W., Hochachka, W.M., Koehncke, A., Meiri, S., Riva, I.D., 2016.
660 Patterns and biases in climate change research on amphibians and reptiles: a systematic
661 review. *Royal Society Open Science*, 3, 160158. <http://dx.doi.org/10.1098/rsos.160158>.

662


663 Xing, Y., Zhang, C., Fan, E., Zhao, Y., 2016. Freshwater fishes of China: species
664 richness, endemism, threatened species and conservation. *Diversity and Distributions*,
665 22, 358-370. <https://doi.org/10.1111/ddi.12399>.

666

667 Zhang, H., Hu, Y., Zhang, Y., Li, W., 2015. Evidence of the Matthew effect in scientific
668 research on mammals in the Chinese First-class National Protected Animals list.
669 *Biodiversity and Conservation*, 24, 2883-2886. [https://doi.org/10.1007/s10531-015-](https://doi.org/10.1007/s10531-015-0983-8)
670 0983-8.

671


672



673
674
675
676
677
678
679


Figure 1. Frequency distribution of amphibian species per number of articles (transformed in log) from Web of Science. Publications cover the period between 1945 and February 2018.

680
681



682
683
684
685
686
687
688

Figure 2. The 10 most studied amphibian species in “Biodiversity and Conservation” WoS thematic area, considering all categories (a) and only threat level (b) according IUCN Red List. These 10 most studied species for all IUCN categories were classified as Least Concern.



689
690

691 **Figure 3.** Coefficient estimates (95% confidence intervals), showing direction and
692 magnitude of effects of explanatory variables on conservation scientific production for
693 all amphibian species, for zero and count Hurdle models (a and b). We perform analysis
694 for each amphibian order separately (c-f), but for Caudata and Gymnophiona, which had
695 a very low amount of zeros, we make common regression model (blue and red symbols
696 represent positive and negative effects, respectively; grey represents no effect).

697 *Color should be used in this figure.*