

<https://helda.helsinki.fi>

Transfusion strategies in bleeding critically ill adults : A clinical practice guideline from the European Society of Intensive Care Medicine: Endorsement by the Scandinavian Society of Anaesthesiology and Intensive Care Medicine

Moller, Morten Hylander

2022-05

Moller , M H , Sigurosson , M I , Olkkola , K T , Rehn , M , Yli-Hankala , A & Chew , M S
2022 , ' Transfusion strategies in bleeding critically ill adults : A clinical practice guideline from the European Society of Intensive Care Medicine: Endorsement by the Scandinavian Society of Anaesthesiology and Intensive Care Medicine ' , Acta Anaesthesiologica Scandinavica , vol. 66 , no. 5 , pp. 638-639 . <https://doi.org/10.1111/aas.14047>

<http://hdl.handle.net/10138/343294>

<https://doi.org/10.1111/aas.14047>

publishedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

CLINICAL PRACTICE GUIDELINE

Transfusion strategies in bleeding critically ill adults: A clinical practice guideline from the European Society of Intensive Care Medicine: Endorsement by the Scandinavian Society of Anaesthesiology and Intensive Care Medicine

Morten Hylander Møller¹  | Martin Ingi Sigurðsson^{2,3} | Klaus T. Olkkola⁴ | Marius Rehn^{5,6,7}  | Arvi Yli-Hankala^{8,9}  | Michelle S. Chew¹⁰

¹Department of Intensive Care, Copenhagen University Hospital Rigshospitalet, Copenhagen, Denmark

²Division of Anesthesia and Intensive Care Medicine, Landspítali-The National University Hospital of Iceland, Reykjavik, Iceland

³Faculty of Medicine, University of Iceland, Reykjavik, Iceland

⁴Department of Anaesthesiology, Intensive Care and Pain Medicine, University of Helsinki and Helsinki University Hospital, Helsinki, Finland

⁵Division of Prehospital Services, Air Ambulance Department, Oslo University Hospital, Oslo, Norway

⁶The Norwegian Air Ambulance Foundation, Oslo, Norway

⁷Faculty of Health Sciences, University of Stavanger, Stavanger, Norway

⁸Department of Anaesthesia, Tampere University Hospital, Tampere, Finland

⁹Faculty of Medicine and Health Technology, Tampere University, Tampere, Finland

¹⁰Department of Anaesthesia and Intensive Care, Biomedical and Clinical Sciences, Linköping University, Linköping, Sweden

Correspondence

Morten Hylander Møller, Department of Intensive Care, Copenhagen University Hospital Rigshospitalet, DK-2100 Copenhagen, Denmark.
Email: mortenhylander@gmail.com

Funding information

Funding was provided from the Scandinavian Society of Anaesthesiology and Intensive Care Medicine and institutional and/or departmental sources.

Abstract

The Clinical Practice Committee of the Scandinavian Society of Anaesthesiology and Intensive Care Medicine endorses the clinical practice guideline *Transfusion strategies in bleeding critically ill adults: a clinical practice guideline from the European Society of Intensive Care Medicine*. This trustworthy clinical practice guideline serves as a useful decision aid for Nordic anaesthesiologists caring for critically ill patients with bleeding.

KEYWORDS

AGREE II, bleeding, clinical practice guideline, critically ill, ICU, transfusion

1 | BACKGROUND

Critically ill patients, including those in the intensive care unit (ICU) frequently experience bleeding, and this is associated with increased morbidity and mortality.^{1,2}

The management of bleeding in critically ill patients is challenging and complex, and often involve multiple concurrent strategies for monitoring coagulopathy, transfusing blood products, and administering medications to support coagulation.³

The clinical practice guideline *Transfusion strategies in bleeding critically ill adults: a clinical practice guideline from the European*

Society of Intensive Care Medicine provides evidence-based recommendations for transfusion of bleeding in critically ill patients in the ICU.⁴

2 | METHODS

It was decided by the Clinical practice committee (CPC) of the Scandinavian Society of Anaesthesiology and Intensive Care Medicine (SSAI) to assess the clinical practice guideline *Transfusion strategies in bleeding critically ill adults: a clinical practice guideline from the European*

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *Acta Anaesthesiologica Scandinavica* published by John Wiley & Sons Ltd on behalf of Acta Anaesthesiologica Scandinavica Foundation.

*Society of Intensive Care Medicine*⁴ for possible endorsement. The Appraisal of Guidelines for REsearch and Evaluation (AGREE) II tool⁵ was used. Details on the endorsement process are available elsewhere.⁶

3 | RESULTS

All six SSAI CPC members completed the appraisal. The individual domain totals were: Scope and Purpose 92%; Stakeholder Involvement 72%; Rigour of Development 79%; Clarity of Presentation 77%; Applicability 69%; Editorial Independence 90%; Overall Assessment 89% (Figure 1).

The breakdown of the individual appraisers (de-identified) is available in the Supporting information.

4 | DISCUSSION

Agreement between the SSAI CPC appraisers was high, and the overall assessment of the guideline was good. There were minor issues related to stakeholder involvement, as there were no patient representatives on the panel, but patient values and preferences were sought obtained through a literature review. Also, applicability to low resource settings and mass casualties may be limited. Of note, the body of evidence was limited for many questions, why no recommendation was issued in 11 out of 26 questions.

The guideline can be used in daily clinical practice in the Nordic countries without major adaptation or modification.

The clinical practice guideline *Transfusion strategies in bleeding critically ill adults: a clinical practice guideline from the European Society*

*of Intensive Care Medicine*⁴ serves as a useful decision aid for Nordic anaesthesiologists caring for critically ill patients with bleeding.

5 | CONCLUSION

The SSAI CPC endorses the clinical practice guideline *Transfusion strategies in bleeding critically ill adults: a clinical practice guideline from the European Society of Intensive Care Medicine*.⁴

CONFLICTS OF INTEREST

No Clinical Practice Committee member had direct or indirect conflicts of interest.

AUTHOR CONTRIBUTIONS

All authors drafted, revised and approved the final manuscript.

ORCID

Morten Hylander Møller  <https://orcid.org/0000-0002-6378-9673>

Marius Rehn  <https://orcid.org/0000-0001-9519-241X>

Arvi Yli-Hankala  <https://orcid.org/0000-0002-5029-9181>

REFERENCES

- Russell L, Holst LB, Kjeldsen L, Stensballe J, Perner A. Risks of bleeding and thrombosis in intensive care unit patients with haematological malignancies. *Ann Intensive Care*. 2017;7:119.
- Lauzier F, Arnold DM, Rabbat C, et al. Risk factors and impact of major bleeding in critically ill patients receiving heparin thromboprophylaxis. *Intensive Care Med*. 2013;39:2135-2143.
- Pham HP, Shaz BH. Update on massive transfusion. *Br J Anaesth*. 2013;111:i71-i82.
- Vlaar APJ, Dionne JC, de Bruin S, et al. Transfusion strategies in bleeding critically ill adults: a clinical practice guideline from the European Society of Intensive Care Medicine. *Intensive Care Med*. 2021. doi:10.1007/s00134-021-06531-x
- Brouwers MC, Kho ME, Browman GP, et al.; Consortium ANS. AGREE II: advancing guideline development, reporting and evaluation in health care. *J Clin Epidemiol*. 2010;63:1308-1311.
- Rehn M, Chew MS, Olkkola KT, Örn Sverrisson K, Yli-Hankala A, Møller MH. Endorsement of clinical practice guidelines by the Scandinavian Society of Anaesthesiology and Intensive Care Medicine. *Acta Anaesthesiol Scand*. 2019;63:161-163.

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.




FIGURE 1 Summary of the appraisal of guidelines for REsearch and Evaluation (AGREE) II assessment⁵ [Colour figure can be viewed at wileyonlinelibrary.com]

How to cite this article: Møller MH, Sigurðsson MI, Olkkola KT, Rehn M, Yli-Hankala A, Chew MS. Transfusion strategies in bleeding critically ill adults: A clinical practice guideline from the European Society of Intensive Care Medicine: Endorsement by the Scandinavian Society of Anaesthesiology and Intensive Care Medicine. *Acta Anaesthesiol Scand*. 2022;66:638–639. doi:10.1111/aas.14047