

Survival, morphological variability, and performance of *Opuntia ficus-indica* in a semi-arid region of India

Sunil Kumar, Dana Ram Palsaniya, Tirumala Kiran Kumar, Asim Kumar Misra, Shahid Ahmad, Arvind Kumar Rai, Ashutosh Sarker, Mounir Louhaichi, Sawsan Hassan, Giorgia Liguori, Probir Kumar Ghosh, Prabhu Govindasamy, Sonu Kumar Mahawer & Hulgathur Appaswamy Bhargavi

To cite this article: Sunil Kumar, Dana Ram Palsaniya, Tirumala Kiran Kumar, Asim Kumar Misra, Shahid Ahmad, Arvind Kumar Rai, Ashutosh Sarker, Mounir Louhaichi, Sawsan Hassan, Giorgia Liguori, Probir Kumar Ghosh, Prabhu Govindasamy, Sonu Kumar Mahawer & Hulgathur Appaswamy Bhargavi (2022): Survival, morphological variability, and performance of *Opuntia ficus-indica* in a semi-arid region of India, Archives of Agronomy and Soil Science, DOI: [10.1080/03650340.2022.2031998](https://doi.org/10.1080/03650340.2022.2031998)

To link to this article: <https://doi.org/10.1080/03650340.2022.2031998>



© 2022 International Center for Agricultural Research in the Dry Areas (ICARDA).
Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 20 Feb 2022.



Submit your article to this journal [↗](#)



Article views: 214








View related articles [↗](#)



View Crossmark data [↗](#)

Survival, morphological variability, and performance of *Opuntia ficus-indica* in a semi-arid region of India

Sunil Kumar ^a, Dana Ram Palsaniya^a, Tirumala Kiran Kumar^b, Asim Kumar Misra^c, Shahid Ahmad^d, Arvind Kumar Rai^e, Ashutosh Sarker^f, Mounir Louhaichi ^{g,h}, Sawsan Hassan ⁱ, Giorgia Liguori ^j, Probir Kumar Ghosh^k, Prabhu Govindasamy ^a, Sonu Kumar Mahawer^a and Hulgathur Appaswamy Bhargavi^d

^aICAR-Indian Grassland and Fodder Research Institute, Division of Crop Production, Jhansi-284003, UP, India; ^bICAR-Central Tobacco Research Institute, Division of Crop Production, Rajamundry-533 105, A.P., India; ^cICAR-Indian Grassland and Fodder Research Institute, Division of Plant and Animal relationship, Jhansi-284003, UP, India; ^dShahid Ahmad, ICAR-Indian Grassland and Fodder Research Institute, Division of Crop Improvement, Jhansi-284003, UP, India; ^eICAR-Central Soil Salinity Research Institute, Division of Soil and Crop Management, Karnal-132001, Haryana, India; ^fInternational Center for Agricultural Research in the Dry Areas, Biodiversity & Crop Improvement Program, New Delhi-110012, India; ^gMounir Louhaichi, International Center for Agricultural Research in the Dry Areas, Resilient Agricultural Livelihood Systems Program, Tunis, Ariana 2049, Tunisia; ^hDepartment of Animal and Rangeland Sciences, Oregon State University, Corvallis, OR 97331, USA; ⁱInternational Center for Agricultural Research in the Dry Areas, Resilient Agricultural Livelihood Systems Program, Amman 11195, Jordan; ^jUniversity of Palermo, Department of Agricultural Food and forest Sciences Viale delle Scienze, Ed. 4, 90128 Palermo, Italy; ^kICAR -National Institute of Biotic Stress Management, Raipur, 493 225, India

ABSTRACT

Cactus pear (*Opuntia ficus-indica* (L.) Mill.) can survive extreme environmental condition and is known for its fodder potential in many parts of the world. The morphological diversity of 15 introduced accessions was evaluated at Jhansi, Uttar Pradesh, India. The plants were established in 2013. Survival and nutrient status were evaluated after two years. Above-ground plant height, biomass, primary and secondary cladode numbers, primary and secondary cladode lengths and below-ground root length, weight, and surface area measurements were done six years after cladode planting. Yellow San Cono, White Roccapalumba, and Seedless Roccapalumba survived 100%. The discriminant traits according to principal component analysis were: primary cladodes plant⁻¹ (component loading, 0.87), primary cladodes biomass (0.95), secondary cladodes plant⁻¹ (0.83), canopy width (0.84), and plant biomass (0.92). Hierarchical cluster analysis grouped 15 accessions into two main clusters based on 17 morphological traits. Cluster I showed favorable values for many above- and below-ground morphological traits while Cluster II showed higher performance for root system width, height, and biomass, and primary and secondary cladode numbers. The results indicate that cactus pear accessions have considerable morphological variability and genetic diversity suitable for promotion as alternative fodder resources in semi-arid regions of India.

ARTICLE HISTORY

Received 8 June 2021
Accepted 17 January 2022

KEYWORDS

Cactus pear; cladode; fodder; principal component analysis; root architecture

Introduction

Cactus pear (*Opuntia ficus-indica*) is a drought-tolerant cactus and exhibits crassulacean acid metabolism (CAM). Cactus pear is mainly grown in arid and semi-arid regions of the world with less than 250 and 250–450 mm annual precipitation (Stintzing and Carle 2005; Prasad et al. 2021). With considerable genetic variability, cactus pear grows in Italy, Spain, Mexico, Brazil, Chile, Argentina, and California (Inglese et al. 2002; Louhaichi et al. 2018). The genus *Opuntia* consists of two subgenera: *Platyopuntia* (cactus pear) characterized by flattened pads (cladodes) and used for animal and human consumption. *Cylindropuntia* (cholla cactus) has cylindrical pads and is generally used for ornamental purposes. Most cactus pear plant biomass is pad material rather than fruits. Therefore, cactus pear is widely used as animal feed in arid and semi-arid tropical regions, especially during droughts. Consumption of the water-rich cactus pads reduces animal drinking water requirements (Tegegne et al. 2007).

The genus *Opuntia* has special morphological, physiological, and biochemical attributes that enable these plants to grow under adverse climate conditions. The taxonomic classification of *Opuntia* has been reported as complex, mainly due to limited information about its morphological descriptors (Samah et al. 2016). However, *Opuntia* shows high morphological variation within species, which is influenced by many factors such as geographical distribution, population size, and pollen and seed dispersal (Rayamajhi and Sharma 2018). This provides scope for morphological descriptors as an effective tool for identification and classification of genotypes in *Opuntia*.

The CAM photosystem helps cactus pear survive in hot (arid) and semi-hot (semi-arid) areas of the world. Han and Felker (1997) reported that cactus pear has a greater water-to-biomass conversion ratio than C_3 and C_4 plants, which helps produce more biomass in hot regions (Felker et al. 2005; Inglese et al. 2017). High atmospheric temperature is not a limiting factor for cactus pear growth (Nobel et al. 1986) but temperatures below -5°C can impair growth or kill plants (Inglese et al. 1995). Cactus pear can thrive in areas with average annual rainfall of 400–600 mm and in regions with 200 mm (Acevedo et al. 1983). However, supplemental irrigation during summer and unfavorable climatic conditions (e.g. dry spells during monsoon periods) is highly advantageous for better biomass yield (Bakali 2013).

Cactus pear can produce 1 kg of dry matter (DM) per 162 kg of water. This is a physiological advantage allowing it to thrive in hot areas (Han and Felker 1997). Guevara et al. (2011) found that the conversion efficiency of water to biomass was five times that of C_3 and three times that of C_4 plants. The water content varies with the age of the cladodes. Generally, younger cladodes have more water than mature and older cladodes (90.8%, 89.1%, and 83.4% respectively; Flores et al. 1995). Similarly, nutrient content (nitrogen, phosphorus, potassium, manganese, zinc, and sodium) also decreases with age, except for magnesium (Lopez et al. 1988). Cactus pear can survive with rainfall that is too low to benefit other fodder crops (Snyman 2013). In semi-arid regions, the mean biomass yield of cactus pear ranged from 10–20 t ha⁻¹ (Santos et al. 2000; Dubeux et al. 2006; Gebretsadik et al. 2013). Dev et al. (2018) reported that the greatest green plant biomass was 1,007 g plant⁻¹ (25% shade) and 1,086 g plant⁻¹ (50% shade) two years after establishment. Similarly, Pareek et al. (2003) reported green biomass of 268 to 1,300 g plant⁻¹ in 51 cactus pear accessions in hot arid regions of India.

There are few studies on the performance of cactus pear accessions in semi-arid regions of India. Therefore, to make more efficient use of the cactus germplasm available in the country, it would be beneficial to study the genetic diversity, the above- and below-ground morphological variability in germplasm accessions and determine the relationships among the traits that contribute to this variability. There is also a need to understand the genetic relationships among cactus pear accessions and the contribution of different above- and below-ground traits to multivariate polymorphism. We therefore conducted a study to assess the performance of different accessions to i) identify the morphological attributes of cactus pear grown in a semi-arid environment in India and ii), to use above- and below-ground morphological variability and their relationships among different accessions to investigate the genetic diversity and performance of cactus pear.

Materials and methods

Experimental site description

The cactus pear accessions we examined were established at the Central Research Farm of ICAR Indian Grassland and Fodder Research Institute, Jhansi, India. A total of 15 accessions were received from the University of Palermo, Italy in April 2013 (Algerian, Blue Motto, Morado, Red Roccapalumba, Red San Cono, Roly Poly, Seedless Roccapalumba, Trunzara Red Bronte, White Roccapalumba, White San Cono, White Santa Margherita, Yellow Roccapalumba, Yellow San Cono, Yellow Santa Margherita Belice, and Zastron).

Five cladodes of each accession were obtained through the ICARDA India office, New Delhi and subjected to quarantine at the national facility at ICAR National Bureau of Plant Genetic Resources, New Delhi. The introduced cladodes from ICARDA were multiplied in a nursery. Cladodes collected from the nursery were planted for this experiment in April 2013. The plot size of individual accessions was 13 m × 6 m and with an interplant spacing of 1 m and 2 m between rows (5,000 plants ha⁻¹). Three rows of cladodes were planted with 39 plants in each block. Cactus pear cladodes were planted after antifungal treatment with Bavistin 50% WP (2 g L⁻¹). At the time of planting, 3–5 kg plant⁻¹ of well-composted farmyard manure was applied (4.0 g N, 1.16 g P and 3.1 g K kg⁻¹). Newly planted cladodes were irrigated twice during the summer of the first year and survived without further irrigation. Weeds were managed using both mechanical and chemical methods. A targeted spray of glyphosate (0.8 kg a.i. ha⁻¹) was done before the rainy season of each year to control early-season weeds. The matured cladodes (2-years old) were harvested to remove crowded and dense stands. Approximately 60% of the cladodes per plant were removed in a single harvest.

The site soil type was clay loam soil, containing 47.86% sand, 20% silt, and 32.14% clay at 0 to 15 cm soil depth. The pH and organic carbon content of the soil were 7.26 and 0.69%, respectively. The long-term (1969–2018) yearly average rainfall in the study location was 950 mm. The maximum yearly average rainfall was recorded in 2013 (1,286 mm) and the minimum in 2017 (690 mm, [Figure 1](#)). The yearly average maximum and minimum temperatures were 32.95°C and 19.08°C. At this location, the soil moisture was generally high (107–168 mm) from July to November due to the annual monsoon rains ([Figure 1](#)).

Survival rate of cactus pear accessions

The survival percentage of accessions was recorded two years after planting. The survival rate was measured by dividing the number of live plants by the total number of plants and expressed as a percentage.

Nutrient status

The proximate nutrient values were estimated by collecting cladodes from 2-year-old accessions. From each accession, mature cladodes (three plants and three cladodes plant⁻¹) were chopped into small pieces and oven-dried at 63°C for 72 hours or until the weight was constant. Oven-dried cladodes were ground using a plant sample grinder. One sample per accession (1 g) was used to estimate crude protein (CP; [AOAC 2007](#)) and cell wall constituents NDF (neutral detergent fiber), ADF (acid detergent fiber), and lignin contents ([Van Soest et al. 1991](#)). *In vitro* dry matter digestibility (IVDMD) was determined using the standard method of [Tilley and Terry \(1963\)](#). In this case, 0.5 g of dry sample was incubated in 50 mL of digestion solution containing 40 mL of CO₂ saturated phosphate carbonate buffer and 10 mL of strained sheep-rumen liquor.




Figure 1. Weather parameters during 2013–2019 at Jhansi (UP), India.

Above- and below-ground growth analysis

Observations on above- and below-ground growth parameters were recorded in 2019 after six years of growth. Above-ground parameters were recorded from three randomly selected plants per plot (plant height and width, number of primary and secondary cladodes per plant, primary and secondary cladode length and width, and biomass of primary and secondary cladodes, Kumar et al. 2021). Plant height was measured from the bottom of the plant to the uppermost cladode. Width was measured by selecting the maximum swath width (middle of the plant). Primary and secondary cladode length and width were measured vertically and horizontally in three randomly selected cladodes per plant. The average fresh weight of three random cactus pear plants was considered as a biomass yield plant^{-1} . Root system depth, width, and weight were manually recorded using three randomly selected plants plot^{-1} . For extracting the whole root system of a cactus pear plant, a circular pit was made around the plant and the pit was filled with water for 24 hours (1.5–2.0 m diameter). The saturated soil was removed carefully from around the plant and the roots removed. The thoroughly washed root system was used for further analysis. The entire root system length and width were measured using a ruler, and then the secondary and finer roots were separated from the primary root system for analysis of root length, surface area, and volume using a Biovis PSM root scanner (Expert Vision Lab Pvt. Ltd., Dadar West, Mumbai, India). Observations on primary and secondary roots were recorded using a digital Vernier caliper. The root surface area and volume measured using the Biovis PSM root system and the Vernier caliper were added together to represent the total root surface area and volume.




Figure 2. Length of primary cladodes in accessions of cactus pear after six years. Note: Al, Algerian; Bm, Blue Motto; Mo, Morado; Rr, Red Roccapalumba; Rs, Red San Cono; Rp, Roly Poly; Sr, Seedless Roccapalumba; Tr, Trunzara Red Bronte; Wr, White Roccapalumba; Ws, White San Cono; Wm, White Santa Margherita; Yb, Yellow Santa Margherita Belice; Yr, Yellow Roccapalumba; Ys, Yellow San Cono; Za, Zastron.*means with different alphabetical letters are significant at $\alpha = 0.05$.

Data handling and statistical analysis

Analysis of variance (ANOVA) was performed for above- and below-ground morphological traits using the PROC GLIMMIX procedure in SAS (v9.3, SAS Institute 2011). A randomized complete block design was used in this experiment. All the experimental units were replicated three times. Normality within the residuals was performed by PROC UNIVARIATE and residual plots in SAS using the Shapiro-Wilk test and visually confirmed with a normal probability plot. Data on above- and below-ground parameters were normally distributed, thus, data transformation was not required. However, data after two years of establishment were not normally distributed (survival, cladode length, cladode width, moisture content, yield, dry matter, NDF, ADF, cellulose, lignin, ash, CP, and IVDMD), therefore a square root transformation ($\sqrt{X + 0.5}$) was done to fulfill the assumptions of ANOVA. In all ANOVAs, accession was a fixed effect and replication and year were random effects. The post-hoc test was performed using the least significant difference test described by Fisher ($\alpha = 0.05$). Pearson's correlation was performed using PROC CORR to determine the relationship between biomass and morphological traits. Combined principal component analysis (PCA) was applied to 17 above-ground and below-ground morphological traits using the factoextra package (Kassambara and Mundt 2017) in R (v3.6.1, R Core Team 2019). For grouping based on the 17 morphological traits, a hierarchical cluster analysis (HCA) was performed using the Ward.D2 package in R. We used the k-1 cut off to keep the clusters at a constant height. Figures in the manuscript were made using Sigmaplot (v14, Sigmaplot 1996).

Results and discussion

Survival and growth performance of cactus pear accessions

The survival and growth performance are shown in Table 1. The survival rates of all accessions ranged from 30 to 100%. The Yellow San Cono, White Roccapalumba, and Seedless Roccapalumba had a 100% survival rate. The range in mean yield was 15.3 to 48 kg plant⁻¹ year⁻¹. The highest was for Yellow San Cono (48 kg plant⁻¹ year⁻¹) and the lowest for Roly Poly (15.3 kg plant⁻¹ year⁻¹; Table 1). This indicates a relationship between accession survival rate and biomass yield. Cladode lengths and widths also varied among accessions. The greatest length (34.5 cm) and width (15.5 cm) were for Roly Poly and Yellow Santa Margherita Belice, respectively. Overall, Yellow San Cono can survive better in a semi-arid region compared to the other accessions.

Nutritional profile of cactus pear accessions

Plant sample analysis data at two years after establishment are presented in Table 2. Nutritional profile data confirmed that the major constituent of cactus pear plants was water (93.1–97.1%, Table 1). The DM range was 2.9 to 6.9%. Misra et al. (2018) reported an average DM of 7.56% in feeding trials in India. The low percentage of DM indicates that cactus pear alone cannot meet the DM requirement for animals (eg. 4% of animal body weight). Therefore, mixing it with other fodder would be an effective option to support livestock nutrition needs. However, the low fiber fractions (NDF, ADF, cellulose, and lignin) of the plants (31.3 to 40.9%, 17.2 to 23.1%, 10.6 to 18.5%, and 3.7 to 5.5%, respectively) indicated that cactus pear is nutritive and digestible (Misra et al. 2018) and is also the reason for its higher crude protein level (CP, 9.0–13.8%). Our results are consistent with those of Gregory and Felker (1992), who reported 4–12% CP content in eight contrasting cactus pear accessions. Similarly, Sirohi et al. (1997) reported 9.2% mean CP content in cactus pear. Therefore, cactus pear would seem to be a suitable alternative fodder where soil moisture and rainfall are major constraints to the cultivation of other fodder crops.

Table 1. Survival and growth performance of cactus pear novel accessions following two years of growth.

Accessions	Survival (%)	Cladode length (cm)	Cladode width (cm)	Moisture content (%)	Yield (kg plant ⁻¹ year ⁻¹)
Algerian	8.2(60)d	5.8 (28.5)†e	3.9 (12.2)i *	10.2 (95.5)g	7.3 (46.6)b
Blue Motto	6.2 (33)g	5.3 (24.0)l	3.7 (10.2)k	10.1 (93.4)m	4.5 (16.4)k
Morado	8.6 (67)c	5.7 (27.7)f	4.3 (14.7)d	10.2 (95.2)j	4.9 (20.0)i
Red Roccapalumba	6.2 (33)g	5.7 (27.5)g	4.2 (14.2)e	10.2 (95.6)f	4.9 (20.0)i
Red San Cono	5.9 (30)h	5.5 (25.0)k	4.1 (13.5)g	10.3 (95.8)e	5.9 (29.8)d
Roly Poly	8.6 (67)c	6.3 (34.5)a	4.2 (14.2)e	10.3 (96.3)c	4.4 (15.3)l
Seedless Roccapalumba	10.5 (100)a	5.5 (25.5)j	3.7 (10.2)k	10.2 (95.4)h	4.7 (18.0)j
Trunzara Red Bronte	9.4 (80)b	6.2 (33.0)c	4.4 (15.0)c	10.3 (96.3)c	5.0 (20.8)h
White Roccapalumba	10.5 (100)a	5.6 (26.7)i	4.1 (13.0)h	10.1 (93.1)n	7.1 (43.7)c
White San Cono	7.5 (50)e	5.6 (27.0)h	3.9 (12.0)j	10.2 (95.1)k	5.5 (25.2)e
White Santa Margherita	6.2 (33)g	5.8 (28.5)e	4.2 (13.7)f	10.2 (95.3)i	5.3 (23.2)f
Yellow Roccapalumba	6.8 (40)f	5.8 (28.5)e	4.4 (15.2)b	10.3 (96.1)d	5.2 (22.4)g
Yellow San Cono	10.5 (100)a	6.1 (32.0)d	4.4 (15.0)c	10.3 (97.1)a	7.4 (48.0)a
Yellow Santa Margherita Belice	8.2 (60)d	5.5 (25.5)j	4.4 (15.5)a	10.2 (94.5)l	7.1 (43.6)c
Zastron	6.8 (40)f	6.2 (33.5)b	3.9 (12.0)j	10.3 (96.4)b	7.3 (46.5)b
p-value	<.0001	<.0001	<.0001	<.0001	<.0001
CV (%)	19.19	5.13	5.15	0.54	18.84

‡Square root transformed values ($\sqrt{X + 0.5}$); †values inside the parenthesis are original values; *means with different alphabetical letters are significant at $\alpha = 0.05$

Table 2. Nutritional profile of cactus pear novel accessions following two years of growth.

Accessions	Dry matter (%)	NDF (%)	ADF (%)	Cellulose (%)	Lignin (%)	Ash (%)	CP† (%)	IVDMD‡ (%)
Algerian	2.6 (4.4)‡b	6.5 (36.6)‡*	4.9 (19.4)h	4.2 (14.1)d	2.6 (4.4)j	6.4(35.3)j	3.7(10.6)g	8.5 (65.2)g
Blue Motto	3.0 (6.6)k	6.7 (39.1)d	5.1 (21.3)b	4.3 (14.6)c	2.8 (5.5)b	6.4(35.3)j	3.8(11.3)d	8.5 (64.9)h
Morado	2.6 (4.7)l	6.4 (35.4)j	4.9 (19.9)e	4.3 (14.6)c	2.7 (4.9)e	6.3(34.3)k	3.8(11.0)e	7.9 (55.6)k
Red Roccapalumba	2.5 (4.4)l	6.8 (40.9)a	5.3 (23.1)a	4.8 (18.5)a	2.4 (3.7)m	6.2(33.4)m	3.8(11.5)c	9.0 (73.1)b
Red San Cono	2.5 (4.2)d	6.7 (39.0)e	4.9 (19.5)g	4.1 (13.3)f	2.7 (5.0)d	6.7(38.9)c	3.6(10.1)j	8.7 (68.7)d
Roly Poly	2.4 (3.7)l	6.5 (37.1)g	4.8 (18.7)k	4.0 (12.6)h	2.7 (5.1)c	6.5(36.8)f	3.5(9.0)m	8.9 (70.1)c
Seedless Roccapalumba	2.6 (4.6)j	6.3 (34.7)l	4.7 (17.7)m	4.0 (12.5)l	2.6 (4.6)h	6.7(38.5)d	3.6(9.9)j	8.7 (67.3)f
Trunzara Red Bronte	2.4 (3.7)h	6.5 (36.8)h	4.8 (18.8)j	4.1 (13.0)g	2.6 (4.5)j	6.7(39.2)b	3.5(9.6)l	8.2 (59.6)j
White Roccapalumba	3.1 (6.9)c	6.8 (40.6)b	5.0 (20.5)c	4.3 (14.6)c	2.6 (4.7)g	6.3(33.9)l	3.7(10.8)h	8.3 (61.9)l
White San Cono	2.7 (4.9)e	6.5 (36.6)j	4.6 (20.3)d	4.4 (15.5)b	2.5 (4.0)k	6.4(35.9)h	3.7(10.5)h	9.0 (73.9)a
White Santa Margherita	2.6 (4.7)l	6.4 (35.0)k	4.6 (17.2)n	3.8 (11.5)j	2.6 (4.7)g	6.5(36.5)g	3.9(11.6)b	8.7 (68.1)e
Yellow Roccapalumba	2.4 (3.9)g	6.6 (38.3)f	4.9 (19.8)f	4.2 (13.9)e	2.6 (4.4)j	6.4(35.1)j	3.7(10.6)g	8.8 (69.9)d
Yellow San Cono	2.2 (2.9)a	6.1 (32.4)m	4.7 (17.7)m	3.7 (10.6)k	2.8 (5.4)b	6.5(37.2)e	3.8(11.3)d	NA
Yellow Santa Margherita	2.8 (5.5)c	6.0 (31.3)n	4.8 (19.1)j	4.3 (14.6)l	2.4 (3.8)l	6.3(34.3)k	4.2(13.8)a	4.5 (16.3)m
Zastron	2.3 (3.6)b	6.8 (40.0)c	4.7 (17.8)l	4.0 (12.5)l	2.6 (4.8)f	6.8(40.0)a	3.6(9.8)k	7.9 (54.8)l
p-value	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001
CV (%)	9.51	3.52	3.59	5.83	5.79	2.62	4.63	13.34

†ADF, acid detergent fiber; NDF, neutral detergent fiber; CP, crude protein; IVDMD, *in vitro* dry matter digestibility; NA, not available

‡Square root transformed values (√(X + 0.5)); values inside the parenthesis are original values; *means with different alphabetical letters are significant at α = 0.05

Above-ground morphological traits

The plant height, biomass, primary and secondary cladode numbers, and primary and secondary cladode lengths at six years after establishment are presented in Table 3 and Figures 2 and 3. Accessions differed in terms of above-ground morphological traits ($p < 0.05$). The range in plant height was 98 to 266 cm (Table 3). Accessions Red San Cono (266 cm) and Algerian (265 cm) had the greatest height ($p < 0.0001$), followed by White Roccapalumba (254.33 cm) and Morado (233 cm), with the lowest for Red Roccapalumba (98 cm) and White Santa Margherita (115.33 cm). The greater-height accessions can be useful as both fodder and a wind barrier. In contrast, Oelofse et al. (2006) reported that Zastron was the highest (223 cm) and Algerian the lowest (173 cm) among 10 studied accessions in a study conducted in a sub-tropical region in South Africa. This may be due to the differences in climate and soil type of the two study regions. However, the reported plant height range in this study is within the range found by Oelofse et al. (2006). Nieto-Garibay et al. (2011) also noted that *Opuntia tapona* was taller when grown in the California Gulf region of Mexico than in the Pacific Ocean region because of less wind. Taller cactus pear accessions can be useful as fodder shrubs (higher yield), wind barriers, and live fences in semi-arid regions.

The primary cladode numbers were greater ($p < 0.05$) for Yellow San Cono (18 plant⁻¹), Red San Cono (15 plant⁻¹), White Roccapalumba (15 plant⁻¹), Seedless Roccapalumba (14 plant⁻¹), Trunzara Red Bronte (14 plant⁻¹), Morado (13 plant⁻¹), and Yellow Rocca Palumba (13 plant⁻¹) (Table 3). However, the trend was different for the secondary cladode number. The highest were Red San Cono (86 plant⁻¹), Yellow San Cono (66 plant⁻¹), Yellow Santa Margherita Belice (61 plant⁻¹), and White Roccapalumba (52 plant⁻¹). The trend change could be due to the individual difference in genetic potential rather than location and management. This agrees with the results of Lima et al. (2016), who documented the average numbers of primary cladodes in the range of 8 to 20 plant⁻¹ in a semi-arid region of Brazil. The number of primary cladodes was affected by year (i.e. weather variables) and cutting interval (Lima et al. 2016). Similarly, Kumar et al. (2017) and Hassan et al. (2020) reported that the average number of cactus pear primary cladodes was 3 to 16 (Italian origin) at 24 months after planting in Italy and 0.6 to 5.0 (Mexico origin), 1.0 to 6.2 (Chile origin), 2.2 (Brazil origin), 2.0 to 6.0 (Argentina origin), and 5.6 (Algeria origin), at 16 months after planting in India. Greater numbers of cladodes per plant increase photosynthetic activity (carbon dioxide capture) and productivity (Nobel 2001). Therefore, accessions with more cladodes and plant height can yield higher biomass than those with fewer cladodes (Table 1).

In this study, primary and secondary cladode lengths were in the range of 19.33 to 39.00 and 10.66–19.17 cm respectively (Figures 2 and 3). The maximum lengths ($p = 0.0015$) of primary cladodes were for White Roccapalumba (39.00 cm), Trunzara Red Bronte (37.00 cm), Seedless Roccapalumba (37.00 cm), Yellow San Cono (36.33 cm), Red San Cono (36.16 cm), Zastron (36.00 cm), and Algerian (35.83 cm). Similarly, accessions Yellow Roccapalumba (42.83 cm), White Roccapalumba (42.50 cm), and Trunzara Red Bronte (41.67 cm) had the maximum lengths of secondary cladodes ($p < 0.0001$).

Cladode length is also an indicator of high water-use efficiency, photosynthetic activity, and biomass production. Accessions with higher cladode length and width produced higher biomass yield per unit area. Therefore, cladode length could be a potential morphological indicator for selection of accessions in semi-arid regions. The data on primary ($p = 0.02$) and secondary ($p = 0.001$) cladode width were significant. The range of primary and secondary cladode widths were 10.00 to 23.33 cm and 10.00 to 18.66 cm respectively. The highest primary and secondary cladode width were recorded for Yellow San Cono (23.33 cm) and Yellow Roccapalumba (18.66 cm; Table 3).

Plant biomass ranged from 3.83 to 118.66 kg plant⁻¹ (Table 3), with the highest ($p = 0.02$) for Red San Cono (118.66 kg plant⁻¹), followed by Yellow Santa Margherita Belice (91.33 kg plant⁻¹), Seedless Roccapalumba (88 kg plant⁻¹), White Roccapalumba (65.33 kg plant⁻¹), Trunzara Red Bronte (62 kg plant⁻¹), and Yellow San Cono (56.33 kg plant⁻¹). The biomass yield per plant was significantly correlated with primary cladode number ($p < 0.0001$; $r = 0.853$), primary cladode length ($r = 0.619$),

Table 3. Plant height, plant biomass, primary cladodes per plant, secondary cladodes per plant of cactus pear accessions.

Genotypes	Plant height (cm)	Plant biomass (kg plant ⁻¹)	Primary cladodes (no. plant ⁻¹)	Secondary cladodes (no. plant ⁻¹)	Primary cladodes Width (cm)	Secondary cladodes Width (cm)
Algerian	265.00 (± 22.54) [†] a*	42.33 (± 13.16) bc	9 (± 1.0) bcdef	33 (± 6.49) bcd	16.66 (± 1.76) bc	13.33 (± 0.88) cdefg
Blue Motto	162.33 (± 2.96) cdef	10.16 (± 4.60) c	6 (± 1.45) def	11 (± 3.46) d	10.00 (± 1.00) d	11.66 (± 0.33) fg
Morado	233.00 (± 19.65) ab	43.16 (± 15.87) bc	13 (± 3.48) abcd	36 (± 8.56) bcd	14.00 (± 2.64) bcd	14.00 (± 1.15) bcdef
Red Roccapalumba	98.00 (± 15.88) f	5.16 (± 0.93) c	5 (± 0.88) ef	9 (± 0.88) d	13.33 (± 2.90) cd	10.00 (± 0.57) g
Red San Cono	266.00 (± 9.71) a	118.66 (± 45.89) a	15 (± 1.45) ab	86 (± 20.95) a	17.66 (± 0.66) abc	15.66 (± 0.88) abcde
Roly Poly	127.33 (± 20.73) ef	21.66 (± 16.67) c	6 (± 1.53) def	29 (± 19.00) cd	14.00 (± 1.00) bcd	11.33 (± 1.76) fg
Seedless Roccapalumba	222.00 (± 37.04) abc	88.00 (± 46.87) ab	14 (± 5.84) ab	34 (± 7.81) bcd	18.66 (± 3.71) abc	16.66 (± 1.85) abc
Trunzara Red Bronte	222.00 (± 37.04) abc	62.00 (± 41.66) abc	14 (± 6.56) abc	44 (± 26.05) bcd	19.33 (± 3.17) ab	16.33 (± 2.40) abcd
White Roccapalumba	254.33 (± 13.92) ab	65.33 (± 16.04) abc	15 (± 1.00) ab	52 (± 11.78) abc	19.33 (± 0.66) ab	17.00 (± 1.73) ab
White San Cono	160.33 (± 16.45) cdef	15.83 (± 4.74) c	6 (± 1.66) def	25 (± 3.84) cd	14.66 (± 1.20) bcd	13.00 (± 0.57) edfg
White Santa Margherita	115.33 (± 21.60) f	3.83 (± 1.58) c	3 (± 0.66) f	9 (± 1.20) d	10.33 (± 0.88) d	11.66 (± 1.20) fg
Yellow Roccapalumba	195.66 (± 32.27) bcde	28.00 (± 8.66) bc	13 (± 2.08) abcd	25 (± 6.17) cd	17.66 (± 3.38) abc	18.66 (± 1.20) a
Yellow San Cono	200.33 (± 14.90) abcd	56.33 (± 17.52) abc	18 (± 3.78) a	66 (± 30.35) ab	23.33 (± 2.60) a	17.00 (± 2.08) ab
Yellow Santa Margherita Belice	215.33 (± 41.33) abc	91.33 (± 40.34) ab	10 (± 1.33) bcde	61 (± 16.18) abc	18.66 (± 0.88) abc	15.66 (± 0.88) abcde
Zastron	140.66 (± 15.38) def	20.00 (± 3.51) c	7 (± 1.20) cdef	28 (± 8.83) cd	18.66 (± 1.85) abc	12.33 (± 1.45) jefg
p-value	<0.0001	0.02	0.003	0.005	0.0001	0.02
CV (%)	35.12	109.39	59.54	81.50	28.69	22.60

[†]Standard error of mean; *Means with different alphabetical letters are significant at α = 0.05




Figure 3. Length of secondary cladodes in accessions of cactus pear after six years. Note: Al, Algerian; Bm, Blue Motto; Mo, Morado; Rr, Red Roccapalumba; Rs, Red San Cono; Rp, Roly Poly; Sr, Seedless Roccapalumba; Tr, Trunzara Red Bronte; Wr, White Roccapalumba; Ws, White San Cono; Wm, White Santa Margherita; Yb, Yellow Santa Margherita Belice; Yr, Yellow Roccapalumba; Ys, Yellow San Cono; Za, Zastron. *means with different alphabetical letters are significant at $\alpha = 0.05$.

primary cladode width ($r = 0.650$), primary cladode biomass ($r = 0.971$), secondary cladode number ($r = 0.923$), secondary cladode length ($r = 0.470$), secondary cladode width ($r = 0.830$), secondary cladode biomass ($r = 0.992$), plant height ($r = 0.815$), and canopy width ($r = 0.810$; Table 5). Selecting a cultivar with superior above-ground morphological traits would appear to be an appropriate way to increase biomass yield in this semi-arid region.

Below-ground morphological traits

Below-ground morphological traits are presented in Table 4. Root system length, weight, and surface area did not differ significantly among accessions and were in the range of 22.22–63.66 cm, 0.46–2.61 kg, and 226–1,047 cm² respectively. Soil moisture, air, temperature, and texture, the interaction of microorganisms, and plant nutrients influence the root length, weight, and surface area of a plant (Koç and Acar 2015). In this study, all these factors were constant for all the genotypes which may help explain the lack of differences among genotypes.

However, root system width ($p = 0.03$) and volume ($p < 0.005$) significantly differed among accessions. The maximum root system width was for Seedless Roccapalumba (152.33 cm) followed by Red San Cono (121.66 cm), White Santa Margherita (118.33 cm), and Zastron (105.50 cm). Root system volume was higher for Yellow Santa Margherita Belice (10,337 cm³) and Red San Cono (7,874 cm³) than for other accessions. The width and depth of cactus pear root systems will expand with time after establishment. Snyman (2006) reported that the roots of the cactus pear can grow to depths of 5–10 cm and 20 cm laterally from the stem two years after establishment. Snyman (2005) reported that cactus pear can grow only 0–5 cm deep within one year of establishment. In this study, a maximum root system depth of 63 cm and width of 152 cm were observed six years after establishment. However, earlier studies reported that cactus pear plants are generally shallow rooted (15 cm deep; Snyman 2005). Possible reasons for the greater root system depth, width, and volume in this study could be due to soil

Table 4. Root characteristics of cactus pear accessions.

Genotypes	Root weight (kg)	Root volume (cm ³)	Root surface area (cm ²)	Root length (cm)	Root width (cm)
Algerian	1.31 (± 0.43) [†] a*	3321 (± 1481) c	666 (± 143)a	58.33 (± 7.83)a	71.66 (±18.44) bcd
Blue Motto	0.74 (± 0.12)a	2922 (± 423) c	504 (± 61)a	39.66 (± 2.90)a	72.33 (±8.41) bcd
Morado	0.77 (± 0.30)a	4693 (± 2118) bc	983 (± 213)a	63.66 (± 10.73)a	79.33 (±34.07) bcd
Red Roccapalumba	1.52 (± 1.01)a	2630 (± 335) c	338 (± 78)a	53.66 (± 3.28)a	75.66 (±10.41) bcd
Red San Cono	1.09 (± 0.36)a	7874 (± 1361) ab	485 (± 110)a	46.00 (± 7.09)a	121.66 (±41.15) ab
Roly Poly	1.50 (± 0.95)a	3036 (± 2033) c	1034 (± 594)a	35.33 (± 7.21)a	48.33 (±14.33) d
Seedless Roccapalumba	2.61 (± 1.95)a	5690 (± 2857) bc	1047 (± 534)a	53.33 (± 14.85)a	152.33 (±36.97) a
Trunzara Red Bronte	0.48 (± 0.30)a	1931 (± 996) c	331 (± 206)a	38.00 (± 4.73)a	58.66 (±19.32) cd
White Roccapalumba	0.67 (± 0.20)a	1847 (± 644) c	473 (± 155)a	43.00 (± 8.18)a	77.00 (±11.53) bcd
White San Cono	0.38 (± 0.08)a	1801 (± 538) c	226 (± 81)a	55.33 (± 11.66)a	50.66 (±8.96) d
White Santa Margherita	0.83 (± 0.23)a	1864 (± 607) c	288 (± 81)a	30.33 (± 1.76)a	118.33 (±16.82) abc
Yellow Roccapalumba	1.34 (± 0.92)a	2627 (± 163) c	462 (± 195)a	42.67 (± 11.55)a	48.66 (±20.75) d
Yellow San Cono	0.46 (± 0.17)a	2574 (± 495)c	372 (± 28)a	42.67 (± 1.76)a	64.00 (±8.62) bcd
Yellow Santa Margherita Belice	0.83 (± 0.14)a	10,337 (± 1649) a	658 (± 151)a	51.00 (± 6.50)a	41.00 (±7.00) d
Zastron	0.51 (± 0.13)a	1816 (± 183) c	527 (± 97)a	22.20 (± 120)a	105.50 (±9.50) abcd
p-value	0.72	0.005	0.36	0.31	0.03
CV (%)	112.48	107.16	78.01	63.64	55.21

[†]Standard error of mean; *Means with different alphabetical letters are significant at $\alpha = 0.05$

Table 5. Pearson correlation coefficients among 17 above-ground and below-ground traits of cactus pear accessions.

	MCPP	MCL	MCW	MCB	SCPP	SCL	SCW	SCB	PH	CW	PB	RB	RD	RW	RV	RSA	RL
MCPP																	
MCL	0.496																
MCW	0.714*	0.813*															
MCB	0.853*	0.631*	0.704*														
SCPP	0.862*	0.622*	0.602*	0.909*													
SCL	0.487	0.617*	0.486	0.329													
SCW	0.841*	0.599*	0.839*	0.734*	0.667*												
SCB	0.844	0.641*	0.949*	0.944*	0.440	0.807*											
PH	0.787*	0.312	0.406	0.765*	0.326	0.608*	0.794*										
CW	0.733*	0.739*	0.689*	0.864*	0.382	0.680*	0.811*	0.750*									
PB	0.853*	0.619*	0.650*	0.923*	0.470	0.830*	0.992*	0.815*	0.810*								
RB	-0.107	0.113	-0.045	-0.120	0.209	0.138	0.105	0.036	-0.054	0.129							
RD	0.374	-0.046	0.232	0.263	-0.081	0.161	0.321	0.389	0.110	0.372	0.234						
RW	0.102	-0.093	-0.122	0.154	-0.202	0.032	0.272	0.186	-0.032	0.275	0.504	0.314					
RV	0.281	0.038	0.000	0.324	0.272	0.469	0.483	0.384	0.195	0.507	0.258	0.646**	0.504				
RSA	0.197	0.290	0.081	0.308	0.263	0.265	0.443	0.388	0.252	0.414	0.236	0.475	0.204	0.607**			
RL	0.238	0.186	0.002	0.333	0.248	0.313	0.490	0.424	0.193	0.464	0.302	0.571**	0.304	0.746**	0.954**		

MCPP, primary cladode per plant (no.); MCL, primary cladode length; MCW, primary cladode width; MCB, primary cladode biomass; SCPP, secondary cladode per plant (no.); SCL, secondary cladode length; SCW, secondary cladode width; SCB, secondary cladode biomass; PH, plant height; CW, canopy width; PB, plant biomass; RB, root biomass; RD, root diameter; RW, root width; RV, root volume; RSA, root surface area; RL, root length.

*indicates correlation is significant at $\alpha = 0.05$.

type, rainfall, and nutrient availability. Growers can select accessions with maximum root system length, width, and volume for better water and nutrient uptake, water-use efficiency, and yield in semi-arid regions (Felker et al. 1997).

Relationships among above-ground and below-ground traits

In this study, the relationships among above- and below-ground traits were compared for all combinations of traits. The highest correlation coefficients were for plant biomass with above-ground traits such as the number of primary cladodes per plant ($r = 0.853$), primary cladode biomass ($r = 0.971$), secondary cladodes per plant ($r = 0.923$), secondary cladode width ($r = 0.830$), secondary cladode biomass ($r = 0.992$), plant height ($r = 0.815$), and canopy width ($r = 0.810$; Table 5). These results indicate positive relationships between plant biomass and its contributing above-ground biomass traits.

In a similar study, Neder et al. (2013) and Alves et al. (2016) also reported significant positive associations of cladode number and width with green matter yield. However, above-ground traits such as primary cladode width and canopy width were negatively correlated with both below-ground traits (i.e. root system biomass and root width) but positively correlated with root system diameter and root surface area. Additionally, a positive correlation was observed between above-ground traits and plant biomass. The combination of traits with $r > 0.70$ indicates a beneficial association (Skinner et al. 1999). In the present study, there were 31 trait combinations with $r > 0.70$ (Table 5). These traits can be considered for development of an ideal plant accession targeting biomass yield.

Regarding below-ground traits, positive associations among all trait combinations were observed, whereas root traits had the weakest relationships with plant biomass. The below-ground traits of root system volume and length had significant positive correlations with root system biomass. The other root traits (e.g. root system surface area $r = 0.95$ and volume $r = 0.74$) had positive correlations with root system length ($P < 0.05$). Therefore, these traits can be considered for the selection of genotypes that are adaptable to dry conditions (Snyman 2007). These associations between above- and below-ground traits suggest that traits such as the number of primary cladodes per plant, primary cladode biomass, number of secondary cladodes per plant, secondary cladode biomass, plant height, and plant biomass may be considered for developing accessions with high biomass.

Morphological variability for above- and below-ground traits in cactus pear accessions

Hierarchical cluster analysis (HCA)

The HCA based on 17 morphological traits resulted in two main clusters for the 15 cactus pear accessions. Cluster II was the largest and with 12 accessions, while Cluster I had three accessions (Figure 4). Cluster I consisted of accessions Red San Cono, Yellow Roccapalumba, and Yellow Santa Margherita Belice. Cluster II consisted of accessions Algerian, Blue Motto, Morado, Red Roccapalumba, Roly Poly, Seedless Roccapalumba, Trunzara Red Bronte, White Roccapalumba, White San Cono, Yellow San Cono, and Zastron. The HCA grouped the 15 accessions independently of their geographical origin. Bendhifi et al. (2013) reported similar results in their study on genetic diversity of 28 cactus pear ecotypes using morphological traits. The accessions under Cluster I represented greater CP content ($>10\%$), lower fiber content (i.e. NDF, ADF, lignin and cellulose), high plant height, biomass and root volume. All the traits in Cluster I could be useful for higher fodder production.




Figure 4. Ward's clustering dendrogram of 15 cactus pear accessions based on 17 above-ground and below-ground traits.

Principal component analysis

A principal component analysis (PCA) was performed to check the relationship between above- and below-ground traits and the results are presented in [Figure 5](#). The biplot indicates that dimension 1 (PC1), PC2, PC3 and PC4 accounted for 49.70%, 16.00%, 9.60% and 6.40% of the total variance. PC3 and PC4 accounted for negligible variation. In the biplot ([Figure 5](#)), the strength of the different variables is indicated with arrows and illustrate distance from the origin. Above-ground traits such as primary cladode per plant (no.) (MCP), primary cladode length (MCL), primary cladode width (MCW), primary cladode biomass (MCB), secondary cladode per plant (no.) (SCPP), secondary cladode length (SCL), secondary cladode width (SCW), secondary cladode biomass (SCB), plant height (PH), canopy width (CW), and plant biomass (PB) were highly related to the accessions Yellow Santa Margherita Belice, Algerian, Red San Cono and White Roccapalumba. All these traits are closely associated.

Below-ground traits like root biomass (RB), root diameter (RD), root width (RW), root volume (RV), root surface area (RSA), and root length (RL) were closely associated with Seedless Roccapalumba, Morado and Yellow Santa Margherita Belice. The PC1 loadings showed that above-ground traits were important for explaining morphological variability in the accessions (number of primary cladodes per plant: 0.90, primary cladode biomass: 0.90, secondary cladodes per plant: 0.82, secondary cladode width: 0.93, secondary cladode biomass: 0.92, plant height: 0.81, canopy width: 0.80, and plant biomass: 0.90; [Table 6](#)). Gallegos-Vázquez et al. (2012), reported that cladode length had a potential role in distinguishing 21 accessions from Mexico. Alves et al. (2016) also noted cladode fresh weight as a useful trait in explaining multivariate polymorphism in *Opuntia* and *Nopalea* species. It is of interest to observe the separation of accessions based on primary cladode length, primary cladode width, number of secondary cladodes per plant, and canopy width ([Figure 5](#)). Loadings for primary cladode biomass and the number of secondary cladodes per plant were positive for the first four PCs, indicating they play a role in variation in the accessions. The below-ground trait root width (0.70) contributed to the highest variation for PC4, which accounted for 6.43% of the total variation ([Table 6](#)). PC2 and PC3 showed that the traits root system surface area, length, and biomass made




Figure 5. PCA plots of 15 cactus pear accessions (MCPP, primary cladode per plant (no.); MCL, primary cladode length; MCW, primary cladode width; MCB, primary cladode biomass; SCPP, secondary cladode per plant (no.); SCL, secondary cladode length; SCW, secondary cladode width; SCB, secondary cladode biomass; PH, plant height; CW, canopy width; PB, plant biomass; RB, root biomass; RD, root diameter; RW, root width; RV, root volume; RSA, root surface area; RL, root length).

Table 6. Eigenvalues, component loading, and percent variation explained by the first four PC components (PC1-PC4) in cactus accessions.

Characteristics	PC1	PC2	PC3	PC4
Eigen value	8.41	2.70	1.61	1.00
Variation explained (%)	49.71	15.91	9.62	6.40
Primary cladodes per plant (no.)	0.90	0.22	0.11	-0.10
Primary cladode length (cm)	0.72	0.23	-0.50	0.10
Primary cladode width (cm)	0.71	0.31	-0.40	0.21
Primary cladode biomass (g)	0.90	0.20	0.11	0.01
Secondary cladodes per plant (no.)	0.82	0.30	0.32	0.02
Secondary cladode length (cm)	0.61	0.04	-0.51	-0.22
Secondary cladode width (cm)	0.93	0.11	-0.11	-0.04
Secondary cladode biomass (g)	0.92	-0.11	0.31	0.21
Plant height (cm)	0.81	0.11	0.32	-0.22
Canopy width (cm)	0.80	0.31	-0.21	-0.05
Plant biomass (g)	0.90	-0.01	0.32	0.11
Root diameter (cm)	0.40	-0.41	0.31	0.11
Root width (cm)	0.20	-0.61	0.05	0.70
Root volume (cm ³)	0.50	-0.41	0.31	-0.08
Root surface area (cm ²)	0.41	-0.70	-0.22	-0.40
Root length (cm)	0.52	-0.80	-0.04	-0.30
Root biomass (g)	0.21	-0.70	-0.41	0.20

no contribution to variation in cactus accessions. The information on genetic variability will help to identify genetically diverse accessions within species for a range of useful traits (root depth, root volume, plant height, cladode number, biomass, and nutrient composition).

Conclusion

This study identified the most promising cactus pear accessions for fodder production in water-scarce and dry weather conditions of India. The results showed that the new cactus pear accessions introduced had potential morphological variability and genetic diversity, indicating their adaptation ability to the extreme weather conditions of India, based on principal component analysis and hierarchical cluster analysis of morphological traits. The observed above-ground morphological traits, except secondary cladode length, showed medium to very high (0.619–0.992) positive, statistically significant ($P < 0.05$) correlations with plant biomass yield, but weaker (0.129–0.507) statistically insignificant correlations were found between below-ground traits and plant biomass. Hierarchical cluster analysis showed that Cluster I represented greater CP content (>10%), lower fiber content (i.e. NDF, ADF, lignin and cellulose), high plant height, biomass and root volume, thus all the traits in Cluster I could be useful for higher fodder production. The principal component analysis results further showed that the observed separation of accessions based on primary cladode length, primary cladode width, number of secondary cladodes per plant, and canopy width could be of interest for selecting suitable accessions for cultivation. According to the coefficient of variation, traits such as cladode length, cladode width, moisture content, dry matter, neutral detergent fibre, acid detergent fibre, cellulose, lignin, ash, and crude protein classified as traits with the least variation (CV <10%), survival (%), yield and *in vitro* dry matter digestibility as traits with medium variation (CV 10–20%), primary cladode width and secondary cladode width as traits with high variation (CV 20–30%) and plant height, plant biomass, primary cladode numbers, secondary cladode numbers, root weight, root volume, root surface, root length and root width as traits with very high variation (CV >30%).

Disclosure statement

The authors do not have any conflicts of interest regarding this manuscript

Funding

This work was implemented by the Indian Grassland and Forage Research Institute (IGFRI) in collaboration with the International Center for Agricultural Research in the Dry Areas (ICARDA). The funding was provided by the Indian Council of Agricultural Research (ICAR) within the framework of the Consultative Group on International Agricultural Research (CGIAR) under grant No. 200091. The authors extend their appreciation to the CGIAR Research Program on Livestock Agri-Food Systems and the CGIAR.

ORCID

Sunil Kumar  <http://orcid.org/0000-0001-6427-5132>
Mounir Louhaichi  <http://orcid.org/0000-0002-4543-7631>
Sawsan Hassan  <http://orcid.org/0000-0002-5057-8957>
Giorgia Liguori  <http://orcid.org/0000-0002-5713-1445>
Prabhu Govindasamy  <http://orcid.org/0000-0003-4678-7809>

References

- Acevedo E, Badilla I, Nobel PS. 1983. Water relations, diurnal acidity changes, and productivity of a cultivated cactus, *Opuntia ficus-indica*. *Plant Physiol.* 72(3):775–780. doi:10.1104/pp.72.3.775.
- Alves FAL, de Andrade AP, Bruno RDLA, Dos Santos DC, Pereira VLA. 2016. Study of the genetic variability, correlation and importance of phenotypic characteristics in cactus pear (*Opuntia* and *Nopalea*). *Afr J Agric Res.* 11:2849–2859. doi:10.5897/AJAR2016.11011.
- AOAC. 2007. Association of Official Analytical Chemists. Official methods of analysis of AOAC international. 18th ed. Washington DC (USA).

- Bakali A. 2013. Impact of irrigation frequencies on the installation and the production of two forms of *Opuntia ficus-indica*. Acta Horti. 995(995):145–156. doi:10.17660/ActaHortic.2013.995.18.
- Bendhif M, Baraket G, Zourgui L, Souid S, Salhi-Hannachi A. 2013. Assessment of genetic diversity of tunisian barbary fig (*Opuntia ficus indica*) cultivars by RAPD markers and morphological traits. Sci Horti. (158):1–7.
- Dev R, Singh JP, Singh T, Dayal D. 2018. Effect of shade levels on growth, and biomass production of cactus (*Opuntia ficus-indica* (L.) Mill.). Int J Curr Microbiol Appl Sci. 7:3145–3153.
- Dubeux JCB Jr, Dos Santos Job F, Andrade Lirab MV, Dos Santos Mde C, Lima DLE, Ferreira RLC, Ferreira RLC. 2006. Productivity of *Opuntia ficus-indica* (L.) Miller under different N and P fertilization and plant population in north-east Brazil. J Arid Environ. 67(3):357–372. doi:10.1016/j.jaridenv.2006.02.015.
- Felker P, de C RS, Casoliba R, Filippini P, Medina D, Zapata R. 2005. Comparison of *Opuntia ficus-indica* varieties of Mexican and Argentine origin for fruit yield and quality in Argentina. J Arid Environ. 60(3):405–422. doi:10.1016/j.jaridenv.2004.06.003.
- Felker P, Singh GB, Parcek OP. 1997. Opportunities for development of cactus (*Opuntia* spp.) in arid- and semi-arid regions. Ann Arid Zone. 36:267–278.
- Flores HA, Murillo SM, Borrego EF, and Rodríguez OJL. 1995. Variación de la composición química en estratos de la planta de 20 variedades de nopal. In: Pimienta-Barrios E, Neri-Luna C, Muñoz-Urías A, and Huerta FM, editors. Memorias del 6o Congreso Nacional y 4o Internacional sobre et Conocimiento y Aprovechamiento del Nopal. Jalisco: México 110–115. <https://www.redalyc.org/pdf/4555/455545052014.pdf>
- Gallegos-Vázquez C, Scheinvar L, Núñez-Colín CA, Mondragón-Jacobo C. 2012. Morphological diversity of xoconostles (*Opuntia* spp.) or acidic cactus pears: a Mexican contribution to functional foods. Fruits. 67(2):109–120. doi:10.1051/fruits/2012001.
- Gebretsadik G, Anmut G, Tegegne F. 2013. Assessment of the potential of cactus pear (*Opuntia ficus indica*) as livestock feed in Northern Ethiopia. Livest Res Rural Dev. 25:1–10.
- Gregory RA, Felker P. 1992. Crude protein and phosphorus contents of eight contrasting *Opuntia* forage clones. J Arid Environ. 22(4):323–331. doi:10.1016/S0140-1963(18)30574-3.
- Guevara JC, Felker P, Balzarini MG, Páez SA, Estevez OR, Paez MN, Antúnez JC. 2011. Productivity, cold hardiness and forage quality of spineless progeny of the *Opuntia ficus-indica* 1281 x *O. Lindhei merii* 1250 cross in Mendoza plain, Argentina. J Prof Ass Cactus Dev. 13:48–62.
- Han H, Felker P. 1997. Field validation of water-use efficiency of the CAM plant *Opuntia ellisiana* in south Texas. J Arid Environ. 36(1):133–148. doi:10.1006/jare.1996.0202.
- Hassan S, Liguori G, Inglese P, Louhaichi M, Sortino G. 2020. The effect of soil volume availability on *Opuntia ficus-indica* canopy and root growth. Agron. 10(5):635. doi:10.3390/agronomy10050635.
- Inglese P, Barbera G, La Mantia T. 1995. Research strategies for the improvement of cactuspear (*Opuntia ficus-indica*) fruit quality and production. J Arid Environ. 29(4):455–468. doi:10.1016/S0140-1963(95)80018-2.
- Inglese P, Basile F, Schirra M. 2002. Cactus pear fruit production. In: Nobel PS, editor. Cacti Biology and Uses. Berkeley (USA): University of California Press; p. 163–183.
- Inglese P, Mondragon C, Nefzaoui A, and Saenz C. 2017. Crop ecology, cultivation and uses of cactus pear, food and agriculture organization of the United Nations (FAO). Rome (Italy). <https://www.fao.org/3/I7012EN/i7012en.pdf>
- Kassambara A, Mundt F 2017. Factoextra: extract and visualize the results of multivariate data analyses. R package version 1.0.5. <https://CRAN.R-project.org/package=factoextra> .
- Koç N, Acar R 2015. The soil factors which effect on root growth in forage plants. In 2nd ICSAE 2015, International Conference on Sustainable Agriculture and Environment, September 30 2015 Oct 03, Konya, Turkey. Proceedings book, volume I & II (pp. 636–642). Selcuk University.
- Kumar S, Dayal D, Shamsudheen M, Machiwal D, Yadav OP. 2017. Cactus pear (*Opuntia ficus-indica*) in India. Jodhpur: ICAR-Central Arid Zone Research Institute.
- Kumar S, Louhaichi M, Dana Ram P, Tirumala KK, Ahmad S, Rai AK, Sarker A, Hassan S, Liguori G, Probir Kumar G, et al. 2021. Cactus pear (*Opuntia ficus-indica*) productivity, proximal composition and soil parameters as affected by planting time and agronomic management in a semi-arid region of India. Agron. 11(8):1647. doi:10.3390/agronomy11081647.
- Lima GFDC, Rego M, Teles M, Dantas FDG, Lobo RNB, Silva JGMD, Aguiar EMD. 2016. Morphological characteristics and forage productivity of irrigated cactus pear under different cutting intensities. Revista Caatinga. 29(2):481–488. doi:10.1590/1983-21252016v29n226rc.
- Lopez MJL, Cruz HP, López AJ. 1988. Contenidos nutrimentales de tallos y raíces de nopal tunero *Opuntia amyoclaea* Tenore. Revista Chapingo. 62–63:143–148.
- Louhaichi M, Kumar S, Tiwari S, Islam M, Hassan S, Yadav OP, Dayal D, Moyo HP, Dev R, Sarker A. 2018. Adoption and utilization of cactus pear in South Asia—Smallholder farmers’ perceptions. Sustainability. 10(10):3625. doi:10.3390/su10103625.
- Misra AK, Kumar S, Kumar TK, Ahmed S, Palsaniya DR, Ghosh PK, Louhaichi M, Sarker A, Hassan S, Ates S. 2018. Nutrient intake and utilization in sheep fed *Opuntia* [*Opuntia ficus-indica* (L.) Mill.] combined with conventional green and dry fodders. Range Manag Agrofor. 39:97–102.

- Neder DG, Costa FRD, Edvan RL, Souto Filho LT. 2013. Correlations and path analysis of morphological and yield traits of cactus pear accessions. *Crop Breed Appl Biotechnol.* 13(3):203–207. doi:10.1590/S1984-70332013000300009.
- Nieto-Garibay A, Troyo-Diéguez E, Valdez-Cepeda RD, Rueda-Puente E, García-Hernández JL, Flores-Hernández A, Zamora-Salgado S, Orona-Castillo I, Murillo-Amador B. 2011. Plant height and mineral content of *Opuntia tapona* growing along the coasts of Baja California Sur, México. *J Prof Ass Cactus Dev.* 13:77–87.
- Nobel PS. 2001. *Biología Ambiental.* In *Agroecología, Cultivo e Usos da Palma forrageira* (Eds. Barbera G, Inglesa P, Pimienta-Barrios E). Roma: FAO/Sebrae-PB; p. 36–48.
- Nobel PS, Geller GN, Kee SC, Zimmerman AD. 1986. Temperatures and thermal tolerance for cacti exposed to high temperatures near the soil surface. *Plant Cell Environ.* 9:279–287.
- Oelofse RM, Labuschagne MT, Potgieter JP. 2006. Plant and fruit characteristics of cactus pear (*Opuntia* spp.) cultivars in South Africa. *J Sci Food Agric.* 86(12):1921–1925. doi:10.1002/jsfa.2564.
- Pareek OP, Singh RS, Vashishtha BB. 2003. Performance of cactus pear (*Opuntia ficus-indica* (L.) Mill.) clones in hot arid region of India. *J Prof Ass Cactus Dev.* 5:121–130.
- Prasad M, Chaudhary M, Ramakrishnan S, Singh HV, Kumar S, Ghosh PK. 2021. Impact of long term land use changes on carbon dynamics and soil quality in the semi arid bundelkhand region of India. *Fresenius Environ Bull.* 30:2536–2544.
- R Core Team. 2019. R: a language and environment for statistical computing. R foundation for statistical computing. Vienna: R Foundation for Statistical Computing. <https://www.R-project.org>.
- Rayamajhi N, Sharma J. 2018. Genetic diversity and structure of a rare endemic cactus and an assessment of its genetic relationship with a more common congener. *Genetica.* 146(3):329–340. doi:10.1007/s10709-018-0024-0.
- Samah S, Cvdt P, Cruz MAS, Valadez-Moctezuma E. 2016. Genetic diversity, genotype discrimination, and population structure of Mexican *Opuntia* sp., determined by SSR markers. *Plant Mo Biol Rep.* 34(1):146–159. doi:10.1007/s11105-015-0908-4.
- Santos DC, Lira Md A, Dias FM, Farias I, Santos MVF, Santos VF. 2000. Produtividade de cultivares de palma forrageira (*Opuntia e Nopalea*). In: *Proceedings of the 7th North eastern Ruminant Feeding Symposium.* SNPA: Teresina, PI, Brazil, 121–123.
- SAS Institute. 2011. SAS/STAT. version 8.02 for windows. Cary (NC, USA):SAS Institute, Inc.
- Sigmaplot. 1996. Sigma graphs, version 10. San Jose (California, USA):Systat Software, Inc.
- Sirohi SK, Karim SA, Misra AK. 1997. Nutrient intake and utilization in sheep fed with prickly pear cactus. *J Arid Environ.* 36(1):161–166. doi:10.1006/jare.1996.0184.
- Skinner DZ, Bauchan GR, Auricht G, Hughes S. 1999. A method for the efficient management and utilization of large germplasm collections. *Crop Sci.* 39(4):1237–1242. doi:10.2135/cropsci1999.0011183X003900040046x.
- Snyman HA. 2005. A case study on in situ rooting profiles and water-use efficiency of *Opuntia ficus-indica* and *O. robusta*. *J Prof Ass Cactus Dev.* 7:1–21.
- Snyman HA. 2006. Root distribution with changes in distance and depth of two-year-old cactus pears *Opuntia ficus-indica* and *O. robusta* plants. *South Afr J Bot.* 72(3):434–441. doi:10.1016/j.sajb.2005.12.008.
- Snyman HA. 2007. Root studies on cactus pears *Opuntia ficus-indica* and *O. robusta* along a soil–water gradient. *Haseltonia.* 13:64–75. doi:10.2985/1070-0048(2007)13[64:RSOCPO]2.0.CO;2.
- Snyman HA. 2013. Growth rate and water-use efficiency of cactus pears *Opuntia ficus-indica* and *O. robusta*. *Arid Land Res Manag.* 27(4):337–348. doi:10.1080/15324982.2013.771232.
- Stintzing FC, Carle R. 2005. Cactus stems (*Opuntia* spp.): a review on their chemistry, technology, and uses. *Mol Nutr Food Res.* 49(2):175–194. doi:10.1002/mnfr.200400071.
- Tegegne F, Kijora C, Peters KJ. 2007. Study on the optimal level of cactus pear (*Opuntia ficus-indica*) supplementation to sheep and its contribution as source of water. *Small Ruminant Res.* 72(2–3):157–164. doi:10.1016/j.smallrumres.2006.10.004.
- Tilley JMA, Terry RA. 1963. A two-stage technique for the in vitro digestion of forage crops. *Grass Forage Sci.* 18(2):104–111. doi:10.1111/j.1365-2494.1963.tb00335.x.
- Van Soest PV, Robertson JB, Lewis BA. 1991. Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *J Dairy Sci.* 74(10):3583–3597. doi:10.3168/jds.S0022-0302(91)78551-2.