



2011

# Introduction of a 6-cyano group in 14-oxygenated N-methylmorphinans influences in vitro and in vivo pharmacological activities

Valeria Follia  
*University of Innsbruck*

Mario D. Aceto  
*Virginia Commonwealth University, maceto@vcu.edu*

Louis S. Harris  
*Virginia Commonwealth University, harris@vcu.edu*

*See next page for additional authors*

Follow this and additional works at: [http://scholarscompass.vcu.edu/phtx\\_pubs](http://scholarscompass.vcu.edu/phtx_pubs)

© 2011 Follia et al; licensee BioMed Central Ltd. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

---

Downloaded from

[http://scholarscompass.vcu.edu/phtx\\_pubs/5](http://scholarscompass.vcu.edu/phtx_pubs/5)

This Article is brought to you for free and open access by the Dept. of Pharmacology and Toxicology at VCU Scholars Compass. It has been accepted for inclusion in Pharmacology and Toxicology Publications by an authorized administrator of VCU Scholars Compass. For more information, please contact [libcompass@vcu.edu](mailto:libcompass@vcu.edu).

---

**Authors**

Valeria Follia, Mario D. Aceto, Louis S. Harris, Andrew Coop, Helmut Schmidhammer, and Mariana Spetea

MEETING ABSTRACT

Open Access

# Introduction of a 6-cyano group in 14-oxygenated *N*-methylnorphinans influences *in vitro* and *in vivo* pharmacological activities

Valeria Follia<sup>1</sup>, Mario D Aceto<sup>2</sup>, Louis S Harris<sup>2</sup>, Andrew Coop<sup>3</sup>, Helmut Schmidhammer<sup>1</sup>, Mariana Spetea<sup>1\*</sup>

From 17th Scientific Symposium of the Austrian Pharmacological Society (APHAR). Joint meeting with the Hungarian Society of Experimental and Clinical Pharmacology (MFT) Innsbruck, Austria. 29-30 September 2011

## Background

Being a disabling symptom of many medical conditions, effective pain control is one of the most important therapeutic priorities. Morphine and other opioid drugs produce analgesia primarily through  $\mu$  opioid (MOP) receptors, which mediate beneficial but also the non-beneficial actions. Appropriate identification of novel opioid analgesics may reduce complications and improve patient compliance. It was reported that hydrazones, oximes, carbazones and semicarbazone derivatives of morphinan-6-ones, e.g. dihydromorphinone or oxymorphone, exhibit high affinity at the MOP receptor [1]. Since most of these structures show high antinociceptive potency while having less pronounced side effects, it remains a promising task to convert the carbonyl group of morphinan-6-ones into various functionalities. In this study, we aimed to investigate the effect of the replacement of the 6-keto function with a 6-cyano group on *in vitro* and *in vivo* pharmacological profiles.

## Methods

Binding affinities at opioid receptors were determined using competition binding assays in rodent brain membranes. *In vitro* [<sup>35</sup>S]GTP $\gamma$ S functional assays were performed with Chinese hamster ovary (CHO) cell membranes expressing human opioid receptors. Antinociceptive activities were assessed in mice using tail-flick, hot-plate and writhing tests.

## Results

Replacement of the 6-keto group by a 6-cyano substituent in *N*-methylnorphinan-6-ones leads to qualitative and quantitative differences in the interaction with opioid receptors. Consequently, we have conducted a comparison of the biological activities of the 6-cyanomorphinans to those of structurally-related opioids, oxycodone, oxymorphone and of the clinically relevant morphine. The 6-cyanomorphinans displayed high affinity and behaved as agonists at the MOP receptor. When tested *in vivo*, they acted as potent antinociceptive agents after subcutaneous administration, being more active than the 6-keto analogues. The presence of a 14-methoxy or a 14-cinnamyloxy group instead of a hydroxy group not only increased *in vitro* opioid activity at the MOP receptor, but also enhanced the antinociceptive potency.

## Conclusions

Our findings revealed that targeting position 6 in the morphinan skeleton represents a viable approach for tuning the pharmacological properties of this class of opioids. Appropriate molecular manipulations could afford ligands that, besides their scientific value as pharmacological tools, may also have the potential of emerging as novel analgesics with fewer side effects compared to currently available treatments.

## Acknowledgements

Supported by the Austrian Science Fund (FWF: TRP 19-B18) and College on Problems of Drug Dependence of the USA (N01DA-1-7725).

## Author details

<sup>1</sup>Department of Pharmaceutical Chemistry, Institute of Pharmacy and Center for Molecular Biosciences, University of Innsbruck, 6020 Innsbruck, Austria.

<sup>2</sup>Department of Pharmacology and Toxicology, Medical College of Virginia,

\* Correspondence: mariana.spetea@uibk.ac.at

<sup>1</sup>Department of Pharmaceutical Chemistry, Institute of Pharmacy and Center for Molecular Biosciences, University of Innsbruck, 6020 Innsbruck, Austria  
Full list of author information is available at the end of the article

Virginia Commonwealth University, Richmond, VA 23298-0613, USA.  
<sup>3</sup>Department of Pharmaceutical Sciences, University of Maryland, School of Pharmacy, Baltimore, MD 21201, USA.

Published: 5 September 2011

#### Reference

1. Schmidhammer H, Spetea M: **Synthesis of 14-alkoxymorphinan derivatives and their pharmacological actions.** *Top Curr Chem* 2011, **299**:63-91.

doi:10.1186/1471-2210-11-S2-A26

**Cite this article as:** Follia *et al.*: Introduction of a 6-cyano group in 14-oxygenated *N*-methyldmorphinans influences *in vitro* and *in vivo* pharmacological activities. *BMC Pharmacology* 2011 **11**(Suppl 2):A26.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

