

2015

Spatial Analysis of Nest Productivity and Predation in Prothonotary Warblers

Miranda Foster
fostermj2@vcu.edu

Lesley P. Bulluck
VCU, lpbulluck@vcu.edu

Follow this and additional works at: <http://scholarscompass.vcu.edu/uressposters>

© The Author(s)

Downloaded from

Foster, Miranda and Bulluck, Lesley P., "Spatial Analysis of Nest Productivity and Predation in Prothonotary Warblers" (2015).
Undergraduate Research Posters. Poster 167.
<http://scholarscompass.vcu.edu/uressposters/167>

This Book is brought to you for free and open access by the Undergraduate Research Opportunities Program at VCU Scholars Compass. It has been accepted for inclusion in Undergraduate Research Posters by an authorized administrator of VCU Scholars Compass. For more information, please contact libcompass@vcu.edu.

Spatial Analysis of Nest Productivity and Predation in Prothonotary Warblers

M.J. FOSTER² AND LESLEY BULLUCK¹⁻³

¹Department of Biology, ²Center for Environmental Studies, ³Life Sciences, Virginia Commonwealth University

BACKGROUND

- A Prothonotary Warbler is a migratory songbird that breeds in the Eastern US and overwinters in Central and S. America.
- Prothonotaries are secondary cavity nesters, meaning they take advantage of hollows that they do not create on their own. They prefer to nest over water, as it reduces predation risk.
- In the past nest productivity and predation has been analyzed assuming that observations are independent from each other in space
- In larger ecological systems, there can be spatial autocorrelation whereby data collected in nearby areas are not independent of one another which violates the assumption of most statistical tests

OBJECTIVES

- We investigated whether Prothonotary Warbler nest productivity and nest predation are spatially autocorrelated at Deep Bottom Park along the lower James River, VA.
- We analyzed spatial autocorrelation for two main reasons:
 1. To eliminate error, accounting for a possible nuisance variable that could give false positives.
 2. To better understand the spatial structure of the nest boxes at Deep Bottom Park.



METHODS AND RESULTS

- Nest success and predation were monitored at 57 different nest boxes at Deep Bottom Park over a period of four years (2009-2013).
- Spatial Autocorrelation (Moran's I) was run using ArcGIS version 10.2.1, and points were projected using World Geodetic System (WGS) 1984 coordinates.
- Nest success (total number of young fledged) (p value = .345) and predation (p =value.768) were found to be not spatially auto correlated.




Fig. 1
Total number of young fledged At 57 nest boxes at Deep Bottom Park from 2009-2013.

CONCLUSIONS

- Because neither nest success or predation is spatially autocorrelated, our observations are independent from each other in space.
- Spatial variables are not driving nest success or predation and individual bird quality is likely the main driver for these differences.
- This is most likely because of the relative similarity of habitats at the nest boxes as well as the limited size of the site.
- Future studies could compare boxes over land vs. boxes placed over water or at a larger continuous nest site.

ACKNOWLEDGEMENTS

Thanks to all the volunteers on the PROW Project especially Ethan and Ashley. Thanks to Jenna Dodson and Nik Moy the graduate students working on the project. Thanks to HSURP for providing me with the opportunity to continue my research, and finally to Dr. Lesley Bulluck for mentoring me and providing priceless guidance

