

2015

Vegetation Structure of Mangrove Ecosystems in Panama

Christopher L. Arrington

Virginia Commonwealth University, arringtoncl@vcu.edu

Cole Gandee

Virginia Commonwealth University, gandeecc@vcu.edu

Spencer R. Parkinson

Virginia Commonwealth University, parkinsonsr@vcu.edu

Follow this and additional works at: http://scholarscompass.vcu.edu/rice_symp

 Part of the [Terrestrial and Aquatic Ecology Commons](#)

© The Author

Downloaded from

http://scholarscompass.vcu.edu/rice_symp/2

This Poster is brought to you for free and open access by the Rice Rivers Center at VCU Scholars Compass. It has been accepted for inclusion in Rice Rivers Center Research Symposium by an authorized administrator of VCU Scholars Compass. For more information, please contact libcompass@vcu.edu.

Vegetation Structure of Mangrove Ecosystems in Panama

Christopher Arrington, Cole Gandee, Spencer Parkinson
Virginia Commonwealth University, Center for Environmental Studies



Introduction

- Mangroves provide important habitat for terrestrial and marine wildlife
- They buffer shorelines from flooding and sequester excess nutrients and pollutants in runoff before reaching rivers and oceans
- They provide a wintering habitat for migratory bird species
- These habitats are being rapidly lost to coastal development




Objectives

- Quantify habitat differences among mangrove sites by
- Quantifying differences in forest structure and tree species composition
 - Determining if these differences are related to site age
 - Analyzing differences in moisture, salinity, and pH across sites

Methods

- 20 meter transects walked diagonally from center of mist nets. Recorded individual tree species, tree height and diameter at breast height (DBH), standing water, percent canopy cover, and leaf litter
- Ocular tube readings every 2 meters for estimated % area cover
- pH and temperature were taken by Hanna Meter in standing water; holes were dug in drier areas until standing water was reached
- Salinity sampled with refractometer
- Canopy height and DBH recorded using a rangefinder and DBH tape



Results




Figure 1: ANOVA comparing salinity levels in ppt by sites. JD Mature East, JD Young, and PV Young are located on Pacific coast while G Mature is on the Caribbean coast

Color: mature (green), intermediate (blue), young (purple)
G Mature: Galeta (Caribbean/light green)
JD Mature East: Juan Diaz 2014 - 2015
JD Mature West: Juan Diaz 2013 - 2014
JD Intermed: Juan Diaz 2010 - 2013
JD Young: Juan Diaz 2013 - 2014
PV Young: Panama 2010 - 2012, 2013 - 2014





Figure 2 (top right): ANOVA comparing the average canopy height by site

Figure 3 (bottom right): ANOVA comparing the average diameter at breast height of tree trunks by site

Conclusions

- Average canopy height and diameter at breast height were higher in mature sites.
- Canopy cover didn't vary across sites - younger stands have more, smaller trees while mature sites have fewer, but larger trees.
- Standing water was greater at the Caribbean site possibly due to rainfall gradient across Panama from north to south (Caribbean to Pacific). This may also explain the significantly higher salinity of standing water at the Pacific sites compared to the Caribbean site Galeta.

Acknowledgements

- Thanks to:
- Panama Audubon Society
 - VCU Rice Center
 - Lesley Bulluck, Catherine Viverette, Edward Crawford, Nik Moy, Jenna Dodson
 - Guido Berguido and guides



Bienvenidos - Welcome