

Virginia Commonwealth University VCU Scholars Compass

Rice Rivers Center Research Symposium

Rice Rivers Center

2015

Behavioral Observations: Foraging and Aggression of Neotropical Migrants and Resident Birds in Panama

Jessica M. Johnston Virginia Commonwealth University, johnstonjm2@vcu.edu

Samuel B. King Virginia Commonwealth University, kingsb3@vcu.edu

Ryan A. Levering Virginia Commonwealth University, leveringra@vcu.edu

Follow this and additional works at: http://scholarscompass.vcu.edu/rice_symp Part of the <u>Behavior and Ethology Commons</u>, and the <u>Ornithology Commons</u>

© The Author

Downloaded from http://scholarscompass.vcu.edu/rice_symp/6

This Poster is brought to you for free and open access by the Rice Rivers Center at VCU Scholars Compass. It has been accepted for inclusion in Rice Rivers Center Research Symposium by an authorized administrator of VCU Scholars Compass. For more information, please contact libcompass@vcu.edu.

Behavioral Observations: Foraging and Aggression of Neotropical Migrants and Resident Birds in Panama Jessica M. Johnston, Samuel B. King & Ryan A. Levering

Introduction

- Lyons suggests that foraging data in songbirds can be an indicator of food abundance , with higher attack rate= higher habitat quality; higher foraging speed= lower habitat quality (2005)
- Territorial behavior such as interspecific aggression among songbirds may also be related to food availability. Previous studies have documented variation in the extent of territorial and aggressive behaviors among resident and Neotropical migratory species on tropical wintering grounds.






Objectives

- Gather foraging and aggression data at two coastal angroves sites in central Panama, one on the Caribbean Coast (Galeta) and one on the Pacific Coast (Juan Diaz-East)
- Use foraging data as an indicator of habitat quality and food availability
- Perform aggression trials in order to compare territorial behaviors among resident Mangrove Warblers (YWAR) and two Neotropical migrant species, the Prothonotary Warbler (PROW), and the Northern Waterthrush (NOWA).
- Determine whether or not territorial behavior is shown and if habitat quality plays a role in these behaviors

Methods

- Random focal observations were performed to observe feeding behaviors of YWAR, PROW, and NOWA, measuring all movements, maneuvers, and non-foraging movements.
- Foraging speed was calculated by examining the number of movements per minute; attack rate was calculated by examining the number of maneuvers per minute.
- Random focal observations were done to examine territorial behaviors of Neotropical birds, developing a numerical score of 1 to 4 to describe an individual's aggressive response to playback and decoy trials.
- A second aggression score (Score B) was developed to incorporate flocking behaviors into the original overall score.


Virginia Commonwealth University, Department of Biology and Center for Environmental Studies



Conclusions

- Although the attack rate did not vary between sites, the Prothonotary Warbler displayed a higher foraging speed at the Pacific site, which may suggest lower habitat quality.
- Unlike previous studies, our results indicate YWAR, PROW, more research is required to examine if flocking behaviors are in fact aggressive responses indicative of territorial behavior.
- Refinement and standardization of methods for aggression trials will enhance research going forward.

and NOWA display similar behaviors, including flocking, but



Acknowledgements

We'd like to thank: Panama Audubon Society VCU Rice Center • Lesley Bulluck, Catherine Viverette, Edward Crawford, Nik Moy, Jenna Dodson • Guido Berguido and guides







