



Virginia Commonwealth University
VCU Scholars Compass

Theses and Dissertations

Graduate School

2008

Fluoride Varnish Use Among Dentists in Virginia

Amanda Bowen Kuhn
Virginia Commonwealth University

Follow this and additional works at: <http://scholarscompass.vcu.edu/etd>

 Part of the [Pediatric Dentistry and Pedodontics Commons](#)

© The Author

Downloaded from

<http://scholarscompass.vcu.edu/etd/1180>

This Thesis is brought to you for free and open access by the Graduate School at VCU Scholars Compass. It has been accepted for inclusion in Theses and Dissertations by an authorized administrator of VCU Scholars Compass. For more information, please contact libcompass@vcu.edu.

© Amanda Bowen Kuhn 2008

All Rights Reserved

FLUORIDE VARNISH USE AMONG DENTISTS IN VIRGINIA

A thesis submitted in partial fulfillment of the requirements for the degree of Masters of Science in Dentistry at Virginia Commonwealth University.

by

AMANDA BOWEN KUHN

B.S., University of North Carolina at Chapel Hill, 2002

D.M.D., University of Pittsburgh School of Dental Medicine, 2006

Director: TEGWYN H. BRICKHOUSE D.D.S., PH.D

ASSISTANT PROFESSOR, DEPARTMENT OF PEDIATRIC DENTISTRY

Virginia Commonwealth University
Richmond, Virginia
June 2008

Acknowledgement

I would like to thank Drs. Tegwyn H. Brickhouse and Alvin M. Best for their expertise and assistance in making this research project possible. To my fellow residents, thank you for all of your support and humor day in and day out, to my parents, who from the beginning have encouraged me and given me their unwavering support, to my dear friend Mary Claire, whose love of life was infectious, to my beautiful and sweet daughter Carson, thank you for always having a smile to brighten my day and keep me focused on what is truly important, and finally, to my loving husband Todd, you are my rock and my best friend. I could not have accomplished any of this without you.

Table of Contents

| | Page |
|------------------------------|------|
| Acknowledgements..... | ii |
| List of Tables | iv |
| List of Figures | v |
| Abstract..... | vi |
| Chapter | |
| 1 Introduction..... | 8 |
| 2 Materials and Methods..... | 12 |
| Analysis | 13 |
| 3 Results..... | 14 |
| 4 Discussion..... | 18 |
| Literature Cited..... | 21 |
| Appendices | |
| A Survey | 33 |

List of Tables

| | Page |
|--|------|
| Table 1: Description of Those Surveyed. | 25 |
| Table 2: Description of Practice Characteristics..... | 26 |
| Table 3: Application of Topical Fluoride. | 27 |
| Table 4: Unadjusted Relationships with Topical Fluoride Varnish Use..... | 28 |
| Table 5: Unadjusted Relationships with Fluoride Varnish Use..... | 29 |
| Table 6: Perception Score and Varnish Use. | 30 |

List of Figures

| | Page |
|--|------|
| Figure 1: Relationship with Graduation Year..... | 31 |
| Figure 2: Perception Score and Percentage of Varnish Use | 32 |

Abstract

FLUORIDE VARNISH USE AMONG DENTISTS IN VIRGINIA

By Amanda Bowen Kuhn, B.S., D.M.D.

A thesis submitted in partial fulfillment of the requirements for the degree of Masters of Science in Dentistry at Virginia Commonwealth University.

Virginia Commonwealth University, 2008

Major Director: Tegwyn H. Brickhouse, D.D.S., Ph.D
Assistant Professor, Department of Pediatric Dentistry

Purpose: The purpose of this study was to assess fluoride varnish use by dental practitioners in Virginia.

Methods: Using a cross sectional survey design, all dentists in Virginia who are members of the Virginia Dental Association (VDA) were sent an online survey about usage and knowledge of fluoride varnish.

Results: The majority of the respondents were general dentists (79%) followed by pediatric dentists (12%). Fluoride varnish use increased with year of graduation from dental school. Dentists who thought fluoride varnish was more effective and less time

consuming use it more than other topical fluorides. Dentists who thought their patients prefer fluoride varnish use it more than other topical fluorides.

Conclusion: The majority of dentists are not aware of the advantages of fluoride varnish. However, those who are, choose to use it as opposed to foams and gels. Recent graduates, with more exposure to fluoride varnish, use it more frequently.

Introduction

Topical fluoride application has been the standard of care in American dental offices for many years, especially in the pediatric population. Dentists apply fluoride products for the primary prevention of dental caries and to prevent early carious lesions from progressing.¹ Several forms of topical fluoride exist including foams, gels, dentifrices, rinses, and varnish.² The clinical recommendations for the use of professionally applied topical fluoride by the American Dental Association are broken down into age categories. For children under 6 years of age with low risk, topical fluoride may not be beneficial. For children under 6 years of age with moderate risk, fluoride varnish applications should be received at 6 month intervals and higher risk patients should receive fluoride varnish at 3 to 6 month intervals. Patients 6-18 years of age have the same recommendations as the under 6 year old patients with one exception; the moderate and high risk patients may receive fluoride varnish or gel applications. Fluoride gels are not recommended by the ADA for children under 6 years of age due to the risk of inadvertent ingestion.¹

Fluoride varnish has been widely used for over 25 years in Scandinavia and Europe, however, it was only recently introduced in the United States.³ It was approved by the US Food and Drug Administration (FDA) in 1994 for the treatment of dentin hypersensitivity associated with the exposure of root surfaces or as a cavity varnish,

however, it has not been approved yet for caries prevention.^{1,4} Therefore, using it for caries prevention is currently considered “off-label” since it has not been cleared by the FDA for this purpose.⁵ In spite of this, there exists a vast amount of published data verifying its effectiveness, consequently, there is not a legal risk in using it off label.^{3,6,7,8}

Acidulated phosphate fluoride (APF) gels are the most commonly used topical fluoride in the US, but when compared to fluoride varnish the APF gels have several disadvantages; primarily the bitter taste and risk of over ingestion. Many factors can cause over ingestion including inadequate suction, improper placement of the tray, a surplus of gel, or inability of the patient to expectorate the remaining gel.³ If the patient swallows the gel dose they do so in a short amount of time and this will cause a significant increase in the plasma fluoride concentrations.⁹ On the contrary, fluoride ingestion following a varnish treatment has barely detectable effects on plasma fluoride concentrations due to the much smaller dose swallowed over several hours.¹⁰ The varnish adheres to the enamel for an extended period of time preventing systemic consumption of fluoride in large doses. The advantage of APF foam is that less fluoride is applied to the tray, thus decreasing the amount of fluoride that is swallowed. A study by Whitford and colleagues concluded that APF foam and gel are equivalent with respect to fluoride uptake and that only one-fifth the amount of foam is required to sufficiently cover the teeth.¹¹ The two most commonly used gels include APF, which contains 1.23% or 12,300 parts per million fluoride ion and 2% sodium fluoride (NaF) which contains 0.90% or 9,050 ppm fluoride ion. Varnishes typically contain 5% sodium fluoride which equals 2.26% or 22,600 parts per million fluoride ion.¹ The concentration of fluoride in varnish is almost double that of APF gel,

however, as Bawden pointed out, “the amount of fluoride in the mouth of a child as a result of a varnish application is less than 7 mg compared with 30 mg or more with an APF application.”³ The amount of fluoride retained in the oral cavity after treatment with a 1.23% APF foam was reported to be approximately one-half that retained with a gel;¹² however, the in vitro fluoride uptake was not significantly different.¹¹

The American Dental Association produced evidence-based clinical recommendations on professionally applied topical fluoride which found that fluoride varnish applied every six months is effective in preventing caries in the primary and permanent dentition of children and adolescents.¹ It has also been concluded that two or more applications of fluoride varnish per year are effective in preventing caries in high risk populations¹ and decreasing caries in primary teeth.^{13,14} Repeated applications of varnish at more frequent intervals have been shown to result in greater caries reduction.¹⁵ Weintraub et al found in their two-year randomized clinical trial that fluoride varnish can be used to prevent early childhood caries and reduce caries increment in very young children.⁸ Some of the children in this study were younger than 1 year of age (the American Academy of Pediatric Dentistry’s recommended age for the first dental visit) and yet, they had very little difficulty with cooperation of the infants with fluoride varnish. They also concluded that although frequent applications of varnish were more beneficial, one application was better than none for preventing caries.⁸ Fluoride varnishes have been shown to inhibit demineralization^{16,17,18} and to promote remineralization of enamel.¹⁹

Recently, there has been an increase in the awareness of fluoride varnish due to rapidly expanding number of primary prevention fluoride varnish programs aimed at

reducing levels of Early Childhood Caries (ECC) by targeting dentists, physicians, nurses, and Early Head Start programs. Early Childhood Caries is a significant problem in low-income and minority groups. The young children from these high-risk groups are more likely to seek medical care than dental care.²⁰ To take advantage of this fact, programs like the *Smiles for Ohio Fluoride Varnish Program* were created. These programs reimburse primary care providers for assessing oral health and applying fluoride to the teeth of eligible children.²⁰

Fluoride varnish is safe, fast, and easy to apply, thus being a superior choice for use with young children.²¹ Studies have shown children gag less with fluoride varnish and the cost per varnish application are significantly less when you factor in labor costs.²² Despite the large amount of information supporting the use of fluoride varnish to control caries, many general and pediatric dentists do not use them, instead favoring gels and foams.²³ Fiset et al found general dentists in Washington State cited, “lack of awareness, lack of convincing evidence of a favorable cost:benefit ratio, patients’ rejection of the service, and low caries risk among adult patients” as reasons for not using fluoride varnish.²³ The purpose of this study was to assess the use of fluoride varnish by dental practitioners as well as to determine if there were any differences in varnish use between provider type or the practice’s patient profile.

Materials and Methods

This study was a cross-sectional survey of member dentists of a state dental association. The survey was administered on-line and distributed via a list-serve of subscribed member dentists (n=1528) of the Virginia Dental Association. The survey consisted of a 13-item survey regarding fluoride varnish. The survey was sent using the Inquisite 7.0 web survey software.²⁴ The dentists were asked to complete the survey within 60 days. After the 60 days expired, a reminder email was sent. Dentists were given another 30 days to complete the survey. After the total of 90 days expired, the survey was closed on-line.

The first four questions of the survey were used to gather demographic information about the respondents and also to determine particular practice characteristics to create a practice profile. The survey asked the specialty (if applicable) of the dental practitioner and what percentage of the practices' patient population is made up of children. In addition, the percentage of private insurance, public assistance, and self-pay patients that the respondents treat in their practice was documented.

Questions 5-7 asked respondents about their use of fluoride in different age groups (0-3, 4-6, 6-12, 12-18, 18+) and how often they apply fluoride. The survey then asked what type of fluoride was used: foam, gel, varnish, other, or none was also noted. If the

respondent answered “none” they were thanked for their responses and excluded from the survey.

Questions 8-12 tested the respondents’ knowledge of fluoride varnish and how it compares to the other topical fluorides. They were asked about price, effectiveness, efficiency, systemic exposure, and patient preference. Question 13 asked the respondent to insert their email address and comments if they were interested in further research and continuing education.

Analysis

Univariate distributions were obtained for each question. Percents for all items were based on the total number of respondents. The percentages regarding each questionnaire item were analyzed and comparisons made using a chi-squared test. The association between dental practitioner type, practice profile, and fluoride use was tested using chi-square or Fisher’s exact test. The use of fluoride varnish across different age groups was modeled using logistic regression (SAS PROC GENMOD).

Results

Of the 1528 dentists surveyed, a total of 243 surveys were completed yielding a gross response rate of 16%. Of these 243 respondents, 14 filled out the survey twice and one survey was blank. After excluding the duplicate surveys and the blank survey, 228 surveys were analyzed yielding a final response rate of 15%. The distribution of providers were as follows; general dentists at 78.8% (N=178), pediatric dentists 11.5% (N=26), and other respondents were made up of dental specialists including periodontists (N=10), orthodontists (N=6), oral and maxillofacial surgeons (N=5), and an endodontist (N=1). The majority of the respondents (76%) graduated from dental school from 1970-1999 and the average age of respondents was 49 years old. The survey respondents are described in Table 1.

The respondents were asked in the survey to approximate the percentage of their practice population within the following three age groups: 0 to 6 years of age, between 6 and 18 years of age, and older than 18 years of age. The responses ranged from zero to over 75% in each of these three categories. The largest group of practitioners (81.3%) had 50% or more in the 18 years or older range. Of the remaining 18.7% of practices, 16% has more than 20% of their practice devoted to children under the age of 6. The patient populations are described in Table 2.

Topical fluoride use was first assessed by asking “what type of topical fluoride do you mainly use?” and these responses are shown in Table 3. It was noted that $n = 14$ practitioners did not apply topical fluoride and so these were not included in subsequent analyses. Of the remaining $N=211$ who use topical fluoride, foam was the most popular (37%), followed by varnish (28%), then gel (27%), and finally other topical fluorides (8%).

The objective of this study was to determine what characteristics of dental providers are associated with the use of topical varnish (yes/no). First, the provider characteristics were screened to determine which had a potential relationship. Simple two-way contingency tables are shown in Table 4. The area of dental practice was collapsed into three categories and showed no relationship with the use of fluoride varnish ($p > 0.2$), although there was a trend of more varnish use in pediatric dentists. The year of graduation from dental school was collapsed into decades and indicated a relationship ($p < 0.04$); with more recent graduates indicating fluoride varnish is their preferred topical fluoride. When considered as a continuous variable, year of graduation also indicated a possible relationship (LR chi-square = 3.88, $p = 0.0488$). Age was potentially related to varnish use when considered as a continuous variable (LR chi-square = 2.54, $p = 0.1110$), but when broken down into decades (Table 4), there was no longer an indication of a relationship ($p > 0.6$). Of course, age and year of graduation are strongly correlated ($r = -0.93$), and so we chose to use year of graduation as a predictor of varnish use. Practices with more than a third of their patient pool under 18 years of age, did not appear to use varnish more than practitioners with less children in their practice ($p > 0.4$). The same was

found for practices with more than 10% of their patient pool less than 6 years of age, though this was closer to being significant ($p < 0.12$).

The variables in Table 4 that pass a bivariate screening for inclusion in a multivariate regression model ($p < .1$) were the year of the practitioner's graduation and the percentage of youth in the practice who are less than 6 years of age to be at least 10%. When considered in a logistic regression model, the percentage of youth was still not statistically significant (LR chi-square = 2.9, p-value = 0.0881) but year of graduation remained significant (LR chi-square = 4.28, p-value = 0.0385). The relationship with graduation year is shown in Figure 1.

The next analyses centered on the relationship between the practitioners' perceptions of fluoride varnish as opposed to other topical fluorides (questions 8-12, see Table 3). The unadjusted relationship between each of the perception questions and varnish use is shown in Table 5. There does not seem to be a relationship with price ($p > 0.3$), but there is a relationship with effectiveness, time, systemic exposure, and patient preference ($ps < 0.0002$). When these four perception variables are used in a logistic regression model that also includes graduation year, year of graduation is no longer significant ($p > 0.4$) and neither is the perception of systemic exposure ($p > 0.5$). Thus, the final model which predicts the use of fluoride varnish includes the following three terms: effectiveness, length of application time, and patient preference.

The significant terms from the final regression model were used to create a perception score to predict the use of fluoride varnish. The three factors were combined by adding up +1, 0, and -1 scores for each. Specifically, the effectiveness question may be

scored as +1=more effective or -1=other. The time consuming question may be scored as +1=less time consuming, 0=equal, -1=more time consuming. Finally, the patient preference question may be scored as +1=prefer varnish and -1=other preference. The sum of these three may thus range from -3, a negative on all three perception variables, to +3, a positive preference. The relationship between this perception score and the percentage of varnish use is shown in Table 6 and Figure 2. As may be seen, those respondents with a +3 perception score choose varnish 80% of the time while those with a +2 perception score choose varnish 71% of the time. A +1 perception score indicates 46% varnish use, a score of 0 indicates 15% varnish use, and a -1 score designates 9% varnish use. Any score less than -1 did not use fluoride varnish at all.

Discussion

Studies have shown that a minority of dentists use fluoride varnish on a regular basis for caries prevention and control.²³ The various reasons for the low utilization rates of varnish may be due to a lack of awareness, lack of FDA approval, and lack of evidence for a favorable cost:benefit ratio. In this study, the majority of respondents thought that fluoride varnish was more time consuming (43%) and had equal systemic exposure (46%) when compared to foams and gels. In reality, varnish is less time consuming; taking only a minute per application as opposed to the 4 minute application time recommended for gels and foams.¹ Also, varnish has been proven to have less systemic exposure and barely detectable effects on plasma fluoride concentrations due to the much smaller dose swallowed over several hours.¹⁰ The greater part of respondents (44%) also claimed to not know how fluoride varnish compared to other topical fluorides in price. However, those dentists who were aware of the advantages of varnish were more likely to use it as opposed to foams and gels. The respondents who correctly viewed varnish as more effective and less time consuming were more likely to use it on their patients. Naturally, those who perceived patients to prefer varnish were also more likely to use it in their practice.

This study found a positive correlation between the year of graduation from dental school and fluoride varnish use (Figure 1). Not surprisingly, those who have had more exposure to fluoride varnish, use it more frequently. Fiset et al confirmed this theory by

proving that dentists who knew more about fluoride varnish were more likely to implement the technology than those who knew less.²⁵ Fluoride varnish was not available in the United States until the early 1990s; thus, anyone who graduated before its availability was unable to become accustomed to using it during dental school. By contrast, every dental student who graduates from Virginia Commonwealth University School of Dentistry, uses varnish on a daily basis during their rotations in the pediatric dental clinic. Dixon et al demonstrated that diffusion of new technology in the medical profession functions through observation of colleagues and a clinician's own experience in using the new technology.^{25,26}

This study had some limitations in respect to the survey sample and on-line administration. First, it relied on email addresses that were given by the dentists to the Virginia Dental Association (VDA) to be used for their list-serve. Many email accounts have "junk mail" folders that filter email from unidentified senders. Thus, the survey could have been unknowingly discarded by potential respondents if they were blocking VDA-list serve email. Also, many dentists chose not to list their email with the VDA and therefore were automatically prevented from receiving the survey. Unfortunately, not all dentists in the state of Virginia are members of the VDA and therefore those dentists did not receive the survey. Second, the list-serve is not divided by dental specialties. Thus, specialists that rarely use fluoride, such as endodontists and oral surgeons, were included in the survey and consequently the results. This survey also had a modest response rate from dentists compared to recent paper surveys that have been done on this same

population.²⁷ It is probable that the response rate would have increased if a paper survey had been sent in addition to the electronic version.

In conclusion, the majority of dentists are not aware of the advantages of fluoride varnish. However, those who are, choose to use it as opposed to foams and gels. Dentists who have recently graduated from dental school are more likely to use fluoride varnish. Also, dentists who thought fluoride varnish was more effective and less time consuming use it more than other topical fluorides and those who thought their patients prefer fluoride varnish use it more than other topical fluorides. This survey supports the fact that more educational opportunities, both didactic and clinical, should be offered to dentists to familiarize them with the clinical superiority, technique, and advantages of fluoride varnish.

Literature Cited

Literature Cited

1. American Dental Association Council on Scientific Affairs. Professionally applied topical fluoride. *JADA* 2006;137:1151-1159.
2. Hazelrigg C, Dean J, Fontana M. Fluoride varnish concentration gradient and its effect on enamel demineralization. *Pediatr Dent*. 2003;25:119-126.
3. Bawden J. Fluoride varnish: a useful new tool for public health dentistry. *J Public Health Dent*. 1998;58(4):266-9.
4. Chu C, Lo E. A review of sodium fluoride varnish. *Gen Dent*. 2006 Jul-Aug;54(4):247-53.
5. Vaikuntam J. Fluoride varnishes: should we be using them? *Pediatr Dent*. 2000 Nov-Dec;22(6):513-6.
6. Berlin CM, May DG, Notterman DA. American Academy of Pediatrics Committee on Drugs. Unapproved uses of approved drugs: the physician, the package insert, and the Food and Drug Administration. *Pediatr* 1996;98:143-5.
7. Cote CJ, Kauffman RE, Troendle GJ, Lambert GH. Is the “therapeutic orphan” about to be adopted? (commentary). *Pediatr* 1996;98:118-22.
8. Weintraub JA, Ramos-Gomez F, Jue B, Shain S, Hoover CI, Featherstone JD, et al. Fluoride Varnish Efficacy in Preventing Early Childhood Caries. *J Dent Res* 2006;85(2):172-76
9. Ekstrand J, Koch G. Systemic fluoride absorption following fluoride gel application. *J Dent Res* 1980;59:1067.
10. Ekstrand J, Koch G, Petersson LF. Plasma fluoride concentrations and urinary fluoride excretion in children following application of the fluoride-containing varnish Duraphat. *Caries Res* 1980;14:185-9.

11. Whitford GM, Adair SM, Hanes CM, Perdue EC, Russell CM. Enamel uptake and patient exposure to fluoride: comparison of APF gel and foam. *Pediatr Dent* 1995;17(3):199–203
12. Wei SH, Chik FF. Fluoride retention following topical fluoride foam and gel application. *Pediatr Dent* 1990;12(6):368–74.
13. Murray JJ, Winter GB, Hurst CP. Duraphat fluoride varnish: a 2-year clinical trial in 5-year-old children. *Br Dent J* 1977;143:11–7.
14. Holm AK. Effect of fluoride varnish (Duraphat) in preschool children. *Community Dent Oral Epidemiol* 1979;7:241–5.
15. Petersson LG, Arthursson L, Ostberg C, Jonsson G, Glerup A. Caries-inhibiting effects of different modes of Duraphat varnish reapplication: a 3-year radiographic study. *Caries Res* 1991;25:70–3.
16. Derand T, Petersson LG. Effect of fluoride varnishes and Nuva-Seal resin treatment on the formation of artificial carious lesions. *Caries Res* 1981;15:250–5.
17. De Bruyn H, Buskes JA, Arends J. The inhibition of demineralization of human enamel after fluoride varnish application as a function of the fluoride content: an in vitro study under constant composition demineralizing conditions. *J Biol Buccale* 1986;14:133–8.
18. Sorvari R, Meurman JH, Alakuijala P, Frank RM. Effect of fluoride varnish and solution on enamel erosion in vitro. *Caries Res* 1994;28:227–32.
19. Sjöppa L. Effects of a sodium fluoride solution and a varnish with different fluoride concentrations on enamel remineralization in vitro. *Scand J Dent Res* 1988;96:304-9.
20. <http://www.mchoralhealth.org/materials/multiples/smilesforohio/Introduction.pdf>
21. Sjöppa L. Efficacy and safety of fluoride varnishes. *Compend Contin Educ Dent*. 1999;20(1 Suppl):18-26.
22. Hawkins R, Noble J, Locker D, Wiebe D, Murray H, Wiebe P, Frosina C, Clarke M. A comparison of the costs and patient acceptability of professionally applied topical fluoride foam and varnish. *J Public Health Dent*. 2004 Spring;64(2):106-10.

23. Fiset L, Grembowski D, Del Aguila M. Third-party reimbursement and use of fluoride varnish in adults among general dentists in Washington State. *J Am Dent Assoc.* 2000 Jul;131(7):961-8.
24. Inquisite 7.0, Web Survey System. Inquisite, Inc., Ausin, TX, 2007.
25. Fiset L, Grembowski D. Adoption of innovative caries-control services in dental practice: a survey of Washington state dentists. *JADA* 1997;128:337-45.
26. Dixon, AS. The evolution of clinical policies. *Med Care* 1990;28:201-20.
27. Brickhouse TH, Hollowell RL, Unkel JH, Webb MD, Best AM. Articaine use in children among dental practitioners. *Pediatr Dent* (Accepted Feb. 2008).

Table 1: Description of Those Surveyed (N=228)

| Characteristic | N | Percent |
|---------------------------------------|------|---------|
| Area of dental practice | | |
| Endodontics | 1 | 0.4 |
| General Dentistry | 178 | 78.8 |
| Oral and Maxillofacial Surgery | 5 | 2.2 |
| Orthodontics | 6 | 2.7 |
| Pediatric Dentistry | 26 | 11.5 |
| Periodontics | 10 | 4.4 |
| Year of graduation from dental school | | |
| prior to 1969 | 14 | 6.1 |
| 1970s | 62 | 27.2 |
| 1980s | 66 | 28.9 |
| 1990s | 45 | 19.7 |
| 2000s | 40 | 17.5 |
| Age (years) | | |
| Mean | 49.2 | |
| SD | 11.2 | |
| youngest | 24 | |
| oldest | 75 | |

Table 2: Description of Practice Characteristics

| Characteristic | N | Mean | SD | Range |
|---|-----|------|------|-------|
| Percentage of patient population in age groups: | | | | |
| < 6 years | 226 | 11.8 | 16.6 | 0 75 |
| 6-18 years | 226 | 22.0 | 17.0 | 0 90 |
| 18+ years | 226 | 66.2 | 28.4 | 0 100 |
| N Percent | | | | |
| Youth (under 18) patient population 33% or more | | | | |
| N | 153 | 68.0 | | |
| Y | 73 | 32.4 | | |
| Children (under 6) patient population 10% or more | | | | |
| N | 130 | 57.8 | | |
| Y | 96 | 42.7 | | |

Table 3: Application of Topical Fluoride

| Usage | N | Percent |
|--|-----|---------|
| 5. What type of topical fluoride do you mainly use? | | |
| Foam | 77 | 34.2 |
| Gel | 57 | 25.3 |
| Varnish | 60 | 26.7 |
| Other | 17 | 7.6 |
| None | 14 | 6.2 |
| 6. In what age categories of patients are you applying topical fluoride? | | |
| 0-3 Years | 77 | 36.2 * |
| 4-6 Years | 177 | 83.1 * |
| 7-12 Years | 200 | 93.9 * |
| 13-18 Years | 184 | 86.4 * |
| 19+ Years | 105 | 49.3 * |
| 7. In a typical patient, how often do you apply fluoride? | | |
| According to patient's risk/history of decay | 46 | 21.6 |
| Once a year | 25 | 11.7 |
| Twice a year | 142 | 66.7 |
| 8. How do you think fluoride varnish compares in price to other topical fluorides? | | |
| Cheaper | 4 | 1.9 |
| Same | 29 | 13.6 |
| More Expensive | 86 | 40.4 |
| Don't Know | 94 | 44.1 |
| 9. Do you think fluoride varnish is _____ effective than gels and foams? | | |
| More | 138 | 66.0 |
| Equal | 60 | 28.7 |
| Less | 11 | 5.3 |
| 10. Do you think fluoride varnish is _____ time consuming than gels and foams? | | |
| More | 89 | 42.8 |
| Equal | 65 | 31.3 |
| Less | 54 | 26.0 |
| 11. Do you think fluoride varnish has _____ systemic exposure than gels and foams? | | |
| More | 37 | 17.7 |
| Equal | 95 | 45.5 |
| Less | 77 | 36.8 |
| 12. Which product do you think patients would prefer? | | |
| Foam | 63 | 30.0 |
| Gel | 45 | 21.4 |
| Varnish | 84 | 40.0 |
| Other | 18 | 8.6 |

Table 4: Unadjusted Relationships with Topical Fluoride Varnish Use

| Characteristic | Varnish use? | | |
|--|--------------|----|-------|
| | N | Y | % Yes |
| Area of dental practice | | | |
| General Dentistry | 127 | 45 | 26.2 |
| other | 10 | 4 | 28.6 |
| Pediatric Dentistry | 14 | 11 | 44.0 |
| LR chi-square = 3.2, p-value = 0.2038 | | | |
| Year of graduation from dental school | | | |
| prior to 1969 | 8 | 4 | 33.3 |
| 1970s | 46 | 8 | 14.8 |
| 1980s | 42 | 20 | 32.3 |
| 1990s | 33 | 11 | 25.0 |
| 2000s | 22 | 17 | 43.6 |
| LR chi-square = 10.5, p-value = 0.033 | | | |
| Continuous trend, LR chi-square = 3.88, p-value = 0.0488 | | | |
| Age (years) | | | |
| 20s | 4 | 3 | 42.9 |
| 30s | 28 | 16 | 36.4 |
| 40s | 37 | 13 | 26.0 |
| 50s | 53 | 17 | 24.3 |
| 60s and older | 29 | 11 | 27.5 |
| LR chi-square = 2.7, p-value = 0.6035 | | | |
| Continuous trend, LR chi-square = 2.54, p-value = 0.1110 | | | |
| Youth (under 18) patient population 33% or more | | | |
| N | 107 | 38 | 26.3 |
| Y | 44 | 21 | 34.2 |
| Fisher's exact p-value = 0.4074 | | | |
| Continuous trend, LR chi-square = 3.53, p-value = 0.0604 | | | |
| Children (under 6) patient population 10% or more | | | |
| N | 90 | 28 | 23.7 |
| Y | 61 | 31 | 33.7 |
| Fisher's exact p-value = 0.1235 | | | |
| Continuous trend, LR chi-square = 6.21, p-value = 0.0127 | | | |

Table 5: Unadjusted Relationships with Fluoride Varnish Use

| Perception | Varnish use? | | % Yes |
|--|--------------|----|-------|
| | N | Y | |
| 8. How do you think fluoride varnish compares in price to other topical fluorides? | | | |
| Cheaper | 3 | 1 | 25.0 |
| Same | 19 | 9 | 32.1 |
| More Expensive | 56 | 29 | 34.1 |
| Don't Know | 73 | 21 | 22.3 |
| LR chi-square = 3.3, p-value = 0.3462 | | | |
| 9. Do you think fluoride varnish is _____ effective than gels and foams? | | | |
| More | 81 | 55 | 40.4 |
| Equal | 56 | 4 | 6.7 |
| Less | 10 | 1 | 9.1 |
| LR chi-square = 25.3, p-value = <.0001 | | | |
| 10. Do you think fluoride varnish is _____ time consuming than gels and foams? | | | |
| More | 78 | 11 | 12.4 |
| Equal | 47 | 17 | 26.6 |
| Less | 22 | 31 | 58.5 |
| LR chi-square = 34.1, p-value = <.0001 | | | |
| 11. Do you think fluoride varnish has _____ systemic exposure than gels and foams? | | | |
| More | 32 | 4 | 11.1 |
| Equal | 73 | 22 | 23.2 |
| Less | 42 | 34 | 44.7 |
| LR chi-square = 16.8, p-value = 0.0002 | | | |
| 12. Which product do you think patients would prefer? | | | |
| Foam | 58 | 5 | 7.9 |
| Gel | 40 | 5 | 11.1 |
| Other | 16 | 2 | 11.1 |
| Varnish | 34 | 48 | 58.5 |
| LR chi-square = 59.8, p-value = <.0001 | | | |

Table 6: Perception Score and Varnish Use

| Score | Varnish use? | | Total | % |
|-------|--------------|----|-------|-----|
| | N | Y | | Yes |
| -3 | 30 | 0 | 30 | 0 |
| -2 | 11 | 0 | 11 | 0 |
| -1 | 48 | 5 | 53 | 9 |
| 0 | 28 | 5 | 33 | 15 |
| +1 | 15 | 13 | 28 | 46 |
| +2 | 5 | 12 | 17 | 71 |
| +3 | 6 | 24 | 30 | 80 |
| Total | 143 | 59 | 202 | 29 |




Figure 1: Relationship with Graduation Year




Figure 2: Perception Score and Percentage of Varnish Use

APPENDIX A

Survey

1) What is your area of Dental Practice?

- General Dentistry
- Pediatric Dentistry
- Orthodontics
- Periodontics
- Endodontics
- Oral and Maxillofacial Surgery
- Other _____

2) What is your age?

3) What year did you graduate from dental school?

4) Please describe the characteristics of your patient population

| Age | < 6 years | 6-18 years | 18+ years |
|-------------------------|-------------------|-------------------|-----------|
| Approximate Percent (%) | | | |
| Dental Coverage | Private Insurance | Public Assistance | Self-Pay |
| Approximate Percent (%) | | | |

5) What type of topical fluoride do you use?

- Foam
- Gel
- Varnish
- Other _____
- None (This survey focuses on the use of topical fluorides, Thank you for your responses)

6) In what age categories of patients are you applying topical fluoride?

- 0-3 Years
- 4-6 Years
- 6-12 Years
- 12-18 Years
- 18+ Years

7) How often do you apply fluoride?

- Once a year
- Twice a year
- According to patient's risk/history of decay

8) How do you think fluoride varnish compares in price to other topical fluorides?

- Cheaper
- Same
- More Expensive
- Don't Know

9) Do you think fluoride varnish is _____ effective than gels and foams?

- More
- Equal
- Less

10) Do you think fluoride varnish is _____ time consuming than gels and foams?

- More
- Equal
- Less

11) Do you think fluoride varnish has _____ systemic exposure than gels and foams?

- More
- Equal
- Less

12) Which product do you think patients would prefer?

- Foam
- Gel
- Varnish
- Other

13) If you are interested in participating in further research and continuing education related to fluoride varnish please enter your email address here.

VITA

Amanda Bowen Kuhn was born on November 27, 1979 in Indianapolis, Indiana. She graduated from James B. Hunt High School, Wilson, North Carolina in 1998. She received a BS in Biology from The University of North Carolina at Chapel Hill, Chapel Hill, North Carolina in 2002. Dr. Kuhn received her Doctor of Dental Medicine from The University of Pittsburgh School of Dental Medicine, Pittsburgh, Pennsylvania in 2006.