

2015

HIV Protease Inhibitors Sensitize Human Head and Neck Squamous Carcinoma Cells to Radiation by Activating Endoplasmic Reticulum Stress

Runping Liu

Virginia Commonwealth University

Luyong Zhang

China Pharmaceutical University

Jing Yang

China Pharmaceutical University, Virginia Commonwealth University

See next page for additional authors

Follow this and additional works at: http://scholarscompass.vcu.edu/micr_pubs

 Part of the [Medicine and Health Sciences Commons](#)

© 2015 Liu et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Downloaded from

http://scholarscompass.vcu.edu/micr_pubs/41

This Article is brought to you for free and open access by the Dept. of Microbiology and Immunology at VCU Scholars Compass. It has been accepted for inclusion in Microbiology and Immunology Publications by an authorized administrator of VCU Scholars Compass. For more information, please contact libcompass@vcu.edu.

Authors

Runping Liu, Luyong Zhang, Jing Yang, Xiaoxuan Zhang, Ross B. Mikkelsen, Shiyu Song, and Huiping Zhou

RESEARCH ARTICLE

HIV Protease Inhibitors Sensitize Human Head and Neck Squamous Carcinoma Cells to Radiation by Activating Endoplasmic Reticulum Stress

Runping Liu^{1,2}, Luyong Zhang^{1*}, Jing Yang^{1,2}, Xiaoxuan Zhang^{1,2}, Ross Mikkelsen³, Shiyu Song³, Huiping Zhou^{2,4*}

1 Jiangsu Center for Drug Screening, China Pharmaceutical University, Nanjing, 210009, China, **2** Department of Microbiology & Immunology, Virginia Commonwealth University, Richmond, VA, 23298, United States of America, **3** Department of Radiation Oncology, Virginia Commonwealth University, Richmond, VA, 23298, United States of America, **4** McGuire Veterans Affairs Medical Center, Richmond, VA, 23298, United States of America

* hzhou@vcu.edu (HZ); lyzhang@cpu.edu.cn (LZ)



OPEN ACCESS

Citation: Liu R, Zhang L, Yang J, Zhang X, Mikkelsen R, Song S, et al. (2015) HIV Protease Inhibitors Sensitize Human Head and Neck Squamous Carcinoma Cells to Radiation by Activating Endoplasmic Reticulum Stress. PLoS ONE 10(5): e0125928. doi:10.1371/journal.pone.0125928

Academic Editor: Gianfranco Alpini, Texas A&M Health Science Center, UNITED STATES

Received: January 14, 2015

Accepted: March 26, 2015

Published: May 1, 2015

Copyright: © 2015 Liu et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper.

Funding: This work was supported in part by NIH grants R01 DK-057543 and R01AT004148, VA Merit Award 1101BX001390, and National Science Foundation of China Grants 81070245 and 81270489 to HZ. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

Abstract

Background

Human head and neck squamous cell carcinoma (HNSCC) is the sixth most malignant cancer worldwide. Despite significant advances in the delivery of treatment and surgical reconstruction, there is no significant improvement of mortality rates for this disease in the past decades. Radiotherapy is the core component of the clinical combinational therapies for HNSCC. However, the tumor cells have a tendency to develop radiation resistance, which is a major barrier to effective treatment. HIV protease inhibitors (HIV PIs) have been reported with radiosensitizing activities in HNSCC cells, but the underlying cellular/molecular mechanisms remain unclear. Our previous study has shown that HIV PIs induce cell apoptosis via activation of endoplasmic reticulum (ER) stress. The aim of this study was to examine the role of ER stress in HIV PI-induced radiosensitivity in human HNSCC.

Methodology and Principal Findings

HNSCC cell lines, SQ20B and FaDu, and the most commonly used HIV PIs, lopinavir and ritonavir (L/R), were used in this study. Clonogenic assay was used to assess the radiosensitivity. Cell viability, apoptosis and cell cycle were analyzed using Cellometer Vision CBA. The mRNA and protein levels of ER stress-related genes (eIF2 α , CHOP, ATF-4, and XBP-1), as well as cell cycle related protein, cyclin D1, were detected by real time RT-PCR and Western blot analysis, respectively. The results demonstrated that L/R dose-dependently sensitized HNSCC cells to irradiation and inhibited cell growth. L/R-induced activation of ER stress was correlated to down-regulation of cyclin D1 expression and cell cycle arrest under G0/G1 phase.

Conclusion and Significance

HIV PIs sensitize HNSCC cells to radiotherapy by activation of ER stress and induction of cell cycle arrest. Our results provided evidence that HIV PIs can be potentially used in combination with radiation in the treatment of HNSCC.

Introduction

Human head and neck carcinoma includes a heterogeneous group of malignancies of the oral cavity, oropharynx, hypopharynx, larynx, lips, paranasal sinuses and salivary glands [1]. More than 90% of these cancers are squamous cell carcinomas (HNSCC). HNSCC represents the sixth most common malignancy worldwide [2]. The major risk factors include tobacco and alcohol consumption, poor oral hygiene, and infection by human papillomavirus (HPV) [3–5].

The majority of HNSCCs are diagnosed in locally advanced stages. Surgery, radiotherapy and chemotherapy are the current main strategies to treat HNSCC patients. Local recurrence remains the dominant pattern of treatment failure. Recent advances in the understanding of the molecular mechanisms of disease initiation and progression have led to the development of more specific therapies, such as Cetuximab, a monoclonal antibody against the epidermal growth factor receptor (EGFR). Cetuximab has been approved for combinational therapy with radiation in patients with unresectable HNSCC [6]. Overexpression of EGFR was often associated with a poor prognosis in HNSCC [7]. Although enhanced efforts have been put into effect and new therapies have been introduced during the last decade, the morbidity rate of HNSCC has not been reduced significantly [8]. A major challenge of current available therapies (radiation and chemotherapy) is the rapid development of resistance. Therefore, identification of cellular/molecular mechanisms responsible for resistance and development of new therapeutic strategies to overcome the resistance would improve the efficiency of current therapies.

Human immunodeficiency virus protease inhibitors (HIV PIs) are key components of highly active anti-retroviral therapy (HAART) for HIV infection. Previous studies from our laboratory and other laboratories demonstrated that HIV PIs are able to trigger the unfolded protein response (UPR) and endoplasmic reticulum (ER) stress [9, 10]. Three main branches of the UPR have been identified so far, including IRE1, PERK and ATF6 [11, 12]. PERK pathway activation is considered tightly correlated with cell apoptosis and survival. PERK activation-induced phosphorylation of eIF2 α , a key mediator of protein translation, further disrupts translation initiation complexes and subsequently leads to global suppression of protein expression. Phosphorylation of eIF2 α further induces the expression of ATF4, which leads to activation of CHOP, a proapoptotic factor [13–15]. Emerging evidence demonstrated that HIV PI-induced ER stress activation is linked to cell apoptosis in different types of cells [16–20]. Several studies reported that HIV PIs induced apoptosis through activating the STAT3/ERK1/2 pathway in human multiple myeloma cells and prostate cancer cells [21, 22]. HIV PIs are also able to induce cell cycle arrest and cell death by triggering the ER stress response [23, 24]. Recent studies further suggest that HIV PIs could be potential anti-cancer drugs due to their inhibitory effects on the PI3K-Akt signaling pathway, which is considered to be an important survival mechanism in various cancer cells [25]. Several preclinical studies indicated that down-regulation of Akt phosphorylation by HIV PI treatment or by the proteasome inhibitor, Bortezomib, resulted in increased radiosensitivity of cancer cells [20, 26, 27]. Despite the fact that HIV PIs have been identified as potential radio-sensitizers, the underlying mechanisms are still unclear. The current study was aimed at examining whether the most commonly used

HIV PIs (lopinavir and ritonavir, L/R) are able to increase the radiosensitivity in HNSCC and further identify the underlying mechanisms for that increase.

Materials and Methods

Materials

HIV protease inhibitors, lopinavir (L) and ritonavir (R), were obtained from NIH AIDS Research & Reference Reagent Program, Division of AIDS, NIAID, NIH. Thapsigargin (TG) and dimethyl sulfoxide (DMSO) was purchased from Sigma-Aldrich (St. Louis, MO). Cell Counting Kit-8 (CCK-8) was from Dojindo Molecular Technologies (Kumamoto, Japan). Antibodies against p-ERK1/2, ERK1, ERK2, CHOP, ATP-4, XBP-1, Lamin B, p-eIF-2 α , and eIF-2 α were from Santa Cruz Biotechnology (Santa Cruz, CA). Mouse monoclonal antibody against β -actin and polyclonal antibodies against p-AKT, AKT, and cyclin D1 were from Thermo Scientific (Wilmington, DE). IRDye secondary antibodies were purchased from LI-COR Biosciences (Lincoln, NE). Bio-Rad protein assay reagent, Precision Plus Kaleidoscope Standards, iQTM SYBR Green Supermix were obtained from Bio-Rad (Hercules, CA). QIAzol Lysis Reagent was obtained from QIAGEN Sciences (Germantown, MD). High Capacity cDNA Reverse Transcription Kit was from Life Technologies (Grand Island, NY). Cell culture medium and supplement components were products from Invitrogen (Carlsbad, CA, USA)

Cell culture and treatment

Human HNSCC cells, FaDu hypopharyngeal cancer cell line (HTB-43) and SQ20B cell line, were obtained from ATCC. FaDu cells were cultured in MEM medium with 10% Fetal Bovine Serum, penicillin G (100 U/ml), streptomycin (100 μ g/ml) and 1% non-essential amino acids (NEAA). SQ20B cells were cultured in RPMI 1640 medium with 10% Fetal Bovine Serum, penicillin G (100 U/ml) and streptomycin (100 μ g/ml). HIV PIs (L/R, 4/1 ratio) and TG were dissolved in DMSO. Drugs were directly added into the cell culture medium and incubated for different periods as described previously [28–31].

Cell proliferation assay

SQ20B and FaDu cells were seeded into 96-well plates with complete medium and treated with HIV PIs. The viable cell numbers were determined by Cell Counting Kit-8 (CCK8), from Dojindo Molecular Technologies, Inc. (Rockville, MD). Victor³ Multilabel Plate Counter from Perkin Elmer (Waltham, MA) was used to measure the absorbance at 450nm as described previously [32].

Clonogenic survival assay

SQ20B and FaDu cells were cultured in 24-well plates with or without HIV PI treatment for 24 h and followed by irradiation with different doses, 2Gy or 4Gy. After 24 h, cells were detached and plated for colony-formation assays to assess clonogenic survival. Cells were cultured for 12 days, fixed in 10% formaldehyde, and stained with 0.5% violet crystal. Colonies containing 50 or more cells were counted to determine the surviving fractions of clonogenic cells using an Olympus microscope as described previously [33].

Annexin V and Propidium Iodide (PI) staining

SQ20B and FaDu cells were cultured in 6 well plates. After 24h treatment, cells were harvested and resuspended into a density of 10⁶cells/ml. BD Pharmingen FITC Annexin V Apoptosis Detection Kit I (BD Science, San Diego, CA) was used to perform the staining, according to the

protocol recommended by the manufacturer as described previously [30]. The stained cells were then analyzed by Cellometer Vision CBA Analysis System (Nexcelom Bioscience, Lawrence, MA).

Cell cycle assay

SQ20B and FaDu cells were treated with HIV PIs or TG or vehicle control for 24h. Cells were fixed in ice-cold methanol for 15 min and stained with a Cellometer PI Cell Cycle Kit (Nexcelom Bioscience, Lawrence, MA) according to the protocol recommended by the manufacturer. The Cellometer Vision CBA Analysis System was used to analyze the results.

RNA isolation and quantitative real-time RT-PCR

Cells were treated with HIV PIs or vehicle control for 24 h. Total Cellular RNA was isolated using QIAzol Lysis Reagent and reverse transcribed into first-strand cDNA using a High-Capacity cDNA Transcription Kit. The mRNA levels of ATF-4, XBP-1, CHOP, and GAPDH were quantified with real-time RT-PCR as described previously [16, 17, 19, 28–30, 34, 35].

Western Blot analysis

The nuclear proteins and total proteins were isolated as previously described [16]. The protein levels of target genes were detected using specific primary antibodies and IRDye secondary antibodies on an Odyssey Fluorescence Imaging System (LI-COR Biosciences, USA) as described previously [19, 35, 36]. The density of the immunoblots was analyzed using Odyssey V3.0 software

Wound healing assays

SQ20B cells were plated into 12-well plates at a concentration of 5×10^5 cells per well for overnight and allowed to form a confluent monolayer. After that the monolayer was scratched with a sterile pipette tip, cells were washed gently with serum free medium to remove floating and detached cells and photographed (time 0 h) using an Olympus 1x71 microscope using a 10 x objective. Cells were treated with different amounts of L/R (0–25 μ M) or vehicle control (DMSO) or TG (100 nM). After 6h, the wounded area was photographed as described previously [32]. The images acquired for each treatment group were further analyzed using IPLab 4.0.

Statistical analysis

All of the experiments were repeated at least three times and the data were expressed as the mean \pm SD. One-way ANOVA was employed to analyze the differences between sets of data. To confirm the differences that occurred between groups, *post hoc* tests were used for follow-up tests. Statistics were performed using GraphPad Prism 5.0 (GraphPad, San Diego, CA) as described previously [32, 36]. A value of $P < 0.05$ was considered statistically significant.

Results

Effect of HIV PIs on radiosensitivity of HNSCC cells

To evaluate the effect of the most commonly used HIV PIs (L/R) on HNSCC proliferation, SQ20B and FaDu cells were treated with different concentrations of L/R (0, 6.25, 12.5, 25 or 50 μ M) for 48 h. As shown in Fig 1, L/R significantly inhibited cell growth at concentrations of 25 and 50 μ M in both SQ20B and FaDu cells. To further determine whether HIV PIs were able to sensitize HNSCC to radiation, SQ20B and FaDu cells were pretreated with L/R (12.5 or




Fig 1. The effects of HIV PIs on HNSCC cell growth. Human HNSCC cell lines SQ20B and FaDu were cultured in complete medium and treated with different concentrations of HIV PIs (Lopinavir/Ritonavir, L/R, 0–50 μM) or Thapsigargin (TG, 50 nM) for 48h. At the end of treatment, cell viability was assessed using the CCK-8 kit according to the instructions provided by the manufacturer. Relative numbers of viable cells were calculated. (A) SQ20B; (B) FaDu. Values of each group are mean ± S.E. of six independent experiments. *** $P < 0.001$, Statistical significance relative to control group.

doi:10.1371/journal.pone.0125928.g001

25 μM) for 24 h and then irradiated at 2 Gy or 4 Gy. As shown in Fig 2, L/R significantly increased the radiosensitivity and reduced the number of survival colonies in both SQ20B and FaDu cells.

Effect of HIV PIs on ER stress activation and cell apoptosis

Our previous studies have shown that HIV PIs induce ER stress and cell apoptosis in macrophages, hepatocytes, intestinal epithelial cells, and adipocytes [16, 18, 19, 34]. To examine whether L/R also activates ER stress in HNSCC cells, we first examined the effect of L/R on mRNA expression of the key genes involved in the UPR. As shown in Fig 3, L/R dose-dependently increased CHOP, ATF-4, and XBP-1 mRNA levels in both SQ20B and FaDu cells.





Fig 2. Effect of HIV PIs on radiosensitivity of HNSCC cells. Human HNSCC cells (SQ20B and FaDu) were cultured in complete medium and treated with different concentrations of HIV PIs (L/R, 12.5 μM or 25 μM) for 24h. At the end of treatment, cells were exposed to 2Gy or 4Gy of radiation. After irradiation, cells were detached and plated for colony-formation assays to assess clonogenic survival as described in “Methods”. Relative numbers of survival colonies are presented. (A) SQ20B; (B) FaDu; Values are mean ± S.E. of three independent experiments. ** $P < 0.01$, * $P < 0.05$, Statistical significance relative to control group.

doi:10.1371/journal.pone.0125928.g002

A SQ20B cells



B FaDu cells




Fig 3. Effect of HIV PIs on mRNA levels of the UPR genes in HNSCC. HNSCC cells, SQ20B and FaDu, were cultured in complete medium and treated with different concentrations of L/R (0, 12.5, 25, and 50 μM) or TG (50 nM) for 24h. At the end of treatment, total RNAs were isolated and reverse transcribed. The relative mRNA levels of ATF-4, CHOP, spliced XBP-1 (XBP-1s) and unspliced XBP-1 (XBP-1u) were detected by real-time RT-PCR and normalized using GAPDH as an internal control as described under "Methods". Values are mean ± S.E. of three independent experiments. Statistical significance relative to vehicle control, * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$. (A) SQ20B cells; (B) FaDu cells.

doi:10.1371/journal.pone.0125928.g003

TG was the positive control for ER stress induction. As shown in Fig 4, L/R-induced mRNA expression of the UPR genes was correlated to up-regulation of the corresponding protein level. L/R had a more profound effect on ATF-4 expression. To further determine whether L/R-induced UPR activation was able to promote cell apoptosis, Annexin V-PI staining was used to detect apoptotic and necrotic cells after treatment with L/R (25 μ M) or TG (50 nM) for 24 h.




Fig 4. Effect of HIV PIs on protein levels of the UPR genes in HNSCC. HNSCC cells, SQ20B and FaDu, were cultured in complete medium and treated with different concentrations of L/R (0, 12.5, 25, and 50 μ M) or TG (50 nM) for 24h. At the end of treatment, nuclear proteins were isolated. The protein levels of CHOP, ATF-4, XBP-1 and lamin B were determined by Western blotting analysis. Lamin B was used as a loading control. Representative images of immunoblots are shown. Values are mean \pm S.E. of three independent experiments. Statistical significance relative to vehicle control, * P <0.05; ** P <0.01. (A) SQ20B cells; (B) FaDu cells.

doi:10.1371/journal.pone.0125928.g004

As shown in Fig 5, L/R alone did not induce significant apoptosis in SQ20B and FaDu cells. However, pre-treatment with L/R markedly increased radiosensitivity of both HNSCC cell lines. Compared to the control group, L/R reduced cell viability by 60% (data not shown).

Effect of HIV PIs on HNSCC cell migration

Cancer cells spread from the initial site of tumor growth by first invading the surrounding tissue (migration). The ability of tumor cells to migrate is closely linked to tumor growth and invasiveness. To determine the effect of HIV PIs on HNSCC cell migration, we did a wound healing assay in SQ20B cells as described in Methods. As shown in Fig 6, L/R dose-dependently inhibited cell migration. Similar results were obtained with TG, the ER stress inducer.

Effect of HIV PIs on cell cycle distribution

In order to identify the potential mechanisms underlying HIV PI-induced radiosensitivity of HNSCC, we examined the effect of L/R on cell cycle distribution. After treatment with L/R (25 μ M) or TG (50 nM) for 24h, cells were analyzed using the Cellometer PI Cell Cycle Kit from Nexcelom. As shown in Fig 7, L/R significantly increased SQ20B and FaDu cells at G₀/G₁ phase and cells at S/G₂ phases were markedly decreased. These results indicated that L/R increased radiosensitivity of SQ20B and FaDu cells through inducing G₀/G₁ phase arrest rather than promoting apoptosis.

Overexpression of EGFR is detected in 90% of HNSCC. EGFR-mediated activation of PI3K/Akt and mitogen-activated/extracellular signal-regulated kinase (MEK) contributes to activation of NF- κ B and induction of inflammatory response. Activation of Akt also induces activation of the mammalian target of rapamycin (mTOR). Emerging evidence indicates that phosphorylated Akt plays an important role in radio-resistance in cancer cells [37–39]. It also has been reported that the HIV PI, nelfinavir, has an inhibitory effect on Akt activation and mTOR [26, 27]. To determine whether L/R have any effect on Akt activation in HNSCC, we measured the phosphorylation level of Akt in SQ20B and FaDu cells after treatment with L/R for 24 h by western blot analysis. The results indicated that L/R had no significant effect on Akt activation in HNSCC (Data not shown).

Effect of HIV PIs on the expression of cyclin D1

HNSCC exhibits increased expression of cyclin D1, a key moderator in cell cycle progression from G₁ phase to S phase. Overexpression of cyclin D1 correlates to poor prognosis of HNSCC [40]. It has been shown that ER stress-mediated activation of the PERK/eIF2 α /ATF-4 signaling pathway down-regulated expression of cyclin D1 by suppressing global protein synthesis. To further identify the mechanism underlying HIV PI-induced cell cycle arrest in HNSCC, we examined the effect of HIV PIs on PERK/eIF2 α activation and cyclin D1 expression. As shown in Fig 8A and 8B, L/R dose-dependently increased the protein levels of phosphorylated eIF2 α (p-eIF2 α) in both SQ20B and FaDu cells. Furthermore, L/R-induced increase of p-eIF2 α was correlated to the decrease of cyclin D1 protein levels (Fig 8C), suggesting the contribution of activation of ER stress to HIV PI-enhanced radiosensitivity of SQ20B and FaDu cells.

Discussion

HNSCC is still a devastating malignancy worldwide. The overall five-year survival rate is only about 50% despite recent advances in therapy. Radiotherapy is the cornerstone of current therapy for HNSCC. The major challenge to the management of HNSCC is the rapid development of resistance to radiotherapy [41]. HIV PIs have been successfully used to inhibit HIV




Fig 5. Effect of HIV PIs on apoptosis in SQ20B and FaDu cells. Human HNSCC cells SQ20B and FaDu were cultured in complete medium and treated with HIV PIs L/R (25µM) or TG (50 nM) for 24h. At the end of treatment, cells were stained with Annexin V- PI and analyzed using the Cellometer Vision CBA system. FL1 represents Annexin V staining for apoptotic cells and FL2 represents PI staining for necrotic cells. Representative images of flow cytometry analysis and percentages of apoptotic and necrotic cells from three independent experiments are shown: (A) SQ20B; (B) FaDu. Statistical significance relative to vehicle control, * $P < 0.05$.

doi:10.1371/journal.pone.0125928.g005

replication and have significantly reduced the morbidity and mortality of HIV patients in the past two decades. Several studies have reported that HIV PIs, such as nelfinavir, can increase the radiosensitivity of certain cancer cells including HNSCC [20–23, 27, 42–44]. However, the toxicity of nelfinavir is relatively higher than other HIV PIs and the underlying mechanisms still remain unknown.




Fig 6. Effect of HIV PIs on cell migration. HNSCC SQ20B cells were plated on 12-well plates until confluent. Cells were scratched to simulate a wound and images were recorded as “0h”. Cells were treated with different amounts of HIV PIs (L/R, 0–25 μ M) for 6 h. The images of wound areas were recorded as described in Methods. Representative images are shown.

doi:10.1371/journal.pone.0125928.g006

In the present study, we investigated the effects of the most commonly used HIV PIs, L/R, on cell proliferation, migration, and radiosensitivity of HNSCC and further identified the potential mechanisms. The results indicated that L/R significantly inhibited cell growth and migration, and markedly enhanced radiosensitivity. Previous studies from our groups and others’ suggested that HIV PIs induce ER stress and apoptosis in different types of cells [18–20, 45]. It has been reported that ER stress sensitizes cancer cells to radiation [46, 47]. Although L/R did not directly induce apoptosis in HNSCC cells, radiosensitivity was significantly enhanced after pretreatment with L/R. Several studies have reported that nelfinavir sensitized HNSCC cells to radiation mainly by inhibiting the PI3/Akt signaling pathway [27]. The Ras/PI3K/Akt signaling pathway is considered to be an important radio-resistance pathway in cancer cells because of modulation of cell cycle progression, especially at the G_1/S transition, and DNA damage repair after radiation exposure [24, 37–39, 48]. Cells under G_1 phase prepare proteins and enzymes for the following DNA synthesis in S phase. The formation of two critical complexes, Cyclin D1/CDK4 and Cyclin E/CDK2, is the core mechanism of cell cycle progression from G_1 to S phase. Akt is an important regulator of the activation of cyclin D1 and inhibition of P27kip1, a suppressor of the cyclin D1/CDK4 complex, and induces cell proliferation by promoting cells to pass through the G_1 checkpoint [49, 50]. In addition, Akt also increases cancer cell survival after radiation by helping to repair radiation-induced DNA damage [51]. These findings suggested that targeting Akt represents a promising new anti-cancer therapy. However, inhibition of the Akt-signaling pathway alone is not enough to enhance radiosensitivity of HNSCC [27]. In contrast to nelfinavir, which required longer (3-day) treatment, L/R only required a shorter (1-day) treatment to enhance radiosensitivity of HNSCC cells. Also, L/R had no effect on Akt activation, suggesting that another Akt-independent pathway is involved in L/R-mediated radio-sensitization.

ER stress-mediated activation of the UPR has been implicated in cell growth arrest and inhibition of cell migration. In the current studies, we identified that L/R significantly activated the IRE1 α /XBP1 and PERK/eIF2 α /ATF4 in SQ20B and FaDu cells. Although L/R significantly up-regulated CHOP, a proapoptotic factor, there was no significant change in terms of cell




Fig 7. Effect of HIV PIs on cell cycle distribution of HNSCC cells. Human HNSCC cells (A) SQ20B and (B) FaDu were cultured in complete medium and treated with L/R (25µ M) or TG (50 nM) for 24h. At the end of treatment, cells were stained with the Cellometer PI Cell Cycle kit and analyzed by the Cellometer Vision CBA system. Cell population percentages of G₀/G₁, S and G₂/M phases are shown. Values of each group are mean ± S.E. of three independent experiments. **P*<0.05, Statistical significance relative to control group.

doi:10.1371/journal.pone.0125928.g007

apoptosis. These results suggest that an L/R-mediated increase of radiosensitivity of HNSCC cells is independent of suppression of Akt activation and induction of apoptosis.

Regulation of cell cycle checkpoints is important for cell growth. Our results indicated that L/R significantly arrested SQ20B and FaDu cells at the G₀/G₁ phase. Previous studies have shown that ER stress-mediated activation of PERK/eIF2α was correlated to the down-regulation of cyclin D1 by global translation suppression [52, 53]. Overexpression of PERK also induced significant reduction of CDK2 activity [53]. In addition, crosstalk between PERK-induced eIF2α




Fig 8. Effect of HIV PIs on PERK/eIF2 α activation and cyclin D1 expression in HNSCC cells. Human HNSCC cells, SQ20B and FaDu, were cultured in complete medium and treated with different concentrations of L/R (0, 12.5, 25, and 50 μM) or TG (50 nM) for 24h. At the end of treatment, total cellular proteins were isolated. The protein levels of p-eIF2 α , eIF2 α , cyclin D1, and β -actin were determined by Western blotting analysis. β -actin was used as a loading control. Representative images of immunoblots are shown. Values are mean \pm S.E. of three independent experiments. Statistical significance relative to vehicle control, * $P < 0.05$; ** $P < 0.01$. (A) Protein levels of p-eIF2 α and eIF2 α ; (B) Protein levels of cyclin D1.

doi:10.1371/journal.pone.0125928.g008




Fig 9. Proposed mechanism of HIV PI-induced radiosensitivity in HNSCC cells. HIV PIs induce ER stress and activate the UPR in HNSCC cell. Activation of PERK/eIF2 α /ATF-4 will repress global protein translation, reduce cyclin D1 protein level and induce cell cycle arrest. ATF-4 also induces the expression of CHOP, which will inhibit cell growth. PERK, double-stranded RNA-activated protein kinase-like ER kinase; CHOP is also called GADD153 (growth arrest and DNA damage-inducing gene).

doi:10.1371/journal.pone.0125928.g009

phosphorylation and cyclin D1 protein degradation was reported recently [54]. Our studies also showed that L/R-mediated activation of the PERK/eIF2 α pathway was linked to G₀/G₁ cell cycle arrest and down-regulation of cyclin D1 protein levels in SQ20B and FaDu cells. During the last decade, several chemotherapeutic agents, which induce cell cycle arrest, such as flavopiridol and paclitaxel, have been used to enhance radiosensitivity of cancer cells [55]. Our studies suggest that the HIV PIs, L/R, can be used to treat HNSCC in combination with other chemotherapies and radiotherapy.

Conclusions

In summary, our study demonstrated that administration of HIV PIs (L/R) significantly enhanced radiosensitivity of SQ20B and FaDu cells. As illustrated in Fig 9, L/R induced activation of the ER stress response and subsequently activated PERK/eIF2 α /ATF-4 and reduced cyclin D1 protein levels. L/R induced cell cycle arrest mainly through the inhibition of global protein translation and cell growth. Our studies suggest that activation of ER stress response represents one of the key mechanisms underlying HIV PI-induced radiosensitivity. This study provides a rationale for clinical investigation of the combination therapy using the FDA-approved HIV PIs (L/R) with radiotherapy in patients with HNSCC.

Acknowledgments

The authors are grateful to Elaine Kennedy and Yunzhou Li for editing and proofreading the manuscript.

Author Contributions

Conceived and designed the experiments: LZ SS HZ. Performed the experiments: RL XZ JY RM. Analyzed the data: RL XZ JY HZ. Contributed reagents/materials/analysis tools: SS RM HZ. Wrote the paper: RL HZ.

References

1. Suh Y, Amelio I, Guerrero Urbano T, Tavassoli M. Clinical update on cancer: molecular oncology of head and neck cancer. *Cell Death Dis.* 2014; 5:e1018. doi: [10.1038/cddis.2013.548](https://doi.org/10.1038/cddis.2013.548) PMID: [24457962](https://pubmed.ncbi.nlm.nih.gov/24457962/)
2. Bose P, Brockton NT, Dort JC. Head and neck cancer: from anatomy to biology. *Int J Cancer.* 2013; 133(9):2013–23. doi: [10.1002/ijc.28112](https://doi.org/10.1002/ijc.28112) PMID: [23417723](https://pubmed.ncbi.nlm.nih.gov/23417723/)
3. Hashibe M, Brennan P, Chuang SC, Boccia S, Castellsague X, Chen C, et al. Interaction between tobacco and alcohol use and the risk of head and neck cancer: pooled analysis in the International Head and Neck Cancer Epidemiology Consortium. *Cancer Epidemiol Biomarkers Prev.* 2009; 18(2):541–50. doi: [10.1158/1055-9965.EPI-08-0347](https://doi.org/10.1158/1055-9965.EPI-08-0347) PMID: [19190158](https://pubmed.ncbi.nlm.nih.gov/19190158/)
4. Marron M, Boffetta P, Zhang ZF, Zaridze D, Wunsch-Filho V, Winn DM, et al. Cessation of alcohol drinking, tobacco smoking and the reversal of head and neck cancer risk. *Int J Epidemiol.* 2010; 39(1):182–96. doi: [10.1093/ije/dyp291](https://doi.org/10.1093/ije/dyp291) PMID: [19805488](https://pubmed.ncbi.nlm.nih.gov/19805488/)
5. Chaturvedi AK, Anderson WF, Lortet-Tieulent J, Curado MP, Ferlay J, Franceschi S, et al. Worldwide trends in incidence rates for oral cavity and oropharyngeal cancers. *J Clin Oncol.* 2013; 31(36):4550–9. doi: [10.1200/JCO.2013.50.3870](https://doi.org/10.1200/JCO.2013.50.3870) PMID: [24248688](https://pubmed.ncbi.nlm.nih.gov/24248688/)
6. Kim YP, Park D, Kim JJ, Choi WJ, Lee SH, Lee SY, et al. Effective Therapeutic Approach for Head and Neck Cancer by an Engineered Minibody Targeting the EGFR Receptor. *PLoS One.* 2014; 9(12): e113442. doi: [10.1371/journal.pone.0113442](https://doi.org/10.1371/journal.pone.0113442) PMID: [25438047](https://pubmed.ncbi.nlm.nih.gov/25438047/)
7. Su NW, Leu YS, Lee JC, Liu CJ, Cheng CY, Lin JS, et al. EGF and EGFR genetic polymorphisms predict prognosis in locally advanced pharyngolaryngeal squamous cell carcinoma patients receiving post-operative concurrent chemoradiotherapy. *Onco Targets Ther.* 2014; 7:2197–204. doi: [10.2147/OTT.S70188](https://doi.org/10.2147/OTT.S70188) PMID: [25506224](https://pubmed.ncbi.nlm.nih.gov/25506224/)
8. Safdari Y, Khalili M, Farajnia S, Asgharzadeh M, Yazdani Y, Sadeghi M. Recent advances in head and neck squamous cell carcinoma—a review. *Clin Biochem.* 2014; 47(13–14):1195–202.

9. Taura M, Kariya R, Kudo E, Goto H, Iwawaki T, Amano M, et al. Comparative analysis of ER stress response into HIV protease inhibitors: lopinavir but not darunavir induces potent ER stress response via ROS/JNK pathway. *Free Radic Biol Med*. 2013; 65:778–88. doi: [10.1016/j.freeradbiomed.2013.08.161](https://doi.org/10.1016/j.freeradbiomed.2013.08.161) PMID: [23973637](https://pubmed.ncbi.nlm.nih.gov/23973637/)
10. Zhou H. HIV protease inhibitors induce endoplasmic reticulum stress and disrupt barrier integrity in intestinal epithelial cells. *Methods Enzymol*. 2011; 490:107–19. doi: [10.1016/B978-0-12-385114-7.00006-4](https://doi.org/10.1016/B978-0-12-385114-7.00006-4) PMID: [21266246](https://pubmed.ncbi.nlm.nih.gov/21266246/)
11. Hetz C, Chevet E, Harding HP. Targeting the unfolded protein response in disease. *Nat Rev Drug Discov*. 2013; 12(9):703–19. doi: [10.1038/nrd3976](https://doi.org/10.1038/nrd3976) PMID: [23989796](https://pubmed.ncbi.nlm.nih.gov/23989796/)
12. Zhou H, Liu R. ER stress and hepatic lipid metabolism. *Front Genet*. 2014; 5:112. doi: [10.3389/fgene.2014.00112](https://doi.org/10.3389/fgene.2014.00112) PMID: [24847353](https://pubmed.ncbi.nlm.nih.gov/24847353/)
13. Chen Y, Gui D, Chen J, He D, Luo Y, Wang N. Down-regulation of PERK-ATF4-CHOP pathway by Astragaloside IV is associated with the inhibition of endoplasmic reticulum stress-induced podocyte apoptosis in diabetic rats. *Cell Physiol Biochem*. 2014; 33(6):1975–87. PMID: [25012492](https://pubmed.ncbi.nlm.nih.gov/25012492/)
14. Masuda M, Miyazaki-Anzai S, Levi M, Ting TC, Miyazaki M. PERK-eIF2alpha-ATF4-CHOP signaling contributes to TNFalpha-induced vascular calcification. *J Am Heart Assoc*. 2013; 2(5):e000238. doi: [10.1161/JAHA.113.000238](https://doi.org/10.1161/JAHA.113.000238) PMID: [24008080](https://pubmed.ncbi.nlm.nih.gov/24008080/)
15. Fassbender JM, Saraswat-Ohri S, Myers SA, Gruenthal MJ, Benton RL, Whittemore SR. Deletion of endoplasmic reticulum stress-induced CHOP protects microvasculature post-spinal cord injury. *Curr Neurovasc Res*. 2012; 9(4):274–81. PMID: [22873727](https://pubmed.ncbi.nlm.nih.gov/22873727/)
16. Zhou H, Gurley EC, Jarujaron S, Ding H, Fang Y, Xu Z, et al. HIV protease inhibitors activate the unfolded protein response and disrupt lipid metabolism in primary hepatocytes. *Am J Physiol Gastrointest Liver Physiol*. 2006; 291(6):G1071–80. PMID: [16861219](https://pubmed.ncbi.nlm.nih.gov/16861219/)
17. Zha W, Liang G, Xiao J, Studer EJ, Hylemon PB, Pandak WM Jr., et al. Berberine inhibits HIV protease inhibitor-induced inflammatory response by modulating ER stress signaling pathways in murine macrophages. *PLoS One*. 2010; 5(2):e9069. doi: [10.1371/journal.pone.0009069](https://doi.org/10.1371/journal.pone.0009069) PMID: [20161729](https://pubmed.ncbi.nlm.nih.gov/20161729/)
18. Wu X, Sun L, Zha W, Studer E, Gurley E, Chen L, et al. HIV protease inhibitors induce endoplasmic reticulum stress and disrupt barrier integrity in intestinal epithelial cells. *Gastroenterology*. 2010; 138(1):197–209. doi: [10.1053/j.gastro.2009.08.054](https://doi.org/10.1053/j.gastro.2009.08.054) PMID: [19732776](https://pubmed.ncbi.nlm.nih.gov/19732776/)
19. Zha BS, Wan X, Zhang X, Zha W, Zhou J, Wabitsch M, et al. HIV protease inhibitors disrupt lipid metabolism by activating endoplasmic reticulum stress and inhibiting autophagy activity in adipocytes. *PLoS One*. 2013; 8(3):e59514. doi: [10.1371/journal.pone.0059514](https://doi.org/10.1371/journal.pone.0059514) PMID: [23533630](https://pubmed.ncbi.nlm.nih.gov/23533630/)
20. Gills JJ, Lopiccio J, Tsurutani J, Shoemaker RH, Best CJ, Abu-Asab MS, et al. Nelfinavir, A lead HIV protease inhibitor, is a broad-spectrum, anticancer agent that induces endoplasmic reticulum stress, autophagy, and apoptosis in vitro and in vivo. *Clin Cancer Res*. 2007; 13(17):5183–94. PMID: [17785575](https://pubmed.ncbi.nlm.nih.gov/17785575/)
21. Yang Y, Ikezoe T, Takeuchi T, Adachi Y, Ohtsuki Y, Takeuchi S, et al. HIV-1 protease inhibitor induces growth arrest and apoptosis of human prostate cancer LNCaP cells in vitro and in vivo in conjunction with blockade of androgen receptor STAT3 and AKT signaling. *Cancer Sci*. 2005; 96(7):425–33. PMID: [16053514](https://pubmed.ncbi.nlm.nih.gov/16053514/)
22. Ikezoe T, Saito T, Bandobashi K, Yang Y, Koeffler HP, Taguchi H. HIV-1 protease inhibitor induces growth arrest and apoptosis of human multiple myeloma cells via inactivation of signal transducer and activator of transcription 3 and extracellular signal-regulated kinase 1/2. *Mol Cancer Ther*. 2004; 3(4):473–9. PMID: [15078991](https://pubmed.ncbi.nlm.nih.gov/15078991/)
23. Pyrko P, Kardosh A, Wang W, Xiong W, Schonthal AH, Chen TC. HIV-1 protease inhibitors nelfinavir and atazanavir induce malignant glioma death by triggering endoplasmic reticulum stress. *Cancer Res*. 2007; 67(22):10920–8. PMID: [18006837](https://pubmed.ncbi.nlm.nih.gov/18006837/)
24. Jiang W, Mikochik PJ, Ra JH, Lei H, Flaherty KT, Winkler JD, et al. HIV protease inhibitor nelfinavir inhibits growth of human melanoma cells by induction of cell cycle arrest. *Cancer Res*. 2007; 67(3):1221–7. PMID: [17283158](https://pubmed.ncbi.nlm.nih.gov/17283158/)
25. Bernstein WB, Dennis PA. Repositioning HIV protease inhibitors as cancer therapeutics. *Curr Opin HIV AIDS*. 2008; 3(6):666–75. doi: [10.1097/COH.0b013e328313915d](https://doi.org/10.1097/COH.0b013e328313915d) PMID: [19373040](https://pubmed.ncbi.nlm.nih.gov/19373040/)
26. Gupta AK, Lee JH, Wilke WW, Quon H, Smith G, Maity A, et al. Radiation response in two HPV-infected head-and-neck cancer cell lines in comparison to a non-HPV-infected cell line and relationship to signaling through AKT. *Int J Radiat Oncol Biol Phys*. 2009; 74(3):928–33. doi: [10.1016/j.ijrobp.2009.03.004](https://doi.org/10.1016/j.ijrobp.2009.03.004) PMID: [19480971](https://pubmed.ncbi.nlm.nih.gov/19480971/)
27. Yang Y, Ikezoe T, Nishioka C, Bandobashi K, Takeuchi T, Adachi Y, et al. NFV, an HIV-1 protease inhibitor, induces growth arrest, reduced Akt signalling, apoptosis and docetaxel sensitisation in NSCLC cell lines. *Br J Cancer*. 2006; 95(12):1653–62. PMID: [17133272](https://pubmed.ncbi.nlm.nih.gov/17133272/)

28. Chen L, Jarujaron S, Wu X, Sun L, Zha W, Liang G, et al. HIV protease inhibitor lopinavir-induced TNF-alpha and IL-6 expression is coupled to the unfolded protein response and ERK signaling pathways in macrophages. *Biochem Pharmacol*. 2009; 78(1):70–7. doi: [10.1016/j.bcp.2009.03.022](https://doi.org/10.1016/j.bcp.2009.03.022) PMID: [19447225](https://pubmed.ncbi.nlm.nih.gov/19447225/)
29. Cao R, Hu Y, Wang Y, Gurley EC, Studer EJ, Wang X, et al. Prevention of HIV protease inhibitor-induced dysregulation of hepatic lipid metabolism by raltegravir via endoplasmic reticulum stress signaling pathways. *J Pharmacol Exp Ther*. 2010; 334(2):530–9. doi: [10.1124/jpet.110.168484](https://doi.org/10.1124/jpet.110.168484) PMID: [20472667](https://pubmed.ncbi.nlm.nih.gov/20472667/)
30. Zhang X, Cao R, Liu R, Zhao R, Huang Y, Gurley EC, et al. Reduction of the HIV protease inhibitor-induced ER stress and inflammatory response by raltegravir in macrophages. *PLoS One*. 2014; 9(3):e90856. doi: [10.1371/journal.pone.0090856](https://doi.org/10.1371/journal.pone.0090856) PMID: [24625618](https://pubmed.ncbi.nlm.nih.gov/24625618/)
31. Zha W, Wang G, Xu W, Liu X, Wang Y, Zha BS, et al. Inhibition of P-glycoprotein by HIV protease inhibitors increases intracellular accumulation of berberine in murine and human macrophages. *PLoS One*. 2013; 8(1):e54349. doi: [10.1371/journal.pone.0054349](https://doi.org/10.1371/journal.pone.0054349) PMID: [23372711](https://pubmed.ncbi.nlm.nih.gov/23372711/)
32. Liu R, Zhao R, Zhou X, Liang X, Campbell DJ, Zhang X, et al. Conjugated bile acids promote cholangiocarcinoma cell invasive growth through activation of sphingosine 1-phosphate receptor 2. *Hepatology*. 2014; 60(3):908–18. doi: [10.1002/hep.27085](https://doi.org/10.1002/hep.27085) PMID: [24700501](https://pubmed.ncbi.nlm.nih.gov/24700501/)
33. Yin X, Zhang H, Lundgren K, Wilson L, Burrows F, Shores CG. BIIB021, a novel Hsp90 inhibitor, sensitizes head and neck squamous cell carcinoma to radiotherapy. *Int J Cancer*. 2010; 126(5):1216–25. doi: [10.1002/ijc.24815](https://doi.org/10.1002/ijc.24815) PMID: [19662650](https://pubmed.ncbi.nlm.nih.gov/19662650/)
34. Zhou H, Jarujaron S, Gurley EC, Chen L, Ding H, Studer E, et al. HIV protease inhibitors increase TNF-alpha and IL-6 expression in macrophages: involvement of the RNA-binding protein HuR. *Atherosclerosis*. 2007; 195(1):e134–43. PMID: [17531241](https://pubmed.ncbi.nlm.nih.gov/17531241/)
35. Wang Y, Zhang L, Wu X, Gurley EC, Kennedy E, Hylemon PB, et al. The role of CCAAT enhancer-binding protein homologous protein in human immunodeficiency virus protease-inhibitor-induced hepatic lipotoxicity in mice. *Hepatology*. 2013; 57(3):1005–16. doi: [10.1002/hep.26107](https://doi.org/10.1002/hep.26107) PMID: [23080229](https://pubmed.ncbi.nlm.nih.gov/23080229/)
36. Zhang X, Wang G, Gurley EC, Zhou H. Flavonoid apigenin inhibits lipopolysaccharide-induced inflammatory response through multiple mechanisms in macrophages. *PLoS One*. 2014; 9(9):e107072. doi: [10.1371/journal.pone.0107072](https://doi.org/10.1371/journal.pone.0107072) PMID: [25192391](https://pubmed.ncbi.nlm.nih.gov/25192391/)
37. McKenna WG, Muschel RJ, Gupta AK, Hahn SM, Bernhard EJ. The RAS signal transduction pathway and its role in radiation sensitivity. *Oncogene*. 2003; 22(37):5866–75. PMID: [12947393](https://pubmed.ncbi.nlm.nih.gov/12947393/)
38. Steelman LS, Navolanic P, Chappell WH, Abrams SL, Wong EW, Martelli AM, et al. Involvement of Akt and mTOR in chemotherapeutic- and hormonal-based drug resistance and response to radiation in breast cancer cells. *Cell Cycle*. 2011; 10(17):3003–15. PMID: [21869603](https://pubmed.ncbi.nlm.nih.gov/21869603/)
39. Gupta AK, Bakanauskas VJ, Cerniglia GJ, Cheng Y, Bernhard EJ, Muschel RJ, et al. The Ras radiation resistance pathway. *Cancer Res*. 2001; 61(10):4278–82. PMID: [11358856](https://pubmed.ncbi.nlm.nih.gov/11358856/)
40. Dionysopoulos D, Pavlakis K, Kotoula V, Fountzilias E, Markou K, Karasmanis I, et al. Cyclin D1, EGFR, and Akt/mTOR pathway. Potential prognostic markers in localized laryngeal squamous cell carcinoma. *Strahlenther Onkol*. 2013; 189(3):202–14. doi: [10.1007/s00066-012-0275-0](https://doi.org/10.1007/s00066-012-0275-0) PMID: [23400686](https://pubmed.ncbi.nlm.nih.gov/23400686/)
41. Wang F, Arun P, Friedman J, Chen Z, Van Waes C. Current and potential inflammation targeted therapies in head and neck cancer. *Curr Opin Pharmacol*. 2009; 9(4):389–95. doi: [10.1016/j.coph.2009.06.005](https://doi.org/10.1016/j.coph.2009.06.005) PMID: [19570715](https://pubmed.ncbi.nlm.nih.gov/19570715/)
42. Bruning A, Rahmeh M, Friese K. Nelfinavir and bortezomib inhibit mTOR activity via ATF4-mediated sestrin-2 regulation. *Mol Oncol*. 2013; 7(6):1012–8. doi: [10.1016/j.molonc.2013.07.010](https://doi.org/10.1016/j.molonc.2013.07.010) PMID: [23916134](https://pubmed.ncbi.nlm.nih.gov/23916134/)
43. Kraus M, Bader J, Overkleeft H, Driessen C. Nelfinavir augments proteasome inhibition by bortezomib in myeloma cells and overcomes bortezomib and carfilzomib resistance. *Blood Cancer J*. 2013; 3:e103. doi: [10.1038/bcj.2013.2](https://doi.org/10.1038/bcj.2013.2) PMID: [23454896](https://pubmed.ncbi.nlm.nih.gov/23454896/)
44. Pore N, Gupta AK, Cerniglia GJ, Jiang Z, Bernhard EJ, Evans SM, et al. Nelfinavir down-regulates hypoxia-inducible factor 1alpha and VEGF expression and increases tumor oxygenation: implications for radiotherapy. *Cancer Res*. 2006; 66(18):9252–9. PMID: [16982770](https://pubmed.ncbi.nlm.nih.gov/16982770/)
45. Parker RA, Flint OP, Mulvey R, Elosua C, Wang F, Fenderson W, et al. Endoplasmic reticulum stress links dyslipidemia to inhibition of proteasome activity and glucose transport by HIV protease inhibitors. *Mol Pharmacol*. 2005; 67(6):1909–19. PMID: [15755908](https://pubmed.ncbi.nlm.nih.gov/15755908/)
46. Pang XL, He G, Liu YB, Wang Y, Zhang B. Endoplasmic reticulum stress sensitizes human esophageal cancer cell to radiation. *World J Gastroenterol*. 2013; 19(11):1736–48. doi: [10.3748/wjg.v19.i11.1736](https://doi.org/10.3748/wjg.v19.i11.1736) PMID: [23555162](https://pubmed.ncbi.nlm.nih.gov/23555162/)
47. Schonthal AH. Endoplasmic reticulum stress: its role in disease and novel prospects for therapy. *Scientifica (Cairo)*. 2012; 2012:857516. doi: [10.6064/2012/857516](https://doi.org/10.6064/2012/857516) PMID: [24278747](https://pubmed.ncbi.nlm.nih.gov/24278747/)

48. Fokas E, Yoshimura M, Prevo R, Higgins G, Hackl W, Maira S-M, et al. NVP-BEZ235 and NVP-BGT226, dual phosphatidylinositol 3-kinase/mammalian target of rapamycin inhibitors, enhance tumor and endothelial cell radiosensitivity. *Radiation Oncology*. 2012; 7(1):48.
49. Liang J, Slingerland JM. Multiple roles of the PI3K/PKB (Akt) pathway in cell cycle progression. *Cell Cycle*. 2003; 2(4):339–45. PMID: [12851486](#)
50. Testa JR, Bellacosa A. AKT plays a central role in tumorigenesis. *Proceedings of the National Academy of Sciences*. 2001; 98(20):10983–5. PMID: [11572954](#)
51. Toulany M, Lee KJ, Fattah KR, Lin YF, Fehrenbacher B, Schaller M, et al. Akt Promotes Post-Irradiation Survival of Human Tumor Cells through Initiation, Progression, and Termination of DNA-PKcs-Dependent DNA Double-Strand Break Repair. *Molecular Cancer Research*. 2012; 10(7):945–57. doi: [10.1158/1541-7786.MCR-11-0592](#) PMID: [22596249](#)
52. Hamanaka RB. PERK and GCN2 Contribute to eIF2 Phosphorylation and Cell Cycle Arrest after Activation of the Unfolded Protein Response Pathway. *Molecular Biology of the Cell*. 2005; 16(12):5493–501. PMID: [16176978](#)
53. Brewer JW, Diehl JA. PERK mediates cell-cycle exit during the mammalian unfolded protein response. *Proceedings of the National Academy of Sciences*. 2000; 97(23):12625–30. PMID: [11035797](#)
54. Raven JF, Baltzis D, Wang S, Mounir Z, Papadakis AI, Gao HQ, et al. PKR and PKR-like Endoplasmic Reticulum Kinase Induce the Proteasome-dependent Degradation of Cyclin D1 via a Mechanism Requiring Eukaryotic Initiation Factor 2 Phosphorylation. *Journal of Biological Chemistry*. 2007; 283(6): 3097–108. PMID: [18063576](#)
55. George S, Kasimis BS, Cogswell J, Schwarzenberger P, Shapiro GI, Fidias P, et al. Phase I study of flavopiridol in combination with Paclitaxel and Carboplatin in patients with non-small-cell lung cancer. *Clin Lung Cancer*. 2008; 9(3):160–5. doi: [10.3816/CLC.2008.n.024](#) PMID: [18621626](#)