



Draft Genome Sequence of *Paraburkholderia* sp. UYCP14C, a Rhizobium Strain Isolated from Root Nodules of *Calliandra parvifolia*

Mauricio Langleib,^{a,b} Martín Beracochea,^a María Zabaleta,^{a,c} Federico Battistoni,^a José Sotelo-Silveira,^d Elena Fabiano,^a Andrés Iriarte,^b Raúl Platero^a

^aDepartamento de Bioquímica y Genómica Microbianas, Instituto de Investigaciones Biológicas Clemente Estable, MEC, Montevideo, Uruguay

^bLaboratorio de Biología Computacional, Departamento de Desarrollo Biotecnológico, Instituto de Higiene, Facultad de Medicina, Universidad de la República, Montevideo, Uruguay

^cFacultad de Agronomía, CURE, Universidad de la República, Montevideo, Uruguay

^dDepartamento de Genómica, Instituto de Investigaciones Biológicas Clemente Estable, MEC, Montevideo, Uruguay

ABSTRACT Here, we present the draft genome sequence of strain UYCP14C, a rhizobium isolated from *Calliandra parvifolia* nodules. The assembled genome size was around 9.8 million bp, containing 9,031 predicted protein-coding sequences, including several symbiotic and nitrogen fixation genes. UYCP14C appears to be a novel species of the plant growth-promoting *Paraburkholderia* genus.

The genus *Paraburkholderia* is composed of Gram-negative, rod-shaped, straight or slightly curved bacteria that contain one or more flagella. These bacteria are metabolically and morphologically similar to the members of their sister clades *Burkholderia*, *Caballeronia*, and *Robbsia* (1, 2). *Paraburkholderia* comprises diverse environmental and plant-beneficial strains, including dinitrogen-fixing legume symbionts (3–9). In particular, the ability of *Paraburkholderia* spp. to nodulate plants of the genus *Calliandra* has been reported (10). The genome sequences of these rhizobia are valuable resources for understanding the mechanisms and evolution of beneficial plant-microbe interactions (11).

Here, we present the draft genome sequence of the bacterial strain UYCP14C, a rhizobium isolated from root nodules of the native legume *Calliandra parvifolia*, naturally occurring in Rio Negro Department, Uruguay.

Root nodule collection and symbiotic bacterial isolation were performed as previously described (12, 13). UYCP14C identification was performed at EzBioCloud (14) using the PCR-amplified 16S rRNA gene sequence. The most similar sequence (99.49%) corresponded to *Paraburkholderia monticola* strain JC2948^T. Genomic DNA was isolated from stationary-phase cultures in liquid tryptone-yeast (TY) medium (15) using the ZR fungal/bacterial DNA kit (ZymoResearch). The whole-genome, paired-end library was constructed using the TruSeq PCR-free kit (Illumina), and samples were run on a HiSeq 2500 sequencer (Illumina) at Macrogen (South Korea). A total of 2.3 × 10⁷ paired-end reads 101 bp long were produced, with 99.66% of the reads having a Phred quality score over 30.

Sequence trimming was performed using Sickle 1.33 (16). The UYCP14C genome was assembled with SPAdes v3.9.0 (17), keeping paired reads unmerged before assembly. The kmer length parameter was set to 21, 33, 55, and 77, and SPAdes was executed with the “–careful” flag, which activates the BayesHammer read correction module (18). Genome assembly quality was assessed with QUAST v5.0.1 (17). The total assembly sums 9.83 megabase pairs, distributed in 427 contigs, with an N_{50} contig size of 121.2 kb and a maximum length of 381.2 kb. The average coverage reported by the

Citation Langleib M, Beracochea M, Zabaleta M, Battistoni F, Sotelo-Silveira J, Fabiano E, Iriarte A, Platero R. 2019. Draft genome sequence of *Paraburkholderia* sp. UYCP14C, a rhizobium strain isolated from root nodules of *Calliandra parvifolia*. *Microbiol Resour Announc* 8:e00173-19. <https://doi.org/10.1128/MRA.00173-19>.

Editor Frank J. Stewart, Georgia Institute of Technology

Copyright © 2019 Langleib et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

Address correspondence to Andrés Iriarte, ariarte@higiene.edu.uy, or Raúl Platero, rplatero@iibce.edu.uy.

Received 18 February 2019

Accepted 20 March 2019

Published 18 April 2019

assembler is approximately 60×. Genome G+C content was 63.07%, which is within the range reported for *Paraburkholderia* species (2).

The genome was uploaded to the RAST v2.0 annotation server under classic annotation scheme (18). In total, 9,031 protein-coding genes and 67 RNA-coding genes were annotated. Genes related to nodulation, such as *nodCDHII*, and nitrogen-fixation, such as *nifHDK* (among others), were found.

Pairwise average nucleotide identities were calculated using OrthoANI calculator v1.40 (19) by comparing the UYCP14C genome with the 43 genome sequences of organisms of the *Paraburkholderia* genus available in GenBank and analyzed previously (1). In all cases, the average nucleotide identity (ANI) was below the usual threshold of 95% employed as a minimum to designate members of the same species, suggesting that the UYCP14C genome is the first of a new *Paraburkholderia* species. Further studies are in progress to validate this hypothesis.

Data availability. Raw reads are available under the BioSample accession number [SAMN10846787](#). The assembled genome sequence was deposited at GenBank under BioProject accession number [PRJNA517875](#). The version described in this paper is the first version under the accession number [SEL500000000](#).

REFERENCES

- Estrada-de los Santos P, Palmer M, Chávez-Ramírez B, Beukes C, Steenkamp ET, Briscoe L, Khan N, Maluk M, Lafos M, Humm E, Arrabit M, Crook M, Gross E, Simon MF, dos Reis Junior FB, Whitman WB, Shapiro N, Poole PS, Hirsch AM, Venter SN, James EK. 2018. Whole genome analyses suggests that *Burkholderia* sensu lato contains two additional novel genera (*Mycetohabitans* gen. nov., and *Trinickia* gen. nov.): implications for the evolution of diazotrophy and nodulation in the *Burkholderiaceae*. *Genes (Basel)* 9:E389. <https://doi.org/10.3390/genes9080389>.
- Sawana A, Adeolu M, Gupta RS. 2014. Molecular signatures and phylogenomic analysis of the genus *Burkholderia*: proposal for division of this genus into the emended genus *Burkholderia* containing pathogenic organisms and a new genus *Paraburkholderia* gen. nov. harboring environmental species. *Front Genet* 5:429. <https://doi.org/10.3389/fgene.2014.00429>.
- Willems A. 2006. The taxonomy of rhizobia: an overview. *Plant Soil* 287:3–14. <https://doi.org/10.1007/s11104-006-9058-7>.
- Chen W-M, James EK, Coenye T, Chou J-H, Barrios E, de Faria SM, Elliott GN, Sheu S-Y, Sprent JI, Vandamme P. 2006. *Burkholderia mimosarum* sp. nov., isolated from root nodules of *Mimosa* spp. from Taiwan and South America. *Int J Syst Evol Microbiol* 56:1847–1851. <https://doi.org/10.1099/ijss.0.64325-0>.
- Dall'Agnol RF, Costa MR, Ribeiro RA, Delamuta JRM, Chueire LMO, Hungria M. 2016. Genome sequence of *Paraburkholderia nodosa* strain CNPSo 1341, a N₂-fixing symbiont of the promiscuous legume *Phaseolus vulgaris*. *Genome Announc* 4:e01073-16. <https://doi.org/10.1128/genomeA.01073-16>.
- Sheu S-Y, Chou J-H, Bontemps C, Elliott GN, Gross E, James EK, Sprent JI, Young JPW, Chen W-M. 2012. *Burkholderia symbiotica* sp. nov., isolated from root nodules of *Mimosa* spp. native to north-east Brazil. *Int J Syst Evol Microbiol* 62:2272–2278. <https://doi.org/10.1099/ijss.0.037408-0>.
- Chen W-M, Moulin L, Bontemps C, Vandamme P, Béna G, Boivin-Masson C. 2003. Legume symbiotic nitrogen fixation by β-proteobacteria is widespread in nature. *J Bacteriol* 185:7266–7272. <https://doi.org/10.1128/JB.185.24.7266-7272.2003>.
- Martínez-Aguilar L, Salazar-Salazar C, Mendez RD, Caballero-Mellado J, Hirsch AM, Vásquez-Murrieta MS, Estrada-De Los Santos P. 2013. *Burkholderia caballeronis* sp. nov., a nitrogen fixing species isolated from tomato (*Lycopersicon esculentum*) with the ability to effectively nodulate *Phaseolus vulgaris*. *Antonie Van Leeuwenhoek* 104:1063–1071. <https://doi.org/10.1007/s10482-013-0028-9>.
- Gyaneshwar P, Hirsch AM, Moulin L, Chen W-M, Elliott GN, Bontemps C, Estrada-de Los Santos P, Gross E, Dos Reis FB, Jr, Sprent JI, Young JPW, James EK. 2011. Legume-nodulating betaproteobacteria: diversity, host range, and future prospects. *Mol Plant Microbe Interact* 24:1276–1288. <https://doi.org/10.1094/MPMI-06-11-0172>.
- Silva VC, Alves PAC, Rhem MFK, dos Santos JMF, James EK, Gross E. 2018. Brazilian species of *Calliandra* Benth. (tribe Ingeae) are nodulated by diverse strains of *Paraburkholderia*. *Syst Appl Microbiol* 41:241–250. <https://doi.org/10.1016/j.syapm.2017.12.003>.
- Reeve W, Ardley J, Tian R, Eshragi L, Yoon JW, Ngamwisetkun P, Seshadri R, Ivanova NN, Kyrides NC. 2015. A genomic encyclopedia of the root nodule bacteria: assessing genetic diversity through a systematic bio-geographic survey. *Stand Genomic Sci* 10:14. <https://doi.org/10.1186/1944-3277-10-14>.
- Platero R, James EK, Rios C, Iriarte A, Sandes L, Zabaleta M, Battistoni F, Fabiano E. 2016. Novel *Cupriavidus* strains isolated from root nodules of native Uruguayan *Mimosa* species. *Appl Environ Microbiol* 82: 3150–3164. <https://doi.org/10.1128/AEM.04142-15>.
- Taulé C, Zabaleta M, Mareque C, Platero R, Sanjurjo L, Sicardi M, Frioni L, Battistoni F, Fabiano E. 2012. New betaproteobacterial *Rhizobium* strains able to efficiently nodulate *Parapiptadenia rigida* (Benth.) Brenan. *Appl Environ Microbiol* 78:1692–1700. <https://doi.org/10.1128/AEM.06215-11>.
- Yoon S-H, Ha S-M, Kwon S, Lim J, Kim Y, Seo H, Chun J. 2017. Introducing EzBioCloud: a taxonomically united database of 16S rRNA gene sequences and whole-genome assemblies. *Int J Syst Evol Microbiol* 67: 1613–1617. <https://doi.org/10.1099/ijsem.0.001755>.
- Vincent JM. 1970. A manual for the practical study of root-nodule bacteria. Blackwell Scientific Ltd., Oxford, United Kingdom.
- Joshi NA, Fass JN. 2011. Sickle: a sliding-window, adaptive, quality-based trimming tool for FastQ files (Version 1.33).
- Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for genome assemblies. *Bioinformatics* 29:1072–1075. <https://doi.org/10.1093/bioinformatics/btt086>.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.
- Lee I, Kim YO, Park S-C, Chun J. 2016. OrthoANI: an improved algorithm and software for calculating average nucleotide identity. *Int J Syst Evol Microbiol* 66:1100–1103. <https://doi.org/10.1099/ijsem.0.000760>.