

Visualizing Abduction

Fernando Soler Toscano

(Grupo de Lógica, Lenguaje e Información, Universidad de Sevilla)

fsoler@us.es

1 Introduction

1.1 What is abduction?

Traditionally, Logic has focus on *deduction*. Its main objective has been to clarify when a given inference is *valid*, that is, when the conclusion is *entailed* by the premises. One of the pillars of this enterprise is the Aristotelian *Analytics*, where the syllogism is taken as the paradigm for reasoning. The modern notion of *logical consequence*, though is defined in a more abstract and formalised way, ratifies the privileged position of deduction as the centre of the logical universe.

But there exist other kinds of inference. The philosopher C. S. Peirce (1839 - 1914), distinguishes three kinds of reasoning: *deduction*, *induction* and *abduction*. Peirce, contrary to the logical tradition, considers that the most interesting kind of inference is abduction. Peirce characterizes abduction in this schematic way:

The surprising fact, F, is observed;
but if H were true, F would be a matter of course.
Hence, there is reason to suspect that H is true (CP 5.189, 1903).

So abduction is the inference which formulates a hypothesis H that explains a surprising fact F. If we turn back to Aristotle, in the *Prior Analytics* he establishes not only the categorical syllogism, but also other inference patterns which change the order of the propositions. Those patterns, though non deductively valid, are very important in human reasoning. One of them is the *apagoge*. An example of Aristotle:

Animals without bile live many years.
But the man, the horse and the mule live many years.
Hence, the man, the horse and the mule do not have bile.

The similarity between the Aristotelian *apagoge* and the Peircean *abduction* is not a matter of chance. In fact, Peirce himself admits the borrowing of concepts from Aristotle. But for Peirce, abduction is not only a kind of reasoning among others. Peirce thinks that abduction is the most interesting kind of inference because it is the only way of introducing new ideas (EP 2:216, 1903).

Thanks to the Peircean works, abductive reasoning has become a central topic not only in Logic, but also in other disciplines as Linguistics, Epistemology, Artificial Intelligence and, of course, Philosophy of Science.

1.2 Is abduction the logic of discovery?

Philosophy of Science is, exactly, the discipline in which the importance of abduction is more discussed. It is usual the distinction between the *context of discovery* and the *context of justification*. The former is understood as the set of processes which lead to the formulation of a new theory, while the latter refers to the methods used in science to confront the hypothesis with the empirical evidence. P. Lipton (1991) defends that scientific discovery is achieved as an *inference to the best explanation*, so *explanatory reasoning*, and hence abduction, play a central role in the context of discovery. But other authors are critical with these ideas.

S. Paavola (2004) collects the arguments that are commonly given against the characterisation of abduction as a *logic of (scientific) discovery*. The first of them is that in the Peircean formulation of abduction – given above – the only requirement for H is that it would make F true. Authors like P. Achinstein argue that this is too permissive, because it does not exclude some improbable or wild hypotheses if they would explain F. Other criticisms insist in the idea that abduction cannot be understood as a logic of discovery because the explanatory hypothesis H is already included in the premises, because prior to infer H we need to know that H would explain F. Then, as scientific discovery introduces new ideas in science, it cannot be the result of an abductive inference.

To defend abduction as a logic of discovery, S. Paavola follows a distinction borrowed by J. Hintikka from game theory. It is the distinction between *definitory* and *strategic* rules. The former settle the legal moves in a game, whereas the latter are used to decide which is the most suitable rule among all the possible. Logic has been traditionally devoted to definitory rules of the calculi and has barely paid attention to the strategic rules which are able to lead to a good proof, in a strategic sense that goes beyond classical soundness.

In line with this, J. Hintikka (1998) takes from T. Kapitan four theses which sum up the main characteristics of abduction:

Inferential Thesis. Abduction is, or includes, an inferential process or processes.

Thesis of Purpose. The purpose of “scientific” abduction is both (i) to generate new hypotheses and (ii) to select hypotheses for further examination; hence, a central aim of scientific abduction is to “recommend a course of action”.

Comprehension Thesis. Scientific abduction includes *all* the operations whereby theories are engendered.

Autonomy Thesis. Abduction is, or embodies, reasoning that is distinct from, and irreducible to, either deduction or induction.

We will come back to Kapitan theses. For now, it is enough to remark that there are reasons to accept abduction as a logic of discovery. But then, abduction must include some kind of strategic rules and satisfy the above theses which set up abduction as an autonomous inferential process, irreducible to deduction.

2 Abduction by dualizing deduction

2.1 Logic-based abduction

To introduce the formal definitions concerning abduction, let us consider that \mathcal{L} is a propositional language with the habitual connectives, and let \models be defined as the classical logical consequence relation. We use capital Greek letters for sets of formulas and small Greek letters to denote formulas.

Definition 1 (Abductive problem). Given $\Theta \subset \mathcal{L}$ and $\phi \in \mathcal{L}$, we say (Θ, ϕ) is an abductive problem iff (if and only if):

$$\Theta \not\vdash \phi \tag{1}$$

$$\Theta \not\vdash \neg\phi \tag{2}$$

Definition 2 (Abductive solution). Given the abductive problem (Θ, ϕ) , $\alpha \in \mathcal{L}$ is an abductive solution for it iff:

$$\Theta, \alpha \models \phi \tag{3}$$

$$\Theta, \alpha \not\vdash \perp \tag{4}$$

$$\alpha \not\vdash \phi \tag{5}$$

According to definition 1, an *abductive problem* appears whenever there is a formula ϕ such that neither it (1) nor its negation (2) can be derived by only the

background theory Θ . Then, α is an *abductive solution* if it extends the theory in a way such that now $\Theta \cup \{\alpha\}$ entail ϕ (3). Some authors demand only this condition, but following A. Aliseda (2006) we have included in definition 2 requirements (4) and (5) to ensure that α is not a trivial explanation. That is, α is consistent with the theory and it does not entail ϕ by itself without the theory, so it is an explanation *within* the theory. With these strong requirements, abduction becomes an interesting non-monotonic inference very different from deduction. Additional conditions may be added. In fact, we will later concentrate on *conjunctive minimal explanations*, conjunctions of literals such that no proper subset of them is an abductive explanation.

To mechanise the generation of abductive solutions, many calculi have been proposed. Most of them make *abductive uses* of *deductive calculi* by exploiting the equivalence of (3) with:

- $\Theta, \neg\phi \models \neg\alpha$. This is done by most on the logic-based approaches coming from Artificial Intelligence (Kakas *et al.* 1998). Using the resolution calculus, the clausal form of $\Theta \cup \{\neg\phi\}$ is obtained and then resolution is applied. Any dead end of the resolution tree can be taken as the negation of an abductive solution.

- $\Theta, \neg\phi, \alpha \models \perp$. This is what the semantic tableaux approach does (Mayer *et al.*, 1993). The tableaux of $\Theta \cup \{\neg\phi\}$ is obtained and then a formula α which closes all the open branches is searched.

In both approaches $\neg\phi$, the negation of what is intended for explain, is in the starting point of the abductive search. So the abductive process starts by negating the empirical evidence which tries to explain. This is somehow similar to *reductio ad absurdum*, because the explanation α becomes an extension of Θ that makes impossible $\neg\phi$.

Proceeding in this way has moved logical abduction further away from the great expectations coming from Philosophy of Science. It is hardly believable that a logic of discovery proceeds by negating exactly what is trying to explain. It is not possible to take the above procedures as a logical model either of scientific or commonsense reasoning.

2.2 The δ -resolution calculus

Definitions 1 and 2 restrict the scope of abductive reasoning. They do not seem to be appropriate to include «all the operations whereby theories are engendered», as Kapitan's *comprehension thesis* requires. Anyway, though reductive, those definitions can be understood as a scale model of (scientific) explanation. They comprise the

common features of any explanatory process. But to increase their interest a proper abductive calculus should be formulated. That is, a calculus which neither proceeds in an *indirect* way or assimilates explanation to *reductio ad absurdum*.

That is what we have done with the δ -resolution calculus. It is a reformulation of resolution which turns it an abductive calculus which generates explanations in a *direct* way. It works by using the equivalence between (3) and $\alpha \models \Theta \rightarrow \phi$, where Θ denotes the conjunction of its formulas. Now, the observation ϕ is not negated, and we obtain directly abductive explanations, not their negations. Let us see an informal sketch of the process applied to an example of Kakas *et al.* (1998). Let *rained*, *sprinkler*, *grass* and *shoes* represent, respectively “rained last night”, “sprinkler was on”, “grass is wet” and “shoes are wet”. Then, the theory is:

$$\begin{aligned} & \textit{rained} \rightarrow \textit{grass} \\ & \textit{sprinkler} \rightarrow \textit{grass} \\ & \textit{grass} \rightarrow \textit{shoes} \end{aligned}$$

We want the theory to explain that “shoes are wet”, that is:

$$(\textit{rained} \rightarrow \textit{grass}) \wedge (\textit{sprinkler} \rightarrow \textit{grass}) \wedge (\textit{grass} \rightarrow \textit{shoes}) \rightarrow \textit{shoes} \quad (6)$$

If this is a valid formula, then the theory itself explains that the shoes are wet. Otherwise, the theory needs an additional support, that is, an abductive explanation. When is (6) true? A conditional is true when the antecedent is false or the consequent is true:

$$\neg((\textit{rained} \rightarrow \textit{grass}) \wedge (\textit{sprinkler} \rightarrow \textit{grass}) \wedge (\textit{grass} \rightarrow \textit{shoes})) \vee \textit{shoes} \quad (7)$$

This sets the two possible extremes of an abductive process. We can refuse the theory if the observation contradicts it, or we can add the observation to our knowledge base, if there is no possible explanation within the theory. But **Error! Reference source not found.** is equivalent to:

$$(\textit{rained} \wedge \neg\textit{grass}) \vee (\textit{sprinkler} \wedge \neg\textit{grass}) \vee (\textit{grass} \wedge \neg\textit{shoes}) \vee \textit{shoes} \quad (7)$$

Any disjointed term in (7) is a formula which supports (6), by either contradicting the theory or assuming the observation. But, is there any intermediate alternative? Of course, these are the abductive explanations. For example, both $(\textit{grass} \wedge \neg\textit{shoes})$ and *shoes* support (6). So, also *grass* because, whenever it is true, one of $(\textit{grass} \wedge \neg\textit{shoes})$ or *shoes* is too, as a trivial semantic reasoning shows. So *grass* is a possible explanation. It is not the best, as we can continue the abductive process, to explain why the grass is wet. From just obtained *grass* and

($rained \wedge \neg grass$) we get $rained$. Also, $grass$ and ($sprinkler \wedge \neg grass$) produce $sprinkler$. The three obtained explanations, $grass$, $rained$ and $sprinkler$ are abductive solutions.

The previous example shows that δ -resolution is in fact a dual version of resolution¹. In the following, we introduce the most important definitions and results concerning propositional δ -resolution. Formal proofs can be found in (Soler-Toscano *et al.*, 2006) and an extension to predicate logic in (Soler-Toscano *et al.*, 2009).

Definition 3. A δ -clause Σ is a finite set of literals of \mathcal{L} . Given a boolean valuation v , $v \models \Sigma$ iff v satisfies all the literals of Σ . The empty δ -clause, \diamond , is universally valid.

Definition 4. A δ -clausal form A is a finite set of δ -clauses. Given a boolean valuation v , $v \models A$ iff v satisfies at least one δ -clause of A . The empty δ -clausal form is not satisfiable.

It is possible to translate any formula to an equivalent δ -clausal form by obtaining its disjunctive normal form. The following definition introduces two additional restrictions in the requirements of definition 2. We select only minimal conjunctions of literals.

Definition 5 (Set of abductive δ -clauses). Given the abductive problem (Θ, ϕ) , the set of abductive δ -clauses $Abd(\Theta, \phi)$ contains every δ -clause Σ such that:

- The conjunction of the literals of Σ is an abductive solution for (Θ, ϕ) .
- There is no $\Sigma' \subsetneq \Sigma$ such that $\Theta, \Sigma' \models \phi$.

Definition 6 (δ -resolution rule). Given two δ -clauses $\Sigma_1 \cup \{\lambda\}$ and $\Sigma_2 \cup \{\neg\lambda\}$, the δ -resolution rule produces their δ -resolvent $\Sigma_1 \cup \Sigma_2$:

$$\frac{\Sigma_1 \cup \{\lambda\} \quad \Sigma_2 \cup \{\neg\lambda\}}{\Sigma_1 \cup \Sigma_2}$$

Though this rule is presented with the same format that the standard resolution one, they are different since now we are working with δ -clauses. In the standard

¹ Dual versions of the resolution calculus are introduced in (Eder 1991) and (Ligeza 1993). We introduced the abductive possibilities of dual resolution in (Soler-Toscano *et al.* 2006).

resolution calculus (Robinson, 1965), every obtained clause is a logical consequence of the original set. Now, any δ -clausal form which contains $\Sigma_1 \cup \{\lambda\}$ and $\Sigma_2 \cup \{\neg\lambda\}$ is a logical consequence of $\Sigma_1 \cup \Sigma_2$, because any valuation ν which satisfies $\Sigma_1 \cup \Sigma_2$ satisfies λ or $\neg\lambda$, so ν satisfies $\Sigma_1 \cup \{\lambda\}$ or $\Sigma_2 \cup \{\neg\lambda\}$. Then ν satisfies any δ -clausal form with $\Sigma_1 \cup \{\lambda\}$ and $\Sigma_2 \cup \{\neg\lambda\}$.

Definition 7 (Proof by δ -resolution). *The δ -clause Σ is provable by δ -resolution from the δ -clausal form A , what we express with $A \vdash_{\delta} \Sigma$, iff there is a sequence of δ -clauses such that:*

- Each δ -clause in the sequence is either a member of A or a δ -resolvent of previous δ -clauses.
- Σ is the last δ -clause of the sequence.

In deductive logic, soundness and completeness results are important to prove the adequacy of a calculus. Now, these properties are related to abductive adequacy of dual resolution, that is, the δ -resolution process produces every abductive solution, and just them.

Theorem 8 (Soundness). *For every δ -clausal form A and δ -clause Σ , if $A \vdash_{\delta} \Sigma$, then $\Sigma \models A$.*

Theorem 9 (Completeness). *If A is an universally valid δ -clausal form, then $A \vdash_{\delta} \diamond$.*

The following theorem proves the *abductive completeness* of the δ -resolution calculus, that is, all the δ -clauses that satisfy Definition 5 can be proved by δ -resolution.

Theorem 10 (Abductive Completeness). *Let A be the δ -clausal form of $\alpha \in \mathcal{L}_p$. Then $A \vdash_{\delta} \Sigma$ for each satisfiable δ -clause Σ such that:*

- $\Sigma \models \alpha$.
- For every $\Sigma' \subsetneq \Sigma$, $\Sigma' \not\models \alpha$.

Definition 11 (Saturation). Given the δ -clausal form A , the set saturation by δ -resolution from A , that we represent as A^δ , is the minimal set which contains every δ -clause Σ such that

- Σ is satisfiable.
- $A \vdash_\delta \Sigma$.
- There is not $\Sigma' \subsetneq \Sigma$ such that $A \vdash_\delta \Sigma'$.

Given a finite set of δ -clauses A , A^δ can be obtained in a finite number of steps, by successive applications of the δ -resolution rule, and eliminating subsumed² and contradictory δ -clauses.

The following is the fundamental theorem of the δ -resolution calculus as it provides the right way for obtaining all abductive solutions by means of a δ -resolution process.

Theorem 12 (Fundamental theorem). For a given abductive problem $(\{\theta_1, \dots, \theta_n\}, \phi)$, if N_Θ and O are respectively the δ -clausal forms of $\neg(\theta_1 \wedge \dots \wedge \theta_n)$ and ϕ , then

$$Abd(\Theta, \phi) = (N_\Theta^\delta \cup O^\delta)^\delta - (N_\Theta^\delta \cup O^\delta)$$

2.3 An abductive process

By using only δ -resolution operations, an abductive process can be defined, as it is implicit in Theorem 12. Given $\Theta = \{\theta_1, \dots, \theta_n\}$ and ϕ , it follows four steps to determine whether (Θ, ϕ) is an abductive problem and, in the affirmative case, to produce all of its abductive solutions:

Step 1: Theory Analysis. Let N_Θ be the δ -clausal form of $\neg(\theta_1 \wedge \dots \wedge \theta_n)$. Then:

- If N_Θ does not contain any satisfiable δ -clause, then Θ is universally valid, and the process stops, because in case (Θ, ϕ) is an abductive problem it cannot have abductive solutions in the sense of definition 2.

²The δ -clause Σ is *subsumed* by Σ' iff $\Sigma' \subsetneq \Sigma$.

- Else, N_{Θ}^{δ} is obtained, and:
 - If $\diamond \in N_{\Theta}^{\delta}$, then Θ is not satisfiable, and the process stops, because (Θ, ϕ) cannot be an abductive problem.
 - Else,

Step 2: Observation Analysis. Let O be the δ -clausal form of ϕ . Then:

- If O does not contain any satisfiable δ -clause, then ϕ is not satisfiable, and the process stops, because (Θ, ϕ) cannot be an abductive problem.
- Else, O^{δ} is obtained, and:
 - If $\diamond \in O^{\delta}$, then ϕ is universally valid, and the process stops (as $\Theta \models \phi$, (Θ, ϕ) is not an abductive problem).
 - Else,

Step 3: Refutation Search. If for every δ -clause $\Sigma \in O^{\delta}$ there is a $\Sigma' \subseteq \Sigma$ such that $\Sigma' \in N_{\Theta}^{\delta}$, then $\Theta \models \neg\phi$, and the process stops because the observation refutes the theory. Else,

Step 4: Explanations Search. From N_{Θ}^{δ} and O^{δ} , $(N_{\Theta}^{\delta} \cup O^{\delta})$ and then $(N_{\Theta}^{\delta} \cup O^{\delta})^{\delta}$ are obtained. Then,

- If $\diamond \in (N_{\Theta}^{\delta} \cup O^{\delta})^{\delta}$, then $\Theta \models \phi$ and the process stops.
- Else, (Θ, ϕ) is an abductive problem. The process returns:

$$Abd(\Theta, \phi) = (N_{\Theta}^{\delta} \cup O^{\delta})^{\delta} - (N_{\Theta}^{\delta} \cup O^{\delta})$$

Is there a *logic of abduction*? This is a recurrent question with a difficult answer. Abductive reasoning has a double character. It is a product, but also a process. Moreover, the process to obtain an explanation is maybe more interesting than the explanation itself. So, the answer cannot be affirmative if there is not something like an *abductive logic* which integrates abductive process and product. As we argued, traditional logic-based approaches obtain abductive products which fulfil definitions 1 and 2, but their processes can hardly be considered abductive, because of the abuse of deduction and *reductio ad absurdum*.

However, δ -resolution can be considered an abductive logic. Not only its products are correct (theorem 12), but also it is possible to define an abductive process which proceeds only by δ -resolution operations, as we have just shown. The steps of this process can be connected with some ideas coming from the Philosophy of Science. As we show in the next section, it is possible to visualize this process in a

way that makes it very intuitive. So δ -resolution agrees with common arguments for abductive logic.

3. Abductive diagrams

Peirce thinks that diagrammatic elements are fundamental in cognition in general and creativity in particular. So, it makes sense to look for a diagrammatic approach to abduction. In this section, we partly follow the ideas in (San Ginés, 2011). She makes an original representation of propositional logic that allows her to visualize some inference rules. The system has an usability problem that makes difficult reasoning within a two dimensional sheet of paper: it requires a temporal dimension to represent material implication by means of colour-changing squares. But if we simplify the notation and change a little the semantics of the pictures, we can take the approach as a good representation for a sound and complete abductive calculus. In the original formulation, there is a pool of formulas that is interpreted as a conjunction. Now, we take the pool as a δ -clausal form, so its elements (δ -clauses) are interpreted in a disjunction.

First, to represent some proposition we use a colour square. The colour is always the same for a given proposition. If the square is crossed out, it represents the negation of the proposition. As a δ -clause is a set of literals, we join the squares that represent its literals. For example, if the blue square represents p , the green is q and the red one r , the following is a δ -clause that represents $p \wedge \neg q \wedge r$:




When several δ -clauses are represented in a same pool, we interpret it as a disjunction, that is, a δ -clausal form.

The rules of the δ -resolution calculus can be represented with these conventions. To do that, we first need a way for representing a general δ -clause. We do it with a dark rectangle, that means some irrelevant set of literals. As an example, look at the following set of δ -clauses:




The first one represents some δ -clause where a given literal is negated and the second one another δ -clause where the same literal is affirmed. The literals in the dark boxes could be related or not. Note that, as the order of the literals is not relevant, the position of the red squares is arbitrary. And also the position of the δ -clauses, of course.

With these resources we can represent the δ -resolution rule. Note that if the previous two δ -clauses are in the same pool, they constitute two different explanations for a given abductive problem: one where the red proposition is affirmed and another one where is negated. Then the δ -resolution rule allows us to propose a new possible explanation that does not use that proposition, as the following figure shows. Arrows indicate the direction of the inference. As it is an abductive inference it is the opposite direction to that of the logical consequence (look at the Soundness Theorem above).




With these elements we can illustrate the δ -resolution process with an example. We start with the δ -clausal form that corresponds to the *wet shoes* example given at section 2.2. The following picture shows which colour we have chosen to represent each proposition. We have drawn explicitly a border to indicate that all δ -clauses belong to the same δ -clausal form, so they are interpreted as a disjunction and all are possible explanations for the same abductive problem.



Look that there is a natural order in the derivation of new δ -clauses. Number 6 and 7 can only be generated after 5. This corresponds to the intuitive idea that “rained last night” (number 7) or “sprinkler was on” (6) are deeper explanations for the wet shoes than just that “the grass is wet” (5), which of course is also a possible explanation, but not so elaborated. It is easy to visualize the idea of greater reasoning chains in the diagram.

An accurate representation of the abductive process in section 2.3 would require two δ -resolution processes (for the theory and the observation) that are originally separated (steps 1 and 2 of the abductive process) and then they join (steps 3 and 4). One can imagine those breeding cages where birds are originally separated at the beginning of the courtship and only later they are put together. Both stages of the process are necessary for the success. Moreover, additional rules to eliminate contradictory or subsumed δ -clauses can be represented within these diagrams:



The left picture eliminates δ -clauses that contain some literal and its complementary. In the same way, the right rule eliminates a δ -clause whenever there is another one which is a proper subset. That is, the big one becomes subsumed, as there is another simpler explanation.

With these rules, in the previous example, δ -clauses 3, 2 and 1 are eliminated, respectively, when 5, 6 and 7 are generated. The old δ -clauses become subsumed by the new ones. This represents the intuitive idea that some conflicting explanations are discarded, before solving an abductive problem, when a better hypothesis is formulated.

As we have shown, all the resources in the δ -resolution process can be represented by these diagrams. So we can use them as a sound and complete abductive calculus, which allows us to visualize many of the intuitive ideas about the formulation of hypotheses.

4. Conclusions

To conclude, let us evaluate the contributions of the δ -resolution calculus and the proposed diagrams. The four Kapitan's theses draw the objectives that any systematisation of abductive reasoning should attain. These are high objectives which operate as the horizon of abduction. So, we finish this paper revisiting them and evaluating to what extent δ -resolution satisfies them.

First, the *inferential thesis* is completely fulfilled, as δ -resolution is an inferential process, formally characterized, with important logical properties.

The *thesis of purpose* highlights the double task of abduction: the *generation* of new hypotheses and the *selection* of the best ones for further examination. Many

approaches split these processes so first generate a number of hypotheses which satisfy (3) and then select, among them, those that satisfy (4) and (5). Usually, each process is performed in a different way. However, δ -resolution integrates *generation* and *selection*, as the abductive process defined above shows.

The *comprehension thesis* is the most interesting one and, at the same time, the most difficult to satisfy. The δ -resolution calculus is far from including all the operations whereby theories are engendered. But the steps of the abductive process are related to some ideas from Philosophy of Science and Epistemology, as we have commented when showing the diagrammatic approach to abduction at section 3. Indeed, the δ -resolution rule, in its graphical representation, can be interpreted within some cognitive theories, as *Mental Models* (Johnson-Laird, 1983). Johnson-Laird's mental models can be assimilated to sets of literals. Then, if $\Sigma_1 \cup \{\lambda\}$ and $\Sigma_2 \cup \{\neg\lambda\}$ are two mental models which explain a given observation, $\Sigma_1 \cup \Sigma_2$ is another mental model, maybe a smaller and better one, which also explains the observation. Some representations of mental models agree with our diagrams. Also, we underlined in the introduction the relevance of strategic rules for abductive reasoning. That relevance is shown explicitly in the diagrammatic representation, as an intelligent ordering of the applications of the δ -resolution rule can lead to shorter paths to the best explanation.

Finally, the *autonomy thesis* requires that abduction is irreducible to deduction and induction. Abduction is frequently called retroduction or backward deduction. A duality between deduction and abduction is often suggested. In this sense δ -resolution, which is dual to a typical deductive calculus, is somehow dual to deduction. But δ -resolution is also irreducible to deduction. Classical deduction is a monotonic reasoning, while the abductive process by δ -resolution, as it fulfils definition 2, is a non-monotonic³ inference.

³ Details on the non-monotonicity of abduction can be found in (Aliseda, 2006).

References

- Atocha Aliseda, *Abductive Reasoning: Logical Investigations into Discovery and Explanation*. Dordrecht: Springer, 2006.
- Marta Cialdea Mayer and FIORA Pirri, "First order abduction via tableau and sequent calculi". In: *Bulletin of the IGPL*, n. 1, 99-117, 1993.
- Elman Eder, "Consolation and its relation with resolution". In John Mylopoulos and Raymond Reiter (Eds.), *Proceedings of the 12th International Joint Conference on Artificial Intelligence (IJCAI-91)*, Sydney, Morgan Kaufmann, 1991, 132-136.
- Jaakko Hintikka, "What is abduction? The fundamental problem of contemporary epistemology". In: *Transactions of the Charles S. Peirce Society*, n. 34 (3), 503-533, 1998.
- Philip N. Johnson-Laird, *Mental Models: Toward a Cognitive Science of Language, Inference and Consciousness*, Cambridge: Harvard University Press, 1983.
- Antoni Kakas, Robert Kowalski and Francesca Toni, "The role of abduction in logic programming". In: Dov M. Gabbay et al. (Eds.), *Handbook of logic in Artificial Intelligence and Logic Programming*, Oxford: Oxford University Press, 1998, 235-324.
- Antoni Ligeza, "A note on Backward Dual Resolution and its application to proving completeness of rule-based systems". In Ruzena Bajcsy (Ed.), *Proceedings of the 13th International Joint Conference on Artificial Intelligence (IJCAI-93)*, Chambéry, Morgan Kaufmann, 1993, 132-137.
- Peter Lipton, *Inference to the Best Explanation*. New York: Routledge, 1991.
- Sami Paavola, "Abduction as a logic and methodology of discovery: the importance of strategies". In: *Foundations of Science*, n. 9 (3), 267-283, 2004.
- John Alan Robinson, "A Machine-Oriented Logic Based on the Resolution Principle". In: *Journal of the ACM*, n. 12, 23-41, 1965.
- Aránzazu San Ginés, "Visual Tools for Teaching Propositional Logic". In Patrick Blackburn et al. (Eds.), *Tools for Teaching Logic. Third International Congress, TICTTL 2011*, Dordrecht: Springer, 2011.
- Fernando Soler-Toscano, Ángel Nepomuceno-Fernández and Atocha Aliseda-Llera, "Model-Based Abduction via Dual Resolution". In: *Logic Journal of the IGPL*, nº 14 (2), 305-319, 2006.
- Fernando Soler-Toscano, Ángel Nepomuceno-Fernández and Atocha Aliseda-Llera, "Abduction via C-tableaux and δ -resolution". In: *Journal of Applied Non-Classical Logics*, n. 19(2), 211-225, 2009.