

1 ***Bacillus locisalis* sp. nov., a new haloalkaliphilic species from hypersaline and alkaline**
2 **lakes of China, Kenya and Tanzania**

3

4 **M. Carmen Márquez¹, Inmaculada J. Carrasco¹, Rafael R. de la Haba¹, Brian E.**
5 **Jones², William D. Grant³ and Antonio Ventosa¹**

6

7 ¹Department of Microbiology and Parasitology, Faculty of Pharmacy, University of Sevilla,
8 41012 Sevilla, Spain

9 ²Genencor International, Archimedesweg 30, 2333 CN Leiden, The Netherlands

10 ³Department of Infection, Immunity and Inflammation, University of Leicester, Leicester
11 LE1 9HN, UK

12

13 **Running title:** *Bacillus locisalis* sp. nov.

14

15 **Author for correspondence:** M.C. Márquez, Department of Microbiology and
16 Parasitology, Faculty of Pharmacy, University of Sevilla, 41012 Sevilla, Spain. Tel. +34 95
17 455 38 09; Fax +34 95 462 81 62. E-mail: cmarquez@us.es

18

19 The GenBank/EMBL/DDBJ accession numbers for the 16S rRNA gene sequences of strain
20 CG1^T, CG2, CG4, CG6, CG7, 103NT4 and WE1 are FR714930, FR714931, FR714932,
21 FR714933, FR714934, X92163 and X92164, respectively.

22 **Abstract**

23 A polyphasic taxonomic study was performed on seven *Bacillus*-like bacteria isolated from
24 three hypersaline and alkaline lakes located in China, Kenya and Tanzania. All strains were
25 moderately halophilic and alkaliphilic, Gram positive, motile rods. The DNA G+C content
26 from the seven isolates ranged from 42.2 to 43.4 mol% and their major fatty acid was
27 anteiso-C_{15:0}. Strain CG1^T, selected as representative strain of the isolates, possesses *meso*-
28 diaminopimelic acid in the cell wall peptidoglycan, MK-7 as the predominant menaquinone
29 and diphosphatidyl glycerol, phosphatidylglycerol and phosphatidylethanolamine as the
30 major polar lipids. Comparative 16S rRNA gene sequence analysis indicated that the
31 isolates belonged to the genus *Bacillus*. The seven isolates shared 97.7-99.9% 16S rRNA
32 gene sequence similarity, and formed a branch that was distinct from the type strains of the
33 recognized species of the genus *Bacillus*. They were most closely related to *Bacillus*
34 *agaradhaerens* DSM 8721^T (92.6-93.8% 16S rRNA sequence similarity). DNA-DNA
35 hybridization values between the seven isolates were 85-100%. According to the
36 polyphasic characterization, the strains represent a novel species, for which the name
37 *Bacillus locisalis* sp. nov. is proposed. The type strain is CG1^T (CCM 7370^T = CECT 7152^T
38 = CGMCC 1.6286^T = DSM 18085^T).

39

40 **Key words:** *Bacillus locisalis* sp. nov., New species, 16S rRNA gene analysis, Taxonomy,
41 Polyphasic study, Hypersaline lakes, Soda lakes

42 **Scope of the paper:** Systematics

43 Haloalkaliphilic bacteria are extremophilic microorganisms that are widely distributed in
44 different hypersaline and alkaline habitats with a variable (up to saturation) salt
45 concentration and high pH values. The genus *Bacillus* was proposed by Cohn in 1872 [6]
46 and since then it has undergone substantial taxonomic changes. Currently, this genus
47 groups near 200 species [10] with some of them having a moderately halophilic and
48 alkaliphilic/alkalitolerant response, such is the case of *B. oshimensis* (from soil in Japan)
49 [29], *B. saliphilus* (from algal mat from a mineral pool in Italy) [24], *B. chagannorensis*
50 (from a soda lake in China) [3], *B. aurantiacus* (from an extremely shallow soda lake in
51 Hungary) [2], and *Bacillus polygoni* (from indigo balls in Japan) [1].

52 In the present study, we report the discovery of a novel moderately halophilic, alkaliphilic
53 *Bacillus* species during a study of bacterial diversity in hypersaline habitats using a culture-
54 dependent approach. Seven bacterial strains were isolated from water and sediment samples
55 from hypersaline and alkaline lakes located in three different countries: China, Kenya and
56 Tanzania. The taxonomic status of the isolates was determined using a polyphasic
57 approach.

58 Strains 103NT4 and WE1 were isolated in 1988 following the methodology described by
59 Duckworth et al. [8]. Strain 103NT4 was isolated from orange-coloured soda crusts
60 surrounding a warm soda seep brine (35°C) located on the northern shore of Lake Natron
61 (Tanzania) (2°08' S, 36°00' E, pH 10.5, conductivity 35 mS cm⁻¹), while strain WE1 was
62 isolated from a sediment sample from the eastern shore of Lake Elmenteita, in the Kenyan
63 section of the East African Rift Valley (0°25' S, 36°15' E, pH 10.5, conductivity 12.7 mS
64 cm⁻¹) [8]. The other five strains were isolated from water (CG1^T and CG2) and sediment
65 (CG4, CG6 and CG7) samples taken from Lake Chagannor, during an expedition in
66 September 2003. This lake is situated near a soda works, 120 km south of Mandulatu

67 (43°16' N 112°55' E, pH 10.5, conductivity 202 mS cm⁻¹), on the Inner Mongolian steppe,
68 northwest of Beijing, China. The water samples were diluted in sterile 10% (w/v) marine
69 salts (g l⁻¹): NaCl, 78; MgCl₂ x 6H₂O, 13; MgSO₄ x 7H₂O, 20.3; CaCl₂, 0.33; KCl, 2;
70 NaHCO₃, 0.07; NaBr, 0.23 [28], plating on alkaline saline medium and incubating at 37°C
71 aerobically. The alkaline saline isolation medium contained (g l⁻¹): glucose, 10.0; peptone
72 (Difco), 5.0; yeast extract (Difco), 5.0; KH₂PO₄, 2.0; MgSO₄ x 7H₂O, 0.4; NaCl, 80;
73 Na₂CO₃, 20. The salts NaCl and Na₂CO₃ were autoclaved separately and added to the
74 organic components at 60°C. The pH of this medium was adjusted to pH 10. When it was
75 necessary, the medium was solidified by adding 2.0% (w/v) agar. The sediments (0.1g)
76 were suspended in 10% (w/v) marine salts. The suspensions were vortexed for 1 min,
77 allowed to settle, serially diluted in 10% (w/v) marine salts and then spread-plated in
78 duplicate on alkaline saline medium followed by aerobic incubation at 37°C. The strains
79 were subsequently purified three times by plating on the same medium and maintained on
80 the same alkaline saline medium and at -80°C on this medium without agar and
81 supplemented with 30% (v/v) glycerol. In addition to the seven isolates, *Bacillus*
82 *agaradhaerens* DSM 8721^T was obtained from the Deutsche Sammlung von
83 Mikroorganismen und Zellkulturen (DSMZ), Braunschweig, Germany, and cultivated at 37
84 °C on alkaline saline medium. This bacterium was used as reference for comparative
85 phenotypic and chemotaxonomic studies.

86 The phylogenetic position of the seven isolates was determined by complete 16S rRNA
87 gene sequence analysis. Genomic DNAs were prepared using the method described by
88 Marmur [19]. PCR amplifications of the 16S rRNA gene were carried out with the forward
89 primer 16F27 and the reverse primer 16R1488. Sequencing was performed using an
90 automated DNA sequencer model 3130XL (Applied Biosystems). Identification of

91 phylogenetic neighbours and calculation of pairwise 16S rRNA gene sequence similarities
92 were achieved using the EzTaxon server version 2 (<http://www.eztaxon.org/>; [5]). The 16S
93 rRNA gene sequences were aligned with the published sequences of closely related
94 bacteria. The alignment was confirmed and checked against both primary and secondary
95 structures of the 16S rRNA molecule using the alignment tool of the ARB software
96 package [18]. The phylogenetic trees were constructed using three different methods:
97 maximum-likelihood [11], maximum-parsimony [13] and neighbour-joining [25],
98 algorithms integrated in the ARB software for phylogenetic inference. The robustness of
99 the topology in the phylogenetic trees was evaluated by bootstrap analyses [12] of the
100 neighbour-joining method based on 1000 resamplings. The 16S rRNA gene sequences used
101 for phylogenetic comparisons were obtained from the GenBank database and their strain
102 designations and accession numbers are shown in Fig. 1.

103 Almost-complete 16S rRNA gene sequences of the seven isolates (1441 nucleotides) were
104 obtained and used for initial BLAST and EzTaxon searches in GenBank and phylogenetic
105 analysis. Comparative 16S rRNA gene sequence analysis revealed that the seven isolates
106 have the closest phylogenetic affiliation with the genus *Bacillus*. A tree constructed by
107 neighbour-joining analysis clearly showed that the seven isolates grouped together with
108 97.7-99.9% 16S rRNA gene sequence similarity among themselves. This cluster was
109 separated from one formed by some other *Bacillus* species with 100 bootstrap support (Fig.
110 1). The topologies of phylogenetic trees built using the maximum-likelihood and
111 maximum-parsimony algorithms were similar to those of the tree constructed by neighbour-
112 joining analysis (data not shown). The nearest known relative of the isolates was *Bacillus*
113 *agaradhaerens* DSM 8721^T, with values of 16S rRNA gene sequence similarity comprised
114 between 92.6 and 93.8%. For determination of the DNA base composition of the seven

115 isolates the DNAs were extracted and purified by the method of Marmur [19] and the G+C
116 contents of the DNAs were determined in triplicate from the midpoint value of the thermal
117 denaturation profile [20] by using the equation of Owen and Hill [24]. The genomic DNA
118 G+C contents of the seven isolates ranged from 42.2 to 43.4 mol% (Table S1). These
119 values are within the range for *Bacillus* but are higher than that of *Bacillus agaradhaerens*
120 DSM 8721^T (39.5 mol%). DNA-DNA hybridization was carried out to evaluate the
121 genomic DNA relatedness between the seven isolates, following the competition procedure
122 of Johnson [16], described in detail elsewhere [21]. The hybridization temperature was 46.4
123 °C, which was within the limit of validity for the filter method [7] and the percentage of
124 hybridization was calculated according to Johnson [16]. The values presented were based
125 on a minimum of four replicates. The values of DNA-DNA hybridization between strain
126 CG1^T and the other six isolates ranged between 85% and 99%. These values are clearly
127 higher than 70%, cut-off generally accepted for species delineation and support the
128 placement of the seven isolates as the same genotypic species [27].

129 In order to phenotypically characterize the isolates and, following the minimal standards for
130 describing new genera and species of aerobic, endospore-forming bacteria recommended by
131 Logan et al. [17], standard phenotypic tests were performed. The Gram stain reaction was
132 carried out using the method described by Dussault [9]. Cell morphology and motility were
133 studied by phase-contrast microscopy. The morphology of colonies, their size and
134 pigmentation were observed on the alkaline saline solid medium with different salt
135 concentrations after 2 days of incubation. Growth at different concentrations of salts was
136 determined on the alkaline saline medium containing 0, 0.5, 1, 3, 5, 7, 10, 15, 20, 25 or
137 30% (w/v) NaCl. The pH range for growth was determined on the alkaline saline liquid
138 medium at pH values ranging from 7.0 to 12.5, with increments of 0.5 pH units, using the

139 appropriate biological buffers, Na₂HPO₄/NaH₂PO₄ (below pH 8.0), Na₂CO₃/NaHCO₃ (pH
140 8.0-10.0) and Na₂HPO₄/NaOH (pH 11), as described previously [15]. The pH was
141 readjusted after sterilization; growth was scored as optical density at 600 nm. The
142 temperature range for growth was determined at temperatures between 6 and 50°C. Catalase
143 was tested by adding 3% H₂O₂ to culture plates. The oxidase reaction was performed on
144 filter paper moistened with 1% (w/v) aqueous solution of *N, N, N', N'*-tetramethyl-*p*-
145 phenyldiamine. Sporulation was tested on the alkaline saline solid medium supplemented
146 with 5 mg l⁻¹ MnSO₄ (Merck). Utilization of various substrates as sole carbon and energy
147 sources, or carbon, nitrogen and energy sources, were determined using a basal medium
148 with the following composition (g l⁻¹): yeast extract (Difco), 0.01; KNO₃, 1.0; KH₂PO₄,
149 1.0; MgSO₄ x 7H₂O, 0.2; (NH₄)₂HPO₄, 1.0; NaCl, 80; Na₂CO₃, 20. To this liquid medium
150 a 0.1% (w/v) filter-sterilized substrate was added. Carbohydrates were used at a final
151 concentration of 0.2% (w/v). When amino acids were used as substrate the basal medium
152 contained neither KNO₃ nor (NH₄)₂HPO₄. A growth test was considered positive when the
153 OD₆₀₀ reached or exceeded a value of 0.3 after 4 days at 37 °C. Other tests shown in Table
154 S1 or included in the species description were carried out following methodologies
155 described previously [14, 23, 28]. Unless otherwise indicated the tests were carried out in
156 the alkaline saline medium (pH 10) and incubated at 37°C in sealed containers to minimise
157 evaporation. The seven isolates studied in this work were very similar in their phenotypic
158 characteristics, although some differences were observed between them (Table S1 and
159 species description).

160 Fatty acids were determined for the seven isolates, as well as for the reference strain
161 *Bacillus agaradhaerens* DSM 8721^T using the MIDI system (Microbial Identification
162 System). All the strains were grown on alkaline saline medium, pH 10 at 37°C, for 48 h.

163 This analysis was carried out by the Identification and Characterization Service of the
164 CECT (Valencia, Spain). Anteiso-C_{15:0} was the predominant compound although slight
165 variation was observed between the compositions of the seven isolates (Table 1).

166 Analysis of peptidoglycan of the cell wall, quinones and polar lipids content of strain
167 CG1^T, selected as representative strain of the isolates, was carried out by the Identification
168 Service of the DSMZ (Braunschweig, Germany). The cell biomass for these analyses was
169 obtained by cultivation on the alkaline saline medium (pH 10) at 37°C, for 48 h. Strain
170 CG1^T possessed a cell wall peptidoglycan of type A1 γ (*meso*-Dpm, directly cross-linked;
171 A31; http://www.dsmz.de/microorganisms/main.php?content_id=35) and contained MK-7
172 (98%) as the predominant menaquinone, with MK-6 (2%) present in minor amounts. The
173 polar lipids of this strain consisted of diphosphatidylglycerol, phosphatidylglycerol,
174 phosphatidylethanolamine, four phospholipids and an aminophospholipid of unknown
175 structure (Fig. S1). The results obtained from these chemotaxonomic analyses were
176 consistent with the results from the phylogenetic analysis that suggest that our isolates may
177 belong to the genus *Bacillus* [4].

178 The characteristics that differentiate strain CG1^T from *Bacillus agaradhaerens* DSM8721^T
179 are summarized in Table 2. The differences in some features, such as colony pigmentation,
180 growth in anaerobic conditions, range and optimal salt concentration for growth, optimal
181 temperature for growth, hydrolysis of starch, Voges-Proskauer test, as well as the genomic
182 DNA G+C content, can be used to distinguish this strain from *Bacillus agaradhaerens*
183 (Table 2). Therefore, the taxonomic data from polyphasic analysis clearly suggest that our
184 isolates belong to the genus *Bacillus* and represent a new species of this genus, for which
185 the new name *B. locisalis* sp. nov. is proposed.

186 *Description of Bacillus locisalis* sp. nov.

187 *Bacillus locisalis* (lo.ci.sa'lis. L. n. *locus* place, locality; L. gen. n. *salis* of salt; N.L. gen. n.
188 *locisalis* from a salted place).

189 Gram-positive rods, 1.0 by 2.0–5.0 μm . Motile, oval endospores are produced at terminal
190 and subterminal positions in swollen sporangia. Facultatively anaerobic. Colonies are
191 orange, circular, opaque and entire on alkaline saline medium after 2 days of cultivation.
192 Moderately halophilic, growing in a wide range (1 to 25% w/v) of salt concentrations, with
193 optimal growth at 7-10% (w/v) NaCl. Grows at 10-45°C (optimal at 37°C) and pH 8-12
194 (optimal at pH 9-10). Oxidase negative and catalase positive. Acid is produced from D-
195 fructose, D-glucose, D-maltose, D-mannitol, D-melibiose, D-ribose, D-trehalose and D-
196 xylose. Acid is not produced from D-amygdaline, D-arabinose, L-citruline, dulcitol, DL-
197 ethionine, glycerol, inulin, lactose, D-melezitose, *m*-inositol, and xylitol. Casein is not
198 hydrolyzed. Indole production and Voges-Proskauer test are negative. D-fucose, D-fructose
199 and D-glucose are utilized as sole carbon and energy sources. The following compounds
200 are not utilized as sole carbon and energy sources: aesculin, butanol, *m*-inositol, sorbitol,
201 xylitol and citrate. L-alanine and cysteine are utilized as sole carbon, nitrogen, and energy
202 sources. L-phenylalanine and L-glutamine are not utilized as sole carbon, nitrogen, and
203 energy sources. DNA base composition ranges from 42.2 to 43.4 mol%. The cell wall
204 contains peptidoglycan of the *meso*-diaminopimelic acid type. Major isoprenoid quinone is
205 MK-7. The polar lipids are diphosphatidylglycerol, phosphatidylglycerol,
206 phosphatidylethanolamine, four phospholipids and an aminophospholipid of unknown
207 structure. Additional characteristics of the strains are listed in Table S1. Cellular fatty acid
208 composition is given in Table 1.

209 The habitats are saline and alkaline waters and soils.

210 The type strain is CGI^T (CCM 7370^T = CECT 7152^T = CGMCC 1.6286^T = DSM 18085^T),
211 isolated from Lake Chagannor, in Inner Mongolia, China.

212 *Description of the type strain.*

213 The description of the type strain is the same as that of the species. The base composition of
214 its DNA is 42.2 mol% G+C, as determined by the thermal denaturation method. Other
215 characteristics are shown in Supplementary Table 1.

216 **Acknowledgements**

217 This study was supported by grants from the Quality of Life and Management of Living
218 Resources Programme of the European Commission (Project “Multigenome access
219 technology for industrial catalysts”, QLK3-CT-2002-01972), Spanish Ministerio de
220 Educación y Ciencia (BIO2009-10138 and CGL2010-19303) and Junta de Andalucía (PO6-
221 CVI-01829).

222 **References**

223 [1] Aino, K., Hirota, K., Matsuno, T., Morita, N., Nodasaka, Y., Fujiwara, T., Matsuyama,
224 H., Yoshimune, K., Yumoto, I. (2008) *Bacillus polygona* sp. nov., a moderately halophilic,
225 non-motile obligate alkaliphile isolated from indigo balls. *Int. J. Syst. Evol. Microbiol.* 58,
226 120-128.

227 [2] Borsodi, A.K., Márialigeti, K., Szabó, G., Palatinszky, M., Pollák, B., Kéki, Z., Kovács,
228 A.L., Schumann, P., Tóth, E.M. (2008) *Bacillus aurantiacus* sp. nov., an alkaliphilic and
229 moderately halophilic bacterium isolated from Hungarian soda lakes. *Int. J. Syst. Evol.*
230 *Microbiol.* 58, 845-851.

- 231 [3] Carrasco, I.J., Márquez, M.C., Xue, Y., Ma, Y., Cowan, D.A., Jones, B.E., Grant, W.D.,
232 Ventosa, A. (2007) *Bacillus chagannorensis* sp. nov., a moderate halophile from a soda
233 lake in Inner Mongolia, China. *Int. J. Syst. Evol. Microbiol.* 57, 2084-2088.
- 234 [4] Chen, Y.-G., Zhang, Y.-Q., Wang, Y.-X., Liu, Z.-X., Klenk, H.-P., Xiao, H.-D., Tang,
235 S.-K., Cui, X.-L., Li, W.-J. (2009) *Bacillus neizhouensis* sp. nov., a halophilic marine
236 bacterium isolated from a sea anemone. *Int. J. Syst. Evol. Microbiol.* 57, 2084-2088.
- 237 [5] Chun, J., Lee, J.-H., Jung, Y., Kim, M., Kim, S., Kim, B.K., Lim, Y.-W. (2007)
238 EzTaxon: a web-based tool for the identification of prokaryotes based on 16S ribosomal
239 RNA gene sequences. *Int. J. Syst. Evol. Microbiol.* 57, 2259-2261.
- 240 [6] Cohn, F. (1872). *Untersuchungen über Bakterien. Beitr. Biol. Pflanz.* 1, 127-244.
- 241 [7] De Ley, J., Tijtgat, R. (1970) Evaluation of membrane filter methods for DNA-DNA
242 hybridization. *Antonie van Leeuwenhoek* 36, 461-474.
- 243 [8] Duckworth, A.W., Grant, W.D., Jones, B.E., van Steenberg, R. (1996) Phylogenetic
244 diversity of soda lake alkaliphiles. *FEMS Microbiol. Ecol.* 19, 181-191.
- 245 [9] Dussault, H.P. (1955) An improved technique for staining red halophilic bacteria. *J.*
246 *Bacteriol.* 70, 484.
- 247 [10] Euzéby, J.P. (2010) List of Prokaryotic names with Standing in Nomenclature.
248 <http://www.bacterio.cict.fr>.
- 249 [11] Felsenstein, J. (1981) Evolutionary trees from DNA sequences: a maximum likelihood
250 approach. *J. Mol. Evol.* 17, 368–376.

- 251 [12] Felsenstein, J. (1985) Confidence limits on phylogenies: an approach using the
252 bootstrap. *Evolution* 39, 783–791.
- 253 [13] Fitch, W.M. (1971) Toward defining the course of evolution: minimum change for a
254 specific tree topology. *Syst. Zool.* 20, 406–416.
- 255 [14] García, M.T., Ventosa, A., Ruiz-Berraquero, F., M. Kocur. (1987) Taxonomic study and
256 amended description of *Vibrio costicola*. *Int. J. Syst. Bacteriol.* 37, 251-256.
- 257 [15] Gomori, G. (1955) Preparation of buffers for use in enzyme studies. *Methods*
258 *Enzymol.* 1, 138-146.
- 259 [16] Johnson, J. L. (1994) Similarity analysis of DNAs. In: Gerhardt, P., Murray, R.G.E.,
260 Wood, W.A., Krieg, N.R. (Eds.), *Methods for General and Molecular Bacteriology*,
261 American Society for Microbiology, Washington, D.C., pp. 655-681.
- 262 [17] Logan, N.A., Berge, O., Bishop, A.H., Busse, H.-J., De Vos, P., Fritze, D.,
263 Heyndrickx, M., Kämpfer, P., Rabinovitch, L., other authors. (2009) Proposed minimal
264 standards for describing new taxa of aerobic, endospore-forming bacteria. *Int. J. Syst. Evol.*
265 *Microbiol.* 59, 2114-2121.
- 266 [18] Ludwig, W., Strunk, O., Westram, R., Richter, L., Meier, H., Yadhukumar, Buchner,
267 A., Lai, T., Steppi, S., other authors. (2004) ARB: a software environment for sequence
268 data. *Nucleic. Acids Res.* 32, 1363–1371.
- 269 [19] Marmur, J. (1961) A procedure for the isolation of deoxyribonucleic acid from micro-
270 organisms. *J. Mol. Biol.* 3, 208-218.

271 [20] Marmur, J., Doty, P. (1962) Determination of the base composition of deoxyribonucleic
272 acid from its thermal denaturation temperature. *J. Mol. Biol.* 5, 109-118.

273 [21] Márquez, M.C., Carrasco, I.J., Xue, Y., Ma, Y., Cowan, D.A., Jones, B.E., Grant,
274 W.D., Ventosa, A. (2007) *Aquisalimonas asiatica* gen. nov., sp. nov., a moderately
275 halophilic bacterium isolated from an alkaline, saline lake in Inner Mongolia, China. *Int. J.*
276 *Syst. Evol. Microbiol.* 57, 1137-1142.

277 [22] Owen, R.J., Hill, L.R. (1979) The estimation of base compositions, base pairing and
278 genome size of bacterial deoxyribonucleic acids. In: Skinner, F. A., Lovelock, D. W. (Eds.),
279 *Identification Methods for Microbiologists*, 2nd edn, Academic Press, London, pp. 217-296.

280 [23] Quesada, E., Ventosa, A., Ruiz-Berraquero, F., Ramos-Cormenzana, A. (1984) *Deleya*
281 *halophila*, a new species of moderately halophilic bacteria. *Int. J. Syst. Bacteriol.* 40, 287-
282 292.

283 [24] Romano, I., Lama, L., Nicolaus, B., Gambacorta, A., Giordano, A. (2005) *Bacillus*
284 *saliphilus* sp. nov., isolated from a mineral pool in Campania, Italy *Int. J. Syst. Evol.*
285 *Microbiol.* 55, 159-163.

286 [25] Saitou, N., Nei, M. (1987) The neighbour-joining method: a new method for
287 reconstructing phylogenetic trees. *Mol. Biol. Evol.* 4, 406-425.

288 [26] Subow, N.N. (1931) *Oceanographical tables*. Commissariat of agriculture of USSR.
289 Hydro-Meteorological Committee of USSR. Oceanographical Institute of USSR, Moscow.

290 [27] Stackebrandt, E., Ebers, J. (2006) Taxonomic parameters revisited: tarnished gold
291 standards. *Microbiol. Today.* 33, 152-155.

- 292 [28] Ventosa, A., Quesada, E., Rodríguez-Valera, F., Ruiz-Berraquero, F., Ramos-
293 Cormenzana, A. (1982) Numerical taxonomy of moderately halophilic Gram-negative rods.
294 J. Gen. Microbiol. 128, 1959-1968.
- 295 [29] Yumoto, I., Hirota, K., Goto, T., Nodasaka, Y., Nakajima, K. (2005) *Bacillus*
296 *oshimensis* sp. nov., a moderately halophilic, non-motile alkaliphile. Int. J. Syst. Evol.
297 Microbiol. 55, 907-911.

298 **Legend to Figure**

299 **Fig. 1.** Neighbour-joining tree, based on the 16S rRNA gene sequence comparison,
300 showing the relationship of strain CG1^T with related species. The accession numbers of the
301 sequences used in this study are shown in parentheses after the strain designation.
302 *Paenibacillus polymixa* NCDO 1774^T was used as an outgroup. Bootstrap values >70% are
303 shown. The scale bar represents 0.01 substitutions per nucleotide position.

304 **Table 1.** Cellular fatty acid composition of the seven isolate strains and *B. agaradhaerens*
 305 DSM 8712^T grown on alkaline saline medium (pH 10) at 37°C, for 48 h. Data are
 306 percentages of the total fatty acids. -, Values less than 0.5% in all strains; ND, Not detected

Fatty acids	Strains							
	CG1 ^T	CG2	CG4	CG6	CG7	WE1	103NT4	DSM 8712 ^T
Straight chain								
C _{12:0}	-	-	-	-	-	0.7	0.8	0.5
C _{14:0}	1.1	1.0	0.7	0.7	0.7	1.4	1.3	0.7
C _{16:0}	4.3	3.9	5.3	3.9	5.6	2.1	2.4	6.0
C _{18:0}	1.0	-	0.7	0.5	0.8	0.5	ND	-
Branched								
iso- C _{14:0}	3.6	3.1	4.5	3.9	4.6	9.0	5.3	1.0
iso-C _{15:0}	11.4	11.4	11.5	11.1	10.3	15.9	11.5	23.3
anteiso C _{15:0}	54.1	55.9	42.0	44.2	39.4	46.2	61.5	40.9
iso-C _{16:0}	3.8	3.4	6.1	5.3	6.7	4.9	3.7	3.3
iso-C _{17:0}	3.9	3.6	6.7	5.6	7.2	0.9	1.5	7.0
anteiso C _{17:0}	11.5	11.6	13.9	13.9	15.0	4.4	6.2	11.9
iso-C _{18:0}	ND	ND	-	-	0.7	ND	ND	ND
Unsaturated								
C _{16:1} ω7 <i>c</i> alcohol	ND	ND	ND	ND	ND	2.0	ND	ND
C _{16:1} ω11 <i>c</i>	1.3	1.5	2.1	3.0	2.8	4.3	1.7	1.2
C _{17:1} iso ω10 <i>c</i>	1.7	2.0	3.7	4.9	4.0	3.8	1.7	1.5

$C_{18:1}$	$\omega 9c$	1.2	1.0	1.1	1.0	1.0	2.0	1.6	1.4
------------	-------------	-----	-----	-----	-----	-----	-----	-----	-----

307 **Table 2.** Characteristics used to distinguish *Bacillus locisalis* strain CG1^T from *B.*
 308 *agaradhaerens* DSM 8721^T (data from this study). +, positive; -, negative.

Characteristic	<i>Bacillus locisalis</i> strain CG1 ^T	<i>Bacillus agaradhaerens</i> DSM 8721 ^T
Sampling site	Water	Soil
Colony pigmentation	Orange	White
Anaerobic growth	+	-
NaCl range (% w/v)	1-20	0-16
Optimum NaCl (% w/v)	10	0.5
Optimum temperature (°C)	37	30
Hydrolysis of starch	-	+
Voges-Proskauer test	-	+
Growth on ^a :		
Aesculin	-	+
D-Fucose	+	-
D-Melezitose	-	+
D-Raffinose	-	+
D-Ribose	-	+
Salicin	-	+
Sucrose	-	+
D-Trehalose	-	+
Butanol	-	+
Ethanol	+	-
Glycerol	-	+
Propanol	-	+
D-Sorbitol	-	+
Xylitol	-	+
Acetate	-	+
Citrate	-	+
Growth on ^b :		
L-Alanine	+	-
L-Arginine	+	-
L-Aspartate	+	-
L-Cysteine	+	-

L-Glutamine	-	+
L-Methionine	+	-
DNA G+C content (mol%)	42.2	39.2
Major fatty acids	Anteiso-C _{15:0} (53%) Anteiso-C _{17:0} (13%) Iso- C _{15:0} (13%)	Anteiso-C _{15:0} (41%) Iso- C _{15:0} (25%) Anteiso-C _{17:0} (12%)

309 ^a When supplied as the sole source of carbon and energy.

310 ^b When supplied as the sole source of carbon, nitrogen, and energy.