

International Journal of Bifurcation and Chaos
© World Scientific Publishing Company

**(This paper is for the Special Issue edited by
Prof. Gregoire Nicolis , Prof. Marko Robnik, Dr. Vassilis Rothos
and Dr. Haris Skokos)**
**Multibreather and vortex breather stability in Klein–Gordon
lattices: Equivalence between two different approaches**

J. Cuevas

*Nonlinear Physics Group of the University of Sevilla, Departamento de Física Aplicada I,
Escuela Politécnica Superior, C/ Virgen de África 7, 41011 Sevilla, Spain
jcuevas@us.es*

V. Koukouloyannis

*Department of Civil Engineering,
Technological Educational Institute of Serres, 62124 Serres, Greece
and
Department of Physics, Section of Astrophysics, Astronomy and Mechanics,
Aristotle University of Thessaloniki, 54124 Thessaloniki, Greece
vkouk@physics.auth.gr*

P.G. Kevrekidis

*Department of Mathematics and Statistics,
University of Massachusetts, Amherst MA 01003-4515
kevrekid@gmail.com*

J.F.R. Archilla

*Nonlinear Physics Group of the University of Sevilla, Departamento de Física Aplicada I,
ETSI Informática, Avda. Reina Mercedes s/n, 41012 Sevilla, Spain
archilla@us.es*

Received (to be inserted by publisher)

In this work, we revisit the question of stability of multibreather configurations, i.e., discrete breathers with multiple excited sites at the anti-continuum limit of uncoupled oscillators. We present two methods that yield quantitative predictions about the Floquet multipliers of the linear stability analysis around such exponentially localized in space, time-periodic orbits, based on the Aubry band method and the MacKay effective Hamiltonian method and prove that by making the suitable assumptions about the form of the bands in the Aubry band theory, their conclusions are equivalent. Subsequently, we showcase the usefulness of the methods through a series of case examples including one-dimensional multi-breathers, and two-dimensional vortex breathers in the case of a lattice of linearly coupled oscillators with the Morse potential and in that of the discrete ϕ^4 model.

1. Introduction

Over the past two decades, there has been an explosion of interest towards the study of Intrinsic Localized Modes (ILMs), otherwise termed discrete breathers [Flach & Willis, 1998; Flach & Gorbach, 2008]. This activity has been, to a considerable extent, fueled by the ever-expanding applicability of these exponentially localized in space and periodic in time modes. A partial list of the relevant applications includes their emergence in halide-bridged transition metal complexes as e.g. in [Swanson *et al.*, 1999], their potential role in the formation of denaturation bubbles in the DNA double strand dynamics summarized e.g. in [Peyrard, 2004], their observation in driven micromechanical cantilever arrays as shown in [Sato *et al.*, 2006], their investigation in coupled torsion pendula [Cuevas *et al.*, 2009], electrical transmission lines [English *et al.*, 2008, 2010], layered antiferromagnetic samples such as those of a $(\text{C}_2\text{H}_5\text{NH}_3)_2\text{CuCl}_4$ [English *et al.*, 2001b,a], as well in nonlinear optics [Lederer *et al.*, 2008] and possibly in atomic physics of Bose-Einstein condensates [Brazhnyi & Konotop, 2004; Morsch & Oberthaler, 2006] and most recently even in granular crystals [Boechler *et al.*, 2010].

In parallel to the above experimental developments in this diverse set of areas, there has been a considerable progress towards the theoretical understanding of the existence and stability properties of such localized modes summarized in a number of reviews and books; see e.g. [Flach & Willis, 1998; Flach & Gorbach, 2008; Lederer *et al.*, 2008; Aubry, 1997; Kevrekidis, 2009]. Arguably, one of the most important developments in establishing the fundamental relevance of this area in coupled nonlinear oscillator chains has been the work of MacKay and Aubry [MacKay & Aubry, 1994], which established the fact that if a single oscillator has a periodic orbit (and relevant non-resonance conditions are satisfied), then upon inclusion of a non-vanishing coupling between adjacent such oscillators, an ILM type waveform will generically persist.

Given the confirmation of persistence of such modes, naturally, the next question concerns their robustness under the dynamical evolution of the relevant systems, which is critical towards their experimental observability. This proved to be a substantially more difficult question to answer in a quantitative fashion, especially so for ILMs featuring multiple localized peaks, i.e., multi-site breathers (since single-site breathers are typically stable in chains of linearly coupled anharmonic oscillators). Two principal theories were proposed for addressing the stability of such periodic orbits, which correspond to discrete breather states (and for identifying their corresponding Floquet multipliers). Interestingly, these originated independently from the same pioneers which established (jointly) the existence of such modes in [MacKay & Aubry, 1994]. In particular, the first theory was pioneered by Aubry in his seminal work of [Aubry, 1997] and will go under the name Aubry Band (AB) theory, hereafter. The second one is an effective Hamiltonian method which was introduced in a series of papers by MacKay and collaborators [Ahn & Sepulchre, 2001; MacKay & Sepulchre, 2002; MacKay, 2004] (and will be termed accordingly MacKay Effective Hamiltonian method (MEH)). The AB approach was adapted to the stability of discrete breathers and multibreathers in the setting of Klein-Gordon (KG) lattices in the work of [Archilla *et al.*, 2003]; see also [Cuevas *et al.*, 2005, 2011]. The MEH approach was applied to the same setting in the recent work of [Koukouloyannis & Kevrekidis, 2009]; see also [Koukouloyannis *et al.*, 2010].

Our aim in the present work is to unify the two methods by firmly establishing the equivalence of the stability conclusions of the Aubry band and MacKay effective Hamiltonian methods. Subsequently, we illustrate the usefulness and versatility of the methods, we apply them to a range of physically interesting chains of oscillator model examples, such as the Morse potential which arises in the study of DNA bubbles [Peyrard, 2004], as well as the ϕ^4 potential which arises in applications in dusty plasmas [Koukouloyannis & Kourakis, 2007, 2009], as well as in field theory, particle physics and elsewhere; see e.g. the recent discussion of [Cubero *et al.*, 2009] and the earlier review [Belova & Kudryavtsev, 1997] and references therein. Our presentation is structured as follows. In section 2, we compare the two approaches and showcase the equivalence of their conclusions. In particular, in section 2.2 we show that, although the “original” AB method provides only qualitative results about the stability of multibreathers, if we make suitable assumptions about the form of the bands we can also get quantitative results; i.e., we can calculate the corresponding characteristic exponents. In section 3, we illustrate the use of the method by calculating the characteristic exponents of multibreathers in some classic lattice configurations. In particular we study a 1D and 2D Klein-Gordon lattice with Morse on-site potential (subsections 3.1 and 3.2, respectively) and a

ϕ^4 1D chain (subsection 3.3). Finally, in section 4, we summarize our findings and present our conclusions.

2. Comparison between the two approaches

2.1. Preliminaries - Terminology

The relevant system under consideration will be a Klein-Gordon chain of oscillators with nearest-neighbor interaction and Hamiltonian

$$H = H_0 + \epsilon H_1 = \sum_{i=-\infty}^{\infty} \left[\frac{1}{2} p_i^2 + V(x_i) \right] + \frac{\epsilon}{2} \sum_{i=-\infty}^{\infty} (x_i - x_{i-1})^2. \quad (1)$$

As indicated previously, we will examine the two approaches (AB and MEH) for the linear stability of multi-site breathers of this general class of systems. Both approaches are based on the notion of the anti-continuum limit. In this limit ($\epsilon = 0$) we consider n “central” oscillators moving in periodic orbits with the same frequency ω (this will be our “multibreather” for $\epsilon \neq 0$), while the rest lie at the equilibrium $(x, \dot{x}) = (0, 0)$. For $\epsilon \neq 0$ some of these configurations, depending on the phase differences between the oscillators, can be continued in order to provide multibreather solutions. It is interesting/relevant to note here that while the MEH approach provides explicit conditions about which configurations can be continued to finite ϵ (the critical points of the relevant effective Hamiltonian), the AB theory provides only stability information for a given configuration (for which we already know otherwise that it should exist at finite ϵ).

The linear stability of these solutions is determined by the corresponding Floquet multipliers. For a stable multibreather we require that all the multipliers lie on the unit circle. In the anti-continuum limit these multipliers lie in three bundles. The two conjugate ones, that correspond to the non-central oscillators, lie at $e^{\pm i\omega T}$, while the third one lies at $+1$ and consists of n multiplier pairs, corresponding to the central oscillators. Each pair of $+1$'s corresponds to the *phase mode* and *growth mode* of each isolated excited oscillator, meaning that a small change in the initial phase or a small change in frequency leads to an extremely close periodic solution, for the growth mode with slightly larger or smaller amplitude.

For $\epsilon \neq 0$, the non-central corresponding bundles split and their multipliers move along the unit circle to form the phonon band, while the multipliers at unity can move along, either the unit circle (stability), or along the real axis (instability). However, a pair of multipliers always continues at $+1$ corresponding to the phase mode and growth mode of the whole system. Hence, the stability of the multibreather, at least for small values of the coupling, is determined by the multipliers of the central oscillators. For larger values of ϵ , a Hamiltonian-Hopf bifurcation can occur and destabilize an initially stable multibreather.

At this point, it is relevant to make a note in passing about the striking similarities between the discussion above (at and near the anti-continuum limit) and that of the linear stability of standing waves in the discrete nonlinear Schrödinger (DNLS) equation. In that case, due to the monochromatic nature of the solutions and the $U(1)$ invariance of the latter model, it is possible to directly consider the eigenvalues associated with the standing wave solutions. However, there is a direct analogy with the spectrum of the excited sites being associated with the eigenvalues at the origin at the anti-continuum limit and the continuous spectrum lying at a finite distance from the spectral plane origin, and how at finite coupling these zero eigenvalue pairs of the excited oscillators are the ones that may give rise to instability. In fact, it turns out that even the conditions under which instability will ensue for multibreathers of the KG models directly parallel the ones for multi-breathers (or multi-site standing waves) of the DNLS. The latter are analyzed in considerable detail for 1–, 2– and 3– dimensional settings in [Pelinovsky *et al.*, 2005b,a; Lukas *et al.*, 2008]; see also [Kevrekidis, 2009].

Returning to our KG setting, the MEH approach considers the Floquet multipliers given as $\lambda = \exp(\sigma T)$, with $T = 2\pi/\omega$, whereas in the AB approach, $\lambda = \exp(i\theta)$. Then,

$$\sigma = \frac{i\theta}{T} = \frac{i\theta\omega}{2\pi} \quad (2)$$

Due to the symplectic character of the Floquet matrix if λ is a multiplier, so is λ^{-1} , and due to its real character if a λ is non real multiplier, so is λ^* , where the asterisk denotes the complex conjugate.

Therefore, the corresponding multipliers come in complex quadruplets $(\lambda, \lambda^{-1}, \lambda^*, \lambda^{*-1})$ if $|\lambda| \neq 1$ and λ is not real, or in pairs λ, λ^{-1} if λ is real, or λ, λ^* if $|\lambda| = 1$ and not real. In addition, due to the time translation invariance of the system there is always a pair of eigenvalues at $+1$. This has as a result that both σ_i 's and θ_i 's, come also in quadruplets or pairs $(\sigma, -\sigma)$ if σ is real, $((\theta^*, -\theta^*)$ if θ is imaginary) or $(\sigma^*, -\sigma^*)$ if σ is imaginary $((\theta, -\theta)$ if θ is real). In principle, the pairs could collide at a single value $\lambda = 1$ or $\lambda = -1$ if the system undergoes an exponential instability or a period-doubling bifurcation, respectively, although there is always a pair of $+1$'s for the systems under study, as explained above.

2.2. The MEH approach

The MEH approach consists of constructing an effective Hamiltonian, whose critical points are in correspondence with the periodic orbits (in our case multibreathers) of the original system. This method has been originally proposed in [Ahn & Sepulchre, 2001; MacKay & Sepulchre, 2002; MacKay, 2004] and used in the present form in [Koukouloyannis & MacKay, 2005]. The effective Hamiltonian can be constructed as follows.

After considering the central oscillators we apply the action-angle canonical transformation to them. Note that, in the anti-continuous limit, the motion of the central oscillators, in the action-angle variables, is described by $w_i = \omega_i t + w_{i0}$, $J_i = \text{const.}$, for $i = 0, \dots, n-1$. Where w_i is the angle, w_{i0} is the initial phase and J_i the action of the i -th central oscillator. For this kind of systems, the action of an oscillator can be calculated as

$$J_i = \frac{1}{2\pi} \int_0^T p_i dx_i = \frac{1}{2\pi} \int_0^T [\dot{x}_i(t)]^2 dt. \quad (3)$$

Since we are interested in a first order approach, the effective Hamiltonian can be written as $H^{\text{eff}} = H_0 + \epsilon \langle H_1 \rangle$, by neglecting terms which do not contribute to the results in this order of approximation. In this formula, $\langle H_1 \rangle$ is the average value of the coupling term of the Hamiltonian, over an angle in the anticontinuous limit, which is equivalent to the average value of H_1 over a period

$$\langle H_1 \rangle = \frac{1}{T} \oint H_1 dt.$$

This averaging procedure is performed in order to lift the phase degeneracy of the system. For the same reason we introduce a second canonical transformation

$$\begin{aligned} \vartheta &= w_0 & \mathcal{A} &= J_0 + \dots + J_{n-1} \\ \phi_i &= w_i - w_{i-1} & I_i &= \sum_{j=i}^{n-1} J_j & i &= 1, \dots, n-1. \end{aligned} \quad (4)$$

In these variables, the effective Hamiltonian reads

$$H^{\text{eff}} = H_0(I_i) + \epsilon \langle H_1 \rangle(\phi_i, I_i). \quad (5)$$

Note that, since the calculations are performed in the anticontinuous limit, the contribution of the non-central oscillators has disappeared.

As we seldom know the explicit form of the transformation $(x, p) \mapsto (w, J)$, we use the fact that since the motion of the central oscillators for $\epsilon = 0$ is periodic, and possesses the $t \mapsto -t, x \mapsto x, p \mapsto -p$ symmetry, it can be described by a cosine Fourier series $x_i(t) = \sum_{k=0}^{\infty} A_k(J_i) \cos(kw_i)$.

Note that at the anti-continuous limit, the orbits differ only in phase (i.e. $\omega_i = \omega \forall i$), therefore $J_i = J$ and the coefficients A_k 's do not depend on the index i .

So, excluding the constant terms, $\langle H_1 \rangle$ becomes for the KG problem [Koukouloyannis & Kevrekidis, 2009]

$$\langle H_1 \rangle = -\frac{1}{2} \sum_{k=1}^{\infty} \sum_{i=1}^{n-1} A_k^2 \cos(k\phi_i) \quad (6)$$

One of the main features of the MEH approach is that the critical points of this effective Hamiltonian correspond to the multibreather solutions of the system. This fact provides the corresponding persistence conditions, as the simple roots of $\frac{\partial \langle H_1 \rangle}{\partial \phi_i} = 0$. Remarkably, in this setting, similarly to what is known also for the DNLS [Kevrekidis, 2009], it can be proved that the only available multibreather solutions in the one-dimensional case are the ones with relative phase among the excited sites of 0 or π .

The second important fact the MEH approach yields is that the linear stability of these critical points (i.e., the Hessian of the effective Hamiltonian) determines the stability of the corresponding multibreather. In particular, the nonzero characteristic exponents of the central oscillators σ_i (see the discussion in the previous subsection) are given as eigenvalues of the stability matrix $\mathbf{E} = \mathbf{J}D^2H^{\text{eff}}$ where $\mathbf{J} = \begin{pmatrix} \mathbf{O} & -\mathbf{I} \\ \mathbf{I} & \mathbf{O} \end{pmatrix}$ is the matrix of the symplectic structure. By using the form in (5) for the H^{eff} we get:

$$\mathbf{E} = \begin{pmatrix} \mathbf{A} & \mathbf{B} \\ \mathbf{C} & \mathbf{D} \end{pmatrix} = \begin{pmatrix} \epsilon \mathbf{A}_1 & \epsilon \mathbf{B}_1 \\ \mathbf{C}_0 + \epsilon \mathbf{C}_1 & \epsilon \mathbf{D}_1 \end{pmatrix} = \begin{pmatrix} -\epsilon \frac{\partial^2 \langle H_1 \rangle}{\partial \phi_i \partial I_j} & -\epsilon \frac{\partial^2 \langle H_1 \rangle}{\partial \phi_i \partial \phi_j} \\ \frac{\partial^2 H_0}{\partial I_i \partial I_j} + \epsilon \frac{\partial^2 \langle H_1 \rangle}{\partial I_i \partial I_j} & \frac{\partial^2 \langle H_1 \rangle}{\partial \phi_j \partial I_i} \end{pmatrix}. \quad (7)$$

Since the only permitted values of the relative phases are $\phi_i = 0$, or $\phi_i = \pi$, the matrix simplifies considerably acquiring the form:

$$\mathbf{E} = \begin{pmatrix} \mathbf{O} & \mathbf{B} \\ \mathbf{C} & \mathbf{O} \end{pmatrix} = \begin{pmatrix} \mathbf{O} & \epsilon \mathbf{B}_1 \\ \mathbf{C}_0 + \epsilon \mathbf{C}_1 & \mathbf{O} \end{pmatrix}. \quad (8)$$

Then, if we consider only the dominant eigenvalue contributions, we get that $\sigma_i^2 = \epsilon \chi_{BC}$, where χ_{BC} are the eigenvalues of the $(n-1 \times n-1)$ matrix $\mathbf{B}_1 \cdot \mathbf{C}_0$ which reads

$$\mathbf{B}_1 \cdot \mathbf{C}_0 = -\frac{\partial \omega}{\partial J} \mathbf{Z} = -\frac{\partial \omega}{\partial J} \begin{pmatrix} 2f_1 - f_1 & 0 & & & \\ -f_2 & 2f_2 & -f_2 & & 0 \\ & \ddots & \ddots & \ddots & \\ & & 0 & -f_{n-2} & 2f_{n-2} & -f_{n-2} \\ & & & 0 & -f_{n-1} & 2f_{n-1} \end{pmatrix}. \quad (9)$$

In this expression $\omega = \partial H_0 / \partial J$ denotes the frequency, while

$$f_i \equiv f(\phi_i) = \frac{1}{2} \sum_{k=1}^{\infty} k^2 A_k^2 \cos(k\phi_i). \quad (10)$$

This leads to the characteristic exponents (i.e., effective eigenvalues) of the DB in the form:

$$\sigma = \pm \sqrt{-\epsilon \frac{\partial \omega}{\partial J} \chi_z}, \quad (11)$$

with χ_z being the eigenvalues of the $(n-1 \times n-1)$ matrix

$$Z_{i,j} = \begin{cases} Z_{i,i\pm 1} = -f_i \\ Z_{i,i} = 2f_i \\ 0 \text{ otherwise.} \end{cases} \quad (12)$$

2.3. The AB approach

We demonstrate hereby that (11) can be reobtained based on the AB approach, by using the exposition of [Archilla *et al.*, 2003]. To this end, we recall that the aim of the AB approach is to look for the displacement that Aubry's bands [Aubry, 1997] experience when the coupling ϵ is switched on. What we plan to do below is to calculate the Floquet eigenvalues assuming that the bands are parabolic and their shape does not change when the coupling is introduced.

First, we recall the basics of Aubry's band theory with the notation used in [Archilla *et al.*, 2003] adapted to the notation in the present paper, where convenient, for ease of comparison. The Hamilton equations applied to the Hamiltonian of Eq. (1) can be written as:

$$\ddot{x}_n + V'(x_i) + \epsilon \frac{\partial H_1}{\partial x_i} = 0 \quad i = 1, \dots, N, \quad (13)$$

for a generic coupling potential H_1 , or, if it is harmonic:

$$\ddot{x}_n + V'(x_i) + \epsilon \sum_{j=1}^N C_{ij} x_j = 0 \quad i = 1, \dots, N \quad (14)$$

where C is a coupling constant matrix. Let us define $x \equiv [x_1(t), \dots, x_N(t)]^\dagger$ (\dagger meaning the transpose matrix). Defining $V(x) = [V(x_1), \dots, V(x_N)]^\dagger$, $\partial H_1/\partial x = [\partial H_1/\partial x_1, \dots, \partial H_1/\partial x_N]^\dagger$ and so on, Eq. (13) can be written as:

$$\ddot{x} + V'(x) + \epsilon \frac{\partial H_1}{\partial x} = 0. \quad (15)$$

Suppose that $x(t)$ is a time-periodic solution, with period T and frequency ω , its (linear) stability depends on the characteristic equation for the Newton operator \mathcal{N}_ϵ given by

$$\mathcal{N}_\epsilon(u) \xi \equiv \ddot{\xi} + V''(x) * \xi + \epsilon \frac{\partial^2 H_1}{\partial x^2} \xi = E \xi, \quad (16)$$

where $*$ product is the list product, i.e., $f(x) * \xi$ is the column matrix with elements $f(x_i(t)) \xi_i(t)$, and $\partial^2 H_1/\partial x^2$ is the matrix of functions $\partial^2 H_1/\partial x_i \partial x_j$, which depends on t through $x = x(t)$.

If $E = 0$, this equation describes the evolution of small perturbations $\xi = \xi(t)$ of $x = x(t)$, which determines the stability or instability of x . It is however, extremely useful to consider the characteristic equation for any eigenvalue E as it is the cornerstone for Aubry's band theory.

Any solution ξ of Eq. (16) is determined by the column matrix of the initial conditions for positions and momenta $\Omega(0) = [\xi_1(0), \dots, \xi_N(0), \pi_1(0), \dots, \pi_N(0)]^\dagger$, with $\pi_i(t) = \dot{\xi}_i(t)$. A basis of solutions is given by the $2N$ functions with initial conditions $\Omega^\nu(0)$, $\nu = 1, \dots, 2N$, with $\Omega_\nu^\nu(0) = \delta_{\nu l}$.

The Newton operator depends on the T -periodic solution $x(t)$, and therefore, it is also T -periodic and its eigenfunctions can be chosen also as eigenfunctions of the operator of translation in time (a period T). They are the Bloch functions $\xi(\theta_i, t) = \chi(\theta_i, t) \exp(i\theta_i t/T)$, with $\chi(\theta, t)$ being a column matrix of T -periodic functions. The sets $\{\xi(\theta_i, 0), \dot{\xi}(\theta_i, 0)\}$ are also the eigenvectors of the Floquet operator \mathcal{F}_E or monodromy, that maps $\Omega(0)$ into $\Omega(T)$, that is, $\Omega(T) = \mathcal{F}_E \Omega(0)$. Their corresponding eigenvalues are the $2N$ multipliers $\{\lambda_i\} = \exp(\theta_i)$, with $\{\theta_i\}$ being the $2N$ Floquet arguments.

The set of points (θ, E) , with θ being a real Floquet argument of \mathcal{F}_E , has a band structure. As the Newton and Floquet operators are real, the Floquet multipliers come in complex conjugate pairs. Therefore, if (θ, E) belongs to a band (i.e. θ is real), $(-\theta, E)$ does as well, i.e., the bands are symmetric with respect to θ , which implies that $dE/d\theta(0) = 0$. There are always two T -periodic solutions, with Floquet multiplier $\lambda = 1$ ($\theta = 0$) for $E = 0$. One is $\dot{x}(t)$, which represents a change in phase of the solution $x(t)$ and corresponds to the *phase mode*; the other is the *growth mode*, given by $\partial x(t)/\partial \omega$, and represents a change in frequency and consequently in amplitude. The consequence is that there is always a symmetric band tangent to the axis $E = 0$ at $\theta = 0$.

There are at most $2N$ points for a given value of E and, therefore, there are at most $2N$ bands crossing the horizontal axes $E = 0$ in the space of coordinates (θ, E) . The condition for linear stability of $x(t)$ is equivalent to the existence of $2N$ bands crossing the axis $E = 0$ (including tangent points with their multiplicity). If a parameter like the coupling ϵ changes, the bands evolve continuously, and they can lose crossing points with $E = 0$, leading to an instability of the system.

The first item to find out are the bands at the anticontinuous limit, where Eq. (14) reduces to N identical equations:

$$\ddot{x}_i + V'(x_i) = 0. \quad (17)$$

If we consider solutions around a minimum of V , the oscillators can be at rest $x_i = 0$, or oscillating with period T ; the latter are identical except for a change in the initial phase, so they can be written as $x_i(t) = g(\omega t + w_{i0})$ with $g(\omega t)$ being the only T -periodic, time-symmetric solution of Eq. (17) with $g(0) > g(\pi)$. Therefore, the excited oscillators can be written as:

$$x_i(t) = z_0 + 2 \sum_{k=1}^{\infty} z_k \cos[k(\omega t + w_{i0})] = \sum_{k=0}^{\infty} A_k \cos[k(\omega t + w_{i0})] = \sum_{k=0}^{\infty} A_k \cos(kw_i), \quad (18)$$

with $A_k = 2z_k$ if $k > 0$, $A_0 = z_0$ and $w_i = \omega t + w_{i0}$.

Let n be the number of excited oscillators at the anticontinuous limit, labeled $i = 0, \dots, n-1$. Then, there are n identical bands tangent to the axis $E = 0$ at $\theta = 0$ for each excited oscillator, and $N - n$ bands, corresponding to the oscillators at rest, with $2(N - n)$ points intersecting the $E = 0$ axis.

In what follows we will show that, by making the appropriate assumptions for the excited bands, we can use the AB method to calculate the characteristic exponents of the multibreather and the results are equivalent to the MEH method. These assumptions are that the form of the bands is parabolic and remains that way in the first order of approximation. Thus, the excited bands can be approximated around $(\theta, E) = (0, 0)$ by

$$E(\theta) \approx E_0 + \kappa \theta^2, \quad (19)$$

with $E_0 = \epsilon \chi_q$ and χ_q being the eigenvalues of the $(n \times n)$ Q -matrix defined below. Additionally,

$$\kappa = \frac{1}{2} \frac{\partial^2 E}{\partial \theta^2} = -\frac{\omega^2}{4\pi^2 J} \frac{\partial H}{\partial \omega} = -\frac{1}{T^2 J} \frac{\partial H}{\partial \omega} \quad (20)$$

where we have made use of [Archilla *et al.*, 2003, Eq. (B14)]. The factor κ is positive if the on-site potential V is hard and negative if V is soft (a potential is hard if the oscillation amplitude increases with the frequency and soft otherwise). When the coupling is switched on, the bands will move and change shape; the $E = 0$ eigenvalue is degenerate with multiplicity $N - n$ at $\epsilon = 0$, but this degeneracy is generically lifted for $\epsilon \neq 0$ and only one band will continue being tangent at $(\theta, E) = (0, 0)$ due to the phase mode. Applying degenerate perturbation theory to Eq. (16), with ϵH_1 being the perturbation, a perturbation matrix Q can be constructed [Archilla *et al.*, 2003], whose eigenvalues χ_q are those of the perturbed Newton operator. The non-diagonal elements of Q are given by

$$Q_{ij} = \frac{1}{\mu_i \mu_j} \int_0^T \dot{x}_i \frac{\partial^2 H_1}{\partial x_i \partial x_j} \dot{x}_j dt, \quad i \neq j, \quad i = 0 \dots n-1, \quad j = 0 \dots n-1, \quad (21)$$

with $\mu_i = \sqrt{\int_0^T (\dot{x}_i)^2 dt}$. The diagonal elements are

$$Q_{ii} = -\sum_{j \neq i} \frac{\mu_j}{\mu_i} Q_{ij}. \quad (22)$$

If the on-site potential $V(x_i)$ is homogeneous and the coupling is given as in Eq. (1), as is the case in the present paper, $\mu_i = (2\pi J)^{1/2} \forall i$. Let us calculate the derivatives of H_1 , $h_{i,j} = \partial^2 H_1 / \partial x_i \partial x_j$. Because of the way the diagonal elements of Q are constructed, we only need the derivatives with $i \neq j$. It is easy to see that they are zero except for $h_{i-1,i} = h_{i,i-1} = -q_i$ (defined below) for $i = 1, \dots, n-1$. The derivatives $h_{0,n-1}$ and $h_{n-1,0}$ are also zero as the oscillators at the extremes of the multibreather are not coupled between them. Then, the matrix Q becomes:

$$Q_{i,j} = \begin{cases} Q_{i,i-1} = Q_{i-1,i} = -q_i, & \text{for } i = 1 \dots n-1 \\ Q_{i,i} = q_{i-1} + q_i, & \text{for } i = 1 \dots n-2 \\ Q_{0,0} = q_1 \\ Q_{n-1,n-1} = q_{n-1} \\ 0 & \text{otherwise} \end{cases} \quad (23)$$

or, explicitly:

$$Q = \begin{pmatrix} q_1 & -q_1 & 0 & & \\ -q_1 & q_1 + q_2 & -q_2 & & 0 \\ & \ddots & \ddots & \ddots & \\ & 0 & -q_{n-2} & q_{n-2} + q_{n-1} & -q_{n-1} \\ & & 0 & -q_{n-1} & q_{n-1} \end{pmatrix}, \quad (24)$$

with

$$q_i \equiv q(\phi_i) = \frac{\int_0^T \dot{x}_i(t) \dot{x}_{i-1}(t) dt}{\int_0^T [\dot{x}_i(t)]^2 dt} = \frac{\omega}{2J} \sum_{k \geq 1} k^2 A_k^2 \cos(k\phi_i) = \frac{\omega}{J} f_i, \quad i = 1, \dots, n-1, \quad (25)$$

Then, by using [Sandstede, 1998, Lemma 5.4] we see that the matrices Q and $\frac{\omega}{J}Z$ have the same nonzero eigenvalues i.e.

$$\chi_q = \frac{\omega}{J} \chi_z. \quad (26)$$

In addition, Q has also a zero eigenvalue¹.

Some important values of $q(\phi)$ are the following ones:

$$q(0) = 1 \quad (27)$$

$$q(\pi) = \frac{\sum_{k \geq 1} (-1)^k k^2 z_k^2}{\sum_{k \geq 1} k^2 z_k^2} \equiv -\gamma \quad (28)$$

For a Morse potential, $\gamma = \omega$; for an even potential, $\gamma = 1$, as shown in [Archilla *et al.*, 2003].

According to the AB theory [Aubry, 1997], the Floquet multipliers are given by the cuts of the bands with the $E = 0$ axis; thus

$$\theta = \pm \sqrt{-\frac{E_0}{\kappa}} = \sqrt{-\frac{\epsilon \omega}{\kappa J}} \chi_z \quad (29)$$

and, applying the last results

$$\theta = \pm T \sqrt{\epsilon \frac{\partial \omega}{\partial J}} \chi_z, \quad (30)$$

where we have taken into account that²

$$\frac{\partial H}{\partial \omega} = \frac{\partial H}{\partial J} \frac{\partial J}{\partial \omega} = \omega \frac{\partial J}{\partial \omega} \quad (31)$$

Finally, introducing (30) into (2), we get (11), which completes the proof of equivalence of the relevant Floquet multiplier predictions.

¹The fact that the Z matrix does not have a zero eigenvalue is a key point for the MEH method. This method is based in the continuation from the anticontinuous limit, which means that it makes use of the implicit function theorem. The Z matrix is directly related to the continuation matrix. So, if Z had a 0 eigenvalue the continuation would fail since the implicit function theorem would not be valid. This is the the reason we use H^{eff} instead of H .

²The expression $\omega = \partial H / \partial J$ comes from the Hamilton's equations for the action-angle variables, since all the calculations are performed in the uncoupled, and therefore integrable, limit.

3. Some Examples

In what follows we will illustrate the use of the method in order to acquire stability results by calculating the characteristic exponents of some prototypical Klein-Gordon lattice configurations. In section 3.1 we calculate the exponents of a multibreather in a 1D Klein-Gordon chain with a Morse on-site potential. These results were already known from earlier works (e.g. [Koukouloyannis & Ichtiaroglou, 2006]) for the 2-site and 3-site cases. In the present exposition we generalize these results for the n -site breather case. In section 3.2 we show that this method can also be used for calculations in higher dimensional lattices. For this purpose we use a 2D square lattice with Morse on-site potential and focus on the case of vortex-breathers. Finally, in section 3.3 we show how this method can be used in cases where the analytical calculations are too cumbersome to handle. In particular, we use a 1D Klein-Gordon chain with hard ϕ^4 potential and the rotating wave approximation in order to avoid the complicated calculations.

3.1. The 1D Morse Klein-Gordon chain

We now consider some special case examples, starting with a linearly coupled lattice of oscillators subject to the Morse potential, $V(x) = (\exp(-x) - 1)^2$. As indicated previously, the only configurations that may exist in the one-dimensional setting are ones which involve excited oscillators either in-phase (i.e., with $\phi_i = 0$) or out-of-phase (i.e., with $\phi_i = \pi$); see [Cuevas *et al.*, 2005], [Koukouloyannis & Kevrekidis, 2009] and also [Cuevas, 2003] for a detailed discussion. Here, we proceed to perform some explicit calculations for the Floquet multipliers σ in the case of n -site breathers. In what follows we consider only the positive σ . To this end we express (11) making use of (26):³

$$\sigma = \sqrt{-\epsilon \frac{J}{\omega} \frac{\partial \omega}{\partial J} \chi_q(\phi)} \quad (32)$$

where $\chi_q(\phi)$ denotes the Q -matrix eigenvalues for a given ϕ . It is straightforward to show that⁴

$$\chi_q(0) = 4 \sin^2 \frac{m\pi}{2n} \quad m = 1, \dots, n-1 \quad (33)$$

and that

$$\chi_q(\pi) = -\gamma \chi_q(0) \quad (34)$$

For instance, in the case of a 2-site breather, $\chi_q(0) = 2$ and $\chi_q(\pi) = -2\gamma$.

We now focus on the particular case of the Morse potential, since it is a potential for which closed form analytical expressions can be found. [For other types of potentials, some approximations can be made for small and high frequencies; alternatively, the required single-oscillator parameters, such as J and $\partial\omega/\partial J$ can be calculated numerically].

In the Morse case and in order to evaluate J and $\partial\omega/\partial J$, we express J as a function of the Fourier coefficients:

$$J = 2\omega \sum_{k \geq 1} k^2 z_k^2. \quad (35)$$

For this potential,

$$z_0 = \ln \frac{1 + \omega}{2\omega^2}; \quad z_k = \frac{(-1)^k}{k} r^{k/2}, \quad r = \frac{1 - \omega}{1 + \omega}. \quad (36)$$

³In what follows, and in order to fix ideas, given the equivalence of the two methods, we will use the formulation with the Q -matrix.

⁴We are neglecting the 0 eigenvalue, associated with $m = 0$.

Substituting into the action

$$J = 1 - \omega \rightarrow \frac{\partial \omega}{\partial J} = -1. \quad (37)$$

Thus,

$$\sigma(\phi) = \sqrt{\epsilon \frac{1 - \omega}{\omega} \chi_q(\phi)}. \quad (38)$$

In the case of a general phase, we can express $\chi_q(\phi) = q(\phi)\chi_q(0)$ with $q(\phi)$ given by (25). In the special case of the Morse potential, we have

$$q(\phi) = \frac{2\omega}{J} \sum_{k \geq 1} r^k \cos(k\phi). \quad (39)$$

To obtain the relevant sum, we use a simple geometric series formula that can be found e.g. in [Gradshcheyn & Ryzhik, 1965], according to which:

$$q(\phi) = \frac{2\omega}{J} r \frac{\cos \phi - r}{1 - 2r \cos \phi + r^2}. \quad (40)$$

Consequently,

$$\sigma(\phi) = \sqrt{2\epsilon r \frac{\cos \phi - r}{1 - 2r \cos \phi + r^2} \chi_q(0)}. \quad (41)$$

For the relevant values of ϕ for time-reversible multibreathers, we get:

$$\sigma(0) = \sqrt{\epsilon \frac{1 - \omega}{\omega} \chi_q(0)} = 2 \sin \frac{m\pi}{2n} \sqrt{\epsilon \frac{1 - \omega}{\omega}} \quad m = 1, \dots, n - 1 \quad (42)$$

$$\sigma(\pi) = \sqrt{-\epsilon(1 - \omega) \chi_q(0)} = 2 \sin \frac{m\pi}{2n} \sqrt{-\epsilon(1 - \omega)} \quad m = 1, \dots, n - 1 \quad (43)$$

Figures 1 and 2 show, respectively, the analytical eigenvalue predictions (dashed lines) for stable and unstable two-site and three-site breathers with the Morse potential and how they favorably compare to the corresponding numerical results (solid lines), obtained via a fully numerical linear stability analysis (and corresponding computation of the Floquet multipliers). It is clear that the predictions are very accurate close to the anti-continuum limit, and their validity becomes progressively limited for larger values of the coupling parameter ϵ , yet they yield a powerful qualitative and even quantitative (in the appropriate parametric regime) tool for tracking the stability of these localized modes. The figures also illustrate typical profiles of the corresponding two- and three-site ILMs.

3.2. Vortices in square Morse lattices

The methodology can also be extended to lattices of higher dimensionality. We consider below some basic properties of discrete vortex breathers of different integer topological charges, in a square 2D lattice. Firstly, we consider square vortices over a single ‘‘plaquette’’ of the 2D lattice with $S = 1$, i.e., at the anti-continuum limit, the excited sites are $(0, 0)$, $(0, 1)$, $(1, 1)$ and $(1, 0)$ with a phase difference $\phi = \pi/2$ between nearest neighbors. This implies a perturbation matrix given by:




Fig. 1. (Top panels) Profiles of an in phase (left) and an out-of-phase (right) 2-site breather with the Morse potential for $\omega = 0.8$ and $\epsilon = 0.05$. The bottom panels show the value of the characteristic exponents σ of the corresponding configurations, with respect to the coupling parameter ϵ . Dashed lines correspond to the predictions of the stability theorems, while solid ones to full numerical linear stability analysis results.

$$Q = q(\pi/2)\tilde{Q}, \text{ with } \tilde{Q} = \begin{pmatrix} 2 & -1 & 0 & -1 \\ -1 & 2 & -1 & 0 \\ 0 & -1 & 2 & -1 \\ -1 & 0 & -1 & 2 \end{pmatrix} \quad (44)$$

with $q(\pi/2)$ given from (40) which is evaluated as:

$$q(\pi/2) = -\frac{(1-\omega)^2}{J(1+\omega^2)}. \quad (45)$$

This, in turn, upon use of Eq. (32) implies that

$$\sigma = i(1-\omega)\sqrt{\frac{\epsilon}{1+\omega^2}}\tilde{\chi}_q, \quad (46)$$

where $\tilde{\chi}_q$ are the eigenvalues of the \tilde{Q} matrix. This corresponds to the matrix of the normal modes of a 1D chain of 4 linearly coupled oscillators with periodic boundary conditions. Let us recall that for a system of n coupled oscillators, the eigenvalues are given by:




Fig. 2. Same as in Fig. 1, but now for the unstable left-panel configuration of three in-phase excited sites (with two real multiplier pairs as shown in the bottom panel) and the case of the out-of-phase, three-site right-panel configuration, which is stable only in the proximity of to the anti-continuum limit.

$$\tilde{\chi}_q = 4 \sin^2 \frac{m\pi}{n} \quad m = 1, \dots, n-1, \quad (47)$$

in addition to the 0 eigenvalue. In the present case of 4 oscillators with periodic boundary conditions, the nonzero eigenvalues are given by 2 and 4, with the former being doubly degenerate. Thus, we have for $S = 1$ vortices the following spectrum:

$$\sigma = \begin{cases} i(1 - \omega) \sqrt{2 \frac{\epsilon}{1 + \omega^2}} & \text{single eigenvalue} \\ 2i(1 - \omega) \sqrt{\frac{\epsilon}{1 + \omega^2}} & \text{double eigenvalue} \end{cases} \quad (48)$$

which implies stability for $\epsilon > 0$.

This type of analysis can be generalized for arbitrary values of the vorticity S , leading to the conclusion that \tilde{Q} is the matrix of $4S$ coupled oscillators, which implies that vortices with *any* integer topological charge will be stable for $\epsilon > 0$ in the case of a lattice with an on-site Morse potential. For instance, in the case of the $S = 2$ vortex, we obtain the explicit expressions for the eigenvalues:




Fig. 3. The characteristic exponents of vortex configurations with $S = 1$ (left) and $S = 2$ (right), with respect to ϵ , for the Morse potential and $\omega = 0.8$. Dashed lines correspond to the theoretical predictions based on Eqs. (48) and (49), respectively; the full numerical linear stability results are given by solid lines and indicate that all doubly degenerate eigenvalue pairs split due to higher order contributions in the relevant expansions in the coupling constant ϵ .

$$\sigma = \begin{cases} \frac{i(1-\omega)}{2} \sqrt{(2 - 2^{1/2}) \frac{\epsilon}{1+\omega^2}} & \text{single eigenvalue} \\ \frac{i(1-\omega)}{2} \sqrt{(2 + 2^{1/2}) \frac{\epsilon}{1+\omega^2}} & \text{double eigenvalue} \\ i(1-\omega) \sqrt{2 \frac{\epsilon}{1+\omega^2}} & \text{double eigenvalue} \\ 2i(1-\omega) \sqrt{\frac{\epsilon}{1+\omega^2}} & \text{single eigenvalue} \end{cases} \quad (49)$$

It is important to highlight here some interesting differences between the above results and the case of the DNLS (and more generally that of even potentials in KG chains, including the case of the hard ϕ^4 lattice considered below). In the latter class of problems, the vanishing of the odd coefficients in the Fourier expansion of the periodic orbit leads to the conclusion that $q(\pi/2) = 0$ and hence there is no contribution to the eigenvalues to leading order. This is the situation which has been characterized as “super-symmetric” in [Pelinovsky *et al.*, 2005a; Kevrekidis, 2009] and one in which the higher order contributions would be critical in determining the stability. Nevertheless, in the case considered herein, the asymmetry of the Morse potential produces a nonvanishing of $q(\pi/2)$ and offers a corresponding nonzero leading order correction to the eigenvalues at $O(\epsilon^{1/2})$.

Figure 3 shows the dependence of stability eigenvalues for the $S = 1$ and $S = 2$ vortices and their comparison with the obtained fully numerical linear stability results as a function of the coupling ϵ . As can be observed in the figures, the approximation is less accurate in this case, although it is qualitatively correct. The reason for the partial disparity is that higher order contributions to the relevant eigenvalues (whose calculation is considerably more technically involved) lead to the observed splitting of all the doubly degenerate eigenvalue pairs. In the relevant cases, the analytical (dashed line) predictions can be seen to straddle the two observed numerical pairs.

3.3. The 1D hard ϕ^4 Klein-Gordon chain

The time evolution of a single oscillator in the hard ϕ^4 potential, $V(x) = x^2/2 + x^4/4$ is given by:

$$x(t) = \sqrt{\frac{2m}{1-2m}} \operatorname{cn} \left(\frac{t}{\sqrt{1-2m}}, m \right) = \sqrt{\frac{2m}{1-2m}} \operatorname{cn} \left(\frac{2K(m)}{\pi} \omega t, m \right), \quad (50)$$

where cn is a Jacobi elliptic function of modulus m and $K(m)$ is the complete elliptic integral of the first kind defined as $K(m) = \int_0^{\pi/2} [1 - m \sin^2 x]^{-1/2} dx$.

The breather frequency ω is related to the modulus m through:

$$\omega = \frac{\pi}{2\sqrt{1-2m}K(m)}. \quad (51)$$

The elliptic function can be expanded into a Fourier series leading to [Abramowitz & Stegun, 1965]:

$$z_{2\nu+1} = \frac{\pi}{K(m)} \sqrt{\frac{2}{1-2m}} \frac{q^{\nu+1/2}}{1+q^{2\nu+1}}, \quad \nu = 0, 1, 2, \dots \quad (52)$$

where q is the elliptic Nome which is defined as

$$q \equiv q(m) = \exp(-\pi K(1-m)/K(m)). \quad (53)$$

In order to get $\chi_q(\phi)$ and $\partial\omega/\partial J$, we cannot use (10) and (25) because it is not possible to find a closed form expression. Instead, we use the integral expression:

$$f(\phi_i) = \frac{1}{2\pi\omega} \int_0^T \dot{x}_i(t) \dot{x}_{i+1}(t) dt. \quad (54)$$

After some manipulations (where it is crucial to apply [Khare *et al.*, 2003, identity 171]), we obtain:

$$\begin{aligned} f(\phi) &= \frac{8K(m)}{\pi^3\omega(1-2m)} [\operatorname{cs}(a, m)\operatorname{ns}(a, m)[2E(m) - K(m)(1 + \operatorname{dn}^2(a, m))]] \\ &\quad - \frac{8K(m)}{\pi^3\omega(1-2m)} [\operatorname{ds}(a, m)(\operatorname{cs}^2(a, m) + \operatorname{ns}^2(a, m))Z(a, m)] \end{aligned} \quad (55)$$

where $E(m)$ is the complete elliptic integral of the second kind defined as $E(m) = \int_0^{\pi/2} [1 - m \sin^2 x]^{1/2} dx$, $Z(a, m)$ is the Jacobi zeta function and $a = 2K(m)\phi/\pi$.

For the action J , a similar manipulation leads to

$$J = \frac{16K(m)}{3\pi^2} \left[\frac{1-m}{1-2m} K(m) - E(m) \right]. \quad (56)$$

The derivative of this expression is cumbersome to handle. So, in what follows, we will work instead with numerically obtained values of J and $\partial\omega/\partial J$ which are relevant for time-reversible multibreathers and vortex breathers, as for these cases we need $f(0)$, $f(\pi)$ and $f(\pi/2)$. As indicated previously, for every even potential, $z_{2\nu+1} = 0$, and, consequently, $f(0) = J/\omega$, $f(\pi) = -f(0)$ and $f(\pi/2) = 0$. This leads to:

$$\sigma(0) = \sqrt{-\epsilon \frac{J}{\omega} \frac{\partial\omega}{\partial J} \chi_q(0)}, \quad (57)$$

$$\sigma(\pi) = \sqrt{\epsilon \frac{J}{\omega} \frac{\partial\omega}{\partial J} \chi_q(0)}. \quad (58)$$

Figure 4 shows the dependence of J and $\partial\omega/\partial J$ with respect to the frequency. Figures 5 and 6 illustrate subsequently the relevant stability eigenvalues for 2-site and 3-site breathers as obtained from the




Fig. 4. Dependence with respect to ω of the action (left) and $\partial\omega/\partial J$ (right) for the hard ϕ^4 potential. The dashed line corresponds to the prediction of the RWA [Eq. (59)], while the solid one represents the exact numerical result.

expressions above and compare them to the full numerical linear stability results. The agreement in this case is very good (there are no degeneracies and associated higher-order contributions that may deteriorate the quality of the agreement as in the vortex breather case above); in fact, in some of the cases, the curves are almost indistinguishable throughout the considered parameter range.

An important observation concerns, however, the role of the “hard” nature of the potential. In particular, as illustrated in Fig. 4, the quantity $\partial\omega/\partial J$ is positive in this case, i.e., its sign is opposite from the soft case of the Morse potential (where $\partial\omega/\partial J = -1$). This results in the corresponding reversal of the stability conclusions in Figs. 5 and 6, in comparison with Figs. 1 and 2 of the Morse case. That is, in-phase modes are now stable, while out-of-phase ones are unstable (as is true for the defocusing nonlinearity DNLS case also), while the reverse was true in the Morse potential (as well as for the focusing DNLS case). Lastly, we recall that since this is an even potential and thus $f(\pi/2) = 0$, the leading order calculation would yield a vanishing contribution to the eigenvalues for the vortex case and a higher-order calculation is necessary to determine the stability of the latter.

As an aside towards obtaining a fully analytical prediction for this case (as some of the quantities need to be obtained numerically above), we note the following. Although we cannot acquire an exact form for $J(\omega)$, as in the case of the Morse potential, an approximate form for J can be found by using the rotating wave approximation (RWA), i.e. by supposing that $x(t) \approx 2z_1 \cos(\omega t)$. The introduction in the dynamical equations for the single oscillator leads to:

$$z_1 = \sqrt{\frac{\omega^2 - 1}{3}} \quad (59)$$

Thus, $J = 2\omega z_1^2 = 2\omega(\omega^2 - 1)/3$ and $\partial\omega/\partial J = 1/[2(\omega^2 - 1/3)]$, and the corresponding expressions for the eigenvalues read:

$$\sigma(0) \approx \sqrt{-\epsilon \frac{\omega^2 - 1}{3\omega^2 - 1} (\cos \phi) \chi_q(0)} \quad (60)$$

$$\sigma(\pi) \approx \sqrt{\epsilon \frac{\omega^2 - 1}{3\omega^2 - 1} (\cos \phi) \chi_q(0)} \quad (61)$$

A comparison between the numerically acquired values of $J(\omega)$ and $\partial\omega/\partial J$ with the ones calculated from the RWA is shown in figure 4. The agreement is remarkable and attests to the quality of the “single




Fig. 5. (Top panels) Profiles of an in-phase (left) and an out-of-phase (right) 2-site breather with the hard ϕ^4 potential; $\omega = 3$ and $\epsilon = 0.05$. The bottom panels show the dependence of the characteristic exponents σ , of the corresponding configurations, on the coupling parameter ϵ . The dashed lines correspond to the predictions of the stability theorems and dash-dotted lines to the RWA predictions, while the solid ones represent the full numerical result.

frequency” rotating wave approximation. In Figs. 5 and 6 the characteristic exponents calculated numerically (solid lines) as well as using Eqs. (57)-(58) (dashed lines) and via Eqs. (60)-(61) are compared, illustrating the excellent agreement between all three.

4. Conclusions and Perspectives

The results presented in this work underscore the formulation of a toolbox that enables the systematic characterization of both the qualitative and even the quantitative aspects of stability

of multibreather and vortex breather waveforms in these large number of degree of freedom, Hamiltonian lattice systems of the Klein-Gordon variety. A systematic calculation of the corresponding Floquet multipliers is presented and highlights the crucial components that imply stability, namely the proper combination of the sign of the coupling constant, the nature (hard or soft) of the potential and the relative phases between the adjacent excited sites. E.g., for positive couplings, and soft potentials, out-of-phase structures may be stable near the vanishing coupling limit, while in-phase ones are always unstable, resembling the corresponding DNLS predictions [Kevrekidis, 2009]; the nature of the conclusions is reversed for either (small) negative couplings or for hard potentials. The explicit analytical predictions have been tested against numerical results both for symmetric (such as the hard ϕ^4) and asymmetric (such as the Morse) potentials, both for hard and soft ones, and both for simpler, non-degenerate one-dimensional multibreather settings and for more complex and degenerate two-dimensional vortex breathers. In all cases, the




Fig. 6. Same as Fig. 5, but for the three-site in-phase (left) and out-of-phase (right) configuration. Again the dash-dotted lines in the bottom panels represent the (fully-analytical) RWA predictions, which agree well with the semi-analytical dependences (dashed lines) and, in turn, with the full numerical results (solid lines).

two theories whose results were shown to be equivalent herein, namely the Aubry band theory and the MacKay Effective Hamiltonian method yield excellent qualitative and good quantitative agreement with the full numerical linear stability results. The accuracy of the predictions is lowered in degenerate cases where higher order contributions may be critical in breaking the relevant degeneracy (as we saw in the case of the discrete vortices for the Morse model).

Naturally, a number of interesting directions for future consideration hereby arise. Perhaps the canonical one among them would involve a systematic derivation of higher order corrections for prototypical cases where the leading order approach yields vanishing results. For instance, the characterization of the stability of discrete vortices in the “super-symmetric” case of phase difference $\phi = \pi/2$ for even potentials would be a natural example. Another possibility that is also emerging and would be relevant to consider from a mathematical point of view would be to examine models with inter-site nonlinearities, such as ones of the Fermi-Pasta-Ulam type. In these cases, where the potential is a function $V(x_n - x_{n-1})$, it is relevant to point out that upon consideration of the so-called strain variables $r_n = x_n - x_{n-1}$, the problem is reverted to an on-site potential case, for which it would be worthwhile to explore methods similar to the ones analyzed herein. These directions are presently under consideration and will be reported in future publications.

Acknowledgments. PGK gratefully acknowledges support from NSF-DMS-0349023 (CAREER), NSF-DMS-0806762, NSF-CMMI-1000337, from the Alexander von Humboldt Foundation and from the Alexander S. Onassis Public Benefit Foundation. JC and JFRA acknowledge financial support from the MICINN

project FIS2008-04848.

References

- Abramowitz, M. & Stegun, I. [1965] *Handbook of mathematical functions* (Dover, New York).
- Ahn, R., T. MacKay & Sepulchre, J.-A. [2001] “Dynamics of relative phases: generalised multibreathers,” *Nonlinear Dyn.* **25**, 157.
- Archilla, J., Cuevas, J., Sánchez-Rey, B. & Álvarez, A. [2003] “Demonstration of the stability or instability of multibreathers at low coupling,” *Physica D* **180**, 235.
- Aubry, S. [1997] “Breathers in nonlinear lattices: Existence, linear stability and quantization,” *Physica D* **103**, 201.
- Belova, T. & Kudryavtsev, A. [1997] “Solitons and their interaction in classical field theory,” *Phys. Usp.* **40**, 359.
- Boechler, N., Theocharis, G., Job, S., Kevrekidis, P., Porter, M. & Daraio, C. [2010] “Discrete breathers in one-dimensional diatomic granular crystals,” *Phys. Rev. Lett.* **104**, 244302.
- Brazhnyi, V. & Konotop, V. [2004] “Theory of nonlinear matter waves in optical lattices,” *Mod. Phys. Lett. B* **18**, 627.
- Cubero, D., Cuevas, J. & Kevrekidis, P. [2009] “Nucleation of breathers via stochastic resonance in nonlinear lattices,” *Phys. Rev. Lett.* **102**, 205505.
- Cuevas, J. [2003] “Localization and energy transfer in non-homogeneous anharmonic lattices,” PhD thesis, University of Seville, Spain.
- Cuevas, J., Archilla, J. & Romero, F. [2005] “Effect of the introduction of impurities on the stability properties of multibreathers,” *Nonlinearity* **18**, 76.
- Cuevas, J., Archilla, J. & Romero, F. [2011] “Stability of non-time-reversible phonobreathers,” *J. Phys. A.: Math. Theor.* **44**, 035102.
- Cuevas, J., English, L., Kevrekidis, P. & Anderson, M. [2009] “Discrete breathers in a forced-damped array of coupled pendula: Modeling, Computation and Experiment,” *Phys. Rev. Lett.* **102**, 224101.
- English, L., Basu Thakur, R. & Stearett, R. [2008] “Patterns of travelling ilms in a driven electrical line,” *Phys. Rev. E* **77**, 066601.
- English, L., Palmero, F., Sievers, A., Kevrekidis, P. & Barnak, D. [2010] “Traveling and stationary intrinsic localized modes and their spatial control in electrical lattices,” *Phys. Rev. E* **81**, 046605.
- English, L., Sato, M. & Sievers, A. [2001a] “Modulational instability of the nonlinear uniform spin wave mode in easy axis antiferromagnetic chains: II. influence of sample shape on intrinsic localized modes and dynamic spin defects,” *Phys. Rev. B* **67**, 024403.
- English, L., Sato, M. & Sievers, A. [2001b] “Nanoscale intrinsic localized modes in an antiferromagnetic lattice,” *J. Appl. Phys.* **89**, 6706.
- Flach, S. & Gorbach, A. [2008] “Discrete breathers — Advances in theory and applications,” *Phys. Rep.* **467**, 1.
- Flach, S. & Willis, C. [1998] “Discrete breathers,” *Phys. Rep.* **295**, 182.
- Gradshteyn, I. & Ryzhik, I. [1965] *Table of integrals, series, and products* (Academic Press, New York).
- Kevrekidis, P. [2009] *The discrete nonlinear Schrödinger equation* (Springer-Verlag, Berlin).
- Khare, A., Lakshminarayan, A. & Sukhatme, U. [2003] “Local identities involving elliptic functions,” ArXiv:math-ph/0306028. This paper was published as *Pramana* 62 (2004) 1201, but the referred identity only appears in the preprint.
- Koukouloyannis, V. & Ichtiaroglou, S. [2006] “A stability criterion for multibreathers in klein-gordon chains,” *International Journal of Bifurcation and Chaos* **16**, 1823–1827.
- Koukouloyannis, V. & Kevrekidis, P. [2009] “On the stability of multibreathers in klein–gordon chains,” *Nonlinearity* **22**, 2269.
- Koukouloyannis, V., Kevrekidis, P., Law, K., Kourakis, I. & Frantzeskakis, D. [2010] “Existence and stability of multisite breathers in honeycomb and hexagonal lattices,” *J. Phys. A: Math. Theor.* **43**, 235101.
- Koukouloyannis, V. & Kourakis, I. [2007] “Existence of multisite intrinsic localized modes in one-

- dimensional debye crystals,” *Phys. Rev. E* **76**, 016402.
- Koukoulouyannis, V. & Kourakis, I. [2009] “Discrete breathers in hexagonal dusty plasma lattices,” *Phys. Rev. E* **80**, 026402.
- Koukoulouyannis, V. & MacKay, R. [2005] “Existence and stability of 3-site breathers in a triangular lattice,” *J. Phys. A: Math. Gen.* **38**, 1021.
- Lederer, F., Stegeman, G., Christodoulides, D., Assanto, G., Segev, M. & Silberberg, Y. [2008] “Discrete solitons in optics,” *Phys. Rep.* **463**, 1.
- Lukas, M., Pelinovsky, D. & Kevrekidis, P. [2008] “Lyapunov-schmidt reduction algorithm for three-dimensional discrete vortices,” *Physica D* **237**, 339.
- MacKay, R. [2004] “Slow manifolds,” *Energy Localisation and Transfer* (World Scientific), pp. 149–192.
- MacKay, R. & Aubry, S. [1994] “Proof of the existence of breathers for time-reversible or Hamiltonian networks of weakly coupled oscillators,” *Nonlinearity* **7**, 1623.
- MacKay, R. & Sepulchre, J.-A. [2002] “Effective Hamiltonian for traveling discrete breathers,” *J. Phys. A: Math. Gen.* **35**, 3985.
- Morsch, O. & Oberthaler, M. [2006] “Dynamics of bose-einstein condensates in optical lattices,” *Rev. Mod. Phys.* **78**, 179.
- Pelinovsky, D., Kevrekidis, P. & D.J., F. [2005a] “Persistence and stability of discrete vortices in nonlinear schrödinger lattices,” *Physica D* **212**, 20.
- Pelinovsky, D., Kevrekidis, P. & D.J., F. [2005b] “Stability of discrete solitons in nonlinear schrödinger lattices,” *Physica D* **212**, 1.
- Peyrard, M. [2004] “Nonlinear dynamics and statistical mechanics of DNA,” *Nonlinearity* **17**, R1.
- Sandstede, B. [1998] “Stability of multiple-pulse solutions,” *Trans. Am. Math. Soc.* **350**, 429.
- Sato, M., Hubbard, B. & Sievers, A. [2006] “Nonlinear energy localization and its manipulation in micro-mechanical oscillator arrays,” *Rev. Mod. Phys.* **78**, 137.
- Swanson, B., Brozik, J., Love, S., Strouse, G., Shreve, A., Bishop, A., Wang, W.-Z. & Salkola, M. [1999] “Observation of intrinsically localized modes in a discrete low-dimensional material,” *Phys. Rev. Lett.* **22**, 3288.