

1 **Model selection and averaging in the estimation of population parameters of *Bemisia tabaci***
2 **(Gennadius) from stage frequency data in sweet pepper plants**

3 **Authors:** J.E. González-Zamora, R. Moreno

4 **Affiliation:** Department of Ciencias Agroforestales, University of Seville (Spain)

5 Ctra. de Utrera, km 1 E- 41013-Seville (Spain)

6 **Corresponding author:** J. E. González-Zamora

7 zamora@us.es Phone number: (+34) 954 486 459 Fax number: (+34) 954 486 436

8
9 **Abstract**

10 *Bemisia tabaci* is a significant pest for many crops, but there are few population studies of this
11 insect on sweet pepper (*Capsicum annuum*). In this study, stage frequency data were generated with
12 *B. tabaci* in sweet pepper plants in various situations, and the Bellows and Birley method was used
13 to obtain population parameters from the data. The Akaike Information Criterion (AIC) was used to
14 select the best option of the Bellows and Birley method and, in some cases, to estimate the
15 parameters of the population using model averaging. The ratios estimated/observed for each
16 population parameter were calculated to assess bias and were used to correct the estimations if the
17 ratios were different from 1. The effects of different factors on the estimations of population
18 parameters were analysed. The total duration of development was affected by the experimental
19 conditions (laboratory vs. greenhouse) and temperature, but it had the highest precision. The final
20 survival rate was affected by temperature, and the estimation of individuals entering each stage was
21 affected only by the options included in the Bellows and Birley method. AIC helped to detect
22 differences in the daily survival rate among the different experiments between N1 (first instar)
23 (range 0.842-0.923), and the egg (range 0.989-1.0) and N4 (fourth instar) (0.990). The methodology
24 used can be employed in field population studies. For example, the final survival rate in the

25 greenhouse experiments varied between 0.624 and 0.097, depending on if the parasitoids were
26 present or not, and the total development varied between 420.6 and 440.7 degree-days.

27

28 **Keywords:** Akaike Information Criterion, Bellows and Birley method, *Bemisia tabaci*, sweet
29 pepper, model selection, model averaging

30

31

32 **Published in: J Pest Sci (2011) 84:165–177**

33 **DOI 10.1007/s10340-010-0337-y**

34

Final Version

35 **Introduction**

36 The whitefly, *Bemisia tabaci* (Gennadius), is a significant pest in many crops around the world
37 (Oliveira et al. 2001). In regions where there is a great density of vegetable crops cultivated in
38 greenhouses, as is found in the southeast of Spain, the whitefly becomes a serious threat in terms of
39 the increase of its population and the potential transmission of several viral diseases (Navas-Castillo
40 et al. 2000; Segundo et al. 2004; Ruiz et al. 2006). Crops such as sweet peppers, tomatoes, melons,
41 cucumbers, green beans and others may be seriously affected by this pest. Many studies have
42 focused on the biology of this species in different crops and on the analysis of life tables to
43 investigate different parameters of the population or the key factors that regulate its population (Von
44 Arx et al. 1983; Horowitz et al. 1984; Baumgartner et al. 1986; Baumgartner and Yano 1990;
45 Naranjo and Ellsworth 2005; Asiimwe et al. 2007). Several of these studies have compared different
46 models. The biology of *B. tabaci* has been studied in sweet peppers (*Capsicum annuum*) under
47 controlled (laboratory) conditions (González-Zamora and Gallardo 1999; Muñiz 2000; Muñiz et al.
48 2002), but no studies have been presented on the biology of this species with sweet peppers under
49 field conditions.

50
51 Stage frequency data are analysed in different ways to obtain information on populations. One way
52 is to use a model or models, which can be as simple or as complicated as needed under the
53 circumstances (for a review see Manly 1990; Southwood and Henderson 2000). If different models
54 are used to analyse the data, the results must be compared to select the most suitable one. Different
55 biological conclusions may be drawn from the data depending upon the final model selected, and
56 therefore, it is important to have a method that selects the best model and measures the strength of
57 the evidence for each one. The Akaike Information Criterion (AIC) is widely used in biological
58 studies to select the best model due to the advantages it has over other criteria, and it is used to
59 estimate parameters by model averaging (Burham and Anderson 2002; Johnson and Omland 2004;

60 Posada and Buckley 2004). In the field of entomology, the application of AIC or other information
61 criteria is generally used to select models that help explain different aspects of the biology and
62 behaviour of insects and to select models that can be used in the field of crop protection (Luh and
63 Croft 1999; Hansen et al. 2001; Hemerik and van der Hoeven 2003; Umble and Fisher 2003; Sileshi
64 2006; Takeuchi 2006; Saint-Germain et al. 2007). A study undertaken by Sileshi (2006) is one of
65 few examples of the use of AIC for insect count data or applications for life table analyses. Model
66 averaging is applied when none of the set of models is clearly the best, and several can be used. In
67 such case, the parameters of interest are estimated based on the relative importance (or weight) of
68 the models. To date, no examples have been found on the use of model averaging to estimate
69 population parameters.

70

71 This work had different objectives corresponding to the information that can be obtained from stage
72 frequency data of *B. tabaci* in sweet peppers, both from the laboratory and field. We studied the bias
73 generated after using a model (in this case, the Bellows and Birley method), comparing the
74 observed and estimated parameters, and how different factors can influence this bias. The other
75 objective of this study was to demonstrate the application of model selection and averaging with the
76 AIC to accurately estimate population parameters. The Bellows and Birley method produces
77 different parameters from stage frequency data and, with the help of the AIC, it can be of great
78 interest in population studies due the information generated, such as, for example, survival rates,
79 development time, number of entering stages, and others. Finally, population parameters from field
80 studies are presented to show the potential of this methodology.

81

82 **Materials and Methods**

83

84 Experimental conditions

85 The study was carried out in the facilities of the I.F.A.P.A. (Instituto para la Formación Agraria y
86 Pesquera de Andalucía) of "La Mojonera-La Cañada" (36°47'18.57" N and 2°42'13.87" W) in
87 Almería (southeast Spain). The experiments were conducted under laboratory conditions with
88 potted plants and in a plastic greenhouse using sweet pepper plants (*Capsicum annuum*) cv.
89 "Espartaco". The pots had a diameter of 13.8 cm and a volume of 1.2 L. The substratum was
90 coconut fibre, and the plants were periodically fertilised with Multi Poli-Feed[®] (Haifa Chemical).

91

92 A 600-m² plastic greenhouse was used for the greenhouse conditions. The sweet pepper plants were
93 transplanted to the ground in August 1995. The normal agricultural practice in the area for this crop
94 was followed during the period of cultivation, with the spraying of pesticides on the upper part of
95 the plants to control certain diseases and pests, such as powdery mildew (*Leveillula taurica* (Lev.)
96 Arnaud (Perisporales: Erysiphaceae)) with dinocap and bupirimate; beet armyworm (*Spodoptera*
97 *exigua* (Hübner) (Lepidoptera: Noctuidae)) with *Bacillus thuringiensis* and trichlorfon mixed with
98 wheat bran; and broad mite (*Poliphagotarsonemus latus* (Banks) (Acari: Tarsonemidae)) with
99 bromopropylate and avermectin. Care was taken to avoid products harmful to whiteflies and their
100 natural enemies.

101

102 The adults of *B. tabaci* that were used to lay the eggs were collected from a different greenhouse
103 planted with peppers (cv. "Espartaco"), where a colony of *B. tabaci* was constantly reared.

104

105 Three trials were carried out in different situations to generate stage frequency data that could be
106 used for the posterior analysis of model selection and model averaging to obtain population
107 parameters. The experiments were carried out under the following two experimental conditions:
108 controlled temperature (laboratory conditions in Trials 1 and 2) vs. uncontrolled temperature (field

109 conditions in Trial 3). There were also two scales of observation: individual counts (in Trial 1) vs.
110 grouped counts (in Trials 2 and 3). The three trials were as follows:

111

112 1) Trial 1 (individual counts and controlled temperature). Assays were carried out at 20 ± 1 , 25 ± 1
113 and 30 ± 1 °C in a growth chamber (KOXKA model MEC-185/F) with 4,000 lux, a 16:8
114 photoperiod (light:dark) and a relative humidity of $75 \pm 10\%$; and in a breeding chamber at $25 \pm$
115 2°C with 6,000 lux, a 16:8 photoperiod (light:dark) and a relative humidity of $65 \pm 10\%$. Each of
116 the studies consisted of one or two potted plants with six to eight leaves each. The plants were
117 infested with high numbers of *B. tabaci* adults. The adults were confined to one or two leaves per
118 plant by means of a cloth bag for 24 h at the different temperatures defined above. After this time,
119 the adults were eliminated, and the eggs were counted. This time point was considered the initial
120 moment, or zero time point, for the study of development. The eggs were observed and counted
121 daily. When nymphs of the first instar emerged, we waited until they fixed on the leaf, and then
122 their positions were marked with a soft marker (Lumocolor[®], Staedtler, Germany). Daily counts of
123 each individual took place until the whitefly adults emerged.

124

125 2) Trial 2 (grouped counts and controlled temperature). Assays were carried out at 20 ± 1 and 30 ± 1
126 °C in a growth chamber (KOXKA model MEC-185/F) with 4,000 lux, a 16:8 photoperiod
127 (light:dark) and a relative humidity of $75 \pm 10\%$; and in a breeding chamber at $25 \pm 2^\circ\text{C}$ with 6,000
128 lux, a 16:8 photoperiod (light:dark) and a relative humidity of $65 \pm 10\%$. Two replicates were done
129 at each temperature. The plant infestation was initiated following the same procedure as described
130 in Trial 1. Once the adults were eliminated, the leaves where the adults had been confined were
131 observed daily. The eggs and individuals that emerged were counted and grouped according to their
132 developmental stage, but they were not marked. The daily counts took place until the adults
133 emerged.

134

135 3) Trial 3 (grouped counts with uncontrolled temperature in a plastic greenhouse). Three
 136 experiments were carried out at several time periods throughout the year, as follows: September 19,
 137 1995 to October 23, 1995 (experiment 1); December 1, 1995 to February 26, 1996 (experiment 2);
 138 and March 6, 1996 to April 30, 1996 (experiment 3). In each experiment, six to ten plants were
 139 selected, and a leaf from each plant was isolated with a cloth bag. Large numbers of adult *B. tabaci*
 140 were introduced into each bag and left for 24 h. The eggs that were laid were counted, and this was
 141 considered the starting point for the developmental study. The population was counted daily except
 142 in experiment 1, where it was counted every two days until all of the adults had emerged.
 143 Individuals were counted and grouped according to their developmental stage. The temperature and
 144 relative humidity were registered daily during the experiments, with mean temperatures (and
 145 ranges) of 24.9 °C (12 - 37 °C), 15.9 °C (4 - 30 °C) and 18.9 °C (8 - 33 °C) for experiments 1, 2, and
 146 3, respectively. Time was measured using degree days (DDs). The DDs were calculated using the
 147 maximal and minimal temperatures of two periods in the day (from 00.00 to 12.00 h and from 12.00
 148 to 24.00 h), and 10 °C was the minimum development threshold temperature (Zalom et al. 1985).
 149 The following equation was used to calculate DD:

$$150 \quad DD = \frac{\frac{(T_{\max} + T_{\min})_{0-12}}{2} + \frac{(T_{\max} + T_{\min})_{12-24}}{2}}{2} - 10,$$

151 where T is temperature.

152

153 In Trials 1 and 2, the different developmental stages, from egg to fourth instar (N4), were observed
 154 in the laboratory with a stereobinocular microscope (9x and 45x magnification) by turning the leaf
 155 under the microscope. The developmental stages were distinguished as a function of size. The first
 156 instar (N1) was the smallest, and the fourth (N4) was the largest. In Trial 3, the different
 157 developmental stages from egg to N4 were distinguished using a field lens (8x magnification), and

158 they were separated according to their relative size. In the first greenhouse experiment, the nymphs
159 of the first, second and third instars were counted together.

160

161 In the individual counts (Trial 1), the fate of each individual was recorded. Thus, it was possible to
162 calculate different parameters, such as the number of individuals entering each instar, the daily
163 survival rate of each instar, the survival rate of each developmental stage, the duration of each
164 instar, the final survival rate of the population and the total development period (González-Zamora
165 and Gallardo 1996). In contrast, in the grouped counts (Trials 2 and 3), only the number of
166 individuals found at each instar was recorded daily. In the grouped counts, the initial number of
167 eggs and the final number of adults that emerged from the empty pupal cases was known. With
168 these data, most of the previous parameters could not be calculated except for the final survival rate
169 of the population and the total development period. Therefore, the other parameters had to be
170 estimated with the help of a model. With individual counts, it was possible to compare the observed
171 values of all parameters with their estimations and to establish the bias and validity of the model
172 used. With Trials 2 and 3, the bias was identified using fewer parameters, such as individuals
173 entering the egg and adult stage, the final survival rate and the total development.

174

175 Model Fitting

176 The P1f software package (Manly 1994) was used to analyse the life tables, and it was specifically
177 designed to analyse data from stage-structured populations with different models. In model
178 selection, it is important to have a group of models that are relevant to the data and to the objectives
179 of the analysis, representing a plausible research hypothesis (Burham and Anderson 2002). The
180 Bellows and Birley method (Bellows and Birley 1981) is the most flexible method because it allows
181 estimation of the duration of each stage, the unit time survival rate, the final survival rate, and the

182 numbers entering each stage. This method also allows for different assumptions when different
 183 survival parameters for each stage or time of entry in stage 1 are considered (Manly 1990).

184

185 All of the experiments were performed using a single cohort, and therefore, the initial number
 186 entering stage 1 was known, and no entering distribution was necessary. The Weibull distribution
 187 was also used to model the distribution function of each stage. The Weibull distribution function is
 188 as follows:

$$189 \quad f(t) = \frac{\alpha}{\lambda} \left(\frac{t}{\lambda} \right)^{\alpha-1} e^{-\left(\frac{t}{\lambda} \right)^\alpha} \quad (3)$$

190 where alpha (α) is the shape parameter, and lambda (λ) determines the spread of the curve along the
 191 X axis. Lambda is also an estimate of the mean duration of each stage.

192

193 The P1f program allows for different combinations of the unit time survival rate (needed in the
 194 Bellows and Birley method) and the shape parameter (α) of the Weibull distribution, which may be
 195 different for each stage in the Bellows and Birley method. The combinations of these two
 196 parameters produce the following four options in the program: 1) different survival and shape
 197 parameters for each stage; 2) the same survival parameters but different shape parameters for each
 198 stage; 3) different survival parameters but the same shape parameters for the stages; and 4) the same
 199 survival parameters and shape parameters for all stages. These four options of the Bellows and
 200 Birley method produced different estimations of the same population parameters along with fitting
 201 of the model to the data as expressed in the log *likelihood* for each option. For this reason, the four
 202 options were considered in this study as models to be selected with a given criterion.

203

204 The output estimates produced by the Bellows and Birley method were as follows: a) individuals
 205 entering into each stage (egg, N1, N2, N3, N4 and adult); b) the stage-specific survival rate (SSSR),

206 which is the survival of a given stage; and c) duration, which is the developmental time for each
207 stage considered. The P1f program generated the standard deviation for the duration of each stage,
208 but not for the individuals entering the stage or for the SSSR. Two other estimates were calculated
209 using the results of the program as follows: a) the SR_f , which is the final survival rate from egg to
210 adult; and b) the $Duration_t$, which is the duration of the total developmental period, from egg to
211 adult. However, neither of these two last estimates had an associated standard deviation. The shape
212 parameter (α) of the Weibull distribution and the unit time (daily) survival rate (ϕ) were also
213 estimated in the four options of the Bellows and Birley method. Both of these parameters could be
214 considered equal or different for each stage, and both of them had a standard deviation produced by
215 the P1f program.

216

217 Statistical analyses

218 The output estimates produced by the Bellows and Birley method, such as the individuals entering
219 each stage, the SSSR, the duration of each immature stage, the final survival rate (SR_f) and the total
220 duration ($Duration_t$), were compared with the observed values of the same parameters, which were
221 obtained mainly from Trial 1, but also from Trials 2 and 3, to obtain the estimated/observed ratios.
222 The ratios were used to identify the bias of the estimates and to determine if the bias was affected
223 by different factors.

224

225 The previous parameters, expressed by relative values in the ratios, were considered as variables
226 that could be affected by different factors. These factors and their levels were as follows: a) *Scale of*
227 *observation*, with two levels, including individual counts (with data coming from the experiments
228 of Trial 1) and grouped counts (with data coming from the experiments of Trial 2); b) *Experimental*
229 *conditions*, with two levels, including controlled temperatures (with data coming from the
230 experiments of Trial 2) and uncontrolled temperatures (with data coming from the experiments of

231 Trial 3); c) *Temperatures*, with the three temperatures used in controlled temperatures (with data
232 coming from the experiments of Trials 1 and 2); d) *Options*, with the four options of the Bellows
233 and Birley method supported by the P1f program (with the available data from Trials 1, 2 and 3).

234

235 The ratios were not transformed in any way, and they were first analysed to test the homogeneity of
236 the variances within factors using Cochran's C contrast, Bartlett's contrast and Levene's tests. If the
237 probability associated with any of them was less than 0.05, the Kruskal-Wallis test was used to
238 analyse the data. In contrast, if the p -value was greater than 0.05 in all of them, a one-way ANOVA
239 was used to analyse the data. If the p -value of the Kruskal-Wallis statistic was less than 0.05, the
240 means of the different levels within the factor were separated using Mann-Whitney's U test (Steel
241 and Torrie 1988). If the p -value of the ANOVA test was less than 0.05, the means of the different
242 levels within the factor were separated using Tukey's honestly significant difference (HSD) test at p
243 = 0.05. The ratios were then tested to determine whether they differed from 1 using the contrast
244 hypothesis test, with $p = 0.05$. All analyses were performed using the Statgraphics package
245 (Statistical Graphics 2000).

246

247 Model selection

248 There are different ways of comparing models to select the most appropriate one. This study used
249 the AIC, which is a powerful method for model selection and the inference of ecological data
250 (Burham and Anderson 2002). With AIC, the goal is to select the model with the least number of
251 parameters that represents the data adequately (i.e., the principle of parsimony) (Franklin et al.
252 2001; Mazerolle 2004). The AIC was used to select the best option of the Bellows and Birley
253 method, and it is defined as follows:

254

$$AIC = -2 \ln(L) + 2k$$

255 In this equation, K is the number of estimated parameters included in each model (or the options of
256 the Bellows and Birley method in this study). The log-likelihood of the model given the data are
257 readily available in the statistical output, and reflects the overall fit of the model. When there was a
258 comparison of models, the model with the smallest AIC was selected. With count data, as was the
259 case here, it is normal to find overdispersion. Therefore, the AIC was modified to obtain QAIC_c.
260 (Burham and Anderson 2002).

261

262 Two measures, delta AIC (Δ) and Akaike weights (w), associated with the AIC and equally with
263 QAIC_c were used to compare models (Burham and Anderson 2002). The delta AIC is a measure of
264 each model relative to the best model. As a rule of thumb, when Δ_i is less than two, it suggests
265 substantial evidence in support of the model. When the values are between three and seven, it
266 indicates that the model has considerably less support. When Δ_i is greater than ten, it indicates that
267 the model is unlikely. Akaike weights provide another measure of the strength of evidence for each
268 model, and they represent the ratio of delta AIC (Δ_i) values for each model relative to the whole set
269 of candidate models (the four options of the Bellows and Birley model). Akaike weights also
270 indicate the probability that the model is the best among the set of candidate models.

271

272 Selecting a model from a set of candidate models may produce a new problem. When no single
273 model is clearly the best, predictions cannot be based on the model ranked in first place. In some
274 cases, the best model may have competitors for the top rank (e.g., when $\Delta_i < 2$). A solution to this
275 problem is to base the inference on the entire set of models, an approach called “multimodel
276 inference” or “model averaging” (Burham and Anderson 2002; Johnson and Omland 2004; Posada
277 and Buckley 2004). When this situation happened in the present study, a weighted average of the
278 estimates was computed using the Akaike weights.

279

280 To conduct model averaging, the estimate of the parameter for each model was weighted by the
 281 Akaike weights as follows:

$$282 \quad \hat{\theta} = \sum_{i=1}^R w_i \hat{\theta}_i$$

283 where $\hat{\theta}_i$ denotes the estimate for model i . Similarly, the precision (as standard error, SE) of the
 284 model averaged estimate may also be computed and is called the unconditional SE (Burham and
 285 Anderson 2002). In many cases, model averaging reduces bias and increases precision, which are
 286 desirable properties (Burham and Anderson 2002). Once the model averaged estimates and SE were
 287 calculated, confidence intervals were used to assess the magnitude of the effect. After using AIC, or
 288 model averaging if needed, the observed parameters and their final estimates were used to calculate
 289 the coefficients of determination R^2 .

290

291 Results

292

293 Differences among the observed parameters and their estimates from the Bellows and Birley
 294 method in its different options were observed. The discrepancies were measured by calculating the
 295 estimated/observed ratios (Table 1), which were used to identify the bias of the method in the
 296 different parameters and to assess the effect of the factors on estimations of the same parameters.
 297 The ratios were less than one in most cases, but some were close to one. Table 2 shows the statistics
 298 obtained for the analyses and their significance. There were no differences in the ratios of each
 299 variable studied within the *scale of observation* (individual counts vs. grouped counts), although the
 300 *experimental conditions* (controlled temperature vs. uncontrolled temperature) showed a significant
 301 difference in the ratios only in the total duration of development. The *temperature* factor displayed
 302 differences only in the final survival rate and total duration. Finally, the *option* factor, which must
 303 be considered as appertaining to the Bellows and Birley method, demonstrated significant

304 differences within each factor only in the entering individuals. The estimated/observed ratios were
305 used to correct the estimated parameters in each situation, but only when the ratio was significantly
306 different from one, by dividing the estimated parameter by the ratio value. This correction was used
307 to obtain the final values of each parameter, but it was not used to select the best option with the
308 AIC.

309
310 The different parameters of the AIC analysis for each Trial (Table 3) indicated that the data were
311 overdispersed (c between 1 and 5; Burham and Anderson 2002). Each trial comprised its own set of
312 data, and therefore, the results are shown separately. The four options supported by the P1f program
313 with the Bellows and Birley method were compared with the delta AIC (Δ) and the Akaike weights
314 (w). The first option (i.e., different daily survival rates for the stages) was selected as the best with
315 the experiments carried out at the lowest temperature under controlled conditions (20°C in both
316 individual and grouped counts, Trials 1 and 2, respectively). In contrast, the second option (i.e., the
317 same daily survival rate for all stages) was selected as the best at the intermediate temperature
318 (25°C in both individual and grouped counts, Trials 1 and 2, respectively). At the higher temperature
319 (30°C in both individual and grouped counts, Trials 1 and 2, respectively), the selection was not as
320 clear, but the options with the same survival rate for the stages (options 2 and 4) generally had
321 higher weights. Similar selections of options occurred in the experiments carried out in the
322 greenhouse (Trial 3). In the experiment carried out in winter (experiment 2 with a mean temperature
323 of 15.9°C), the first option (i.e., different daily survival rates for the stages) was clearly selected. In
324 experiment 3 (mean temperature of 18.9 °C and at the beginning of spring), the third option (i.e.,
325 different daily survival rate for each stage) was the most important. In experiment 1, under
326 greenhouse conditions (warm temperatures with a mean of 24.9°C and at the beginning of autumn),
327 the second and fourth options (both with the same daily survival rate for each stage) were more
328 important.

329

330 Finally, the final population parameters for each experiment (with their own standard errors), were
331 obtained using multimodel inference when necessary, according to the w factor of Table 3. The
332 coefficients of determination R^2 between the observed and estimated parameters (corrected with the
333 ratio if necessary) were calculated as follows: 0.909 (41), 0.925 (20), 0.988 (20), 0.985 (12) and
334 0.999 (13) for the individuals entering the different stages, the stage specific survival rate (SSSR),
335 the duration of development in the different stages, the final survival rate, and the duration of total
336 development period, respectively (the numbers between brackets are the number of points used to
337 calculate the coefficient of determination in each case).

338

339 In 6 out of 13 experiments, the first and/or third option of the Bellows and Birley method (i.e., in
340 which the daily survival rates were considered differently for each instar) produced higher Akaike
341 weights (Table 4). Three of the experiments demonstrated a clear difference with non-overlapping
342 confidence intervals between the daily survival rates of the egg and N1 stages, and in one
343 experiment between the N1 and N4 stages (Table 4). In several cases, there was only a light overlap
344 in the confidence intervals of the daily survival rate of the N4 with the egg stage. Also, the estimates
345 of the daily survival rate with the different options in the individual counts (Trial 1) were similar to
346 the observed daily survival rates, which were included in the 95% confidence intervals of the
347 estimates (Table 4). The observed values of daily survival rate showed statistical differences in the
348 stage and temperature factors and their interaction ($F_{4,5} = 68.5$, $p < 0.001$; $F_{2,5} = 113.8$, $p < 0.001$;
349 and $F_{8,5} = 16.6$, $p = 0.003$, respectively). The differences between stages within the temperature
350 were consistent with the results from the estimations and their confidence intervals at 20°C (N1 was
351 different from egg and N4; Table 4). At 30°C and 25°C, however, differences were found between
352 the egg stage and the remaining instar stages (Table 4 and data not shown).

353

354 The final estimates obtained for the greenhouse experiments are presented in Table 5, and they
355 show the type of output obtained using the AIC and the multimodel inference. From the results, the
356 similarity of the total duration among the three experiments (ranging from 420.6 to 440.7 DD) was
357 significant, whereas the final survival rates (ranging from 0.097 to 0.624) indicate that the
358 environmental conditions in the three experiments were different.

359

360 Discussion

361

362 The study presented in this paper had two objectives. The first objective was to show the potential
363 of the Bellows and Birley method to estimate parameters from stage frequency data. As in any
364 estimation, the parameters estimated may differ from the observed values of the same parameters.
365 The observed parameters are not always known. In this study, however, the observed parameters
366 were known in most of the cases, and they were used to identify the bias of the Bellows and Birley
367 method as presented by the software P1f program. The second objective was to use a procedure to
368 select a model that provides the best trade-off between bias and accuracy (i.e., AIC selected the
369 most parsimonious model from those used to adjust the data). If different models support the data
370 similarly, a multimodel inference (e.g., model averaging) can be considered to obtain a precision
371 estimator for the different parameters.

372

373 The Bellows and Birley method produced several estimates that were different from the observed
374 values, as shown by the ratios in Table 1. The ratios of the estimated/observed parameters are useful
375 for identifying the bias of the estimates and for determining if this bias is affected by the different
376 factors considered. It is of particular interest that no difference of the ratios within the *scale of*
377 *observation* (individual vs. grouped counts) was observed in the variables estimated. However,
378 there were statistical differences in the ratios within the *experimental conditions* and *temperatures*,

379 which affected variables representing general values (i.e., final survival rate and total duration of
380 development). There were differences among the experiments carried out in the greenhouses and
381 among the different temperatures in the laboratory. However, there was a remarkably high precision
382 in several of the estimated variables, such as the total duration of development, across all factors.

383

384 The *option* factor was associated with the Bellows and Birley method of the P1f program, and it had
385 a significant effect on one of the studied variables. The second option of the Bellows and Birley
386 method (i.e., same survival parameter but different shape parameter for each stage) had a
387 remarkable effect on the estimations of entering individuals, with a ratio close to 0.5 (Table 1).

388

389 With the analysis presented herein, we have a tool to help answer the question of whether there are
390 different daily survival rates at each stage. The options of the Bellows and Birley method with
391 different daily survival rates in each stage (options 1 and 3) obtained better support at lower
392 temperatures (both in controlled conditions and the plastic greenhouse), whereas the options with
393 equal survival rates (options 2 and 4) were selected at medium and higher temperatures. The factors
394 that affect mortality in each situation may be different and may change in other experiments, but
395 their identification and quantification may only be answered with an adequate sampling
396 methodology that identifies them and relates them to the results of the analysis.

397

398 The daily survival rates of the stages was of particular interest (Table 4). The observed values of the
399 daily survival rates in the experiments of Trial 1 fell within the confidence interval of their
400 estimations, reflecting their accuracy. It may be assumed that the estimations in the other
401 experiments shown in Table 4, where no observed values were generated, were equally accurate.
402 Although the daily survival rates were different among some stages within the temperatures, as
403 confirmed by the ANOVA of the observed values (even though these results must be considered

404 with caution due the low number of replications of the observed data), this was not confirmed with
405 the Bellows and Birley method in some cases (e.g., the experiment performed at 25°C, not shown in
406 Table 4, the experiment performed at 30°C in Trial 1, shown in Table 4, or one of the experiments
407 performed at 20°C in Trial 2, shown in Table 4).

408

409 For differences among stages, studies that relate this variation to environmental variables, such as
410 climate, number of predators, number of parasitoids, food availability or any other variable, may
411 help to explain these differences (Manly 1990). In experiments conducted at a controlled
412 temperature at 20°C, the temperature and possible manipulations under the stereomicroscope may
413 be considered as factors affecting the survival of N1. Other factors may have produced these
414 differences in the experiments carried out with uncontrolled temperatures (Trial 3 in the
415 greenhouse).

416

417 The egg and N1 and N4 instar stages had the greatest effect on the *B. tabaci* population in the life
418 table analysis in different locations and crops (Horowitz et al. 1984; Naranjo and Ellsworth 2005;
419 Asiiimwe et al. 2007). Predation (Naranjo and Ellsworth 2005) and parasitisation (Asiiimwe et al.
420 2007) were the principal factors responsible for decreasing populations, affecting mainly the N4
421 instar, and dislodgement was second in most of the previous studies. In general, these conclusions
422 agree with the results obtained in the present study for the greenhouse experiments (although with
423 only three experiments), where N1 and N4 had the highest mortalities, with estimated survival rates
424 of 0.758 and 0.474, respectively, in experiment 2, and 0.372 and 0.515, respectively, in experiment
425 3 (Table 5). In greenhouses, the N1 instar is the most exposed to environmental factors, such as
426 peaks of high temperature and low humidity typical of greenhouses in southeast Spain and low
427 temperatures in some periods of the year. The N1 instar is also subject to other factors related to the
428 plant itself, such as nutritional factors and cuticle thickness (Byrne and Bellows 1991). Predation by

429 lacewings or beetles on whiteflies was not observed in Trial 3 (greenhouses). However,
430 *Eretmocerus mundus* Mercet adults, which are the main parasitoid of *B. tabaci* in greenhouses in
431 Almería (Rodríguez-Rodríguez et al. 1994), and subsequently, parasitised N4, were detected. The
432 high mortality found in N4 (mainly from parasitism but also from direct feeding) and the low final
433 survival rates recorded in experiments 2 and 3 ($SR_f = 0.143$ and $SR_f = 0.097$, respectively) (Table 5)
434 were due to *E. mundus* in contrast to the values obtained in the first experiment in which no *E.*
435 *mundus* was detected.

436
437 The level of parasitism due to *E. mundus* in the N4 instar may be high (González-Zamora et al.
438 1996; Téllez et al. 2003; Stansly et al. 2005). Predation by *E. mundus* adults on different instars of
439 the whitefly *B. tabaci* is well known and has been evaluated (Gerling and Fried 2000; Urbaneja et
440 al. 2007), and is considered an important factor in population regulation (Téllez et al. 2003; Zang
441 and Liu 2008).

442
443 The methodology used in this work allowed the bias of the Bellows and Birley method to be
444 identified in different experimental situations (laboratory vs. greenhouse and individual vs. grouped
445 counts) and with the use of the P1f program. The final estimates obtained from the best option of
446 the Bellows and Birley method selected with the AIC or with model averaging, when it was needed,
447 include the correction with the estimated/observed ratio when the ratios were different from one.
448 The estimations may be used to analyse life tables to, for example, study the key factors that affect
449 the survival of a population with a high enough number of generations (Southwood and Henderson
450 2000). However, this was not the objective of the present work. The final estimates for all of the
451 experiments were compared with the observed values, obtaining, in general, a good agreement
452 between them, as reflected in the coefficient of determination R^2 . It is especially remarkable that the
453 total duration of development had an $R^2 = 0.999$, which reflects the robustness of its estimation in

454 both experimental conditions (Table 1). The duration of development of each stage and the final
455 survival rates also had high values of R^2 (0.988 and 0.985, respectively). The high similarity
456 between the observed and final estimated parameters with the field experiments in greenhouses is
457 shown in Table 5.

458

459 The final estimates for the experiments carried out in the greenhouses (Table 5) are of particular
460 interest for their implications in population studies in field conditions. The final estimation of the
461 total duration of development may be compared with other studies, such as those carried out in
462 cotton, to study the developmental time of *B. tabaci* (Zalom et al. 1985; Zalom and Natwick 1987).
463 The mean generation time of *B. tabaci* in cotton was found to be between 316.0 DD and 369.5 DD,
464 which differs from the values obtained for sweet peppers in the present study. The duration of each
465 instar also indicates the difference among instars. The development times of the egg and N4 stages
466 were longer than the other instars, in agreement with results from laboratory studies on sweet
467 peppers and other crops (González-Zamora and Gallardo 1996, Muñiz 2000).

468

469 In conclusion, the methodology used in this paper allowed the bias of the estimations obtained from
470 the model (the Bellows and Birley method) to be identified. This methodology also permitted the
471 selection of the best model from a set of models (applying model averaging when necessary) to
472 analyse stage frequency data in life tables to shed light on important aspects of a population, which
473 was presented in this study for the whitefly *B. tabaci* in sweet pepper plants.

474

475

476 **Acknowledgements**

477

478 The first author was granted a postdoctoral scholarship from the Junta de Andalucía (Spain) at the
479 I.F.A.P.A. (Instituto para la Formación Agraria y Pesquera de Andalucía) centre “La Mojonera-La
480 Cañada” (Almería, Spain), within the research project “Lucha integrada en cucurbitáceas”. The
481 O.E.C.D. (Organisation for Economic Co-operation and Development) supported the first author
482 with a grant to visit Prof. B.F.J. Manly at Otago University (Dunedin, New Zealand). The authors
483 would like to thank Jose Miguel Gallardo for his help in collecting data.

484

485

486

Final Version

487

References

- 488 Asiimwe P, Ecaat JS, Otim M, Gerling D, Kyamanywa S, Legg JP (2007) Life-table analysis of
489 mortality factors affecting populations of *Bemisia tabaci* on cassava in Uganda. Entomol Exp
490 Appl 122:37-44
- 491 Baumgartner J, Delucchi V, Von Arx R, Rubli D (1986) Whitefly (*Bemisia tabaci* Genn., Stern.:
492 Aleyrodidae) infestation patterns as influenced by cotton weather and *Heliothis*: Hypothesis
493 testing by using simulation models. Agric Ecosystems and Environ 17:49-59
- 494 Baumgartner J, Yano E (1990) Whitefly population dynamics and modelling. In: Gerling D (ed.)
495 Whiteflies: Their Bionomics, Pest Status, and Management. Intercept Ltd., Newcastle, pp. 123-
496 146
- 497 Bellows TS, Birley MH (1981) Estimating development and mortality rates and stage recruitment
498 from insect stage-frequency data. Res Popul Eco 23:232-244
- 499 Burham KP, Anderson DR (2002) Model selection and multimodel inference: A practical
500 information-theoretic approach. Springer-Verlag, New York
- 501 Byrne DN, Bellows TS (1991) Whitefly biology. Annu Rev Entomol 36:431-457
- 502 Franklin AB, Shenk TM, Anderson DR, Burham KP (2001) Statistical model selection: An
503 alternative to null hypothesis testing. In: Shenk TM, Franklin AB (eds.) Modelling in Natural
504 Resource Management: Development, Interpretation, and Application. Island Press,
505 Washington, pp. 75-90
- 506 Gerling D, Fried PM (2000) Biological studies with *Eretmocerus mundus* Mercet (Hymenoptera:
507 Aphelinidae) in Israel. OILB/SROP Bull 23:117-123

- 508 González-Zamora JE, Moreno Vázquez R, Rodríguez Rodríguez MD, Rodríguez Rodríguez MP,
509 Mirasol Carmona E, Lastres Garcia Testón J, Manzanares Ruiz C (1996) Evolución del
510 parasitismo en *Bemisia tabaci* (Genn.) y *Trialeurodes vaporariorum* (West.) (Homoptera;
511 Aleyrodidae) en invernaderos de Almería. Bol San Veg -Plagas 22:373-389
- 512 González-Zamora JE, Gallardo JM (1999) Desarrollo y capacidad reproductiva de *Bemisia tabaci*
513 (Gennadius) (Homoptera; Aleyrodidae) en pimiento a tres temperaturas. Bol San Veg -Plagas
514 25:3-11
- 515 Hansen EM, Bentz BJ, Turner DL (2001) Temperature-based model for predicting univoltine brood
516 proportions in spruce beetle (Coleoptera: Scolytidae). Can Entomol 133:827-841
- 517 Hemerik L, van der Hoeven N (2003) Egg distributions of solitary parasitoids revisited. Entomol
518 Exp Appl 107:81-86
- 519 Horowitz AR, Podoler H, Gerling D (1984) Life table analysis of the tobacco whitefly *Bemisia*
520 *tabaci* (Gennadius) in cotton fields in Israel. Oecol Applic 5:221-233
- 521 Johnson JB, Omland KS (2004) Model selection in ecology and evolution. Trends Ecol Evol
522 19:101-108
- 523 Luh HK, Croft BA (1999) Classification of generalist or specialist life styles of predaceous
524 phytoseiid mites using a computer genetic algorithm, information theory, and life history traits.
525 Environ Entomol 28:915-923
- 526 Manly BFJ (1990) Staged-structured populations. sampling, analysis and simulation. Chapman and
527 Hall Ltd., London

- 528 Manly BFJ (1994) Population Analysis System: P1f, Single Species Stage Frequency Analysis. Ver.
529 3.0. Ecological Systems Analysis. Pullman, Washington. USA.
- 530 Mazerolle MJ (2004) Appendix 1. Making Sense Out of Akaike's Information Criterion (AIC): Its
531 use and Interpretation in Model Selection and Inference from Ecological Data. University of
532 Laval, Canada. <http://www.theses.ulaval.ca/2004/21842/apa.html>. Accessed 29 november 2005
- 533 Muñiz M (2000) Development of the B-biotype of *Bemisia tabaci* (Gennadius, 1889) (Homoptera:
534 Aleyrodidae) on three varieties of pepper at constant temperatures. Bol San Veg -Plagas
535 26:605-617
- 536 Muñiz M, Nombela G, Barrios L (2002) Within-plant distribution and infestation pattern of the B-
537 and Q-biotypes of the whitefly, *Bemisia tabaci*, on tomato and pepper. Entomol Exp Appl
538 104:369-373
- 539 Naranjo SE, Ellsworth PC (2005) Mortality dynamics and population regulation in *Bemisia tabaci*.
540 Entomol Exp Appl 116:93-108
- 541 Navas-Castillo J, Camero R, Bueno M, Moriones E (2000) Severe yellowing outbreaks in tomato in
542 Spain associated with infections of Tomato chlorosis virus. Plant Dis 84:835-837
- 543 Oliveira MRV, Henneberry TJ, Anderson P (2001) History, current status, and collaborative
544 research projects for *Bemisia tabaci*. Crop Prot 2005:709-723
- 545 Posada D, Buckley T (2004) Model selection and model averaging in phylogenetics: advantages of
546 Akaike information criterion and Bayesian approaches over likelihood ratio tests. Syst Biol
547 53:793-808

- 548 Rodríguez-Rodríguez MD, Moreno R, Téllez MM, Rodríguez-Rodríguez MP, Fernández-Fernández
549 R (1994) *Eretmocerus mundus* (Mercet), *Encarsia lutea* (Masi) y *Encarsia transvena*
550 (Timberlake) (Hym., Aphelinidae) parasitoides de *Bemisia tabaci* (Homoptera, Aleyrodidae) en
551 los cultivos hortícolas protegidos almerienses. Bol San Veg -Plagas 20:695-702
- 552 Ruiz L, Janssen D, Martín G, Velasco L, Segundo E, Cuadrado IM (2006) Analysis of the temporal
553 and spatial disease progress of *Bemisia tabaci*-transmitted Cucurbit yellow stunting disorder
554 virus and Cucumber vein yellowing virus in cucumber. Plant Pathol 55:264-275
- 555 Saint-Germain M, Drapeau P, Buddle CM (2007) Occurrence patterns of aspen-feeding wood-
556 borers (Coleoptera: Cerambycidae) along the wood decay gradient: active selection for specific
557 host types or neutral mechanisms? Ecol Entomol 32:712-721
- 558 Segundo E, Martín G, Cuadrado IM, Janssen D (2004) A new yellowing disease in *Phaseolus*
559 *vulgaris* associated with a whitefly-transmitted virus. Plant Pathol 53:517-517
- 560 Sileshi G (2006) Selecting the right statistical model for analysis of insect count data by using
561 information theoretic measures. Bull Entomol Res 96:479-488
- 562 Southwood TRE, Henderson PA (2000) Ecological methods. Blackwell Science Ltd, Oxford UK
- 563 Stansly PA, Calvo J, Urbaneja A (2005) Release rates for control of *Bemisia tabaci* (Homoptera:
564 Aleyrodidae) biotype "Q" with *Eretmocerus mundus* (Hymenoptera: Aphelinidae) in
565 greenhouse tomato and pepper. Biol Control 35:124-133
- 566 Statistical Graphics. 2000. Statgraphics plus for windows. 5.1.5. Rockville, MD

- 567 Steel RGD, Torrie JH (1988) Bioestadística: principios y procedimientos [translation from the 2nd
568 ed. of Principles and Procedures of Statistics. A biometrial approach]. McGraw-Hill
569 Latinoamericana, Bogotá (Colombia)
- 570 Takeuchi H (2006) Estimation of the occurrence of pecky grain damage by a sweeping census of
571 *Leptocorisa chinensis* Dallas (Hemiptera: Alydidae) in rice fields. Jap J Appl Entomol Zool
572 50:137-143
- 573 Téllez MM, Lara L, Stansly P, Urbaneja A (2003) *Eretmocerus mundus* (Hyt.: Aphelinidae),
574 parasitoide autóctono de *Bemisia tabaci* (Hom.: Aleyrodidae): primeros resultados de eficacia
575 en judía. Bol San Veg -Plagas 29:511-521
- 576 Umble JR, Fisher JR (2003) Influence of below-ground feeding by garden symphylans
577 (Cephalostigmata: Scutigrellidae) on plant health. Environ Entomol 32:1251-1261
- 578 Urbaneja A, Sanchez E, Stansly PA (2007) Life history of *Eretmocerus mundus*, a parasitoid of
579 *Bemisia tabaci*, on tomato and sweet pepper. Biocontrol 52:25-39
- 580 Von Arx R, Baumgartner J, Delucchi V (1983) A model to simulate the population dynamics of
581 *Bemisia tabaci* Genn. (Stern. Aleyrodidae) on cotton in the Sudan Gezira. Z Angew Entomol
582 96:341-363
- 583 Zalom FG, Natwick ET, Toscano NC (1985) Temperature regulation of *Bemisia tabaci*
584 (Homoptera: Aleyrodidae) populations in Imperial Valley cotton. J Econ Entomol 78:61-64
- 585 Zalom FG, Natwick ET (1987) Developmental time of sweetpotato whitefly (Homoptera:
586 Aleyrodidae) in small field cages on cotton plants. Fla Entomol 70:427-431

587 Zang LS, Liu TX (2008) Host-feeding of three parasitoid species on *Bemisia tabaci* biotype B and
588 implications for whitefly biological control. *Entomol Exp Appl* 127:55-63

Final Version

Table 1 Mean value \pm the standard error of the estimated/observed ratios of the different population parameters with the factors used

FACTORS	POPULATION PARAMETERS				
	Individuals Entering	SSSR	Duration	SR _f	Duration _t
Scale of observation ¹					
Individual	0.913 \pm 0.026	-	-	1.010 \pm 0.030	0.878 \pm 0.008
Group	0.902 \pm 0.034	-	-	1.037 \pm 0.044	0.891 \pm 0.005
Experimental conditions ²					
Controlled temperature	0.902 \pm 0.034	-	-	1.037 \pm 0.044	0.891 \pm 0.005 a
Greenhouse	1.028 \pm 0.101	-	-	1.253 \pm 0.191	0.931 \pm 0.006 b
Temperature ³					
20° C	0.914 \pm 0.047	1.020 \pm 0.075	0.891 \pm 0.006	0.971 \pm 0.067 a	0.923 \pm 0.001 b
25° C	0.877 \pm 0.027	1.015 \pm 0.027	0.856 \pm 0.015	0.944 \pm 0.023 a	0.866 \pm 0.003 a
30° C	0.955 \pm 0.036	1.039 \pm 0.043	0.825 \pm 0.028	1.189 \pm 0.031 b	0.875 \pm 0.005 a
Option ⁴					
Opt 1	1.024 \pm 0.026 b	1.033 \pm 0.053	0.869 \pm 0.019	1.085 \pm 0.073	0.900 \pm 0.009
Opt 2	0.548 \pm 0.024 a	1.019 \pm 0.054	0.850 \pm 0.023	1.090 \pm 0.104	0.893 \pm 0.010
Opt 3	1.039 \pm 0.027 b	1.025 \pm 0.045	0.857 \pm 0.025	1.046 \pm 0.079	0.897 \pm 0.008
Opt 4	1.084 \pm 0.032 b	1.013 \pm 0.053	0.852 \pm 0.023	1.033 \pm 0.073	0.894 \pm 0.009

Means in the same column within a factor followed by different letter differ significantly (Tukey's honestly significant difference test, $p = 0.05$).

¹ In controlled conditions (laboratory), with Trials 1 and 2.

² With grouped counts, Trials 2 and 3.

³ With the three temperatures used in the laboratory trials, Trials 1 and 2.

⁴ With the four options of the Bellows and Birley method supported in the P1f software package, using Trials 1, 2 and 3.

Table 2 Statistical analyses of the population parameters studied with the factors used and their significance

FACTORS	POPULATION PARAMETERS									
	Individuals entering		SSSR		Duration		SR _f		Duration _t	
	<i>P</i>	<i>P</i>	<i>P</i>	<i>P</i>	<i>P</i>	<i>P</i>	<i>P</i>	<i>P</i>	<i>P</i>	
Scale of observation	F ^a =0.05	0.815	-	-	-	-	K-W ^b =0.37	0.544	F=2.10	0.156
Experimental conditions	K-W=0.22	0.638	-	-	-	-	K-W=0.32	0.572	F=23.24	<0.001
Temperature	K-W=3.02	0.220	K-W=1.16	0.561	K-W=3.86	0.145	K-W=18.84	<0.001	K-W=25.82	<0.001
Option	F=84.69	<0.001	F=0.03	0.990	F=0.14	0.938	F=0.11	0.951	F=0.13	0.941

^a F statistic obtained by one-way ANOVA

^b Kruskal-Wallis statistic

Table 3 Values of the Akaike Information Criterion (AIC) in the different trials and options of the Bellows and Birley method

		N	K	Deviance	Residuals d.f.	c	QAICc	Δ	w
Trial 1 (Individual counts with controlled temperature)									
20 °C	Option 1	72	16	78.8	55	1.43	99.93	0.00	0.999
	Option 2	72	11	121.4	60	2.02	113.81	13.88	0.001
	Option 3	72	12	124.6	59	2.11	119.05	19.12	0.000
	Option 4	72	7	169.2	64	2.64	136.31	36.38	0.000
25°(I)	Option 1	64	16	138.4	47	2.94	93.75	11.84	0.003
	Option 2	64	11	153.2	52	2.95	81.91	0.00	0.997
	Option 3	64	12	325.3	51	6.38	143.46	61.55	0.000
	Option 4	64	7	326.0	56	5.82	129.25	47.34	0.000
25 °(II)	Option 1	57	16	132.7	40	3.32	88.93	16.18	0.000
	Option 2	57	11	139.2	45	3.09	72.74	0.00	1.000
	Option 3	57	12	267.8	44	6.09	114.84	42.10	0.000
	Option 4	57	7	265.5	49	5.42	98.94	26.19	0.000
30 °C	Option 1	57	16	13.7	40	0.34	88.93	0.00	0.737
	Option 2	57	11	20.7	45	0.46	90.99	2.06	0.263
	Option 3	57	12	52.4	44	1.19	186.61	97.68	0.000
	Option 4	57	7	59.0	49	1.20	190.64	101.71	0.000
Trial 2 (Grouped counts with controlled temperature)									
20°C(I)	Option 1	79	16	295.5	62	4.77	105.71	0.00	0.972
	Option 2	79	11	530.3	67	7.91	139.84	34.13	0.000
	Option 3	79	12	398.8	66	6.04	115.09	9.38	0.009
	Option 4	79	7	455.5	71	6.42	113.56	7.85	0.019
20°C(II)	Option 1	91	16	190.3	74	2.57	116.16	0.00	0.999
	Option 2	91	11	260.7	79	3.30	129.25	13.09	0.001
	Option 3	91	12	393.4	78	5.04	183.55	67.39	0.000
	Option 4	91	7	420.2	83	5.06	181.10	64.94	0.000
25°C(I)	Option 1	67	16	172.4	50	3.45	96.00	10.27	0.006
	Option 2	67	11	193.6	55	3.52	85.73	0.00	0.994
	Option 3	67	12	265.6	54	4.92	109.65	23.92	0.000
	Option 4	67	7	291.2	59	4.93	102.83	17.11	0.000
25°C(II)	Option 1	61	16	114.8	44	2.61	91.60	8.15	0.017
	Option 2	61	11	138.9	49	2.83	83.45	0.00	0.983
	Option 3	61	12	191.7	48	3.99	106.89	23.44	0.000
	Option 4	61	7	223.2	53	4.21	104.20	20.76	0.000
30°C(I)	Option 1	67	16	56.1	50	1.12	96.00	12.89	0.002
	Option 2	67	11	60.1	55	1.09	83.11	0.00	0.998
	Option 3	67	12	149.8	54	2.77	166.01	82.90	0.000
	Option 4	67	7	156.2	59	2.65	157.54	74.44	0.000
30°C(II)	Option 1	64	16	178.9	47	3.81	93.75	5.24	0.050
	Option 2	64	11	223.2	52	4.29	88.51	0.00	0.692
	Option 3	64	12	234.5	51	4.60	94.59	6.08	0.033
	Option 4	64	7	275.0	56	4.91	90.76	2.25	0.225
Trial 3 (Grouped counts with uncontrolled temperatures (greenhouse))									
EXP-1	Option 1	46	9	94.8	36	2.63	61.95	6.02	0.032
	Option 2	46	7	101.2	38	2.66	58.13	2.21	0.213
	Option 3	46	8	96.5	37	2.61	59.38	3.45	0.114
	Option 4	46	6	103.0	39	2.64	55.93	0.00	0.641
EXP-2	Option 1	72	16	43.2	55	0.79	99.93	0.00	1.000
	Option 2	72	11	73.7	60	1.23	122.90	22.97	0.000
	Option 3	72	12	176.7	59	2.99	256.82	156.90	0.000
	Option 4	72	7	199.4	64	3.12	271.82	171.89	0.000
EXP-3	Option 1	51	16	82.9	34	2.44	85.49	10.12	0.006
	Option 2	51	11	171.6	39	4.40	102.16	26.79	0.000
	Option 3	51	12	97.6	38	2.57	75.37	0.00	0.994
	Option 4	51	7	185.6	43	4.32	95.38	20.01	0.000

Table 4 Unit time (daily) survival rates estimated and observed, and their confidence intervals, for the experiments in which the Akaike Information Criterion selected the option with different daily survival rates for the stages

		ESTIMATED		OBSERVED	
		Survival	95% CI	Survival	95% CI ¹
Trial 1 (Individual counts with controlled temperature)					
20°C	Egg	1.000	a ² 0.978 - 1.023	0.997	0.991-1.000
	N1	0.880	b 0.829 - 0.931	0.857	0.828-0.885
	N2	0.928	0.755 - 1.101	0.962	0.944-0.976
	N3	0.932	0.767 - 1.098	0.948	0.928-0.964
	N4	0.990	a 0.928 - 1.052	0.987	0.977-0.995
	Adult	0.990	0.868 - 1.112	--	
30°C	Egg	0.983	0.914 - 1.052	0.986	0.975-0.994
	N1	0.960	0.892 - 1.029	0.941	0.920-0.959
	N2	0.963	0.767 - 1.160	0.956	0.937-0.971
	N3	0.936	0.678 - 1.195	0.941	0.920-0.959
	N4	0.941	0.862 - 1.020	0.941	0.921-0.959
	Adult	0.981	0.895 - 1.067	--	
Trial 2 (Grouped counts with controlled temperature)					
20°C(I)	Egg	0.996	0.973 - 1.019		
	N1	0.964	0.902 - 1.027		
	N2	0.952	0.800 - 1.103		
	N3	0.921	0.751 - 1.091		
	N4	0.972	0.899 - 1.044		
	Adult	0.982	(-7.620) - 9.584		
20°C(II)	Egg	0.989	a 0.972 - 1.006		
	N1	0.923	b 0.876 - 0.970		
	N2	0.999	0.870 - 1.128		
	N3	0.977	0.855 - 1.099		
	N4	0.976	0.934 - 1.018		
	Adult	1.018	(-0.404) - 2.440		
Trial 3 (Grouped counts with uncontrolled temperatures (greenhouse))					
EXP-2	Egg	0.992	a? ³ 0.973 - 1.011		
	N1	0.945	0.878 - 1.012		
	N2	0.976	0.873 - 1.078		
	N3	0.972	0.889 - 1.055		
	N4	0.953	b? 0.919 - 0.987		
	Adult	0.830	(-5.179) - 6.839		
EXP-3	Egg	0.995	a 0.972 - 1.018		
	N1	0.842	b 0.756 - 0.927		
	N2	0.890	0.678 - 1.102		
	N3	0.923	0.737 - 1.110		
	N4	0.961	0.884 - 1.038		
	Adult	1.063	0.727 - 1.400		

The AIC (and model averaging if it was needed) was used to obtain the mean (m) and standard error (s.e.) needed to calculate the 95% confidence intervals of the estimated values. The standard error was corrected with the *c* of the most complex model (option) of the Bellows and Birley method.

¹ Standard errors needed to calculate confidence intervals were obtained from pooled data.

² Different letters in each experiment indicates differences between the stages when comparing the intervals, which do not overlap.

³ The confidence intervals of egg and N4 overlap slightly in this experiment.

Table 5 Final estimates of population parameters with their standard errors obtained from the three experiments carried out in the plastic greenhouse

	Individuals entering		SSSR		Duration ^a	
	m	s.e.	m	s.e.	m	s.e.
EXP-1						
Egg					132.3	0.0
N123	145.1	14.1	0.746	0.024	178.6	37.2
N4	107.9	9.7	0.836	0.010	110.0	19.3
Adult	90.2	8.4				
	<i>(97)^b</i>		0.624^c	0.026	420.7^d	0.5
					<i>(420.2)</i>	
EXP-2						
Egg	56.8	0.0	0.901	0.000	128.7	4.9
	<i>(59)</i>					
N1	51.2	0.0	0.758	0.000	47.2	21.5
N2	38.8	0.0	0.892	0.000	43.5	4.8
N3	34.6	0.0	0.865	0.000	48.2	6.8
N4	29.9	0.0	0.474	0.000	153.1	16.5
Adult	14.2	0.0				
	<i>(8)</i>		0.143^c	0.000	420.6^d	0.0
			<i>(0.136)</i>		<i>(422.2)</i>	
EXP-3						
Egg	120.3	0.0	0.933	0.000	142.3	31.7
	<i>(126)</i>					
N1	112.3	0.0	0.372	0.001	56.9	13.4
N2	41.8	0.1	0.625	0.001	38.3	9.8
N3	26.0	0.0	0.640	0.001	50.3	12.1
N4	17.3	0.0	0.515	0.003	152.6	33.9
Adult	8.9	0.0				
	<i>(12)</i>		0.097^c	0.000	440.7^d	0.1
			<i>(0.095)</i>		<i>(440.2)</i>	

The AIC was used to select the best option of the Bellows and Birley method, and model averaging was used to obtain the parameters of the experiment 1, accordingly with the w values of Table 3. The estimated parameters have been corrected with the estimated/observed ratio. The standard error was corrected with the c of the most complex model (option) of the Bellows and Birley method.

^a Duration is expressed in degree days (DD)

^b Numbers in italics between brackets are the observed values.

^c Final survival rate (SR_f).

^d Duration of total development ($Duration_t$). It is also expressed in degree days (DD).