

ON THE PROPERTIES OF SPECIAL FUNCTIONS ON THE LINEAR-TYPE LATTICES

R. ÁLVAREZ-NODARSE AND J. L. CARDOSO

ABSTRACT. We present a general theory for studying the difference analogues of special functions of hypergeometric type on the linear-type lattices, i.e., the solutions of the second order linear difference equation of hypergeometric type on a special kind of lattices: the linear type lattices. In particular, using the integral representation of the solutions we obtain several difference-recurrence relations for such functions. Finally, applications to q -classical polynomials are given.

2000 Mathematics Subject Classification 33D15, 33D45

1. INTRODUCTION

The study of the so-called q -special functions has known an increasing interest in the last years due its connection with several problems in mathematics and mathematical-physics (see e.g. [3, 6, 8, 13, 17]). A systematic study starting from the second order linear difference equation that such functions satisfy was started by Nikiforov and Uvarov in 1983 and further developed by Atakishiyev and Suslov (for a very nice reviews see e.g. [7, 13, 16]). Of particular interest is the so-called q -classical polynomials (see e.g. [5]) introduced by Hahn in 1949 which are polynomials on the lattice q^s .

Our main aim in this paper is to present a constructive approach for generating recurrence relations and ladder-type operators for the difference analogues of special functions of hypergeometric type on the linear-type lattices. Here we will focus our attention on functions defined on the q -linear lattice (for the linear lattice $x(s) = s$ see [4] and references therein, and for the continuous case see e.g. [18]). Therefore we will complete the work started in [16] where few recurrence relations were obtained. In fact we will prove, by using the q -analogue of the technique introduced in [4] for the discrete case (uniform lattice), that the solutions (not only the polynomial ones) of the difference equation on the q -linear lattice $x(s) = c_1 q^s + c_2$ satisfy a very general recurrent-difference relation from where several well known relations (such as the three-term recurrence relation and the ladder-type relations) follow.

The structure of the paper is as follows: In section 2 the needed results and notations from the q -special function theory are introduced. In sections 3 and 4 the general theorems for obtaining recurrences relations are presented. In section 5 the special case of classical q -polynomials are considered in details and some examples are worked out in details.

2. SOME PRELIMINAR RESULTS

Here we collect the basic background [1, 13, 16] on q -hypergeometric functions needed in the rest of the work.

The hypergeometric functions on the non-uniform lattice $x(s)$ are the solutions of the second order linear difference equation of hypergeometric type on non-uniform lattices

$$\begin{aligned} \sigma(s) \frac{\Delta}{\Delta x(s - \frac{1}{2})} \left[\frac{\nabla y(s)}{\nabla x(s)} \right] + \tau(s) \frac{\Delta y(s)}{\Delta x(s)} + \lambda y(s) &= 0, \\ \sigma(s) = \tilde{\sigma}(x(s)) - \frac{1}{2} \tilde{\tau}(x(s)) \Delta x \left(s - \frac{1}{2} \right), \quad \tau(s) = \tilde{\tau}(x(s)), \end{aligned} \tag{1}$$

Key words and phrases. q -hypergeometric functions, difference equations, recurrence relations, q -polynomials.

MAY 15, 2011

where $\Delta y(s) := y(s+1) - y(s)$, $\nabla y(s) := y(s) - y(s-1)$, are the forward and backward difference operators, respectively; $\tilde{\sigma}(x(s))$ and $\tilde{\tau}(x(s))$ are polynomials in $x(s)$ of degree at most 2 and 1, respectively, and λ is a constant. Here we will deal with the linear and q -linear lattices, i.e., lattices of the form

$$x(s) = c_1 s + c_2 \quad \text{or} \quad x(s) = c_1(q)s + c_2(q), \quad (2)$$

respectively, with $c_1 \neq 0$ and $c_1(q) \neq 0$.

We will define the k -order difference derivative of a solution $y(s)$ of (1) by

$$y^{(k)}(s) := \Delta^{(k)}[y(s)] = \frac{\Delta}{\Delta x_{k-1}(s)} \frac{\Delta}{\Delta x_{k-2}(s)} \cdots \frac{\Delta}{\Delta x(s)} [y(s)],$$

where $x_\nu(s) = x(s + \frac{\nu}{2})$. It is known [13] that $y^{(k)}(s)$ also satisfy a difference equation of the same type. Moreover, for the solutions of the difference equation (1) the following theorem holds

Theorem 2.1. [12, 16] *The difference equation (1) has a particular solution of the form*

$$y_\nu(z) = \frac{C_\nu}{\rho(z)} \sum_{s=a}^{b-1} \frac{\rho_\nu(s) \nabla x_{\nu+1}(s)}{[x_\nu(s) - x_\nu(z)]^{(\nu+1)}}, \quad (3)$$

if the condition

$$\left. \frac{\sigma(s) \rho_\nu(s) \nabla x_{\nu+1}(s)}{[x_{\nu-1}(s) - x_{\nu-1}(z+1)]^{(\nu+1)}} \right|_a^b = 0,$$

is satisfied, and of the form

$$y_\nu(z) = \frac{C_\nu}{\rho(z)} \int_C \frac{\rho_\nu(s) \nabla x_{\nu+1}(s)}{[x_\nu(s) - x_\nu(z)]^{(\nu+1)}} ds, \quad (4)$$

if the condition

$$\int_C \Delta_s \frac{\sigma(s) \rho_\nu(s) \nabla x_{\nu+1}(s)}{[x_{\nu-1}(s) - x_{\nu-1}(z+1)]^{(\nu+1)}} = 0, \quad (5)$$

is satisfied. Here C is a contour in the complex plane, C_ν is a constant, $\rho(s)$ and $\rho_\nu(s)$ are the solution of the Pearson-type equations

$$\begin{aligned} \frac{\rho(s+1)}{\rho(s)} &= \frac{\sigma(s) + \tau(s) \Delta x(s - \frac{1}{2})}{\sigma(s+1)} = \frac{\phi(s)}{\sigma(s+1)}, \\ \frac{\rho_\nu(s+1)}{\rho_\nu(s)} &= \frac{\sigma(s) + \tau_\nu(s) \Delta x_\nu(s - \frac{1}{2})}{\sigma(s+1)} = \frac{\phi_\nu(s)}{\sigma(s+1)}, \end{aligned} \quad (6)$$

where

$$\tau_\nu(s) = \frac{\sigma(s+\nu) - \sigma(s) + \tau(s+\nu) \Delta x(s + \nu - \frac{1}{2})}{\Delta x_{\nu-1}(s)}, \quad (7)$$

ν is the root of the equation

$$\lambda_\nu + [\nu]_q \left\{ \alpha_q(\nu-1) \tilde{\tau}' + [\nu-1]_q \frac{\tilde{\sigma}''}{2} \right\} = 0, \quad (8)$$

and $[\nu]_q$ and $\alpha_q(\nu)$ are the q -numbers

$$[\nu]_q = \frac{q^{\nu/2} - q^{-\nu/2}}{q^{1/2} - q^{-1/2}}, \quad \alpha_q(\nu) = \frac{q^{\nu/2} + q^{-\nu/2}}{2}, \quad \forall \nu \in \mathbb{C}, \quad (9)$$

respectively. The generalized powers $[x_k(s) - x_k(z)]^{(\nu)}$ are defined by

$$[x_k(s) - x_k(z)]^{(\nu)} = (q-1)^\nu c_1^\nu q^{\nu(k-\nu+1)/2} q^{\nu z} \frac{\Gamma_q(s-z+\nu)}{\Gamma_q(s-z)}, \quad \nu \in \mathbb{R}, \quad (10)$$

for the q -linear (exponential) lattice $x(s) = c_1 q^s + c_2$ and

$$[x_k(s) - x_k(z)]^{(\nu)} = c_1^\nu \frac{\Gamma(s-z+\mu)}{\Gamma(s-z)}, \quad \nu \in \mathbb{R},$$

for the linear lattice $x(s) = c_1 s + c_2$, respectively. For the definitions of the Gamma and the q -Gamma functions see, for instance, [6].

Remark 2.2. For the special case when $\nu \in \mathbb{N}$, the generalized powers become

$$\begin{aligned} [x_k(s) - x_k(z)]^{(n)} &= (-1)^n c_1^n q^{-n(n-1)/2} q^{n(z+k/2)} (q^{s-z}; q)_n, \\ [x_k(s) - x_k(z)]^{(n)} &= c_1^n (s-z)_n, \end{aligned}$$

for q -linear and linear lattices, respectively.

We will need the following straightforward proposition which proof we omit here (see e.g. [1, 16])

Proposition 2.3. Let μ and ν be complex numbers and m and k be positive integers with $m \geq k$. For the q -linear lattice $x(s) = c_1 q^s + c_2$ we have

$$\begin{aligned} (1) \quad & \frac{[x_\mu(s) - x_\mu(z)]^{(m)}}{[x_\nu(s) - x_\nu(z)]^{(m)}} = q^{\frac{m(\mu-\nu)}{2}}, \\ (2) \quad & \frac{[x_\mu(s) - x_\mu(z)]^{(m)}}{[x_\mu(s) - x_\mu(z)]^{(k)}} = [x_\mu(s) - x_\mu(z-k)]^{(m-k)}, \\ (3) \quad & \frac{[x_\mu(s) - x_\mu(z)]^{(m)}}{[x_\nu(s) - x_\nu(z)]^{(k)}} = q^{\frac{k(\mu-\nu)}{2}} [x_\mu(s) - x_\mu(z-k)]^{(m-k)}, \\ (4) \quad & \frac{[x_\mu(s) - x_\mu(z)]^{(m+1)}}{[x_{\mu-1}(s+1) - x_{\mu-1}(z)]^{(m)}} = x_{\mu-m}(s) - x_{\mu-m}(z), \\ (5) \quad & \frac{[x_\mu(s) - x_\mu(z)]^{(m+1)}}{[x_{\mu-1}(s) - x_{\mu-1}(z)]^{(m)}} = x_{\mu-m}(s+m) - x_{\mu-m}(z). \end{aligned}$$

To obtain the result for the linear lattice one only has to put in the above formulas $q = 1$.

3. THE GENERAL RECURRENCE RELATION IN THE LINEAR-TYPE LATTICES

In this section we will obtain several recurrence relations for the solutions (3) and (4) of the difference equation (1) in the linear-type lattices (2). Since the equation (1) is linear we can restrict ourselves to the canonical cases $x(s) = q^s$ and $x(s) = s$.

Let us define the functions¹

$$\Phi_{\nu,\mu}(z) = \sum_{s=a}^{b-1} \frac{\rho_\nu(s) \nabla x_{\nu+1}(s)}{[x_\nu(s) - x_\nu(z)]^{(\mu+1)}} \quad (11)$$

and

$$\Phi_{\nu,\mu}(z) = \int_C \frac{\rho_\nu(s) \nabla x_{\nu+1}(s)}{[x_\nu(s) - x_\nu(z)]^{(\nu+1)}} ds. \quad (12)$$

Notice that the functions y_ν and the functions $\Phi_{\nu,\mu}$ are related by the formula

$$y_\nu(z) = \frac{C_\nu}{\rho(z)} \Phi_{\nu,\nu}(z). \quad (13)$$

Lemma 3.1. For the functions $\Phi_{\nu,\mu}(z)$ the following relation holds

$$\nabla_z \Phi_{\nu,\mu}(z) = [\mu + 1]_q \nabla x_{\nu-\mu}(z) \Phi_{\nu,\mu+1}(z), \quad (14)$$

where $[t]_q$ denotes the symmetric q -numbers (9).

¹Obviously the functions (3) correspond to the functions (11), whereas the functions y_ν given by (4) correspond to those of (12).

Proof. We will prove it for the functions (11). The other case is analogous. Using (10), one gets

$$\begin{aligned}
\nabla_z \Phi_{\nu,\mu}(z) &= \sum_{s=a}^{b-1} \nabla_z \left(\frac{\rho_\nu(s) \nabla x_{\nu+1}(s)}{[x_\nu(s) - x_\nu(z)]^{(\mu+1)}} \right) \\
&= \sum_{s=a}^{b-1} \left(\frac{\rho_\nu(s) \nabla x_{\nu+1}(s)}{[x_\nu(s) - x_\nu(z)]^{(\mu+1)}} - \frac{\rho_\nu(s) \nabla x_{\nu+1}(s)}{[x_\nu(s) - x_\nu(z-1)]^{(\mu+1)}} \right) \\
&= \sum_{s=a}^{b-1} \frac{\rho_\nu(s) \nabla x_{\nu+1}(s)}{[x_\nu(s) - x_\nu(z-1)]^{(\mu)}} \left(\frac{1}{x_\nu(s) - x_\nu(z)} - \frac{1}{x_\nu(s) - x_\nu(z-1-\mu)} \right) \\
&= \sum_{s=a}^{b-1} \frac{\rho_\nu(s) \nabla x_{\nu+1}(s)}{[x_\nu(s) - x_\nu(z-1)]^{(\mu)}} \frac{x_\nu(z) - x_\nu(z-1-\mu)}{(x_\nu(s) - x_\nu(z))(x_\nu(s) - x_\nu(z-1-\mu))} \\
&= \sum_{s=a}^{b-1} \frac{\rho_\nu(s) \nabla x_{\nu+1}(s)}{[x_\nu(s) - x_\nu(z)]^{(\mu+2)}} (x_\nu(z) - x_\nu(z-1-\mu))
\end{aligned}$$

Since $x(s) - x(s-t) = [t]_q \nabla x \left(s - \frac{t-1}{2} \right)$ we then have

$$\begin{aligned}
\nabla_z \Phi_{\nu,\mu}(z) &= \sum_{s=a}^{b-1} \frac{\rho_\nu(s) \nabla x_{\nu+1}(s)}{[x_\nu(s) - x_\nu(z)]^{(\mu+2)}} [\mu+1]_q \nabla x_\nu \left(z - \frac{\mu}{2} \right) \\
&= [\mu+1]_q \nabla x_{\nu-\mu}(z) \Phi_{\nu,\mu+1}(z)
\end{aligned}$$

which is (14). \square

From (14) follows that

$$\Delta_z \Phi_{\nu,\mu}(z) = [\mu+1]_q \Delta x_{\nu-\mu}(z) \Phi_{\nu,\mu+1}(z+1).$$

Next we prove the following lemma that is the discrete analog of the Lemma in [14, page 14].

Lemma 3.2. . *Let $x(z)$ be $x(z) = q^z$ or $x(z) = z$. Then, any three functions $\Phi_{\nu_i,\mu_i}(z)$, $i = 1, 2, 3$, are connected by a linear relation*

$$\sum_{i=1}^3 A_i(z) \Phi_{\nu_i,\mu_i}(z) = 0, \quad (15)$$

with non-zero at the same time polynomial coefficients on $x(z)$, $A_i(z)$, provided that the differences $\nu_i - \nu_j$ and $\mu_i - \mu_j$, $i, j = 1, 2, 3$, are integers and that the following condition holds²

$$\left. \frac{x^k(s) \sigma(s) \rho_{\nu_0}(s)}{[x_{\nu_0-1}(s) - x_{\nu_0-1}(z)]^{(\mu_0)}} \right|_{s=a}^{s=b} = 0, \quad k = 0, 1, 2, \dots, \quad (16)$$

when the functions Φ_{ν_i,μ_i} are given by (11) and

$$\int_C \Delta_s \frac{x^k(s) \sigma(s) \rho_{\nu_0}(s) ds}{[x_{\nu_0-1}(s) - x_{\nu_0-1}(z)]^{(\mu_0)}} = 0, \quad k = 0, 1, 2, \dots, \quad (17)$$

when Φ_{ν_i,μ_i} are given by (12). Here ν_0 is the ν_i , $i = 1, 2, 3$, with the smallest real part and μ_0 is the μ_i , $i = 1, 2, 3$, with the largest real part.

Proof. Since in [4] we have proved the case when $x(s) = s$ (the uniform lattice) we will restrict here to the case of the q -linear lattice $x(s) = c_1 q^s + c_2$. Moreover, we will give the proof for the case of functions of the form (11), the other case is completely equivalent. Using the identity

$$\nabla x_{\nu_i+1}(s) = q^{\frac{\nu_i - \nu_0}{2}} \nabla x_{\nu_0+1}(s),$$

²In some cases this condition is equivalent to the condition $x(s)^k \sigma(s) \rho_{\nu_0}(s) \Big|_{s=a}^{s=b} = 0$, $k = 0, 1, 2, \dots$

as well as (3) of Proposition 2.3, we have

$$\begin{aligned}
\sum_{i=1}^3 A_i(z) \Phi_{\nu_i, \mu_i}(z) &= \sum_{i=1}^3 A_i(z) \sum_{s=a}^{b-1} \frac{\rho_{\nu_i}(s) \nabla x_{\nu_i+1}(s)}{[x_{\nu_i}(s) - x_{\nu_i}(z)]^{(\mu_i+1)}} \\
&= \sum_{s=a}^{b-1} \sum_{i=1}^3 A_i(z) \frac{\rho_{\nu_i}(s) \nabla x_{\nu_i+1}(s)}{[x_{\nu_i}(s) - x_{\nu_i}(z)]^{(\mu_i+1)}} = \sum_{s=a}^{b-1} \frac{1}{[x_{\nu_0}(s) - x_{\nu_0}(z)]^{(\mu_0+1)}} \times \\
&\quad \left(\sum_{i=1}^3 A_i(z) q^{\frac{(\mu_i+1)(\nu_0-\nu_i)}{2}} [x_{\nu_0}(s) - x_{\nu_0}(z - \mu_i - 1)]^{(\mu_0-\mu_i)} \rho_{\nu_i}(s) \nabla x_{\nu_i+1}(s) \right) \\
&= \sum_{s=a}^{b-1} \frac{\rho_{\nu_0}(s) \nabla x_{\nu_0+1}(s)}{[x_{\nu_0}(s) - x_{\nu_0}(z)]^{(\mu_0+1)}} \times \\
&\quad \left(\sum_{i=1}^3 A_i(z) q^{\frac{\mu_i(\nu_0-\nu_i)}{2}} [x_{\nu_0}(s) - x_{\nu_0}(z - \mu_i - 1)]^{(\mu_0-\mu_i)} \frac{\rho_{\nu_i}(s)}{\rho_{\nu_0}(s)} \right).
\end{aligned}$$

Using the Pearson-type equation (6) we obtain

$$\rho_{\nu_i}(s) = \phi(s + \nu_0) \phi(s + \nu_0 + 1) \dots \phi(s + \nu_i - 1) \rho_{\nu_0}(s), \quad (18)$$

so

$$\sum_{i=1}^3 A_i(z) \Phi_{\nu_i, \mu_i}(z) = \sum_{s=a}^{b-1} \frac{\rho_{\nu_0}(s) \nabla x_{\nu_0+1}(s)}{[x_{\nu_0}(s) - x_{\nu_0}(z)]^{(\mu_0+1)}} \Pi(s)$$

where

$$\begin{aligned}
\Pi(s) &= \sum_{i=1}^3 A_i(z) q^{\frac{\mu_i(\nu_0-\nu_i)}{2}} [x_{\nu_0}(s) - x_{\nu_0}(z - \mu_i - 1)]^{(\mu_0-\mu_i)} \times \\
&\quad \phi(s + \nu_0) \phi(s + \nu_0 + 1) \dots \phi(s + \nu_i - 1).
\end{aligned} \quad (19)$$

Let us show that there exists a polynomial $Q(s)$ in $x(s)$ (in general, $Q \equiv Q(z, s)$ is a function of z and s) such that

$$\begin{aligned}
\frac{\rho_{\nu_0}(s) \nabla x_{\nu_0+1}(s)}{[x_{\nu_0}(s) - x_{\nu_0}(z)]^{(\mu_0+1)}} \Pi(s) &= \Delta \left[\frac{\rho_{\nu_0}(s-1)}{[x_{\nu_0-1}(s) - x_{\nu_0-1}(z)]^{(\mu_0)}} Q(s) \right] \\
&= \Delta \left[\frac{\sigma(s) \rho_{\nu_0}(s)}{[x_{\nu_0-1}(s) - x_{\nu_0-1}(z)]^{(\mu_0)}} Q(s) \right].
\end{aligned} \quad (20)$$

If such polynomial exists, then, taking the sum in s from $s = a$ to $b - 1$ and using the boundary conditions (16) we obtain (15).

To prove the existence of the polynomial $Q(s)$ in the variable $x(s)$ in (20) we write

$$\begin{aligned}
&\frac{\sigma(s+1) \rho_{\nu_0}(s+1)}{[x_{\nu_0-1}(s+1) - x_{\nu_0-1}(z)]^{(\mu_0)}} Q(s+1) - \frac{\sigma(s) \rho_{\nu_0}(s)}{[x_{\nu_0-1}(s) - x_{\nu_0-1}(z)]^{(\mu_0)}} Q(s) = \\
&\frac{\rho_{\nu_0}(s)}{[x_{\nu_0}(s) - x_{\nu_0}(z)]^{(\mu_0+1)}} \left[\sigma(s+1) \frac{\rho_{\nu_0}(s+1)}{\rho_{\nu_0}(s)} \frac{[x_{\nu_0}(s) - x_{\nu_0}(z)]^{(\mu_0+1)}}{[x_{\nu_0-1}(s+1) - x_{\nu_0-1}(z)]^{(\mu_0)}} Q(s+1) - \right. \\
&\quad \left. \sigma(s) \frac{[x_{\nu_0}(s) - x_{\nu_0}(z)]^{(\mu_0+1)}}{[x_{\nu_0-1}(s) - x_{\nu_0-1}(z)]^{(\mu_0)}} Q(s) \right].
\end{aligned}$$

From (4) and (5) of Proposition 2.3, and using (6), the above expression becomes

$$\begin{aligned}
&\frac{\rho_{\nu_0}(s)}{[x_{\nu_0}(s) - x_{\nu_0}(z)]^{(\mu_0+1)}} \{ \phi_{\nu_0}(s) [x_{\nu_0-\mu_0}(s) - x_{\nu_0-\mu_0}(z)] Q(s+1) - \\
&\quad \sigma(s) [x_{\nu_0-\mu_0}(s + \mu_0) - x_{\nu_0-\mu_0}(z)] Q(s) \}.
\end{aligned}$$

Thus

$$(\sigma(s) + \tau_{\nu_0}(s)\nabla x_{\nu_0+1}(s)) [x_{\nu_0-\mu_0}(s) - x_{\nu_0-\mu_0}(z)] Q(s+1) - \sigma(s) [x_{\nu_0-\mu_0}(s+\mu_0) - x_{\nu_0-\mu_0}(z)] Q(s) = \nabla x_{\nu_0+1}(s)\Pi(s). \quad (21)$$

Since $\nabla x_{\nu_0+1}(s)$ is a polynomial of degree one in $x(s)$, $x_k(s)$ and $\tau_{\nu_0}(s)$ are polynomials of degree at most one in $x(s)$, and $\sigma(s)$ is a polynomial of degree at most two in $x(s)$, we conclude that the degree of $Q(s)$ is, at least, two less than the degree of $\Pi(s)$, i.e., $\deg Q \geq \deg \Pi - 2$. Moreover, equating the coefficients of the powers of $x(s) = q^s$ on the two sides of the above equation (21), we find a system of linear equations in the coefficients of $Q(s)$ and the coefficients $A_i(z)$ which have at least one unknown more than the number of equations. Notice that the coefficients of the unknowns are polynomials in q^z , so that after one coefficient is selected the remaining coefficients are rational functions of q^z , therefore after multiplying by the common denominator of the $A_i(z)$ we obtain the linear relation with polynomial coefficients on $x \equiv x(z) = q^z$. This completes the proof. \square

The above Lemma when $q \rightarrow 1$ and $x(s) = s$ leads to the corresponding result on the uniform lattice $x(s)$ [4].

3.1. Some representative examples. In the following examples, and for the sake of simplicity, we will use the notation

$$\sigma(s) = aq^{2s} + bq^s + c, \quad \tau(s) = dq^s + e, \quad \phi_\nu(s) = \sigma(s) + \tau_{\nu-1}(s)\nabla x_\nu(s) = fq^{2s} + gq^s + h. \quad (22)$$

Example 3.3. *The following relation holds*

$$A_1(z)\Phi_{\nu,\nu-1}(z) + A_2(z)\Phi_{\nu,\nu} + A_3(z)\Phi_{\nu+1,\nu}(z) = 0,$$

where the coefficients A_1 , A_2 and A_3 , are polynomials in $x \equiv x(z) = q^z$, given by

$$A_1(z) = -eq^{\frac{z}{2}} + \frac{b+e\left(q^{\frac{1}{2}} - q^{-\frac{1}{2}}\right)}{a+d\left(q^{\frac{1}{2}} - q^{-\frac{1}{2}}\right)} \left(dq^{\frac{z}{2}} + a[\nu]_q\right) + \left(dq^\nu + a[2\nu]_q\right)q^{\frac{z}{2}+z},$$

$$A_2(z) = \frac{c(dq^\nu + a[2\nu]_q)}{a+d\left(q^{\frac{1}{2}} - q^{-\frac{1}{2}}\right)} + \frac{b+e\left(q^{\frac{1}{2}} - q^{-\frac{1}{2}}\right)}{q^{\frac{1}{2}} - q^{-\frac{1}{2}}} \left(q^\nu + \frac{a}{q^\nu\left(a+d\left(q^{\frac{1}{2}} - q^{-\frac{1}{2}}\right)\right)}\right)q^z + \left(dq^\nu + a[2\nu]_q\right)q^{2z},$$

$$A_3(z) = -\frac{dq^{\frac{z}{2}} + a[\nu]_q}{a+d\left(q^{\frac{1}{2}} - q^{-\frac{1}{2}}\right)},$$

where a , b , c , d , and e , are the coefficients of σ and τ (22).

Proof. Using the notations of Lemma 3.2 we have $\nu_1 = \nu$, $\nu_2 = \nu$, $\nu_3 = \nu + 1$, $\mu_1 = \nu - 1$, $\mu_2 = \nu$ and $\mu_3 = \nu$, thus $\nu_0 = \nu$ and $\mu_0 = \nu$. By (19)

$$\Pi(s) = A_1\left(q^{s+\frac{z}{2}} - q^{z-\frac{z}{2}}\right) + A_2 + A_3q^{-\frac{z}{2}} \left[\left(a+d\left(q^{\frac{1}{2}} - q^{-\frac{1}{2}}\right)\right)q^{2\nu+2s} + \left(b+e\left(q^{\frac{1}{2}} - q^{-\frac{1}{2}}\right)\right)q^{\nu+s} + c\right]. \quad (23)$$

On the other hand, from (21) and because $Q(s) = k$ is a constant –notice that $\deg(\Pi) = 2$ – we have

$$\nabla x_{\nu_0+1}(s)\Pi(s) = k \left\{ \left[\left(a+d\left(q^{\frac{1}{2}} - q^{-\frac{1}{2}}\right)\right)q^{2\nu+2s} + \left(b+e\left(q^{\frac{1}{2}} - q^{-\frac{1}{2}}\right)\right)q^{\nu+s} + c\right] (q^s - q^z) - (aq^{2s} + bq^s + c)(q^{\nu+s} - q^z) \right\} \quad (24)$$

where k is an arbitrary constant. Introducing (23) in (24), using the identity

$$\nabla x_{\nu_0+1}(s) = q^{\frac{z}{2}} \left(q^{\frac{1}{2}} - q^{-\frac{1}{2}}\right) q^s$$

and comparing the coefficients of the powers of $x(s) = q^s$ we get a linear system of three equations with four variables A_1 , A_2 , A_3 and k . Choosing $k = 1$ and solving the corresponding system we get, after some simplifications, the coefficients A_1 , A_2 and A_3 . \square

In the next examples, since the technique is similar to the previous one we will omit the details.

Example 3.4. *The following relation holds*

$$A_1(z)\Phi_{\nu,\nu}(z) + A_2(z)\Phi_{\nu,\nu+1}(z) + A_3(z)\Phi_{\nu+1,\nu+1}(z) = 0,$$

where the coefficients A_1 , A_2 and A_3 , are polynomials in $x \equiv x(z) = q^z$, given by

$$A_1(z) = f(a - f q^{2\nu})q^z + a g q - f b q^{\nu+1},$$

$$A_2(z) = q^{-\frac{z}{2}-1} (a - f q^{2\nu}) (f q^{2z} + g q^{z+1} + h q^2),$$

$$A_3(z) = \sqrt{q} (a q - f q^\nu),$$

where a, b, c, f, g and h , are the coefficients of σ and ϕ_ν (22).

Example 3.5. *The following relation holds*

$$A_1(z)\Phi_{\nu-1,\nu-1}(z) + A_2(z)\Phi_{\nu,\nu-1}(z) + A_3(z)\Phi_{\nu,\nu}(z) = 0,$$

where the coefficients A_1 , A_2 and A_3 , are polynomials in $x \equiv x(z) = q^z$, given by

$$\begin{aligned} A_1(z) = & q^{-\frac{1}{2}-\nu} \left\{ f q^{2z} \left[-a^2 h q^4 + a g b q^{\nu+4} - q^{2\nu+2} (a g^2 q - 2 f a h + f b^2) \right. \right. \\ & \left. \left. - f g b q^{3\nu+1} (q^2 - q - 1) + f q^{4\nu} (g^2 (q - 1) q - f h) \right] + \right. \\ & g q^{z+1} \left[-a^2 h q^5 + a q^{\nu+2} (g b q^3 + f h q^2 - f h) - q^{2\nu+2} ((f a h + f g b + a g^2) q^2 - \right. \\ & \left. f (2 a h - b^2 + g b) q - f a h) + f q^{3\nu} (q^2 (g^2 q - f h + g b - g^2) + f h) + f^2 h q^{4\nu} (q^2 - q - 1) \right] \\ & \left. - a^2 h^2 q^6 + a g h q^{\nu+5} (b q + g q - g) + f g h q^{3\nu+4} (g q + b - g) - f^2 h^2 q^{4\nu+2} \right. \\ & \left. - h q^{2\nu+3} (a g^2 q^3 + f g b q^2 + f g^2 q^2 - 2 f a h q + f b^2 q - 2 f g^2 q - f g b + f g^2) \right\}, \\ A_2(z) = & (q^{-\frac{z}{2}} - q^{\frac{z}{2}}) (f q^{2z} + g q^{z+1} + h q^2) (f q^z (f q^{2\nu} - a q^2) + f q^{\nu+1} (g q + b - g) - a g q^3), \\ A_3(z) = & f (f q^\nu - a q) \left[(f q^{2z} + h q^2) (f q^{2\nu} - a q^2) + g q^{z+1} (f q^\nu (q^\nu + q - 1) - a q^3) \right], \end{aligned}$$

where a, b, c, f, g and h , are the coefficients of σ and ϕ_ν (22).

Example 3.6. *The following relation holds*

$$A_1(z)\Phi_{\nu-1,\nu-1}(z) + A_2(z)\Phi_{\nu,\nu}(z) + A_3(z)\Phi_{\nu,\nu+1}(z) = 0,$$

where the coefficients A_1 , A_2 and A_3 , are polynomials in $x \equiv x(z) = q^z$, given by

$$A_1(z) = a^2 h q^4 - a g b q^{\nu+3} + q^{2\nu+2} (f b^2 - 2 f a h + a g^2) - f g b q^{3\nu+1} + f^2 h q^{4\nu},$$

$$A_2(z) = q^{-\frac{1}{2}} (f q^\nu - a q^2) (f q^{z+2\nu} - a q^{z+2} + g q^{2\nu+1} - b q^{\nu+2}),$$

$$A_3(z) = -q^{\frac{\nu-3}{2}} (f q^{2\nu} - a q^2) (q^{\nu+1} - 1) (g q^{z+1} + f q^{2z} + h q^2),$$

where a, b, c, f, g and h , are the coefficients of σ and ϕ_ν (22).

Example 3.7. *The relation*

$$A_1(z)\Phi_{\nu,\nu-1}(z) + A_2(z)\Phi_{\nu,\nu}(z) + A_3(z)\Phi_{\nu+1,\nu+1}(z) = 0,$$

is verified when the polynomial coefficients A_1 , A_2 and A_3 , in the variable $x \equiv x(z) = q^z$, are given by

$$A_1(z) = q^{\frac{\nu+1}{2}} \left(f q^{z+\nu} (f q^{2\nu} - g q^\nu + b - a) - f (h - b) q^{2\nu+1} - a q (g q^\nu - h) \right),$$

$$A_2(z) = q^{-z+\nu+\frac{1}{2}} \left(q^z (f q^{2\nu} - a) + q^\nu (g q^\nu - b) \right) (f q^{2z} + g q^{z+1} + h q^2),$$

$$\begin{aligned} A_3(z) = & q^{2z} (f q^\nu - a q) + q^{z+\nu} \left(q (g q^\nu - a q - b) - f q^\nu (q^{\nu+1} - q - 1) \right) + \\ & q^{\nu+1} ((h - b) q^{\nu+1} + g q^\nu - h), \end{aligned}$$

where a, b, c, f, g and h , are the coefficients of σ and ϕ_ν (22).

4. RECURRENCES INVOLVING THE SOLUTIONS y_ν

In [16] the following relevant relation was established

$$\Delta^{(k)}y_\nu(s) = \frac{C_\nu^{(k)}}{\rho_k(s)}\Phi_{\nu, \nu-k}(s), \quad (25)$$

where

$$C_\nu^{(k)} = C_\nu \prod_{m=0}^{k-1} \left[\alpha_q(\nu + m - 1)\tilde{\tau}' + [\nu + m - 1]_q \frac{\tilde{\sigma}''}{2} \right].$$

This relation is valid for solutions of the form (3) and (4) of the difference equation (1).

In the following, $y_n^{(k)}(s)$ denotes the k -th differences $\Delta^{(k)}y_n(s)$.

Theorem 4.1. *In the same conditions as in Lemma 3.2, any three functions $y_{\nu_i}^{(k_i)}(s)$, $i = 1, 2, 3$, are connected by a linear relation*

$$\sum_{i=1}^3 B_i(s)y_{\nu_i}^{(k_i)}(s) = 0, \quad (26)$$

where the $B_i(s)$, $i = 1, 2, 3$, are polynomials.

Proof. From Lemma 3.2 we know that there exists three polynomials $A_i(s)$, $i = 1, 2, 3$ such that

$$\sum_{i=1}^3 A_i(s)\Phi_{\nu_i, \nu_i-k_i}(s) = 0,$$

then, using the relation (25), we find

$$\sum_{i=1}^3 A_i(s)(C_\nu^{(k)})^{-1}\rho_{k_i}(s)y_{\nu_i}^{(k_i)}(s) = 0.$$

Now, dividing the last expression by $\rho_{k_0}(s)$, where $k_0 = \min\{k_1, k_2, k_3\}$, and using (18) we obtain

$$\sum_{i=1}^3 B_i(s)y_{\nu_i}^{(k_i)}(s) = 0, \quad B_i(s) = A_i(s)(C_\nu^{(k)})^{-1}\phi(s + k_0) \cdots \phi(s + k_i - 1),$$

which completes the proof. \square

Corollary 4.2. *In the same conditions as in Lemma 3.2, the following three-term recurrence relation holds*

$$A_1(s)y_\nu(s) + A_2(s)y_{\nu+1}(s) + A_3(s)y_{\nu-1}(s) = 0,$$

with polynomial coefficients $A_i(s)$, $i = 1, 2, 3$.

Proof. It is sufficient to put $k_1 = k_2 = k_3 = 0$, $\nu_1 = \nu$, $\nu_2 = \nu + 1$ and $\nu_3 = \nu - 1$ in (26). \square

Corollary 4.3. *In the same conditions as in Lemma 3.2, the following Δ -ladder-type relation holds*

$$B_1(s)y_\nu(s) + B_2(s)\frac{\Delta y_\nu(s)}{\Delta x(s)} + B_3(s)y_{\nu+m}(s) = 0, \quad m \in \mathbb{Z}, \quad (27)$$

with polynomial coefficients $B_i(s)$, $i = 1, 2, 3$.

Proof. It is sufficient to put $k_1 = k_3 = 0$, $k_2 = 1$, $\nu_1 = \nu_2 = \nu$ and $\nu_3 = \nu + m$ in (26). \square

Notice that for the case $m = \pm 1$ (27) becomes

$$B_1(s)y_\nu(s) + B_2(s)\frac{\Delta y_\nu(s)}{\Delta x(s)} + B_3(s)y_{\nu+1}(s) = 0, \quad (28)$$

$$\tilde{B}_1(s)y_\nu(s) + \tilde{B}_2(s)\frac{\Delta y_\nu(s)}{\Delta x(s)} + \tilde{B}_3(s)y_{\nu-1}(s) = 0, \quad (29)$$

with polynomial coefficients $B_i(s)$ and $\tilde{B}_i(s)$, $i = 1, 2, 3$. The above relations are usually called raising and lowering operators, respectively, for the functions y_ν .

Let us now obtain a raising and lowering operators for the functions y_ν but associated to the $\nabla/\nabla x(s)$ operators.

We start applying the operator $\nabla/\nabla x(s)$ to (13)

$$\begin{aligned} \frac{\nabla}{\nabla x(s)}y_\nu(s) &= \frac{\nabla}{\nabla x(s)} \left[\frac{C_\nu}{\rho(s)}\Phi_{\nu,\nu}(s) \right] \\ &= \frac{1}{\nabla x(s)} \left[C_\nu\Phi_{\nu\nu}(s) \left(\frac{1}{\rho(s)} - \frac{1}{\rho(s-1)} \right) + \frac{C_\nu}{\rho(s-1)}\nabla\Phi_{\nu\nu}(s) \right], \end{aligned}$$

or, equivalently,

$$\frac{\nabla\Phi_{\nu\nu}}{\nabla x(s)} = \frac{\rho(s-1)}{C_\nu} \frac{\nabla y_\nu(s)}{\nabla x(s)} - \frac{\Phi_{\nu\nu}(s)}{\nabla x(s)} \left[\frac{\rho(s-1)}{\rho(s)} - 1 \right].$$

By Lemma (3.2) with $\nu_1 = \mu_1 = \nu_2 = \nu$, $\mu_2 = \nu + 1$ and $\nu_3 = \mu_3 = \nu + m$, there exist polynomial coefficients on $x(s)$, $A_i(s)$, $i = 1, 2, 3$, such that

$$A_1(s)\Phi_{\nu,\nu}(s) + A_2(s)\Phi_{\nu,\nu+1}(s) + A_3(s)\Phi_{\nu+m,\nu+m}(s) = 0.$$

From (14)

$$\Phi_{\nu,\nu+1}(s) = \frac{1}{[\nu+1]_q} \frac{\nabla\Phi_{\nu,\nu}}{\nabla x(z)} = \frac{1}{[\nu+1]_q} \frac{\nabla\Phi_{\nu,\nu}}{\nabla x(z)}.$$

Therefore

$$\begin{aligned} A_1(s)\Phi_{\nu,\nu} + \frac{A_2(s)}{[\nu+1]_q} \left[\frac{\rho(s-1)}{C_\nu} \frac{\nabla y_\nu}{\nabla x(s)} - \frac{\Phi_{\nu\nu}(s)}{\nabla x(s)} \left(\frac{\rho(s-1)}{\rho(s)} - 1 \right) \right] \\ + A_3\Phi_{\nu+m,\nu+m} = 0. \end{aligned}$$

Using now the Pearson equation (6) and dividing by $\rho(s)$ we get

$$\begin{aligned} A_1(s)y_\nu(s) + \frac{A_2(q)}{[\nu+1]_q} \left[\frac{\sigma(s)}{\phi(s-1)} \frac{\nabla y_\nu}{\nabla x(s)} - \frac{y_\nu(s)}{\nabla x(s)} \left(\frac{\sigma(s)}{\phi(s-1)} - 1 \right) \right] \\ + A_3 \frac{C_\nu}{C_{\nu+m}} y_{\nu+m}(s) = 0. \end{aligned}$$

Multiplying both sides by $[\nu+1]_q\phi(s-1)$,

$$\begin{aligned} A_1(s)[\nu+1]_q\phi(s-1)y_\nu(s) + A_2(s)\sigma(s)\frac{\nabla y_\nu}{\nabla x(s)} - \\ A_2(s)\frac{\sigma(s) - \phi(s-1)}{\nabla x(s)}y_\nu(s) + [\mu+1]_q C_\nu C_{\nu+m}^{-1} A_3\phi(s-1)y_{\nu+m}(s) = 0. \end{aligned}$$

Thus we have proven the following

Theorem 4.4. *In the same conditions as in Lemma 3.2, the following ∇ -ladder-type relation holds*

$$C_1(s)y_\nu(s) + C_2(s)\frac{\nabla y_\nu(s)}{\nabla x(s)} + C_3(s)y_{\nu+m}(s) = 0, \quad m \in \mathbb{Z}, \quad (30)$$

with polynomial coefficients $C_i(s)$, $i = 1, 2, 3$.

Notice that for the case $m = \pm 1$ (30) becomes

$$C_1(s)y_\nu(s) + C_2(s)\frac{\nabla y_\nu(s)}{\nabla x(s)} + C_3(s)y_{\nu+1}(s) = 0, \quad (31)$$

$$\tilde{C}_1(s)y_\nu(s) + \tilde{C}_2(s)\frac{\nabla y_\nu(s)}{\nabla x(s)}y_\nu(s) + \tilde{C}_3(s)y_{\nu-1}(s) = 0, \quad (32)$$

with polynomial coefficients $C_i(s)$ and $\tilde{C}_i(s)$, $i = 1, 2, 3$. The above relations are usually called raising and lowering operators, respectively, for the functions y_n . Eq. (31) was first obtained in [16, Eq. (3.4)].

To conclude this section let us point out that from formula (25) and the examples 3.3, 3.5, and 3.7 follow the relations

$$\begin{aligned} B_1(s)y_\nu^{(1)}(s) + B_2(s)y_\nu(s) + B_3(s)y_{\nu+1}^{(1)}(s) &= 0, \\ B_1(s)y_\nu^{(1)}(s) + B_2(s)y_{\nu-1}(s) + B_3(s)y_\nu(s) &= 0, \\ B_1(s)y_\nu^{(1)}(s) + B_2(s)y_\nu(s) + B_3(s)y_{\nu+1}(s) &= 0, \end{aligned} \quad (33)$$

respectively, being the last two expressions the lowering and raising operators for the functions y_ν . Moreover, combining the explicit values of A_1 , A_2 and A_3 with formula (25), one can obtain the explicit expressions for the coefficients B_1 , B_2 and B_3 in (33).

5. APPLICATIONS TO q -CLASSICAL POLYNOMIALS

In this section we will apply the previous results to the q -classical orthogonal polynomials [2, 10, 11] in order to show how the method works. We first notice that these polynomials are instances of the functions y_ν on the lattice $x(s) = q^s$ defined in (4). In fact we have [13, 16]

$$P_n(x(s)) = \frac{[n]_q! B_n}{\rho(s) 2\pi i} \int_C \frac{\rho_n(z) \nabla x_{n+1}(z)}{[x_n(z) - x_n(s)]^{(n+1)}} dz, \quad (34)$$

where B_n is a normalizing constant, C is a closed contour surrounding the points $x = s, s-1, \dots, s-n$ and it is assumed that $\rho_n(s) = \rho(s+n) \prod_{m=1}^n \sigma(s+m)$ and $\rho_n(s+1)$ are analytic inside C (ρ is the solution of the Pearson equation (6)), i.e., the condition (5) holds.

A detailed study of the q -classical polynomials, including several characterization theorems, was done in [2, 9, 11]. In particular, a comparative analysis of the q -Hahn tableau with the q -Askey tableau [9] and Nikiforov-Uvarov tableau [15] was done in [5]. In the following we use the standard notation for the q -calculus [8]. In particular by $(a; q)_k = \prod_{m=0}^{k-1} (1 - aq^m)$, we denote the q -analogue of the Pochhammer symbol.

Since the q -classical polynomials are defined by (34) where the contour C is closed and ν is a non-negative integer, then the condition (17) is automatically fulfilled, so Lemma 3.2 holds for all of them. Moreover, the Theorem 4.1 holds and there exist the non-vanishing polynomials B_1 , B_2 and B_3 of (26).

In the following we will assume that the three-term recurrence relation is known, i.e.,

$$\begin{aligned} x(s)P_n(x(s)) &= \alpha_n P_{n+1}(x(s)) + \beta_n P_n(x(s)) + \gamma_n P_{n-1}(x(s)) = 0, \quad n \geq 0 \\ P_{-1}(x(s)) &= 0, \quad P_0(x(s)) = 1, \quad x(s) = q^s. \end{aligned} \quad (35)$$

where the coefficients α_n , β_n and γ_n can be computed using the coefficients σ , τ and $\lambda \equiv \lambda_n$ of (1), being λ_n given by (8) and (9) with $\nu = n$. For more details see, e.g., [1, 11].

Since the TTRR and the differentiation formulas for the q -polynomials are very well known (see e.g. [9, 11, 16]) we will obtain here two recurrent-difference relations involving the q -differences of the polynomials and the polynomials themselves.

5.1. The first difference-recurrence relation. If we choose $\nu_1 = n - 1$, $\nu_2 = n$, $\nu_3 = n + 1$, $k_1 = 1$, $k_2 = 1$ and $k_3 = 0$, in Theorem 4.1 one gets

$$A_1(s)\Delta^{(1)}P_{n-1}(x(s)) + A_2(s)\Delta^{(1)}P_n(x(s)) + A_3(s)P_{n+1}(x(s)) = 0.$$

Using [1, Eq. (6.14), page 193]

$$[\sigma(s) + \tau(s)\Delta x(s - 1/2)]\Delta^{(1)}P_n(x(s)) = \widehat{\alpha}_n P_{n+1}(x(s)) + \widehat{\beta}_n P_n(x(s)) + \widehat{\gamma}_n P_{n-1}(x(s)),$$

where

$$\widehat{\alpha}_n = \frac{\lambda_n}{[n]_q} \left[q^{-\frac{n}{2}} \alpha_n - \frac{B_n}{\tau'_n B_{n+1}} \right], \quad \widehat{\beta}_n = \frac{\lambda_n}{[n]_q} \left[q^{-\frac{n}{2}} \beta_n + \frac{\tau_n(0)}{\tau'_n} - c_3(q^{-\frac{n}{2}} - 1) \right],$$

$$\widehat{\gamma}_n = \frac{\lambda_n q^{-\frac{n}{2}} \gamma_n}{[n]_q},$$

to compute $\Delta^{(1)}P_n(x(s)) = \frac{\Delta P_n(x(s))}{\Delta x(s)}$ we get

$$\begin{aligned} & \left[A_2(s) \frac{\lambda_n}{[n]_q} \left(q^{-\frac{n}{2}} \alpha_n - \frac{B_n}{\tau'_n B_{n+1}} \right) + (\sigma(s) + \tau(s)\Delta x(s - \frac{1}{2})) A_3(s) \right] P_{n+1} + \\ & \left[A_1(s) \frac{\lambda_{n-1}}{[n-1]_q} \left(q^{-\frac{n-1}{2}} \alpha_{n-1} - \frac{B_{n-1}}{\tau'_{n-1} B_n} \right) + A_2(s) \frac{\lambda_n}{[n]_q} \left(q^{-\frac{n}{2}} \beta_n + \frac{\tau_n(0)}{\tau'_n} \right) \right] P_n + \\ & \left[A_1(s) \frac{\lambda_{n-1}}{[n-1]_q} \left(q^{-\frac{n-1}{2}} \beta_{n-1} + \frac{\tau_{n-1}(0)}{\tau'_{n-1}} \right) + A_2(s) \frac{\lambda_n q^{-\frac{n}{2}} \gamma_n}{[n]_q} \right] P_{n-1} + \\ & A_1(s) \frac{\lambda_{n-1} q^{-\frac{n-1}{2}} \gamma_{n-1}}{[n-1]_q} P_{n-2} = 0, \end{aligned}$$

By (35) we may write

$$P_{n-2}(x(s)) = \frac{x(s) - \beta_{n-1}}{\gamma_{n-1}} P_{n-1}(x(s)) - \frac{\alpha_{n-1}}{\gamma_{n-1}} P_n(x(s))$$

so the above equality becomes

$$\begin{aligned} & \left[\frac{\lambda_n}{[n]_q} \left(q^{-\frac{n}{2}} \alpha_n - \frac{B_n}{\tau'_n B_{n+1}} \right) A_2(s) + (\sigma(s) + \tau(s)\Delta x(s - \frac{1}{2})) A_3(s) \right] P_{n+1}(x(s)) + \\ & \left[-\frac{\lambda_{n-1}}{[n-1]_q} \frac{B_{n-1}}{\tau'_{n-1} B_n} A_1(s) + \frac{\lambda_n}{[n]_q} \left(q^{-\frac{n}{2}} \beta_n + \frac{\tau_n(0)}{\tau'_n} \right) A_2(s) \right] P_n(x(s)) + \\ & \left[\frac{\lambda_{n-1}}{[n-1]_q} \left(\frac{\tau_{n-1}(0)}{\tau'_{n-1}} + q^{-\frac{n-1}{2}} x \right) A_1(s) + \frac{\lambda_n}{[n]_q} q^{-\frac{n}{2}} \gamma_n A_2(s) \right] P_{n-1}(x(s)) = 0. \end{aligned} \quad (36)$$

Comparing the above equation with the TTRR (35) one can obtain the explicit values of A_1 , A_2 , and A_3 .

5.1.1. Some examples. Since we are working in the q -linear lattice $x(s) = q^s$, for the sake of simplicity, we will use the letter x to denote the variable of the polynomials [9, 11]. We will consider monic polynomials, i.e., those with the leading coefficient equal to 1. In the following we need the value of $\tau_n(x)$ for each family, which can be computed using (7).

Al-Salam-Carlitz I q -polynomials. For the Al-Salam-Carlitz I monic polynomials $U_n^{(a)}(x; q)$ we have (see [1, see table 6.5, p.208] or [11])

$$\begin{aligned} \sigma(x) &= (1-x)(a-x), \quad \tau_n(x) = \frac{q^{\frac{1-n}{2}}}{1-q} (x - (1+a)), \\ \tau(x) &= \tau_0(x), \quad \lambda_n = -\frac{q^{\frac{3-n}{2}}(1-q^n)}{(1-q)^2}, \end{aligned}$$

and

$$\alpha_n = 1, \quad \beta_n = (1+a)q^n, \quad \gamma_n = -aq^{n-1}(1-q^n).$$

The constant B_n is given by [1, Eq. (5.57), p. 147], $B_n = q^{\frac{1}{4}n(3n-5)}(1-q)^n$. Introducing these values into the equation (36) it becomes

$$\begin{aligned} & \left[q \left(q^{-\frac{n}{2}} - 1 \right) A_2(x) + a(1-q)q^n A_3(x) \right] U_{n+1}^{(a)}(x; q) + \\ & \left[q^{-\frac{n}{2}-\frac{5}{2}} A_1(x) + q^{1+\frac{n}{2}}(1+a) \left(1 - q^{\frac{n}{2}} \right) A_2(x) \right] U_n^{(a)}(x; q) + \\ & \left[\left(q^{\frac{n+3}{2}}(1+a) - q^{2-n}x \right) A_1(x) + aq^n(1-q^n) A_2(x) \right] U_{n-1}^{(a)}(x; q) = 0. \end{aligned}$$

Comparing with the TTRR (35) for the Al-Salam I polynomials we obtain a linear system for getting the unknown coefficients A_1 , A_2 and A_3

$$\begin{aligned} & q \left(q^{-\frac{n}{2}} - 1 \right) A_2(x) + a(1-q)q^n A_3(x) = 1, \\ & q^{-\frac{n}{2}-\frac{5}{2}} A_1(x) + q^{1+\frac{n}{2}}(1+a) \left(1 - q^{\frac{n}{2}} \right) A_2(x) = (1+a)q^n - x, \\ & \left(q^{\frac{n+3}{2}}(1+a) - q^{2-n}x \right) A_1(x) + aq^n(1-q^n) A_2(x) = aq^{n-1}(q^n - 1). \end{aligned}$$

The solution of the above system is

$$\begin{aligned} A_1(x) &= \frac{aq^n(1+q^{\frac{n}{2}})(1+a)q^{-\frac{n}{2}}x}{aq^{-\frac{5}{2}}(1+q^{\frac{n}{2}}) - q(1+a)\left(q^{\frac{n+3}{2}}(1+a) - q^{2-n}x\right)}, \\ A_2(x) &= \frac{-aq^{-\frac{7}{2}}(1-q^n) - ((1+a)q^n - x)\left(q^{\frac{3}{2}}(1+a) - q^{2-\frac{3n}{2}}x\right)}{\left(1 - q^{\frac{n}{2}}\right) \left[aq^{-\frac{5}{2}}(1+q^{\frac{n}{2}}) - q(1+a)\left(q^{\frac{n+3}{2}}(1+a) - q^{2-n}x\right) \right]}, \\ A_3(x) &= \frac{a+q^{\frac{11}{2}-2n}x^2 + q^{-\frac{n}{2}}(a - (1+a)q^5x)}{a(1-q)\left[aq^n + q^{\frac{3n}{2}}(a - (1+a)^2q^5) + (1+a)q^{\frac{11}{2}}x \right]}. \end{aligned} \quad (37)$$

Then, the Al-Salam I q -polynomials satisfy the the following relation

$$A_1(x)\Delta^{(1)}U_{n-1}^{(a)}(x; q) + A_2(x)\Delta^{(1)}U_n^{(a)}(x; q) + A_3(x)U_{n+1}^{(a)}(x; q) = 0, \quad (38)$$

where the coefficients A_1 , A_2 and A_3 are given by (37).

Notice that the coefficients A_1 , A_2 and A_3 are rational functions on x . Therefore, multiplying (38) by and appropriate factor it becomes a linear relation with polynomials coefficients.

Alternative q -Charlier polynomials. In this case (see [1, table 6.6, p.209])

$$\begin{aligned} \sigma(x) &= q^{-1}x(1-x), \quad \tau_n(x) = -\frac{q^{-\frac{n+1}{2}}}{1-q} \left((1+aq^{1+2n})x - 1 \right), \\ \tau(x) &= \tau_0(x), \quad \lambda_n = \frac{q^{\frac{1}{2}-n}(1-q^n)(1+aq^n)}{(1-q)^2}, \end{aligned}$$

and, for the monic case, $\alpha_n = 1$

$$\beta_n = \frac{q^n(1+aq^{n-1}+aq^n-aq^{2n})}{(1+aq^{2n-1})(1+aq^{2n+1})}, \quad \gamma_n = \frac{aq^{3n-2}(1-q^n)(1+aq^{n-1})}{(1+aq^{2n-2})(1+aq^{2n-1})^2(1+aq^{2n})}.$$

The corresponding normalizing constant B_n is given by

$$B_n = \frac{(-1)^n q^{\frac{1}{4}n(3n-1)}(1-q)^n}{(-aq^n; q)_n}.$$

Following the same procedure as before we obtain the following relation for the alternative Charlier q -polynomials:

$$A_1(x)\Delta^{(1)}K_{n-1}(x; a; q) + A_2(x)\Delta^{(1)}K_n(x; a; q) + A_3(x)K_{n+1}(x; a; q) = 0,$$

with the coefficients

$$A_1(x) = \frac{a(1+aq^{\frac{n}{2}})((1+aq^{2n+1})x - q^{-\frac{n}{2}})x}{q^2(1+aq^{2n-2})(1+aq^{2n-1})(1+aq^{2n})(1+aq^{2n+1})},$$

$$A_2(x) = \frac{-q^{\frac{3n+1}{2}}(1+aq^n)x + (1+aq^{2n})\left(q^{1+\frac{n}{2}}(1+aq^{2n+1}) + aq^{2n+\frac{1}{2}}(1+q) + q^{\frac{3n}{2}}(1-aq^{2n})\right)x^2}{q^{3n}(1+aq^n)(1+aq^{2n})(1+aq^{2n+1})} - \frac{q^{\frac{3}{2}}(1+aq^{2n-1})(1+aq^{2n+1})x^3}{q^{3n}(1+aq^n)(1+aq^{2n})(1+aq^{2n+1})},$$

$$A_3(x) = \frac{q^{\frac{n+1}{2}} + aq^{2n}\left(q^{\frac{n}{2}} + 1 + q^{\frac{1}{2}}\right) - q^{\frac{3}{2}}\left(1 - aq^{\frac{3n}{2}}\right)(1+aq^{2n-1})x}{q^{\frac{9n}{2}}(1+aq^n)}.$$

Big q -Jacobi polynomials. In this case (see [1, see table 6.2, p.204] or [11])

$$\sigma(x) = q^{-1}(x - aq)(x - cq), \lambda_n = -q^{\frac{1}{2}-n} \frac{(1 - abq^{1+n})(1 - q^n)}{(1 - q)^2},$$

$$\tau_n(x) = \frac{q^{\frac{1-n}{2}}}{1 - q} \left(\frac{1 - abq^{2+2n}}{q} x + a(b + c)q^{1+n} - (a + c) \right), \tau(x) = \tau_0(x),$$

and, for the monic case $\alpha_n = 1$,

$$\beta_n = \frac{c+a^2bq^n\left((1+b+c)q^{1+n}-q-1\right)+a\left(1+b+c-q^n\left(b(1+q)+c(1+q+b+bq-bq^{1+n})\right)\right)}{q^{-1-n}(1-abq^{2n})(1-abq^{2n+2})},$$

$$\gamma_n = -\frac{a(1-q^n)(1-aq^n)(1-bq^n)(1-cq^n)(c-abq^n)}{q^{-1-n}(1-abq^{2n-1})(1-abq^{2n})^2(1-abq^{2n+1})}.$$

The corresponding normalizing constant is

$$B_n = \frac{(1 - q)^n q^{\frac{1}{4}n(3n-1)}}{(abq^{1+n}; q)_n}.$$

The big q -Jacobi polynomials satisfy the following relation

$$A_1(x)\Delta^{(1)}p_{n-1}(x; a, b, c; q) + A_2(x)\Delta^{(1)}p_n(x; a, b, c; q) + A_3(x)p_{n+1}(x; a, b, c; q) = 0,$$

with the coefficients A_1 , A_2 and A_3 given by

$$A_1(x) = \frac{aq^{-\frac{1}{2}+n}(1-abq^{n+1})(1-x)(c-bx)\left(c-(b+c)x+bx^2\right)}{1-abq^{2n-1}} \times$$

$$\left\{ (1-q)q^{\frac{n}{2}}(1-abq^{2n+2}) \left[\frac{c+a\left(1+b+c+b(c+a(1+b+c))q^{2n+1}-(c+b(1+a+c))q^n(1+q)\right)}{q^{-(n+1)}(1-abq^{2n})(1-abq^{2n+2})} - x \right] D(x) - \right.$$

$$(1-q)q^n(1-abq^{2n}) \left[(1-abq^{2n})(-c+a(-1+(b+c)q^{n+1})) + \right.$$

$$\left. \left. q^{\frac{n}{2}}\left(c+a\left(1+b+c+b(c+a(1+b+c))q^{2n+1}-(c+b(1+a+c))q^n(1+q)\right)\right) N(x) \right] \right\},$$

$$A_2(x) = a(1-q)q^n(1-abq^{2n})^2(1-abq^{2n+2})(1-x)(c-bx)\left(c-(b+c)x+bx^2\right)N(x),$$

$$A_3(x) = (1-abq^{n+1})(1-abq^{2n+2})(1-x)(c-bx)D(x) +$$

$$q^{-1-\frac{n}{2}}\left(1-q^{\frac{n}{2}}\right)\left(1+abq^{1+\frac{3n}{2}}\right)(1-abq^{2n})^2(1-abq^{2n+2})\left(c-(b+c)x+bx^2\right)N(x),$$

where the polynomials $N(x)$ and $D(x)$ are given by

$$N(x) = \frac{aq^2(1-q^n)(1-aq^n)(1-bq^n)(1-cq^n)(c-abq^n)}{(1-abq^{2n})^2(1-aq^{2n+1})} - \left[\frac{q(-c+a(-1+(b+c)q^n))}{1-abq^{2n}} + q^{\frac{1-n}{2}}x \right] \times$$

$$\left[\frac{c+a^2bq^n(-1-q+(1+b+c)q^{n+1})+a\left(1-(b+c)(-1+q^n+q^{n+1})-bcq^n(1+q-q^{n+1})\right)}{q^{-n-1}(1-abq^{2n})(1-abq^{2n+2})} - x \right]$$

and

$$D(x) = \frac{aq(1-q^n)(1-aq^n)(1-bq^n)(1-cq^n)(c-abq^n)}{1-abq^{2n+1}} + \frac{1-q^{\frac{n}{2}}}{1-abq^{2n+2}} \times \\ \left\{ -c + a^2bq^{\frac{3n}{2}}(-1-q+(b+c)q^{n+1}-q^{1+\frac{n}{2}}) + a \left[-1 + (b+c)(q^{\frac{n}{2}} + q^n + q^{n+1}) - \right. \right. \\ \left. \left. bc(q^{\frac{3n}{2}} + q^{1+\frac{3n}{2}} + q^{2n+1}) \right] \right\} \left[(c+a)q^{1+\frac{n}{2}} - a(b+c)q^{1+\frac{3n}{2}} - q^{\frac{1}{2}}(1-abq^{2n})x \right],$$

respectively.

5.2. The second difference-recurrence relation. If we choose $\nu_1 = n - 1$, $\nu_2 = n$, $\nu_3 = n + 1$, $k_1 = 0$, $k_2 = 0$ and $k_3 = 1$ in Theorem 4.1, and proceeding as in the previous case one gets

$$A_1(x)P_{n-1}(x; q) + A_2(x)P_n(x; q) + A_3(x)\Delta^{(1)}P_{n+1}(x; q) = 0, \quad (39)$$

where the coefficients A_1 , A_2 and A_3 , satisfy the linear relation

$$A_3(x) \left[\left(q^{-\frac{n+1}{2}} - \frac{B_{n+1}}{\alpha_{n+1}\tau'_{n+1}B_{n+2}} \right) (x - \beta_{n+1}) + \left(q^{-\frac{n+1}{2}} \beta_{n+1} + \frac{\tau_{n+1}(0)}{\tau'_{n+1}} \right) \right] P_{n+1} + \\ \left[A_3(x) \frac{B_{n+1}}{\alpha_{n+1}\tau'_{n+1}B_{n+2}} \gamma_{n+1} + \left(\sigma(x) + \tau(x)\Delta x \left(s - \frac{1}{2} \right) \right) \frac{[n+1]_q}{\lambda_{n+1}} A_2(x) \right] P_n + \\ \left(\sigma(x) + \tau(x)\Delta x \left(s - \frac{1}{2} \right) \right) \frac{[n+1]_q}{\lambda_{n+1}} A_1(x)P_{n-1} = 0.$$

Comparing the above relation with the three-term recurrence relation (35) one can obtain the explicit expressions for the coefficients A_1 , A_2 and A_3 in (39).

5.2.1. Some examples.

Al-Salam and Carlitz I polynomials. Using the main data for the Al-Salam and Carlitz I polynomials we obtain the relation

$$A_1(x)U_{n-1}^{(a)}(x; q) + A_2(x)U_n^{(a)}(x; q) + A_3(x)\Delta^{(1)}U_{n+1}^{(a)}(x; q) = 0$$

where

$$A_1(x) = aq^{n-1}(1-q^n)x, \quad A_2(x) = \left[a \left(1 + q^{\frac{n+1}{2}} \right) q^n - \left((1+a)q^n - x \right) x \right], \\ A_3(x) = -a \frac{1-q}{1-q^{\frac{n+1}{2}}} q^{\frac{3n+1}{2}}.$$

Alternative q -Charlier polynomials. In this case, one gets

$$A_1(x)K_{n-1}(x; a; q) + A_2(x)K_n(x; a; q) + A_3(x)\Delta^{(1)}K_{n+1}(x; a; q) = 0, \\ A_1(x) = \frac{a(1-q^n)(1+aq^{n-1}) \left\{ aq^n(1-q^{n+1}) + q^{-\frac{n+1}{2}}(1+aq^{2n+1}) \left[(1+aq^{n+1}) - q^{-\frac{n+1}{2}}(1+aq^{2n+2}) \right] x \right\}}{q^{2-3n}(1+aq^{2n-2})(1+aq^{2n-1})(1+aq^{2n})}, \\ A_2(x) = -x \left\{ aq^n(1-q^{n+1}) + q^{-\frac{n+1}{2}}(1+aq^{2n+1}) \left[(1+aq^{n+1}) - q^{-\frac{n+1}{2}}(1+aq^{2n+2}) \right] x \right\} + \\ \frac{a^2q^{3n-1}(1-q^n)(1-q^{n+1}) + q^{\frac{n-1}{2}}(1+aq^{n-1}+aq^n-aaq^{2n})(1+aq^{2n+1}) \left[(1+aq^{n+1}) - q^{-\frac{n+1}{2}}(1+aq^{2n+2}) \right] x}{(1+aq^{2n-1})(1+aq^{2n+1})}, \\ A_3(x) = a(1-q)q^{\frac{n+1}{2}}(1+aq^{2n+1})x^2$$

Concluding remarks. In this paper we present a constructive approach for finding recurrence relations for the hypergeometric-type functions on the linear-type lattices, i.e., the solutions of the hypergeometric difference equation (1) on the linear-type lattices. Important instances of “discret” functions are the celebrated Askey-Wilson polynomials and q -Racah polynomials. Such functions are defined on the non-uniform lattice of the form $x(s) = c_1(q)q^s + c_2(q)q^{-s} + c_2(q)$ with $c_1c_2 \neq 0$, i.e., a non linear-type lattice and therefore they require a more detailed study (some preliminar general results can be found in [16]).

Acknowledgements. The authors thank J. S. Dehesa and J.C. Petronilho for interesting discussions. The authors were partially supported by DGES grants MTM2009-12740-C03; PAI grant FQM-0262 (RAN) and Junta de Andalucía grants P09-FQM-4643, Spain; and CM-UTAD from UTAD (JLC).

REFERENCES

- [1] R. Álvarez-Nodarse, *Polinomios hipergométricos y q -polinomios*. Monografías del Seminario Matemático “García de Galdeano” Vol. **26**. Prensas Universitarias de Zaragoza, Zaragoza, Spain, 2003.
- [2] R. Álvarez-Nodarse, On characterizations of classical polynomials, *J. Comput. Appl. Math.* **196** (2006), 320-337.
- [3] R. Álvarez-Nodarse, N. M. Atakishiyev, and R. S. Costas-Santos, Factorization of hypergeometric-type difference equations on nonuniform lattices: dynamical algebra. *J. Phys. A: Math. Gen.* **38** (2005) 153-174.
- [4] R. Álvarez-Nodarse and J. L. Cardoso, Recurrence relations for discrete hypergeometric functions. *J. Difference Eq. Appl.* **11** (2005), 829-850.
- [5] R. Álvarez-Nodarse y J. C. Medem, q -Classical polynomials and the q -Askey and Nikiforov-Uvarov Tableaus. *J. Comput. Appl. Math.* **135** (2001), 157-196.
- [6] G. E. Andrews, R. Askey, and R. Roy, *Special functions*. Encyclopedia of Mathematics and its Applications, Cambridge University Press, Cambridge, 1999.
- [7] N. M. Atakishiyev, M. Rahman, and S. K. Suslov, Classical Orthogonal Polynomials, *Constr. Approx.* **11** (1995) 181-226.
- [8] G. Gasper and M. Rahman, *Basic Hypergeometric Series. (2nd Ed.)* Encyclopedia of Mathematics and its Applications **96**, Cambridge University Press, Cambridge, 2004.
- [9] R. Koekoek, Peter A. Lesky, and R.F. Swarttouw, *Hypergeometric orthogonal polynomials and their q -analogues*, Springer Monographs in Mathematics, Springer-Verlag, Berlin-Heidelberg, 2010.
- [10] T. H. Koornwinder, Compact quantum groups and q -special functions. En: *Representations of Lie groups and quantum groups*. V. Baldoni y M.A. Picardello (Eds.) Pitman Research Notes in Mathematics, Series 311, Longman Scientific & Technical (1994), 46-128.
- [11] J. C. Medem, R. Álvarez-Nodarse, and F. Marcellán, On the q -polynomials: A distributional study. *J. Comput. Appl. Math.* **135** (2001), 197-223.
- [12] A. F. Nikiforov, S. K. Suslov, and V. B. Uvarov, Classical orthogonal polynomials of a discrete variable on nonuniform grids. *Soviet Math. Dokl.* **34** (1987), no. 3, 576-579.
- [13] A. F. Nikiforov, S. K. Suslov, and V. B. Uvarov, *Classical Orthogonal Polynomials of a Discrete Variable*. Springer Series in Computational Physics. Springer-Verlag, Berlin, 1991.
- [14] A.F. Nikiforov and V.B. Uvarov, *Special Functions of Mathematical Physics*. Birkhäuser, Basel, 1988.
- [15] A. F. Nikiforov and V. B. Uvarov, Polynomial Solutions of hypergeometric type difference Equations and their classification. *Integral Transform. Spec. Funct.* **1** (1993), 223-249.
- [16] S. K. Suslov, The theory of difference analogues of special functions of hypergeometric type. *Uspekhi Mat. Nauk.* **44:2** (1989), 185-226). (Russian Math. Survey **44:2** (1989), 227-278.)
- [17] N. Ja. Vilenkin y A.U. Klimyk, *Representations of Lie Groups and Special Functions. Vol. I, II, III*. Kluwer Academic Publishers. Dordrecht, 1992.
- [18] R.J. Yañez, J.S. Dehesa, and A.F. Nikiforov, The three-term recurrence relations and the differentiation formulas for functions of hypergeometric type. *J. Math. Anal. Appl.* **185** 1994, 855-866.

IMUS & DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, FACULTAD DE MATEMÁTICA, UNIVERSIDAD DE SEVILLA. APDO. POSTAL 1160, SEVILLA, E-41080, SEVILLA, SPAIN

E-mail address: address ran@us.es

CM-UTAD & DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE DE TRÁS-OS-MONTES E ALTO DOURO. APARTADO 202, 5001 - 911 VILA REAL, PORTUGAL

E-mail address: jluis@utad.pt