

BRUNN-MINKOWSKI AND ZHANG INEQUALITIES FOR CONVOLUTION BODIES

DAVID ALONSO-GUTIÉRREZ*, C. HUGO JIMÉNEZ^{†‡}, AND RAFAEL VILLA[†]

ABSTRACT. A quantitative version of Minkowski sum, extending the definition of θ -convolution of convex bodies, is studied to obtain extensions of the Brunn-Minkowski and Zhang inequalities, as well as, other interesting properties on Convex Geometry involving convolution bodies or polar projection bodies. The extension of this new version to more than two sets is also given.

Zhang inequality, Brunn-Minkowski inequality, Convolution Body.

1. INTRODUCTION AND MOTIVATION

The Minkowski sum of two sets $A, B \subseteq \mathbb{R}^n$ is defined as the set

$$A + B = \{x \in \mathbb{R}^n : A \cap (x - B) \neq \emptyset\}.$$

The essential sum in terms of measure is defined as

$$A +_e B = \{x \in \mathbb{R}^n : |A \cap (x - B)| > 0\},$$

for $A, B \subseteq \mathbb{R}^n$ measurable sets, when $|\cdot|$ denotes the Lebesgue measure in \mathbb{R}^n . A quantitative version of this definition, involving the proportional measure of the intersections, gives the following subset of $A + B$

$$A +_\theta B = \{x \in A + B : |A \cap (x - B)| \geq \theta M(A, B)\}$$

for $\theta \in [0, 1]$, whenever $M(A, B) := \sup_{x \in A+B} |A \cap (x - B)|$ is finite. This set is called the θ -convolution set of A and B . Note that $A +_0 B$ is the usual Minkowski sum $A + B$. This set is studied for symmetric convex bodies¹ in [1, 2, 3, 4] and [5], where the term *convolution body* is first introduced. However, our notation differs from the one used there, in order to emphasize the connection with the standard Minkowski sum. Properties of θ -convolution bodies are given in Section 2.

[†] Authors partially supported by the Spanish Ministerio de Economía y Competitividad grant MTM2012-30748 and by the Junta de Andalucía, grant P08-FQM-03543.

[‡] Author partially supported by CONACyT.

* Spanish grant MTM2010-16679 and "Programa de Ayudas a Grupos de Excelencia de la región de Murcia", Fundación Séneca, 04540/GERM/06.

¹A convex body is a compact convex subset of \mathbb{R}^n with non-empty interior.

Recently, there is an increasing interest in finding extensions of the classical integral geometry of the motion group in Euclidean spaces to the group of translations (see [6] and the references therein), motivated by possible applications to the stochastic geometry of homogeneous random geometric structures.

Our purpose is to find volume estimates, from above and below, of the θ -convolution of two sets. In what follows we will motivate our interest in studying the volume of this family of sets.

The celebrated Brunn-Minkowski inequality states that

$$|K + L|^{\frac{1}{n}} \geq |K|^{\frac{1}{n}} + |L|^{\frac{1}{n}}$$

for two convex bodies $K, L \subset \mathbb{R}^n$, with equality if and only if K and L are homothetic. Recall that K and L are called *homothetic* if $L = z + \lambda K$ for some $z \in \mathbb{R}^n$ and $\lambda > 0$. Two classical references in this topic are [7] and [8].

Brunn-Minkowski inequality has been widely applied to solve a large number of problems involving geometrical quantities such as volume, surface area, and mean width. In the last thirty years the Brunn-Minkowski inequality has become an essential analytical tool to develop the so-called Local Theory of Normed Spaces and Convex Geometric Analysis [9, 8, 7, 10]. Extensions of this inequality for non convex sets, even for non-measurable sets, have been also studied. We refer only to [11] for details and references.

In Section 3 a generalization of the Brunn-Minkowski inequality is studied. Even though extensive work with this inequality as backbone has emerged both within the class of convex bodies [12, 13, 14] and under other settings [15, 16, 17, 18], we pursue something closer in spirit to [19, 20]. See [21] for a comprehensive survey on the Brunn-Minkowski inequality including extensions, applications and its relation to other analytical inequalities.

Namely, we pose the problem of finding the best function $\varphi_n(\theta)$ such that

$$(1.1) \quad |K +_{\theta} L|^{\frac{1}{n}} \geq \varphi_n(\theta)^{\frac{1}{n}} (|K|^{\frac{1}{n}} + |L|^{\frac{1}{n}})$$

for any convex bodies $K, L \subset \mathbb{R}^n$. It is proved that $\varphi_n(\theta) = (1 - \theta^{\frac{1}{n}})^n$ satisfies (1.1). Some particular cases are also studied.

Following the work of Kiener [1], Schmuckensläger [2] proved that for any convex body K of volume 1,

$$(1.2) \quad (1 - \theta)\Pi^*(K) \subseteq K +_{\theta} (-K) \subseteq \log \frac{1}{\theta} \Pi^*(K)$$

where $\Pi^*(K)$ is the polar projection body of K , the unit ball of the norm $\|x\|_{\Pi^*(K)} = |x| |P_{x^\perp} K|$. Here P_{x^\perp} denotes the orthogonal projection on the hyperplane orthogonal to x .

These inclusions imply $|K| \Pi^*(K) = \lim_{\theta \rightarrow 1^-} \frac{|K +_{\theta} (-K)|}{1 - \theta}$ in the Hausdorff metric for any convex body K .

In Section 4 we modify the argument to improve the estimate (1.2) (see Proposition 4.1)

$$(1 - \theta)|K|\Pi^*(K) \subseteq K +_\theta(-K) \subseteq n(1 - \theta^{\frac{1}{n}})|K|\Pi^*(K).$$

The most famous inequality concerning the volume of the polar projection body of a convex body $K \subset \mathbb{R}^n$ is Petty projection inequality,

$$|K|^{n-1}|\Pi^*(K)| \leq \left(\frac{\omega_n}{\omega_{n-1}}\right)^n$$

where ω_n denotes the volume of the n -dimensional Euclidean ball. The equality is attained provided K is an ellipsoid (See [22]). A different proof using convolutions can be found in [4].

In [23], Zhang proved a reverse form of this inequality

$$(1.3) \quad |K|^{n-1}|\Pi^*(K)| \geq \frac{1}{n^n} \binom{2n}{n}$$

for any convex body, with equality if and only if K is a simplex. Zhang inequality can be written as

$$(1.4) \quad \left| \lim_{\theta \rightarrow 1^-} \frac{K +_\theta(-K)}{1 - \theta} \right| \geq \frac{1}{n^n} \binom{2n}{n} |K|.$$

It is worth mentioning that Tsolomitis studies in [5] the behavior of limiting convolution bodies

$$(1.5) \quad \lim_{\theta \rightarrow 1^-} \frac{K +_\theta L}{(1 - \theta)^\alpha}$$

for symmetric convex bodies K and L , and some exponent α , giving some regularity conditions under which (for some specific α) the limit (1.5) is non-degenerated, denoted by $C(K, L)$.

In [24], Rogers and Shephard obtained the inequality

$$(1.6) \quad |K - K| \leq \binom{2n}{n} |K|$$

for any convex body K , with equality if and only if K is a simplex. Throughout the proof it is showed that

$$(1.7) \quad K +_\theta(-K) \supseteq (1 - \theta^{\frac{1}{n}})(K - K)$$

with equality if and only if K is a simplex. They also showed in [25] the extension for two different convex bodies

$$(1.8) \quad |K - L||K \cap L| \leq \binom{2n}{n} |K||L|.$$

The last part of Section 4 is devoted to generalize the inclusions stated in (1.2) and Zhang inequality (1.3) for limiting convolutions of different convex bodies. This generalization is a consequence of Corollary 2.5, from which (1.8) can be obtained (see Proposition 4.7).

Given that some classical geometric inequalities are recovered and extended through the use of convolution bodies, it would be natural to consider the extension of the convolution body of more than two bodies in order to generalize these inequalities. In Section 5 we study such an extension obtaining similar inequalities when we consider more than two bodies. Surprisingly, it turns out when studying the equality cases, that these inequalities can only be sharp when convoluting two convex bodies and not when considering three or more bodies.

2. PROPERTIES OF THE θ -CONVOLUTION OF CONVEX BODIES

In this section we give some properties of the θ -convolution of two convex bodies, from which the Brunn-Minkowski-type inequality for the θ -convolution of convex bodies $|K +_{\theta} L|^{\frac{1}{n}} \geq (1 - \theta^{\frac{1}{n}}) \left(|K|^{\frac{1}{n}} + |L|^{\frac{1}{n}} \right)$ will follow. However, this bound is not sharp, as we will see below.

We now list some basic properties of $M(K, L)$ and the θ -convolution in the following

Proposition 2.1. *Let $K, L \subset \mathbb{R}^n$ be compact sets, $\lambda \geq 0$, $x \in \mathbb{R}^n$ and $T \in GL_n(\mathbb{R}) = \{T : \mathbb{R}^n \rightarrow \mathbb{R}^n : T \text{ is linear}\}$.*

- (a1) $M(K, L) = M(L, K)$.
- (a2) $M(x + K, L) = M(K, L)$.
- (a3) $M(\lambda K, \lambda L) = \lambda^n M(K, L)$.
- (a4) $M(TK, TL) = |\det T| M(K, L)$.
- (a5) *If $K = -L$ or K, L are symmetric, then $M(K, L) = |K \cap (-L)|$.*

Let $\theta \in [0, 1]$.

- (b1) $(\lambda K) +_{\theta} (\lambda L) = \lambda(K +_{\theta} L)$.
- (b2) $K +_{\theta} L = L +_{\theta} K$.
- (b3) $(x + K) +_{\theta} L = x + (K +_{\theta} L)$.
- (b4) $TK +_{\theta} TL = T(K +_{\theta} L)$.

A first question about this θ -convolution is its convexity, provided that K and L are both convex. The affirmative answer is a consequence of the following result. In what follows, using (b3) in Proposition 2.1 above, we will assume without loss of generality, that

$$(2.1) \quad M(K, L) = |K \cap (-L)|.$$

Proposition 2.2. *Let $K, L \subset \mathbb{R}^n$ be convex bodies satisfying (2.1). Then for every $\theta_1, \theta_2, \lambda_1, \lambda_2 \in [0, 1]$ such that $\lambda_1 + \lambda_2 \leq 1$ we have that*

$$(2.2) \quad \lambda_1(K +_{\theta_1} L) + \lambda_2(K +_{\theta_2} L) \subseteq K +_{\theta} L,$$

where $1 - \theta^{\frac{1}{n}} = \lambda_1(1 - \theta_1^{\frac{1}{n}}) + \lambda_2(1 - \theta_2^{\frac{1}{n}})$.

Proof. Let $x_1 \in K +_{\theta_1} L$ and $x_2 \in K +_{\theta_2} L$. From the general inclusion

$$K \cap (\lambda_0 A_0 + \lambda_1 A_1 + \lambda_2 A_2) \supseteq \lambda_0 K \cap A_0 + \lambda_1 K \cap A_1 + \lambda_2 K \cap A_2$$

where K is convex and $\lambda_0 + \lambda_1 + \lambda_2 = 1$, and using the convexity of L , we have

(2.3)

$$K \cap (\lambda_1 x_1 + \lambda_2 x_2 - L) \supseteq (1 - \lambda_1 - \lambda_2)[K \cap (-L)] + \lambda_1 [K \cap (x_1 - L)] + \lambda_2 [K \cap (x_2 - L)].$$

Taking volumes, using the classical Brunn-Minkowski inequality and the fact that $x_i \in K +_{\theta_i} L$ we have

$$(2.4) \quad |K \cap (\lambda_1 x_1 + \lambda_2 x_2 - L)| \geq [1 - \lambda_1(1 - \theta_1^{\frac{1}{n}}) - \lambda_2(1 - \theta_2^{\frac{1}{n}})]^n M(K, L),$$

which proves that $\lambda_1 x_1 + \lambda_2 x_2 \in K +_{\theta} L$ for $\theta = [1 - \lambda_1(1 - \theta_1^{\frac{1}{n}}) - \lambda_2(1 - \theta_2^{\frac{1}{n}})]^n$. \square

Taking $\theta_1 = \theta_2$ and $\lambda_2 = 1 - \lambda_1$ we have

Corollary 2.3. *Let $K, L \subset \mathbb{R}^n$ be convex bodies and $\theta \in [0, 1]$. Then $K +_{\theta} L$ is convex.*

The following result on $K +_1 L$ will be used later on, and it is a consequence of Proposition 2.2.

Corollary 2.4. *Let $K, L \subset \mathbb{R}^n$ be convex bodies. For any $x \in K +_1 L$, $K \cap (x - L)$ is a translation of $K \cap (-L)$.*

Proof. Taking $\theta_1 = \theta_2 = 1$ in Proposition 2.2, we get $\theta = 1$, and then Brunn-Minkowski inequality (2.4), obtained from (2.3), holds with equality. Then (see equality cases in Brunn-Minkowski inequality in [26]) the four sets involved in (2.3),

$$K \cap (\lambda_1 x_1 + \lambda_2 x_2 - L), \quad K \cap (-L), \quad K \cap (x_1 - L), \quad K \cap (x_2 - L),$$

are all homothetic. As they all have the same volume (equals to $M(K, L)$), homotheties are indeed translations. \square

Taking $\theta_1 = \theta_2$ leads us to the following

Corollary 2.5. *Let $K, L \subset \mathbb{R}^n$ be convex bodies satisfying (2.1). Then, for every $0 \leq \theta_0 \leq \theta < 1$ we have*

$$\frac{K +_{\theta_0} L}{1 - \theta_0^{\frac{1}{n}}} \subseteq \frac{K +_{\theta} L}{1 - \theta^{\frac{1}{n}}}.$$

Proof. Taking $\theta_1 = \theta_2 = \theta_0$ in the above proposition, for any $\lambda_1, \lambda_2 \in [0, 1]$ such that $\lambda_1 + \lambda_2 \leq 1$

$$\lambda_1(K +_{\theta_0} L) + \lambda_2(K +_{\theta_0} L) \subseteq (\lambda_1 + \lambda_2)(K +_{\theta_0} L) \subseteq K +_{\theta} L,$$

with $1 - \theta^{\frac{1}{n}} = (\lambda_1 + \lambda_2)(1 - \theta_0^{\frac{1}{n}})$. Since $\lambda_1 + \lambda_2 = \frac{1 - \theta^{\frac{1}{n}}}{1 - \theta_0^{\frac{1}{n}}}$,

$$\frac{1 - \theta^{\frac{1}{n}}}{1 - \theta_0^{\frac{1}{n}}}(K +_{\theta_0} L) \subseteq K +_{\theta} L$$

whenever $\lambda_1 + \lambda_2 \leq 1$, which means $0 \leq \theta_0 \leq \theta \leq 1$. \square

The following extension of (1.7) from Rogers and Shephard's work will be used to get a first Brunn-Minkowski-type inequality.

Corollary 2.6. *Let $K, L \subset \mathbb{R}^n$ be convex bodies. Then*

$$(2.5) \quad \theta^{\frac{1}{n}}(K +_1 L) + (1 - \theta^{\frac{1}{n}})(K + L) \subseteq K +_{\theta} L.$$

Proof. Take $\theta_1 = 1$, $\theta_2 = 0$, $\lambda_1 = \theta^{\frac{1}{n}}$, and $\lambda_2 = (1 - \theta^{\frac{1}{n}})$ in Proposition 2.2 to obtain the desired result. \square

Note that the condition $0 \in K +_1 L$ is equivalent to $M(K, L) = |K \cap (-L)|$, and that is verified under the assumptions of Proposition 2.1 (a5).

We will use the following description of the boundary of $K +_{\theta} L$.

Lemma 2.7. *Let $K, L \subset \mathbb{R}^n$ be convex bodies, and $\theta \in [0, 1)$. Then*

$$\partial(K +_{\theta} L) = \{x \in K + L : |K \cap (x - L)| = \theta M(K, L)\}.$$

In particular, for $\theta = 0$,

$$\partial(K + L) = \{x \in K + L : |K \cap (x - L)| = 0\}.$$

Consequently, for any $x \in K + L$, $x \notin K +_1 L$, there exists a unique $\theta \in [0, 1)$ such that $x \in \partial(K +_{\theta} L)$.

Proof. Let $f : \mathbb{R}^n \rightarrow [0, +\infty)$ be given by $f(x) = |K \cap (x - L)|^{\frac{1}{n}}$. Using properties of the Lebesgue measure and Brunn-Minkowski inequality, it can be deduced that f is continuous on \mathbb{R}^n and concave on $K + L$. It is then clear that $K +_{\theta} L$ is closed.

The first assertion is equivalent to $\text{int}(K +_{\theta} L) = \{x \in K + L : f(x)^n > \theta M(K, L)\}$. Since the right-hand set is open and it is contained in $K +_{\theta} L$, it remains to be shown the inclusion $\text{int}(K +_{\theta} L) \subseteq \{x \in K + L : f(x)^n > \theta M(K, L)\}$.

Let us take any $x \in K + L$, with $f(x)^n = \theta M(K, L)$. It is left to show that $x \notin \text{int}(K +_{\theta} L)$.

We will assume without loss of generality that (2.1) is satisfied. For any $\lambda \in (0, 1)$, write $x = \lambda(\lambda^{-1}x) + (1 - \lambda) \cdot 0$. If $\lambda^{-1}x \in K + L$, using the concavity of f on $K + L$,

$$(\theta M(K, L))^{\frac{1}{n}} = f(x) \geq \lambda f(\lambda^{-1}x) + (1 - \lambda)f(0) = \lambda f(\lambda^{-1}x) + (1 - \lambda^{-1})M(K, L)^{\frac{1}{n}}$$

and therefore

$$f(\lambda^{-1}x)^n \leq \frac{\theta^{\frac{1}{n}} - (1 - \lambda^{-1})}{\lambda} M(K, L)^{\frac{1}{n}} < (\theta M(K, L))^{\frac{1}{n}}.$$

Then $\lambda^{-1}x \notin K +_{\theta} L$ for any $\lambda < 1$ (for $\lambda^{-1}x \notin K + L$ it is trivial). Hence $x \notin \text{int}(K +_{\theta} L)$.

To obtain the last assertion, just take $\theta = \frac{|K \cap (x - L)|}{M(K, L)} \in [0, 1)$. \square

Equality cases in Proposition 2.2 and Corollary 2.6 are stated in the following result.

Proposition 2.8. *Let $K, L \subset \mathbb{R}^n$ be convex bodies. The following conditions are equivalent.*

(i) *For every $\theta_1, \theta_2, \lambda_1, \lambda_2 \in [0, 1]$ such that $\lambda_1 + \lambda_2 = 1$, we have*

$$\lambda_1(K +_{\theta_1} L) + \lambda_2(K +_{\theta_2} L) = K +_{\theta} L,$$

$$\text{where } \theta^{\frac{1}{n}} = \lambda_1 \theta_1^{\frac{1}{n}} + \lambda_2 \theta_2^{\frac{1}{n}}.$$

(ii) *For every $\theta \in [0, 1]$, $\theta^{\frac{1}{n}}(K +_1 L) + (1 - \theta^{\frac{1}{n}})(K + L) = K +_{\theta} L$.*

(iii) *K and $-L$ are homothetic n -simplices.*

For the proof, we will use the following result from [27] (we state it here in the form it will be used in this paper).

Theorem 2.9. (Soltan) *Let $K, L \subset \mathbb{R}^n$ be convex bodies. The following conditions are equivalent:*

(i) *K and L are homothetic n -simplices.*

(ii) *The n -dimensional intersections $K \cap (z - L)$, $z \in \mathbb{R}^n$, are all homothetic to $K \cap (-L)$.*

Proof of Proposition 2.8. Using a translation we may assume that (2.1) is satisfied.

Conditions (i) and (ii) are equivalent. Indeed, (i) trivially implies (ii).

Suppose (ii) holds. Then, using the equality for θ_1 , θ_2 and θ successively, and the convexity of the convolution sets,

$$\begin{aligned} & \lambda_1(K +_{\theta_1} L) + \lambda_2(K +_{\theta_2} L) \\ &= \lambda_1 \left(\theta_1^{\frac{1}{n}}(K +_1 L) + (1 - \theta_1^{\frac{1}{n}})(K + L) \right) \\ & \quad + \lambda_2 \left(\theta_2^{\frac{1}{n}}(K +_1 L) + (1 - \theta_2^{\frac{1}{n}})(K + L) \right) \\ &= (\lambda_1 \theta_1^{\frac{1}{n}} + \lambda_2 \theta_2^{\frac{1}{n}})(K +_1 L) + (\lambda_1(1 - \theta_1^{\frac{1}{n}}) + \lambda_2(1 - \theta_2^{\frac{1}{n}}))(K + L) \\ &= K +_{\theta} L, \end{aligned}$$

and (i) follows.

Suppose (ii) holds, and take $x \in K + L$, $x \notin \partial(K + L)$. If $x \in K +_1 L$, Corollary 2.4 shows that $K \cap (x - L)$ is a translation of (so homothetic to) $K \cap (-L)$.

If $x \notin K +_1 L$, by Lemma 2.7, there is a $\theta \in (0, 1)$ such that $x \in \partial(K +_\theta L)$. Using (ii), $x = \theta^{\frac{1}{n}}x_1 + (1 - \theta^{\frac{1}{n}})x_2$, for some $x_1 \in K +_1 L$, $x_2 \in K + L$.

Now, inclusion (2.3) and inequality (2.4) are both equalities (since $|K \cap (x - L)| = \theta M(K, L)$), and then $K \cap (x - L)$, $K \cap (x_1 - L)$ and $K \cap (x_2 - L)$ are all homothetic (see equality cases in Brunn-Minkowski inequality in [26]). Since $x_1 \in K +_1 L$, they are all homothetic to $K \cap (-L)$.

Then, all the n -dimensional intersections $K \cap (x - L)$ are homothetic to the same body $K \cap (-L)$. Now it follows from Theorem 2.9 that K , $-L$ and $K \cap (-L)$ are homothetic simplices.

Now suppose condition (iii) holds. After an affine transformation we may assume that

$$K = \left\{ t \in \mathbb{R}^n : t_i \geq 0, \sum_{i=1}^n t_i \leq 1 \right\}$$

and $-L = \lambda K$ with $0 < \lambda \leq 1$. For any $x \in \mathbb{R}^n$,

$$K \cap (x - L) = \left\{ t \in \mathbb{R}^n : t_j \geq x_j^+, \sum_{i=1}^n t_i \leq \min\{1, \lambda + \sum_{i=1}^n x_i\} \right\},$$

where $r^+ = \max\{r, 0\}$. Then, $K \cap (x - L) = z(x) + \lambda(x)K$, where $z_j(x) = x_j^+$, and

$$\lambda(x) = \min\{1, \lambda + \sum_{i=1}^n x_i\} - \sum_{i=1}^n x_i^+$$

whenever $\lambda(x) \geq 0$. For those $x \in \mathbb{R}^n$, we have $|K \cap (x - L)| = \lambda(x)^n |K|$. It is easy to see that $\lambda(x) \leq \lambda$, and equality holds if and only if $x_j \geq 0$ for all j and $\lambda + \sum_{i=1}^n x_i \leq 1$. Then $M(K, L) = \lambda^n |K|$, and $K +_1 L = (1 - \lambda)K$.

Using absolute values, $\lambda(x)$ can be rewritten as

$$\lambda(x) = \frac{1}{2} \left(1 + \lambda - \left| 1 - \lambda - \sum_{i=1}^n x_i \right| - \sum_{i=1}^n |x_i| \right).$$

Then

$$K +_\theta L = \left\{ x \in \mathbb{R}^n : \left| 1 - \lambda - \sum_{i=1}^n x_i \right| + \sum_{i=1}^n |x_i| \leq 1 + \lambda(1 - 2\theta^{\frac{1}{n}}) \right\}.$$

In particular, letting $\theta = 0$, we obtain an expression for $K + L$. In order to prove (ii), it is enough to prove the inclusion $K +_\theta L \subseteq \theta^{\frac{1}{n}}(K +_1 L) + (1 - \theta^{\frac{1}{n}})(K + L)$ for every $\theta \in (0, 1)$. Any $x \in K +_\theta L$ satisfies

$$(2.6) \quad \left| 1 - \lambda - \sum_{i=1}^n x_i \right| + \sum_{i=1}^n |x_i| \leq \theta^{\frac{1}{n}}(1 - \lambda) + (1 - \theta^{\frac{1}{n}})(1 + \lambda).$$

Recall that $\left|1 - \lambda - \sum_{i=1}^n a_i\right| + \sum_{i=1}^n |a_i| = 1 - \lambda$ for any $a \in K +_1 L$. Consequently, if

$$(2.7) \quad x = \theta^{\frac{1}{n}} a + (1 - \theta^{\frac{1}{n}}) b,$$

with $a \in K +_1 L$, then $b \in K + L$ provided that the left hand side in (2.6)

$$\left| \theta^{\frac{1}{n}} (1 - \lambda - \sum_{i=1}^n a_i) + (1 - \theta^{\frac{1}{n}}) (1 - \lambda - \sum_{i=1}^n b_i) \right| + \sum_{i=1}^n |\theta^{\frac{1}{n}} a_i + (1 - \theta^{\frac{1}{n}}) b_i|$$

equals

$$\theta^{\frac{1}{n}} \left| 1 - \lambda - \sum_{i=1}^n a_i \right| + (1 - \theta^{\frac{1}{n}}) \left| 1 - \lambda - \sum_{i=1}^n b_i \right| + \theta^{\frac{1}{n}} \sum_{i=1}^n |a_i| + (1 - \theta^{\frac{1}{n}}) \sum_{i=1}^n |b_i|.$$

Considering equality cases in triangle inequality, this happens provided that $1 - \lambda - \sum_{i=1}^n a_i$ has the same sign as $1 - \lambda - \sum_{i=1}^n b_i$, and for any j , a_j has the same sign as b_j (here r, s have the same sign iff $r \cdot s \geq 0$).

If $\sum_{i=1}^n x_i^+ \leq 1 - \lambda$, it is enough to consider $a \in K +_1 L$ in (2.7), so that $a_j = x_j^+$. Then $b_j = (1 - \theta^{\frac{1}{n}})^{-1} x_j$ if $x_j < 0$ and $b_j = x_j$ if $x_j \geq 0$. In any case, b_j has the same sign as a_j , and therefore $b \in K + L$.

If $\sum_{i=1}^n x_i^+ > 1 - \lambda$, it is enough to take $a \in K +_1 L$ in (2.7), so that $a_j = (1 - \lambda) \frac{x_j^+}{\sum_{i=1}^n x_i^+}$. Then $b \in K + L$ provided that $b_j \geq 0$ for those j so that $x_j > 0$. But in that case

$$x_j = \theta^{\frac{1}{n}} \frac{(1 - \lambda)}{\sum_{i=1}^n x_i^+} x_j + (1 - \theta^{\frac{1}{n}}) b_j$$

Then $b_j \geq 0$ provided that

$$\theta^{\frac{1}{n}} \frac{(1 - \lambda)}{\sum_{i=1}^n x_i^+} \leq 1,$$

which is assumed to be true. In any case, we can find a decomposition as (2.7), which proves (ii). \square

Now we can deduce equality cases in Corollary 2.5.

Proposition 2.10. *Let $K, L \subset \mathbb{R}^n$ be convex bodies satisfying (2.1). The following conditions are equivalent.*

- (i) For any $0 \leq \theta_0 \leq \theta \leq 1$, $(1 - \theta^{\frac{1}{n}})(K +_{\theta_0} L) = (1 - \theta_0^{\frac{1}{n}})(K +_{\theta} L)$.
- (ii) For every $\theta \in [0, 1]$, $(1 - \theta^{\frac{1}{n}})(K + L) = K +_{\theta} L$.
- (iii) $K = -L$ is an n -simplex.

Proof. Again, conditions (i) and (ii) are easily seen to be equivalent.

Now, suppose (ii) holds; then $K +_1 L = \{0\}$, and consequently (ii) in Proposition 2.8 holds. Then K and $-L$ are homothetic n -simplices.

As in the proof of Proposition 2.8, we have $K +_1 L = (1 - \lambda)K$, which implies $\lambda = 1$, since $K +_1 L = \{0\}$. Consequently, $K = -L$.

Condition (iii) implies (ii) in Proposition 2.8, and since $K +_1 L = \{0\}$, we get (ii). \square

3. BRUNN-MINKOWSKI TYPE INEQUALITY FOR θ -CONVOLUTION BODIES

From the previous study on convolution of two sets, the following natural question arises: what kind of Brunn-Minkowski-type inequality for θ -convolutions

$$(3.1) \quad |K +_{\theta} L|^{\frac{1}{n}} \geq \varphi_n(\theta)^{\frac{1}{n}} (|K|^{\frac{1}{n}} + |L|^{\frac{1}{n}}).$$

does it hold?

As in the classical case, the homogeneity allows one to formulate the inequality in different equivalent forms.

Proposition 3.1. *The following statements are all equivalent:*

(i) For K, L measurable sets in \mathbb{R}^n

$$|K +_{\theta} L|^{\frac{1}{n}} \geq \varphi_n(\theta)^{\frac{1}{n}} (|K|^{\frac{1}{n}} + |L|^{\frac{1}{n}}).$$

(ii) For K, L measurable sets in \mathbb{R}^n and $0 < \lambda < 1$

$$|\lambda K +_{\theta} (1 - \lambda)L|^{\frac{1}{n}} \geq \varphi_n(\theta)^{\frac{1}{n}} (\lambda |K|^{\frac{1}{n}} + (1 - \lambda) |L|^{\frac{1}{n}}).$$

(iii) For K, L measurable sets in \mathbb{R}^n and $0 < \lambda < 1$

$$|\lambda K +_{\theta} (1 - \lambda)L| \geq \varphi_n(\theta) (|K|^{\lambda} \cdot |L|^{1-\lambda}).$$

(iv) For K, L measurable sets in \mathbb{R}^n and $0 < \lambda < 1$

$$|\lambda K +_{\theta} (1 - \lambda)L| \geq \varphi_n(\theta) \min\{|K|, |L|\}.$$

(v) For K, L measurable sets in \mathbb{R}^n such that $|K| = |L| = 1$ and $0 < \lambda < 1$

$$|\lambda K +_{\theta} (1 - \lambda)L| \geq \varphi_n(\theta).$$

Proof. (i) \rightarrow (ii) and (iii) \rightarrow (iv) \rightarrow (v) are immediate. The proof of (ii) \rightarrow (iii) is obtained by taking logarithm and using its concavity.

Finally, apply (v) with $\overline{K} = |K|^{-\frac{1}{n}}K$, $\overline{L} = |L|^{-\frac{1}{n}}L$ and $\lambda = \frac{s|K|^{\frac{1}{n}}}{s|K|^{\frac{1}{n}} + t|L|^{\frac{1}{n}}}$, and use the homogeneity of the convolution (Proposition 2.1 (b1)) to get (i). \square

A first inequality in this direction for convex bodies is obtained from Corollary 2.6.

Corollary 3.2. *Let $K, L \subset \mathbb{R}^n$ be convex bodies. Then*

$$|K +_{\theta} L|^{\frac{1}{n}} \geq (1 - \theta^{\frac{1}{n}})(|K|^{\frac{1}{n}} + |L|^{\frac{1}{n}}).$$

Equivalently, $\varphi_n(\theta) \geq (1 - \theta^{\frac{1}{n}})^n$ in (3.1).

Proof. Taking volumes in (2.5)

$$(3.2) \quad |K +_{\theta} L|^{\frac{1}{n}} \geq (1 - \theta^{\frac{1}{n}})|K + L|^{\frac{1}{n}}$$

and applying Brunn-Minkowski inequality

$$(3.3) \quad (1 - \theta^{\frac{1}{n}})|K + L|^{\frac{1}{n}} \geq (1 - \theta^{\frac{1}{n}})(|K|^{\frac{1}{n}} + |L|^{\frac{1}{n}})$$

we obtain the desired result. \square

In order to have equality in Corollary 3.2, we need to have equality in (3.2) and in Brunn-Minkowski inequality (3.3). However, by Proposition 2.10, equality in (3.2) holds if and only if $K = -L$ is an n -dimensional simplex, and in that case there is not equality in Brunn-Minkowski inequality (unless $n = 1$). See examples at the end of the section for details.

The following result improves the inclusion

$$(3.4) \quad (1 - \theta^{\frac{1}{n}})(K + L) \subseteq K +_{\theta} L$$

providing a new set between them. A good estimate for the volume of this new set would lead to a better estimate for $|K +_{\theta} L|$.

Theorem 3.3. *Let $K, L \subset \mathbb{R}^n$ be convex bodies such that $M(K, L) = |K \cap (-L)|$. Then for all $\theta \in [0, 1]$,*

$$\begin{aligned} K +_{\theta} L &\supseteq \left\{ a + b : a \in K, b \in L, \frac{|(1 - \|a\|_K)K \cap (1 - \|b\|_L)(-L)|}{|K \cap (-L)|} \geq \theta \right\} \\ &\supseteq (1 - \theta^{\frac{1}{n}})(K + L). \end{aligned}$$

Proof. Let $x \in K + L$, then $x = a + b$ with $a \in K$ and $b \in L$. From the convexity of K

$$(1 - \|a\|_K)K + \|a\|_K \frac{a}{\|a\|_K} \subseteq K.$$

Also, since $\| -b \|_{-L} = \|b\|_L$ and L is convex

$$(1 - \|b\|_L)(-L) + \|b\|_L \frac{-b}{\|b\|_L} + x \subseteq x - L.$$

Since $x - b = a$, we have $(1 - \|a\|_K)K + a \subseteq K$ and $(1 - \|b\|_L)(-L) + a \subseteq x - L$. Thus,

$$a + (1 - \|a\|_K)K \cap (1 - \|b\|_L)(-L) \subseteq K \cap (x - L)$$

and then $|K \cap (x - L)| \geq |(1 - \|a\|_K)K \cap (1 - \|b\|_L)(-L)|$. Consequently,

$$K +_{\theta} L \supseteq \left\{ a + b : a \in K, b \in L, \frac{|(1 - \|a\|_K)K \cap (1 - \|b\|_L)(-L)|}{|K \cap (-L)|} \geq \theta \right\}.$$

This set trivially contains the set

$$\{a + b : \inf\{(1 - \|a\|_K)^n, (1 - \|b\|_L)^n\} \geq \theta\} = (1 - \theta^{\frac{1}{n}})(K + L). \quad \square$$

In order to get a more accurate idea of how good the bound in Corollary 3.2 is, we estimate the quotient $\frac{|K +_\theta L|^{\frac{1}{n}}}{|K|^{\frac{1}{n}} + |L|^{\frac{1}{n}}}$ for some particular pairs of bodies.

Examples:

- 1) For K, L cubes whose sides are parallel to the coordinate hyperplanes, it is not hard to see that the quotient is minimized when $K = L = [-1/2, 1/2]^n$, and its value equals

$$\left[1 - \theta \sum_{k=0}^{n-1} \frac{(-\log \theta)^k}{k!} \right]^{\frac{1}{n}}.$$

- 2) For $K = L$ the unit Euclidean ball, the quotient equals $R_n(\theta)$ given by the equality

$$2\omega_{n-1} \int_{R_n(\theta)}^1 (1 - s^2)^{\frac{n-1}{2}} ds = \theta\omega_n$$

where ω_n denotes the volume of the n -dimensional unit Euclidean ball.

- 3) As it was mentioned above, in [24] it was proved that, for $K = L$ the simplex, the quotient equals

$$(1 - \theta^{\frac{1}{n}}) \frac{\binom{2n}{n}^{\frac{1}{n}}}{2} \sim 2(1 - \theta^{\frac{1}{n}}).$$

Comparing these three cases, it seems that the minimum value for the quotient is attained in a different case depending on θ . This fact makes difficult to find a family of bodies in which the minimum is attained.

4. A CONNECTION WITH PROJECTION BODIES AND ZHANG INEQUALITY

This section is devoted to generalize the inclusions (1.2) and Zhang inequality for convolution of different convex bodies.

The following result generalizes the right hand side inclusion in (1.2). We extend the ideas used in [2]

Proposition 4.1. *Let $K, L \subset \mathbb{R}^n$ be convex bodies satisfying (2.1). Then, for every $\theta \in (0, 1)$*

$$K +_\theta L \subseteq \left\{ x \in \mathbb{R}^n : |x| \left| \frac{d^+}{dt} \left| K \cap \left(t \frac{x}{|x|} - L \right) \right|_{t=0} \right| \leq n(1 - \theta^{\frac{1}{n}})M(K, L) \right\}.$$

Proof. The concavity of the function $x \mapsto |K \cap (x - L)|^{\frac{1}{n}}$ implies

$$\begin{aligned} |K \cap (\lambda x - L)| &\geq \left((1 - \lambda)M(K, L)^{\frac{1}{n}} + \lambda|K \cap (x - L)|^{\frac{1}{n}} \right)^n \\ &= M(K, L) \left[1 + \lambda \left(\frac{|K \cap (x - L)|^{\frac{1}{n}}}{M(K, L)^{\frac{1}{n}}} - 1 \right) \right]^n \\ &\geq M(K, L) \left[1 + \lambda n \left(\frac{|K \cap (x - L)|^{\frac{1}{n}}}{M(K, L)^{\frac{1}{n}}} - 1 \right) \right] \end{aligned}$$

for $\lambda \in [0, 1]$ and $x \in K + L$. On the other hand,

$$\begin{aligned} |K \cap (\lambda x - L)| &= M(K, L) + \int_0^{\lambda|x|} \frac{d^+}{dt} \left| K \cap \left(t \frac{x}{|x|} - L \right) \right| dt \\ &\leq M(K, L) + \lambda|x| \max_{t \in [0, \lambda|x|]} \frac{d^+}{dt} \left| K \cap \left(t \frac{x}{|x|} - L \right) \right| \end{aligned}$$

again using the concavity of $x \mapsto |K \cap (x - L)|^{\frac{1}{n}}$. Comparing these two inequalities, and letting $\lambda \rightarrow 0^+$, we obtain

$$nM(K, L) \left(\frac{|K \cap (x - L)|^{\frac{1}{n}}}{M(K, L)^{\frac{1}{n}}} - 1 \right) \leq |x| \frac{d^+}{dt} \left| K \cap \left(t \frac{x}{|x|} - L \right) \right|_{t=0}.$$

Since the lateral derivative is non positive, we get the desired inclusion. \square

Remark 4.2.

- 1) If $L = -K$, then the right-hand side set is exactly $n(1 - \theta^{\frac{1}{n}})|K| \Pi^* K$ which improves the right hand side inclusion in (1.2).
- 2) From Corollary 2.5, the family of sets $\frac{K + \theta L}{1 - \theta^{\frac{1}{n}}}$ is increasing with respect to θ , and using the equivalence $1 - \theta \sim n(1 - \theta^{\frac{1}{n}})$, the existence of the limiting convolution set with $\alpha = 1$,

$$C_1(K, L) = \lim_{\theta \rightarrow 1^-} \frac{K + \theta L}{1 - \theta}$$

follows. However, there are cases in which this set is unbounded. We refer to Example 3.15 in [5] for a detailed construction of an example where the limiting convolution is \mathbb{R}^n . In that paper sufficient conditions for $C_1(K, L)$ to be bounded are also given.

- 3) The righten set in Proposition 4.1 is $n(1 - \theta^{\frac{1}{n}})C_1(K, L)$. The previous result can be deduced from Corollary 2.5 letting $\theta_0 \rightarrow 1^-$ (see the proof of Theorem 4.6 below).
- 4) Nevertheless, a general inclusion $K + \theta L \subseteq n(1 - \theta^{\frac{1}{n}})C$, for some body C independent from θ can not be proved, since it was shown in [5] that the limiting convolution body with $\alpha = 1$ could be non compact.

The left-hand side inclusion in (1.2) is generalized with the following Proposition. Recall that $\text{conv}(A)$ denotes the convex hull of a set A .

Proposition 4.3. *Let $K, L \subset \mathbb{R}^n$ be convex bodies satisfying (2.1). Then, for every $\theta \in (0, 1)$*

$$K +_{\theta} L \supseteq (1 - \theta)M(K, L) \text{conv}(\Pi^*K \cup \Pi^*L).$$

Proof. In [3] it is proved that, given two convex bodies K, L , and $u \in S^{n-1}$, the function $f(r) = |K \cap (ru + L)|$ verifies $f'(0) = |C_u^+(1, 2)| - |C_u^-(2, 1)|$, where

$$\begin{aligned} C_u^+(1, 2) &= P_u(K \cap L) \cap \{\psi_K^+ > \psi_L^+ \geq \psi_K^- > \psi_L^-\} \\ C_u^+(2, 1) &= P_u(K \cap L) \cap \{\psi_L^+ \geq \psi_K^+ > \psi_L^- \geq \psi_K^-\} \end{aligned}$$

with

$$\psi_K^+(y) = \max\{t : tu + y \in K\}$$

and

$$\psi_K^-(y) = \min\{t : tu + y \in K\}.$$

Thus $\frac{d}{d^v} |K \cap (x - L)| \geq -|P_v^+K \cap (x - L)| \geq -\min\{|P_v^+K|, |P_v^+L|\}$. Consequently,

$$|K \cap (x - L)| =$$

$$M(K, L) + \int_0^{|x|} \frac{d}{dt} \left| K \cap \left(t \frac{x}{|x|} - L \right) \right| \geq M(K, L) - |x| \min\{|P_v^+K|, |P_v^+L|\}.$$

Hence, if $M(K, L) - |x| \min\{|P_v^+K|, |P_v^+L|\} \geq \theta M(K, L)$ then $x \in K +_{\theta} L$ and this holds if and only if $\min\{\|x\|_{\Pi^*K}, \|x\|_{\Pi^*L}\} \leq (1 - \theta)M(K, L)$.

So, $(1 - \theta)M(K, L)(\Pi^*K \cup \Pi^*L) \subseteq K +_{\theta} L$. The convexity of the set $K +_{\theta} L$ yields the desired result. \square

Remark 4.4. Taking $L = -K$ and $|K| = 1$, we recover the left hand side inclusion in (1.2).

Remark 4.5. Applying Zhang inequality we deduce that

$$\min\{|K|^{n-1}, |L|^{n-1}\} \left| \frac{C_1(K, L)}{M(K, L)} \right| \geq \binom{2n}{n} \frac{1}{n^n}$$

which extends Zhang inequality (1.4). Nevertheless, a stronger extension of Zhang inequality can be proved using Corollary 2.5.

Theorem 4.6. *Let $K, L \subset \mathbb{R}^n$ be convex bodies such that $M(K, L) = |K \cap (-L)|$. Then*

$$(4.1) \quad |C_1(K, L)| \geq \binom{2n}{n} \frac{1}{n^n} \frac{|K||L|}{M(K, L)}$$

Equality holds if and only if $K = -L$ is a simplex.

Proof. From Corollary 2.5 we have that for every $0 \leq \theta_0 \leq \theta < 1$

$$(4.2) \quad \frac{K +_{\theta_0} L}{1 - \theta_0^{\frac{1}{n}}} \subseteq \frac{K +_{\theta} L}{1 - \theta^{\frac{1}{n}}}.$$

Thus, letting $\theta \rightarrow 1^-$ we obtain that for every $\theta_0 \in [0, 1)$

$$\frac{K +_{\theta_0} L}{1 - \theta_0^{\frac{1}{n}}} \subseteq \lim_{\theta \rightarrow 1^-} \frac{1 - \theta}{1 - \theta^{\frac{1}{n}}} \frac{K +_{\theta} L}{1 - \theta} = nC_1(K, L),$$

and taking volumes

$$|K +_{\theta_0} L| \leq n^n (1 - \theta_0^{\frac{1}{n}})^n |C_1(K, L)|$$

for $\theta_0 \in [0, 1)$. Integrating over $[0, 1]$ yields

$$\int_0^1 |K +_{\theta_0} L| d\theta_0 \leq n^n |C_1(K, L)| \int_0^1 (1 - \theta_0^{\frac{1}{n}})^n d\theta_0 = n^n |C_1(K, L)| \binom{2n}{n}^{-1}.$$

Integrating by parts and using Fubini's Theorem, the first integral equals

$$\begin{aligned} \int_0^1 |K +_{\theta_0} L| d\theta_0 &= \int_0^1 \left| \left\{ x \in K + L : \frac{|K \cap (x - L)|}{M(K, L)} \geq \theta_0 \right\} \right| d\theta_0 \\ &= \int_{K+L} \frac{|K \cap (x - L)|}{M(K, L)} dx = \frac{|K||L|}{M(K, L)} \end{aligned}$$

from which the desired inequality follows.

If equality holds in (4.1), then (4.2) holds also with equality for any $0 \leq \theta_0 \leq \theta < 1$. Letting $\theta_0 = 0$,

$$(4.3) \quad (1 - \theta^{\frac{1}{n}})(K + L) = K +_{\theta} L$$

for any $\theta \in [0, 1)$. Now Corollary 2.6 implies

$$\theta^{\frac{1}{n}}(K +_1 L) + (1 - \theta^{\frac{1}{n}})(K + L) \subseteq K +_{\theta} L = (1 - \theta^{\frac{1}{n}})(K + L).$$

A compactness argument shows that $K +_1 L = \{0\}$, so equality (4.3) holds for every $0 \leq \theta \leq 1$, and then (ii) in Proposition 2.10 implies that $K = -L$ is a simplex. \square

Finally, Corollary 2.5 allows us to recover Rogers-Shephard inequality (1.8). We also solve the problem of characterizing equality cases posed in [25].

Proposition 4.7. *Let $K, L \subset \mathbb{R}^n$ be convex bodies. Then*

$$|K + L|M(K, L) \leq \binom{2n}{n} |K||L|.$$

Equality holds if and only if $K = -L$ is a simplex.

Proof. By a translation we may assume that (2.1) is satisfied. A similar argument to that used in Theorem 4.6, taking $\theta_0 = 0$ in Corollary 2.5, applying volumes and integrating in θ , shows the desired result.

If equality holds, then (ii) in Proposition 2.10 is satisfied, and therefore $K = -L$ is a simplex. \square

5. CONVOLUTION OF m BODIES

In this section we will extend the definition of θ -convolution bodies to more than two sets. The θ -convolution is not associative (as a simple computation with Euclidean balls of different radius shows) so a definition of an m -fold convolution can not be made inductively. Nevertheless, since $|K \cap (x - L)| = \chi_K * \chi_L(x)$ and the convolution is associative, it seems natural to make the following extension of θ -convolution bodies:

Definition 5.1. Let K_1, \dots, K_m be m measurable sets in \mathbb{R}^n and let $\theta \in [0, 1]$. We define their θ -convolution as the set

$$K_1 +_{\theta} \dots +_{\theta} K_m = \{x \in K_1 + \dots + K_m : \chi_{K_1} * \dots * \chi_{K_m}(x) \geq \theta M(K_1, \dots, K_m)\}$$

when $M(K_1, \dots, K_m) = \max_{x \in \mathbb{R}^n} \chi_{K_1} * \dots * \chi_{K_m}(x)$ is finite.

For $\theta = 0$ the set $K_1 +_0 \dots +_0 K_m$ is just the support of the function $\chi_{K_1} * \dots * \chi_{K_m}$, the usual Minkowski sum $K_1 + \dots + K_m$.

The commutative and associative properties of the convolution imply trivially that

- (a1) $\chi_{K_1} * \dots * \chi_{K_m} = \chi_{K_{\sigma(1)}} * \dots * \chi_{K_{\sigma(m)}}$ for any σ permutation of $\{1, \dots, m\}$.
- (a2) $\chi_{z+K_1} * \dots * \chi_{K_m}(x) = \chi_{K_1} * \dots * \chi_{K_m}(x - z)$
- (a3) $\chi_{TK_1} * \dots * \chi_{TK_m}(x) = |\det T|^{m-1} \chi_{K_1} * \dots * \chi_{K_m}(T^{-1}x)$ for any $T \in GL_n(\mathbb{R})$.

Consequently we have the following result, analogous to Proposition 2.1.

Proposition 5.2. Let K_1, \dots, K_m be compact sets in \mathbb{R}^n , $\lambda \in \mathbb{R}, \theta \in [0, 1]$, $x \in \mathbb{R}^n$ and $T \in GL_n(\mathbb{R})$. Then:

- (b1) $(\lambda K_1) +_{\theta} \dots +_{\theta} (\lambda K_m) = \lambda(K_1 +_{\theta} \dots +_{\theta} K_m)$
- (b2) $K_1 +_{\theta} \dots +_{\theta} K_m = K_{\sigma(1)} +_{\theta} \dots +_{\theta} K_{\sigma(m)}$ for any σ permutation of $\{1, \dots, m\}$.
- (b3) $(x + K_1) +_{\theta} K_2 +_{\theta} \dots +_{\theta} K_m = x + (K_1 +_{\theta} \dots +_{\theta} K_m)$
- (b4) $TK_1 +_{\theta} \dots +_{\theta} TK_m = T(K_1 +_{\theta} \dots +_{\theta} K_m)$

The convexity is transmitted to the θ -convolution of m convex bodies.

Proposition 5.3. Let K_1, \dots, K_m be convex bodies in \mathbb{R}^n . Then $K_1 +_{\theta} \dots +_{\theta} K_m$ is a convex body.

Proof. The characteristic function of each convex body K_i is log-concave. The convolution of log-concave functions is log-concave, and the level sets of log-concave functions are convex. \square

Corollary 2.5 is proved by using that $\chi_{K_1} * \chi_{K_2}$ is $\frac{1}{n}$ -concave in its support. In order to generalize this result, we have to prove that the convolution of more than two characteristic functions is s -concave for some s . We get this result for $s^{-1} = (m-1)n$ by considering sets in dimension $(m-1)n$. As in the case $m=2$, we may assume, without loss of generality, that

$$(5.1) \quad M(K_1, \dots, K_m) = \chi_{K_1} * \dots * \chi_{K_m}(0).$$

Proposition 5.4. *Let $K_1, \dots, K_m \subset \mathbb{R}^n$ be convex bodies satisfying (5.1).*

Then, for any $\theta_1, \theta_2, \lambda_1, \lambda_2 \in [0, 1]$ such that $\lambda_1 + \lambda_2 \leq 1$ we have

$$(5.2) \quad \lambda_1(K_1 + \theta_1 \dots + \theta_1 K_m) + \lambda_2(K_1 + \theta_2 \dots + \theta_2 K_m) \subseteq K_1 + \theta \dots + \theta K_m,$$

where $1 - \theta^{\frac{1}{(m-1)n}} = \lambda_1(1 - \theta_1^{\frac{1}{(m-1)n}}) + \lambda_2(1 - \theta_2^{\frac{1}{(m-1)n}})$.

Proof. First at all, notice that for any $x \in \mathbb{R}^n$,

$$\begin{aligned} & \chi_{K_1} * \dots * \chi_{K_m}(x) \\ &= \int_{\mathbb{R}^n} \chi_{K_1} * \dots * \chi_{K_{m-1}}(t_{m-1}) \chi_{K_m}(x - t_{m-1}) dt_{m-1} \\ &= \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \chi_{K_1} * \dots \\ & \quad \dots * \chi_{K_{m-2}}(t_{m-2}) \chi_{K_{m-1}}(t_{m-1} - t_{m-2}) \chi_{K_m}(x - t_{m-1}) dt_{m-2} dt_{m-1} \\ &= \dots \\ &= \int_{\mathbb{R}^n} \dots \int_{\mathbb{R}^n} \chi_{K_1}(t_1) \chi_{K_2}(t_2 - t_1) \dots \\ & \quad \dots \chi_{K_{m-1}}(t_{m-1} - t_{m-2}) \chi_{K_m}(x - t_{m-1}) dt_1 \dots dt_{m-1} \\ &= |\Omega_{m-1}(x)| \end{aligned}$$

where

$$\begin{aligned} \Omega_{m-1}(x) = \{ & (t_1, \dots, t_{m-1}) \in \mathbb{R}^{(m-1)n} : t_1 \in K_1, t_2 - t_1 \in K_2, \dots \\ & \dots t_{m-1} - t_{m-2} \in K_{m-1}, x - t_{m-1} \in K_m\}. \end{aligned}$$

The convexity of K_1, \dots, K_m gives

$$(5.3) \quad \Omega_{m-1}(\lambda_1 x_1 + \lambda_2 x_2) \supseteq (1 - \lambda_1 - \lambda_2) \Omega_{m-1}(0) + \lambda_1 \Omega_{m-1}(x_1) + \lambda_2 \Omega_{m-1}(x_2)$$

for any $x_1, x_2 \in \mathbb{R}^n$ and $\lambda_1, \lambda_2 \geq 0$ such that $\lambda_1 + \lambda_2 \leq 1$. In particular, if $x_1 \in K_1 + \theta_1 \dots + \theta_1 K_m$ and $x_2 \in K_1 + \theta_2 \dots + \theta_2 K_m$, Brunn-Minkowski inequality in $\mathbb{R}^{(m-1)n}$ implies

$$(5.4) \quad |\Omega_{m-1}(\lambda_1 x_1 + \lambda_2 x_2)| \geq ((1 - \lambda_1 - \lambda_2) + \lambda_1 \theta_1^{\frac{1}{(m-1)n}} + \lambda_2 \theta_2^{\frac{1}{(m-1)n}})^{(m-1)n} |\Omega_{m-1}(0)|,$$

which shows that $\lambda_1 x_1 + \lambda_2 x_2 \in K_1 + \theta \dots + \theta K_m$ with

$$1 - \theta^{\frac{1}{(m-1)n}} = \lambda_1(1 - \theta_1^{\frac{1}{(m-1)n}}) + \lambda_2(1 - \theta_2^{\frac{1}{(m-1)n}}). \quad \square$$

Remark 5.5. The concavity of the function

$$x \in K_1 + \cdots + K_m \mapsto \chi_{K_1} * \cdots * \chi_{K_m}(x)^{\frac{1}{(m-1)n}}$$

allows us to write, for any $\theta \in [0, 1)$, the boundary of $K_1 + \theta \cdots + \theta K_m$ as

$$\partial(K_1 + \theta \cdots + \theta K_m) = \{x \in K_1 + \cdots + K_m : \chi_{K_1} * \cdots * \chi_{K_m}(x) = \theta M(K_1, \dots, K_m)\}.$$

In particular, for $\theta = 0$,

$$\partial(K_1 + \cdots + K_m) = \{x \in K_1 + \cdots + K_m : \chi_{K_1} * \cdots * \chi_{K_m}(x) = 0\}.$$

That implies that for any $x \in K_1 + \cdots + K_m$, $x \notin K_1 + \theta \cdots + \theta K_m$, there exists a (unique) $\theta \in [0, 1)$ such that $x \in \partial(K_1 + \theta \cdots + \theta K_m)$.

Remark 5.6. Taking $\theta_1 = \theta_2 = 1$, and following the proof of Proposition 5.4, we get equality in the inclusion (5.3) and in the Brunn-Minkowski inequality (5.4). Consequently, $\Omega_{m-1}(x)$ are all homothetic to $\Omega_{m-1}(0)$ for any $x \in K_1 + \theta \cdots + \theta K_m$. Using that they all have the same volume, we get that they are translations of $\Omega_{m-1}(0)$.

In particular, taking $\theta_1 = \theta_2$ and $\lambda_2 = 1 - \lambda_1$ in Proposition 5.4, we get the convexity of $K_1 + \theta \cdots + \theta K_m$. Also, taking $\theta_1 = \theta_2 = \theta_0$, we get the version of Corollary 2.5 to m bodies.

Corollary 5.7. Let $K_1, \dots, K_m \subset \mathbb{R}^n$ be convex bodies such that (5.1) is satisfied. Then for any $0 < \theta_0 \leq \theta < 1$

$$\frac{K_1 + \theta_0 \cdots + \theta_0 K_m}{1 - \theta_0^{\frac{1}{(m-1)n}}} \subseteq \frac{K_1 + \theta \cdots + \theta K_m}{1 - \theta^{\frac{1}{(m-1)n}}}.$$

We can easily get the Brunn-Minkowski type inequality for m bodies, a generalization of Corollary 3.2.

Corollary 5.8. Let $K_1, \dots, K_m \subset \mathbb{R}^n$ be convex bodies. Then

$$|K_1 + \theta \cdots + \theta K_m|^{\frac{1}{n}} \geq \left(1 - \theta^{\frac{1}{(m-1)n}}\right) (|K_1|^{\frac{1}{n}} + \cdots + |K_m|^{\frac{1}{n}}).$$

Finally, we get Zhang and Roger-Shephard type inequalities for m bodies. For $K_1, \dots, K_m \subseteq \mathbb{R}^n$, let

$$C_1(K_1, \dots, K_m) = \lim_{\theta \rightarrow 1} \frac{K_1 + \theta \cdots + \theta K_m}{1 - \theta}.$$

As in the case $m = 2$, the existence of the previous limit follows from Corollary 5.7.

Corollary 5.9. Let $K_1, \dots, K_m \subseteq \mathbb{R}^n$ be convex bodies. Then

$$|K_1 + \cdots + K_m| \leq \binom{m n}{n} \frac{|K_1| \cdots |K_m|}{M(K_1, \dots, K_m)} \leq (m-1)^n n^n |C_1(K_1, \dots, K_m)|$$

Proof. The proof runs as in Theorem 4.6 and Proposition 4.7 resp., using Corollary 5.7, instead of Corollary 2.5. \square

Regarding the study of equality cases, we will show that equality never occur in (5.2) for all λ_1, λ_2 such that $\lambda_1 + \lambda_2 = 1$ unless $m = 2$ or $n = 1$. That implies that extensions of Zhang, Roger-Shephard and Brunn-Minkowski type inequalities are not sharp for $m > 2$ and $n > 1$.

The following result can be proved as in Proposition 2.8.

Proposition 5.10. *Let $K_1, \dots, K_m \subset \mathbb{R}^n$ be convex bodies satisfying (5.1) and such that for any $\theta_1, \theta_2, \lambda_1, \lambda_2 \in [0, 1]$ such that $\lambda_1 + \lambda_2 = 1$ we have*

$$(5.5) \quad \lambda_1(K_1 +_{\theta_1} \dots +_{\theta_1} K_m) + \lambda_2(K_1 +_{\theta_2} \dots +_{\theta_2} K_m) = K_1 +_{\theta} \dots +_{\theta} K_m,$$

where $\theta^{\frac{1}{(m-1)n}} = \lambda_1 \theta_1^{\frac{1}{(m-1)n}} + \lambda_2 \theta_2^{\frac{1}{(m-1)n}}$. Then for every $x \in K_1 + \dots + K_m$ $\Omega_{m-1}(x)$ is homothetic to $\Omega_{m-1}(0)$.

Then we will show that this consequence can not occur for $m \geq 3$.

Proposition 5.11. *Let $m \geq 3$ and $K_1, \dots, K_m \subset \mathbb{R}^n$ be convex bodies satisfying (5.1). Then it is not possible that $\Omega_{m-1}(x)$ is homothetic to $\Omega_{m-1}(0)$ for every $x \in K_1 + \dots + K_m$.*

For the proof, we will use the following fact on sum of simplices, which is of independent interest.

Lemma 5.12. *Let $K, L \subset \mathbb{R}^n$ be n -dimensional convex bodies. If $K + L$ is an n -dimensional simplex, then K and L are both n -dimensional homothetic simplices.*

Proof. Denote by $h_C(x) = \max\{\langle x, y \rangle : y \in C\}$ the support function of a compact set $C \subset \mathbb{R}^n$.

Let us write $K + L = \text{conv}\{w_0, \dots, w_n\}$. We will show that L is a simplex.

Each w_i is an extreme point of $K + L$. Then $w_i = u_i + v_i$ where $u_i \in K$ and $v_i \in L$ are extreme points of K and L resp. In particular, $v_i = w_i - u_i \in L$ and so $\text{conv}\{w_0 - u_0, \dots, w_n - u_n\} \subseteq L$. Also

$$\begin{aligned} h_K(x) + h_L(x) &= h_{K+L}(x) = \max_{0 \leq i \leq n} \langle x, w_i \rangle \\ &= \max_{0 \leq i \leq n} (\langle x, u_i \rangle + \langle x, w_i - u_i \rangle) \\ &\leq \max_{0 \leq i \leq n} \langle x, u_i \rangle + \max_{0 \leq i \leq n} \langle x, w_i - u_i \rangle \\ &\leq h_K(x) + \max_{0 \leq i \leq n} \langle x, w_i - u_i \rangle \end{aligned}$$

Thus $h_L(x) \leq \max_{0 \leq i \leq n} \langle x, w_i - u_i \rangle$, so $L \subseteq \text{conv}\{w_0 - u_0, \dots, w_n - u_n\}$, and L is a simplex.

Now, write $K = \text{conv}\{u_0, \dots, u_n\}$ and $L = \text{conv}\{v_0, \dots, v_n\}$. We will prove that they are homothetic simplices.

Let $A_i = \{s \in S^{n-1} : h_K(s) = \langle x, u_i \rangle\}$ and $B_i = \{s \in S^{n-1} : h_L(s) = \langle x, v_i \rangle\}$. K and L are homothetic if and only if both partitions of S^{n-1}

$$S^{n-1} = \bigcup_{i=0}^n A_i = \bigcup_{i=0}^n B_i$$

are identical. Assume they are not the same, then the partition $S^{n-1} = \bigcup_{i,j=0}^n C_{i,j}$ where $C_{i,j} = \{s \in S^{n-1} : h_{K+L}(s) = \langle x, u_i + v_j \rangle\}$ has more than $n+1$ elements, since

$$h_{K+L}(s) = h_K(s) + h_L(s) = \max_{0 \leq i \leq n} \langle x, u_i \rangle + \max_{0 \leq j \leq n} \langle x, v_j \rangle = \max_{0 \leq i, j \leq n} \langle x, u_i + v_j \rangle.$$

Then $K+L$ is not a simplex. \square

Remark 5.13. An argument similar to that used in the remark before Theorem 3.2.3 in [8] leads to the same claim. However, we have included a direct proof of it for the sake of completeness.

Proof of Proposition 5.11. Notice that

$$\Omega_{m-1}(x) = \{(t_1, \dots, t_{m-1}) \in \mathbb{R}^{(m-1)n} : (t_1, \dots, t_{m-2}) \in \Omega_{m-2}(t_{m-1}), \\ t_{m-1} \in (x - K_m) \cap (K_1 + \dots + K_{m-1})\}.$$

and that $\Omega_{m-2}(t_{m-1})$ is non-empty if and only if $t_{m-1} \in K_1 + \dots + K_{m-1}$. Then the projection onto the t_{m-1} coordinate is

$$P_{m-1}(\Omega_{m-1}(x)) = (x - K_m) \cap (K_1 + \dots + K_{m-1}).$$

Suppose that $\Omega_{m-1}(x)$ is homothetic to $\Omega_{m-1}(0)$ for every $x \in K_1 + \dots + K_m$. Then their projections are also homothetic. Then $(x - K_m) \cap (K_1 + \dots + K_{m-1})$ is homothetic to $(-K_m) \cap (K_1 + \dots + K_{m-1})$ for any $x \in K_1 + \dots + K_m$ and Soltan's Theorem 2.9 implies that $K_1 + \dots + K_{m-1}$ and $-K_m$ are homothetic simplices.

On the other hand, the projection onto the t_{m-2} coordinate is

$$\begin{aligned} P_{m-2}(\Omega_{m-1}(x)) &= \bigcup_{t_{m-1} \in x - K_m} P_{m-2}(\Omega_{m-2}(t_{m-1})) \\ &= \bigcup_{t_{m-1} \in x - K_m} (t_{m-1} - K_{m-1}) \cap (K_1 + \dots + K_{m-2}) \\ &= (x - (K_{m-1} + K_m)) \cap (K_1 + \dots + K_{m-2}) \end{aligned}$$

and so we have that for every $x \in K_1 + \dots + K_m$, $(x - (K_{m-1} + K_m)) \cap (K_1 + \dots + K_{m-2})$ is homothetic to $(-(K_{m-1} + K_m)) \cap (K_1 + \dots + K_{m-2})$. Soltan's Theorem 2.9 shows again that $K_1 + \dots + K_{m-2}$ and $-(K_{m-1} + K_m)$ are homothetic simplices.

If K_m and $K_{m-1} + K_m$ are both simplices, by Lemma 5.12, K_{m-1} is also a simplex homothetic to K_m . But $(K_1 + \dots + K_{m-1}) + K_m$ is a simplex, so

Lemma 5.12 shows that $K_1 + \cdots + K_{m-1}$ and K_m are homothetic. But then K_m is a simplex homothetic to $-K_m$, a contradiction, unless $n = 1$. \square

ACKNOWLEDGEMENTS

Part of this work was done while the first two named authors were attending the Thematic Program on Asymptotic Geometric Analysis in Fall 2010 at the Fields Institute in Toronto, where the first named author enjoyed the J. Marsden's postdoctoral fellowship. We appreciate the hospitality. We are indebted to Prof. A. Giannopoulos for many helpful discussions and for pointing out to us several references at the early stages of this work. We also would like to thank Steven Taschuk for pointing out to us the alternative proof of Lemma 5.12 exposed in Remark 5.13. Finally, the authors thank the referee for suggesting the study of equality cases and other improvements in exposition.

REFERENCES

- [1] K. Kiener, "Extremalität von Ellipsoiden und die Faltungsungleichung von Sobolev," *Archiv der Mathematik*, vol. 46, pp. 162–168, 1986. 10.1007/BF01197494.
- [2] M. Schmuckenschläger, "The distribution function of the convolution square of a convex symmetric body in \mathbb{R}^n ," *Israel Journal of Mathematics*, vol. 78, pp. 309–334, 1992. 10.1007/BF02808061.
- [3] M. Meyer, S. Reisner, and M. Schmuckenschläger, "The volume of the intersection of a convex body with its translates," *Mathematika*, vol. 40, no. 02, pp. 278–289, 1993.
- [4] M. Schmuckenschläger, "Petty's projection inequality and Santaló's affine isoperimetric inequality," *Geom. Dedicata*, vol. 57, no. 3, pp. 285–295, 1995.
- [5] A. Tsolomitis, "Convolution bodies and their limiting behavior," *Duke Math. J.*, vol. 87, no. 1, pp. 181–203, 1997.
- [6] R. Schneider, "Mixed polytopes," *Discrete Comput. Geom.*, vol. 29, no. 4, pp. 575–593, 2003.
- [7] T. Bonnesen and W. Fenchel, *Theory of convex bodies*. Moscow, ID: BCS Associates, 1987. Translated from the German and edited by L. Boron, C. Christenson and B. Smith.
- [8] R. Schneider, *Convex bodies: the Brunn-Minkowski theory*, vol. 44 of *Encyclopedia of Mathematics and its Applications*. Cambridge: Cambridge University Press, 1993.
- [9] R. J. Gardner, *Geometric tomography*, vol. 58 of *Encyclopedia of Mathematics and its Applications*. Cambridge: Cambridge University Press, second ed., 2006.
- [10] V. D. Milman and G. Schechtman, *Asymptotic theory of finite-dimensional normed spaces*, vol. 1200 of *Lecture Notes in Mathematics*. Berlin: Springer-Verlag, 1986. With an appendix by M. Gromov.
- [11] H. Hadwiger, *Vorlesungen über Inhalt, Oberfläche und Isoperimetrie*. Berlin: Springer-Verlag, 1957.
- [12] W. Firey, " p -means of convex bodies," *Math. Scand.*, vol. 10, pp. 17–24, 1962.
- [13] E. Lutwak, "The Brunn-Minkowski-Firey theory. I. Mixed volumes and the Minkowski problem," *J. Differential Geom.*, vol. 38, no. 1, pp. 131–150, 1993.
- [14] E. Lutwak, "The Brunn-Minkowski-Firey theory. II. Affine and geominimal surface areas," *Adv. Math.*, vol. 118, no. 2, pp. 244–294, 1996.
- [15] C. Borell, "The Brunn-Minkowski inequality in Gauss space," *Invent. Math.*, vol. 30, no. 2, pp. 207–216, 1975.

- [16] R. Vitale, "The Brunn-Minkowski inequality for random sets," *J. Multivariate Anal.*, vol. 33, no. 2, pp. 286–293, 1990.
- [17] B. Uhrin, "Curvilinear extensions of the Brunn-Minkowski-Lusternik inequality," *Adv. Math.*, vol. 109, no. 2, pp. 288–312, 1994.
- [18] H. J. Brascamp and E. H. Lieb, "On extensions of the Brunn-Minkowski and Prékopa-Leindler theorems, including inequalities for log concave functions, and with an application to the diffusion equation," *J. Functional Analysis*, vol. 22, no. 4, pp. 366–389, 1976.
- [19] F. Barthe, "Restricted Prékopa-Leindler inequality," *Pacific J. Math.*, vol. 189, no. 2, pp. 211–222, 1999.
- [20] S. Szarek and D. Voiculescu, "Volumes of restricted Minkowski sums and the free analogue of the entropy power inequality," *Comm. Math. Phys.*, vol. 178, no. 3, pp. 563–570, 1996.
- [21] R. J. Gardner, "The Brunn-Minkowski inequality," *Bull. Amer. Math. Soc. (N.S.)*, vol. 39, no. 3, pp. 355–405, 2002.
- [22] C. M. Petty, "Isoperimetric problems," in *Proceedings of the Conference on Convexity and Combinatorial Geometry (Univ. Oklahoma, Norman, Okla., 1971)*, pp. 26–41, Dept. Math., Univ. Oklahoma, Norman, Okla., 1971.
- [23] G. Y. Zhang, "Restricted chord projection and affine inequalities," *Geom. Dedicata*, vol. 39, no. 2, pp. 213–222, 1991.
- [24] C. A. Rogers and G. C. Shephard, "The difference body of a convex body," *Arch. Math. (Basel)*, vol. 8, pp. 220–233, 1957.
- [25] C. A. Rogers and G. C. Shephard, "Convex bodies associated with a given convex body," *J. London Math. Soc.*, vol. 33, pp. 270–281, 1958.
- [26] H. Brunn, "Referat über eine Arbeit: Exacte Grundlagen für eine Theorie der Ovale," *S.-B. Bayer. Akad. Wiss.*, pp. 93–111, 1894.
- [27] V. Soltan, "A characterization of homothetic simplices," *Discrete Comput. Geom.*, vol. 22, no. 2, pp. 193–200, 1999.

C. HUGO JIMÉNEZ AND RAFAEL VILLA, DEPT. DE ANÁLISIS MATEMÁTICO, FACULTAD DE MATEMÁTICAS, UNIVERSIDAD DE SEVILLA, PO BOX 1160, 41080 SEVILLA, SPAIN

E-mail address: carloshugo@us.es, villa@us.es

DAVID ALONSO-GUTIÉRREZ, DEPT. OF MATH. AND STATS., 632 CENTRAL ACADEMIC BUILDING, UNIVERSITY OF ALBERTA, EDMONTON, AB, CANADA T6G 2G1

E-mail address: alonsogu@ualberta.ca