

1 **Extrapolating base-line trunk shrinkage reference equations across olive orchards**

2

3

4 **Corell, M.^{a,*}, Girón, I.F.^b, Moriana, A.^a, Dell'Amico, J.^c, Morales, D.^c, Moreno, F.^b**

5

6 ^aEscuela Técnica Superior de Ingeniería Agronómica. University of Seville, Carretera

7 de Utrera Km 1, 41013 Sevilla, Spain

8 ^bInstituto de Recursos Naturales y Agrobiología de Sevilla (IRNAS-CSIC), P.O. Box

9 1052, E-41080 Sevilla, Spain.

10 ^c Instituto Nacional de Ciencias Agrícolas, Cuba

11

12 *Corresponding author: mcorell@us.es Phone: (+34)954487298; Fax: (+34)954486436

13

14

15

16

17

18

19

20

21 **Abstract**

22 Maximum daily trunk shrinkage is a common measurement in irrigation scheduling of
23 fruit trees. But the strong relationship between these measurements and the environment
24 severely limit field applications. Reference baselines are the solution for understanding
25 the influence of environmental conditions. Nevertheless, the extrapolation out of the
26 original conditions is not clear. The aim of this study was to compare several
27 approaches to estimate a reference baseline in an olive orchard where there were no
28 previous data from other seasons. Two orchards, separated 60 m, with different tree
29 density were used. Orchard 1 had greater tree density than orchard 2, though the age and
30 the cultivar were the same. Trunk diameters of both orchards were similar but the crown
31 volume of orchard 2 was slightly lower than orchard 1. The current reference baselines
32 of maximum daily trunk diameter in both orchards were not significantly different
33 between them ($p < 0.05$). In orchard 1, the previous reference baseline was calculated in
34 a 5-year study (the so called multi-seasons approach). The multi-seasons approach was
35 not significantly different in slope but it was in the y-interception to the current
36 reference baselines in both orchards ($p < 0.05$). This approach over-estimated the values
37 in both orchards. Two additional approaches were tested. These latter approaches used
38 data before massive pit hardening to estimate the current reference baseline. One of
39 them used the early data to estimate a complete reference baseline (the so-called early
40 approach). The other (the so-called y-early approach) used the same data only to
41 estimate the y-interception and assumed that the slope was the same as in the multi-
42 seasons approach. The early approach under-estimated the value of maximum daily
43 trunk shrinkage. The early-y approach provided a satisfactory estimation of the

44 reference baseline and improved those obtained with the multi-seasons approach. The
45 limitations and uses in irrigation scheduling are also discussed.

46

47 **Keyword:** Irrigation scheduling, plant water status measurements, trunk diameter
48 fluctuations, water relations.

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67 **1. Introduction**

68 The Olive (*Olea europaea* L.) is a traditional rain-fed fruit tree in the Mediterranean
69 basin. In the last 20 years, the intensification of olive cultivation has advanced greatly.
70 These new orchards are irrigated and have higher tree density than traditional ones. The
71 increase in cultivation has coincided with an increase in water demand for other uses
72 (Fereres and Evans, 2006) and this strongly limits the amount of water available for
73 irrigation purposes.

74 Regulated deficit irrigation (RDI) is a common irrigation scheduling technique
75 in fruit trees, which saves water with low or null variations in yield (Behboudian and
76 Mills, 1997; Naor, 2006). When RDI is applied, plant water status measurements are
77 needed to control the intensity of the plant water stress imposed (Fereres and González-
78 Dugo, 2009). In deficit irrigation scheduling, techniques of continuous measurement of
79 plant water status allow adequate daily watering for control of water stress level (Ortuño
80 et al., 2010). Trunk diameter fluctuations have been suggested in several fruit trees as a
81 very efficient tool for RDI (i.e. almond, Goldhamer and Fereres, 2004; peaches,
82 Conejero et al., 2011), but not in others such as olive trees (Moriana et al., 2003;
83 Moriana et al., 2010). Trunk diameter fluctuations are a daily cycle of shrinkage (from
84 the beginning of the day) and swelling (from mid-afternoon) which occurs in all plants
85 (Klepper et al., 1971). This daily cycle provides two parameters which are used in
86 irrigation scheduling: maximum daily trunk shrinkage and trunk growth rate.

87 Goldhamer and Fereres (2001) suggested the first approach for irrigation
88 scheduling with a trunk diameter fluctuations parameter. In a young orchard, these
89 authors suggested the comparison with maximum daily diameter to estimate trunk
90 growth. However, although this parameter is presented in several studies (i.e.
91 Goldhamer et al 1999) extrapolation to other locations is not easy. Moriana and Fereres

92 (2002), in olive trees, reported that maximum daily diameter would be difficult to
93 extrapolate to other conditions and suggested trunk growth rate (the slope of maximum
94 daily diameter) as easier to use in irrigation scheduling. However, only in young olive
95 trees, a good relationship between TGR and temperature has been reported (Pérez-
96 López et al., 2008). No strong relationship between either of these parameters and a
97 meteorological variable has been found in mature orchards of any fruit species. This
98 lack of results is probably related to the strong relationship between fruit development
99 and both parameters (olive, Moriana et al., 2003; plum, Intrigliolo and Castel, 2007;
100 olive, Pérez-López et al., 2008).

101 Maximum daily trunk shrinkage is the indicator derived from trunk diameter
102 fluctuations most widely suggested in irrigation scheduling in several fruit trees (Ortuño
103 et al., 2010; Fernández and Cuevas 2010). The increase in maximum daily trunk
104 shrinkage has traditionally been associated with water stress conditions (Ortuño et al.,
105 2010), though it is also strongly related to evaporative demand (Herzog et al., 1995). A
106 reference is therefore needed in order to separate the effect of evaporative demand and
107 soil water deficit. Reference baseline is a simple regression equation that estimates
108 maximum daily trunk shrinkage in fully irrigated conditions from one meteorological
109 parameter. Parameters such as temperature or vapor pressure deficit would provide an
110 estimation of maximum daily trunk shrinkage in fully irrigated conditions and thus
111 irrigation could be scheduled according to the water stress level previously decided.
112 Goldhamer and Fereres (2001) suggested the maximum daily trunk diameter signal,
113 which is the ratio between the measured and the estimated values, as an indicator of
114 water stress level.

115 Previous studies conducted on olive trees have concluded that maximum daily
116 trunk shrinkage is not a reliable water status indicator in moderate water deficit

117 conditions (Moriana and Fereres, 2002; Moriana et al., 2010; Cuevas et al., 2012).
118 However, Moriana et al (2000) reported a maximum daily trunk shrinkage vs. stem
119 water potential relationship in olives that estimated a decrease in maximum daily trunk
120 shrinkage in severe water deficit conditions. Olive trees are a very drought resistant
121 species, in which severe water stress conditions during massive pit hardening (mid-
122 summer) do not affect or only slightly reduce yield (Goldhamer, 1999). A reference
123 baseline would therefore be useful in order to impose such stress conditions during this
124 phenological period.

125 Several authors have reported reference baselines of maximum daily trunk
126 shrinkage in relation to different meteorological data. The most commonly reported are
127 vapour pressure deficit (VPD) and air temperature (amongst others, almond, Goldhamer
128 and Fereres, 2004; lemon, Ortuño et al., 2009; mandarin, Pagán et al., 2008; peach,
129 Conejero et al., 2011; plum, Intrigliolo and Castel, 2007). There are only three studies
130 of olives that estimated a reference baseline (Moriana and Fereres, 2004; Moreno et al.,
131 2006; Moriana et al., 2011). Moriana and Fereres (2004), with one season's data,
132 suggested VPD as the meteorological variable but without comparison to other
133 meteorological data. Moreno et al (2006), with one season's data, compared four
134 different meteorological measurements (VPD, temperature, radiation and reference
135 evapotranspiration) and reported that VPD and temperature presented the best
136 agreement with maximum daily trunk shrinkage data. Moriana et al (2011), in a five-
137 year study, showed that maximum daily temperature was between others (VPD and
138 temperatures at different time) the best fit with maximum daily shrinkage. These
139 authors also considered that maximum temperature was easier to obtain than VPD and
140 temperature measurements at different time of the day.

141 One of the main problems in the management of the reference baseline is
142 validation in other places or seasons. In order to eliminate the influence of inter-season
143 changes most studies use data from several seasons (i.e. Ortuño et al., 2009; Conejero et
144 al., 2011; Moriana et al., 2011). However, Goldhamer and Fereres (2004) reported a
145 reference baseline estimated only with the data of the beginning of the irrigation season.
146 The extrapolation of the reference baseline to other places (with different cultivars
147 or/and tree spaces) is more related to factors such as fruit load or tree dimensions and,
148 from our knowledge, there are no studies in the literature. Fruit load has been reported
149 as a significant factor that slightly varied the equations obtained (Intrigliolo and Castel
150 2007; Conejero et al., 2010; Moriana et al., 2011). However, such differences are small
151 and, in a commercial orchard, fruit load could be not considered (Moriana et al., 2011).
152 The influences of tree age (Moriana and Fereres, 2004) or trunk diameter (Genard et al,
153 2001; Intrigliolo and Castel, 2006) were also significant. Reference baseline would
154 therefore probably need a previous local calibration.

155 The aim of this study is to compare several approaches to estimating a reference
156 baseline in an orchard where there are no previous data. This reference baseline will be
157 used during the massive pit hardening period. Two approaches use current season data
158 before massive pit hardening. And a third approach uses a previous reference baseline
159 (Moriana et al., 2011), calculated on the same experimental farm, but in an olive
160 orchard with closer tree spacing.

161

162 **2. Material and Methods**

163 The experiment was performed in two orchards during the summer of 2011 at La
164 Hampa, the experimental farm of the Instituto de Recursos Naturales y Agrobiología de
165 Sevilla (IRNAS-CSIC). These orchards are located at Coria del Río near Seville (Spain)

166 (37° 17'N, 6° 3'W, 30 m altitude). The sandy loam soil (about 2 m deep) of the
167 experimental site was characterized by a volumetric water content of 0.33 m³ m⁻³ at
168 saturation, 0.21 m³m⁻³ at field capacity and 0.1 m³m⁻³ at permanent wilting point, and
169 1.30 (0-10 cm) and 1.50 (10-120 cm) g cm⁻³ bulk density.

170 The two olive orchards (*Olea europaea* L cv Manzanillo) were irrigated during
171 the previous season with no water limitation. Orchard 1 was 41 years old and the tree
172 spacing was 7 x 5 m. This orchard was the same where the reference baseline of
173 Moriana et al (2011) was calculated, but the trees used in the present study were
174 different. Orchard 2 was also 41 years old and tree spacing was 7 m x 7m. This orchard
175 was beside orchard 1 and separated by around 60 m. The crown volume and trunk
176 diameter of the experimental trees were not significantly different between orchards
177 (p<0.05, Table 1). However, the trees in orchard 2 had markedly lower values of crown
178 volume than orchard 1 (around 25% less) and lower ground cover (orchard 1 40%;
179 orchard 2 24%). The beginning of the massive pit hardening was estimated according to
180 Gijón et al (2010) at day of the year (DOY) 157 in the orchard 1 and 164 in the orchard
181 2.

182 Pest control, pruning and fertilization practices were those commonly used by
183 growers and no weeds were allowed to develop in the orchard. Irrigation was carried out
184 during the night by drip using one lateral pipe per tree row and five emitters per plant,
185 delivering 8 L h⁻¹ each. Micrometeorological 30 min data, namely air temperature,
186 solar radiation, relative humidity of air and wind speed at 2 m above the soil surface
187 were collected by an automatic weather station located some 40 m from the
188 experimental site. Daily reference evapotranspiration (ET_o) was calculated using the
189 Penman-Monteith equation (Allen et al., 1998).

190 Irrigation requirements were determined as the difference between crop
191 evapotranspiration (ET_c) and rainfall. Soil moisture was not considered in the water
192 balance in order to obtain no water stress conditions. ET_c was determined according to
193 daily reference evapotranspiration (ET_o) and a crop factor based on the time of the year
194 and the percentage of ground area shaded by the tree canopy (Ferreles and Goldhamer,
195 1990). The crop coefficient values (K_c) considered were 0.76 in May, 0.70 in June, 0.63
196 in July and August, 0.72 in September and 0.77 in October (Fernández et al., 2006). The
197 values of the coefficient in relation to the percentage of ground covered by the crop (K_r)
198 were 0.8 in orchard 1 and 0.48 in orchard 2. The values of crop evapotranspiration (ET_c)
199 and total amount of applied water (rainfall not included) during the experimental period
200 (from the end of April until the middle September) are shown in Table 2.

201 Trunk diameter fluctuations (TDF) are a daily cycle of shrinkage and swelling.
202 TDF were measured throughout the experimental periods, using a set of linear variable
203 displacement transducers (LVDT) (model DF±2.5 mm, accuracy ±10 µm, Solartron
204 Metrology, Bognor Regis, UK) attached to the main trunk, with a special bracket made
205 of Invar, an alloy of Ni and Fe with a thermal expansion coefficient close to zero
206 (Katerji et al., 1994). Measurements in 6 trees in orchard 1 and 5 trees in orchard 2 were
207 taken every 10 s and the datalogger (model CR10X with AM 416 multiplexer,
208 Campbell Sci. Ltd., Logan, USA) was programmed to report 15 min means. Maximum
209 daily trunk shrinkage (MDS) was calculated as the difference between the maximum
210 daily diameter, which occurs at the beginning of the day, and the minimum daily
211 diameter, which occurs at mid-afternoon (Goldhamer et al., 1999).

212 The determination of the reference baseline supposed fully irrigated conditions.
213 Several parameters were measured in order to establish the water status of the tree (soil
214 moisture, maximum trunk diameter, stem water potential). Soil moisture was measured

215 with a portable FDR sensor (HH2, Delta-T, U.K.) with a calibration previously
216 obtained. The access tubes for the FDR sensor were placed in the irrigation line around
217 30 cm from the emitter. The data were obtained at 1 m depth and 10 cm intervals.

218 Maximum daily diameter of TDF daily cycle showed the trunk growth
219 (Goldhamer et al 1999) and it was initially suggested as an indicator in water status
220 measurements (Goldhamer and Fereres, 2001). However, trunk growth rate (TGR), the
221 slope of maximum daily diameter, was considered a better parameter to describe the
222 cycle of stress and re-watering (Moriana and Fereres, 2002). TGR in day “n” was
223 calculated as the difference between the maximum daily diameter of day “n+1” minus
224 the ones from the day “n” (Cuevas et al 2010). In order to characterize the trunk growth,
225 maximum daily diameter was represented and also average TGR values were presented.

226 The stem water potential was measured at midday in one leaf per tree, using the
227 pressure chamber technique (Scholander et al., 1965). Leaves near the main trunk were
228 covered with aluminium foil at least one hour before measurements were taken.

229 The reference baseline of maximum daily trunk shrinkage would be used during
230 pit hardening. The aim of this study was to establish a methodology for estimating a
231 reference baseline in a current season before the massive pit hardening period.
232 Reference baseline was estimated with maximum temperature. According to the
233 literature maximum temperature is the best meteorological parameter (Moriana et al.,
234 2011). Three different approaches were used to estimate the reference baseline:

235 1) Multi-seasons approach. This approach used the 5-year reference baseline
236 calculated previously in orchard 1 (Moriana et al. 2011). Moriana et al (2011) reported
237 that the low fruit load season’s equations were significantly different to the high fruit
238 load season’s equations. However, according to this author such differences were small

239 and a general equation with all the seasons is suggested for commercial purposes. This
240 general equation is the one used in this approach in both orchards.

241 2) Early approach. This approach was based on the methodology suggested in
242 Goldhamer and Fereres (2004). The hypothesis is that data at the beginning of the
243 season would allow the estimation of the reference baseline for the current season. The
244 reference baselines in each orchard were estimated with data measured in the current
245 season before the beginning of massive pit hardening.

246 3) Early-y approach. This approach is a mix of approaches 1 and 2. Moriana et al
247 (2011) reported differences in the y-interception of the reference baseline due to fruit
248 load, but not in the slope. The hypothesis is that changes in the reference baseline would
249 be only in y-interception and not in the slope. Therefore, in this approach, data before
250 massive pit hardening would estimate y-interception of the reference baseline. The slope
251 used was the same as the multi-seasons approach.

252 In all three cases reference baseline was obtained by linear regression analysis,
253 between the two variables. Differences between regression lines were determined with
254 T-test of the slope and y-intercept. The comparisons between early-y approach and the
255 current reference baseline and the early approach were done with confidence intervals
256 (95%) of the slope and y-intercept. Equations were validated with the measured data of
257 Maximum daily trunk shrinkage. And statistical differences with the 1:1 line were
258 determined with T-test.

259

260 **3. Results.**

261 The experiment was performed from day of the year (DOY) 100 to 260 in 2011.
262 Environmental variables fluctuated widely during this period, as is customary in the
263 area. Mean daily air temperature (T_m) and maximum air temperature (T_{max}) presented a

264 similar trend, reaching maximum values in August (Fig. 1A). Average T_m and average
265 T_{max} were 25.6 and 33.5°C respectively. The pattern of daily ET_o fluctuated widely,
266 showing maximum values in late June and early July, and minimum values in early
267 September (Fig. 1B). Total ET_o during the experimental period was 637 mm (Fig. 1B).
268 Rainfall was very scarce, 87.1 mm during the experimental period, and occurred in June
269 and late August (Fig. 1B). During the experimental period the volumetric soil water
270 content in the profile (0-1m) was almost constant in the two orchards, with values close
271 to field capacity content (Fig 1C). Soil water content in orchard 2 was always slightly
272 higher than in orchard 1, average values were 0.23 m^3m^{-3} and 0.26 m^3m^{-3} in orchard 1
273 and 2 respectively (Fig 1C).

274 Stem water potential presented a similar pattern in both orchards, with an almost
275 steady pattern throughout the experimental period (Fig. 2A). In orchard 1, stem water
276 potential varied from -0.8 MPa to around -1.2 MPa, the average value was -1.13 MPa.
277 In orchard 2, values varied from -0.7 MPa to -1.4 MPa, the average value was -1.17
278 MPa. Maximum daily trunk shrinkage (MDS) showed a similar pattern in both
279 orchards, as shown in Figure 2B. Maximum daily trunk shrinkage increased slightly
280 from the beginning of the experiment. The highest values of maximum daily trunk
281 shrinkage in both orchards were observed during August (around 779 μm). Maximum
282 daily diameter increased during the season in both orchards; Orchard 2 had the highest
283 growth (Fig. 2C). The periods of sharp increase and decrease of maximum daily
284 diameter (orchard 1, DOY 118-129; DOY 138-143; DOY 241-250; orchard 2, 136-143;
285 DOY 237-250) were related to rainfall events (Fig. 1B), probably strongly influenced by
286 the increased trunk moisture content. If these periods of rain are excluded, trunk growth
287 was almost constant with a slight increase from DOY 187 in orchard 1 and at DOY 183
288 in orchard 2. In orchard 1, trunk growth rate (TGR), the slope of the maximum daily

289 diameter figure, was lower than in orchard 2. (TGR was 8.4 ± 2.3 in orchard 1 and
290 $33.7 \pm 4.6 \mu\text{m day}^{-1}$, in orchard 2).

291 Crop evapotranspiration (ET_c) and applied water in orchard 1 was 30% higher
292 than in orchard 2 (around 30% in both parameters), due to tree spacing (Table 2). Yields
293 were very low in both orchards: with a yield of 2.5 t ha^{-1} in orchard 1 and 0.6 t ha^{-1} in
294 orchard 2.

295 The relationships between maximum daily trunk shrinkage (MDS) and
296 maximum air temperature (T_{max}) for both orchards are shown at Fig. 3. The best fit,
297 when all the data are considered, was $\text{MDS} = -667 + 34T_{\text{max}}$ (equation 1), in orchard 1 and
298 $\text{MDS} = -757 + 37T_{\text{max}}$ (equation 2) in orchard 2 (Fig 3). The last two, equations 1 and 2,
299 were therefore the reference baselines for this season. The equations were not
300 significantly different ($p < 0.05$). Equations 1 and 2 were slightly displaced compared to
301 the multi-seasons approach (Fig. 3a; $\text{MDS} = -640 + 36T_{\text{max}}$, Moriana et al., 2011). The y-
302 interception was significantly different between equations 1 and 2 and the multi-seasons
303 approach ($p < 0.05$). However, there were no significant differences in the slope between
304 the two equations and the multi-seasons approach ($p < 0.05$).

305 The early approach estimated the reference baseline only with the early data
306 (before massive pit hardening, solid symbols at Figure 3b). The best fit with the early
307 approach was $\text{MDS} = -137 + 13T_{\text{max}}$ in orchard 1 and $\text{MDS} = -478 + 26T_{\text{max}}$ in orchard 2.
308 Both latter equations obtained with the early approach were significantly different from
309 equations 1 and 2 respectively ($p < 0.05$).

310 Finally, the Early-y approach estimated only the y-interception with the data
311 before massive pit hardening (Fig. 3c). The equation estimated was $\text{MDS} = -792 + 36T_{\text{max}}$
312 in orchard 1 and $\text{MDS} = -760 + 36T_{\text{max}}$ in orchard 2. In both orchards, the slope and the y-
313 interception of the early-y approach were in the confidence interval (95%) of equations

314 1 and 2, respectively. In orchard 1, the slope was also in the confidence interval (95%)
315 of the early approach but not in orchard 2.

316 The equations calculated in the three approaches were validated with the data
317 measured from the beginning of massive pit hardening (Fig. 4). In both orchards, all the
318 approaches were significantly different to the 1:1 line ($p < 0.05$). The multi-seasons
319 approach clearly over-estimated in all the ranges of maximum daily trunk shrinkage
320 (Fig. 4a). The early approach underestimated the measured values (Fig. 4b). In the early
321 approach, the differences were greater in orchard 1 than in orchard 2 (Fig. 4b). The
322 early-y equation is the nearest to the 1:1 line in both orchards, especially with maximum
323 daily trunk shrinkage values lower than $500 \mu\text{m}$ (Fig 4 c).

324

325 **4. Discussion**

326 Deficit irrigation scheduling based on water status measurements such as trunk diameter
327 fluctuations had a very significant advantage in comparison with the traditional water
328 balance (Ortuño et al., 2010). This is that the control of water stress level instead of
329 applied water permit an easier extrapolation to different conditions. However, the great
330 variability of the indicators of trunk diameter fluctuations (Naor et al., 2006) and the
331 strong relationship with the environment probably limit the use in places different to
332 those where the experiments were performed. According to the present study, tree
333 spacing was not an important factor and almost the same reference baseline could be
334 used (Fig. 3). Although an important parameter as trunk diameter were similar (Table
335 1), the changes in tree environment, mainly radiation, due to the tree spacing (ground
336 covers were clearly different, Table 1) would be enough to produce significant changes
337 in the maximum daily trunk shrinkage. Trunk diameter fluctuations are mainly
338 produced by hydration and dehydration of the bark (Brough et al., 1986) and had been

339 associated with changes in trunk water content (Simmoneau et al., 1993). Therefore,
340 maximum daily trunk shrinkage has been considered a good indicator in fully irrigated
341 conditions of tree transpiration (Herzog et al., 1995). So according to the reference
342 baseline of orchard 1 and 2, tree transpiration was similar in both orchards. However,
343 the interception radiation of the trees was different and, therefore, the canopy
344 transpiration was also likely different. On the other hand, the soil allotted per tree is
345 larger in the wider tree spacing orchard, probably the more water transpired by the
346 canopy is compensated by the larger root water uptake capacity. Then maximum daily
347 shrinkage would not be associated to transpiration as strong as other authors suggest
348 and, in fact, estimates the difference between root uptake and canopy transpiration.

349 Fruit load is a factor that, in olives trees, could also affect the extrapolation of
350 the reference baseline. Alternate bearing could be produced by climatic and biotic
351 conditions, or, which is most common in table olives, excessive pruning. The multi-
352 season approach did not consider this effect, though fruit load produced significant
353 changes in water relations and transpiration in olive trees (Martín-Vertedor et al., 2011).
354 This approach over-estimated maximum daily trunk shrinkage mainly with values lower
355 than 500 μm (Fig. 4). These variations were probably produced by the very low fruit
356 load conditions. Moriana et al (2011) suggested that the variations when fruit load were
357 not considered would be small. However, according to the present study deviations
358 greater than 25% than the measured value would be obtained, mainly with low values of
359 maximum daily trunk shrinkage (Fig. 4). Intrigliolo and Castel (2007) reported a similar
360 decrease (around 34%) in maximum daily trunk diameter due to fruit load in plums. The
361 yield in the present study was very low and should be considered even as null (Table 2).
362 Martín-Vertedor et al (2011) in olive trees reported that the significant variations in
363 stomata conductance occurred between off-season (when no yield was recorded) and the

364 rest of seasons (with medium and high yield). Therefore, the multi-season approach
365 could be useful in conditions of significant yield.

366 Although significant yield would be expected an estimation of the reference
367 baseline with current season data and with the local conditions would be a better
368 approach than the multi-season approach. The approach suggested by Goldhamer and
369 Fereres (2004) in almonds (the so-called early approach) was used in this way.
370 However, the early approach was the worst in both orchards (Figs. 3 and 4). These
371 disagreements with the results in almonds of Goldhamer and Fereres (2004) are
372 probably related to the differences between the water relations of both fruit species. The
373 daily cycle of stomatal leaf conductance in full irrigated conditions is not limited by
374 evaporative demand in almond (Marsal and Girona, 1997) while in olive trees there is a
375 reduction at midday (Angelopoulos et al., 1996). Such differences in the daily pattern
376 stomata could suggest different respond to high transpiration conditions. According to
377 the present study, maximum daily trunk shrinkage greater than 400 μm was under-
378 estimated with the early approach (Fig. 4). Therefore, the greatest conditions of
379 transpiration likely changed the amount of water that the trunk transferred to the
380 transpiration stream. In olive, the tree could increases the amount of water from the
381 trunk more than in almond and then maximum daily shrinkage was higher than
382 expected, Such increase could be related to the greater capacity of dehydration in olive
383 trees (Fereres, 1984).

384 Maximum daily trunk shrinkage is affected by several factors which would
385 change the pattern of its relationship with temperature in the long term, water status
386 conditions (Genard et al., 2001) and wood composition (Drew and Dones, 2009) are
387 probably the most important. Such changes would probably vary the reference baseline
388 even though fully-irrigated conditions were performed during the current or previous

389 season. The present study confirmed that these changes occurred, but only in the y-
390 interception, while the slope of the equation was not affected (Fig. 4). The early-y
391 approach made it possible to obtain the reference baseline before massive pit hardening
392 with only the estimation of the y-interception. The extrapolation to other Manzanillo
393 orchards is probably possible, but further work is needed to study the influence of the
394 cultivar. Moriana and Fereres (2004) reported reference baseline using vapor pressure
395 deficit (VPD) with different cultivars and age orchards, but similar environmental
396 conditions. When VPD is considered for the same fruit load an almost equal slope was
397 found between mature cv Manzanillo (Moriana et al. 2011) and mature cv Picual
398 (Moriana and Fereres, 2004). The influence of tree age is probably different, young cv
399 Arbequino showed a different slope and y-interception than mature cv Picual (Moriana
400 and Fereres, 2004) and mature cv Manzanillo (Moriana et al., 2011). Goldhamer and
401 Fereres (2001) suggested that the maximum daily trunk shrinkage would be smaller in
402 young trees than in mature because of the greater growth in the former. In almonds, the
403 reference baseline reported in several studies for young orchards was similar with
404 different cultivars (Fereres and Goldhamer, 2003; Goldhamer and Fereres, 2004). Egea
405 et al. (2009) with three season's data in a young almond orchard suggested that the
406 reference baseline could be estimated every 1-2 years with the data of the beginning of
407 the season, though slight differences due to phenological stage were reported.

408

409 **5. Conclusions**

410 The extrapolation of reference baseline of maximum daily trunk shrinkage to other
411 orchards was possible in mature cultivar Manzanillo. The reference baselines were
412 similar between orchards, though spacing and ground cover were different. Slight
413 variations in the y-interception were found in both reference equations. The multi-

414 season approach, a general equation previously calculated, over-estimated both
415 reference baselines. Such results were probably related to the extremely low fruit load in
416 both orchards in comparison with the ones obtained when the multi-season approach
417 was calculated. A multi-season approach would be useful in conditions of significant
418 yield. An early approach that estimated the complete (slope and y-interception)
419 reference baseline was not possible because significant variations in the slope were
420 found with the increase in maximum daily trunk shrinkage. This result is not consistent
421 with the ones reported in almond. The differences in the stomata leaf conductance
422 pattern between both species were probably related to the lack of results. The reference
423 baseline in all the estimations was similar in slope but different in y-interception. The
424 early-y approach that only estimated this latter component of the reference baseline
425 presented a good fit between observed and measured data during the period of pit
426 hardening. According to the present results and literature the early-y approach would be
427 useful in the estimation of reference baseline in mature orchards even when different
428 cultivars were considered.

429

430 **Acknowledgements**

431 This research was supported by the Spanish Ministerio de Economía y Competitividad
432 (MINECO), (AGL2010-19201-CO4-03 project). The stay of D. Morales and J.
433 Dell'Amico was supported by the AECID project D/030431/10. Thanks are due to A.
434 Montero and J. Rodríguez for help with field measurements and to professors M. Cera
435 and C. Cortés for help in the statistical analysis.

436

437

438 **Reference**

- 439 Allen, R.G., Pereira, L.S., Raes, D., Smith, M. 1998. Crop evapotranspiration.
440 Guideline for computing crop water requirements. FAO irrigation and drainage
441 paper n° 56. Roma.FAO.
- 442 Angelopoulos, K., Dichio, B., Xiloyannis, C. 1996. Inhibition of photosynthesis in olive
443 trees (*Olea europaea* L.) during water stress and rewatering. J Exp Bot 47, 1093-
444 110.
- 445 Behboudian, M.H., Mills, T.M., 1997. Deficit irrigation in deciduous orchards. Hort.
446 Rev. 21, 105-131.
- 447 Brough, D.W., Jones, H.G., Grace, J. 1986. Diurnal changes in water content of the
448 stems of apple trees, as influenced by irrigation. Plant Cell Environ. 9, 1-7.
- 449 Conejero, W., Mellisho, C.D., Ortuño, M.F., Galindo, A., Pérez-Sarmiento, F.,
450 Torrecillas, A. 2010. Establishing maximum daily trunk shrinkage and midday
451 stem water potential reference equations for irrigation scheduling of early
452 maturing peach trees. Irrig. Sci. 29, 299-309.
- 453 Conejero, W., Mellisho, C. D., Ortuno, M. F., Moriana, A., Moreno, F., Torrecillas, A.
454 2011. Using trunk diameter sensors for regulated deficit irrigation scheduling in
455 early maturing peach trees. Environ. Exp. Bot. 71, 409-415.
- 456 Cuevas MV, Martín-Palomo MJ, Díaz-Espejo A, Torres-Ruíz JM, Rodríguez-
457 Dominguez CM, Pérez-Martín A, Pino-Mejías R, Fernández JE (2012).
458 Assessing water stress in a hedgerow olive orchard from sap flow and trunk
459 diameter measurements. Irrig. Sci. doi 10.1007/s00271-0.12-0357-x.
- 460 Drew, D.M., Downes, G.M. 2009. The use of precision dendrometers in research on
461 daily stem size and wood property variation: A review. Dendrochronologia 27,
462 159-172.

463 Fereres, E. 1984. Variability in adaptive mechanisms to water deficits in annual and
464 perennial crop plants. *Bull. Soc. Bot. France.* 131, 17-32.

465 Fereres, E., Evans, R.G. 2006. Irrigation of fruit trees and vines: an introduction. *Irrig.*
466 *Sci.* 24, 55-57.

467 Fereres, E., Goldhamer, D. 1990. Deciduous fruit and nut trees. In: Stewart, B.A.,
468 Nielsen D.R. (Eds), *Irrigation of Agricultural Crops. Agronomy Monograph No.*
469 *30. American Society of Agronomy, Madison, pp. 987-1017.*

470 Fereres, E., Goldhamer, D. 2003. Suitability of stem diameter variations and water
471 potential as indicators for irrigation scheduling of almond trees. *J. Hort. Sci.*
472 *Biotech.* 78,139-144.

473 Fereres, E., González-Dugo, V. 2009. Improving productivity to face water scarcity in
474 irrigated agriculture. In: *Crop Physiology: applications for genetic improvements*
475 *and agronomy. Sadras, V. and Calderini, D. (eds). Pp 122-143. Amsterdam,*
476 *Academic Press.*

477 Fernández, J. E.; Cuevas, M. V. 2010. Irrigation scheduling from stem diameter
478 variations: A review. *Agr. Forest Meteorol.*, 150, 135-151.

479 Genard, M., Fishman, S., Vercambre, G., Huguet, J.G., Bussi, C., Besset, J., Habib, R.
480 2001. A biophysical analysis of stem and root diameter variations in woody
481 plants. *Plant Physiol.* 126, 188-202.

482 Gijón, M.C., Pérez-López, D., Guerrero, J., Couceiro, J.F., Moriana, A. 2010. Riego
483 deficitario controlado en olivo y pistachero. *Agricultura* 930, 458-462

484 Goldhamer, D.A. 1999. Regulated deficit irrigation for California canning olives. *Acta*
485 *Hortic.* 474, 369-372.

486 Goldhamer D.A.; Fereres E. 2001. Irrigation scheduling protocols using continuously
487 recorded trunk diameter measurements. *Irrig. Sci.* 20, 115-125

488 Goldhamer, DA; Fereres, E 2004. Irrigation scheduling of almond trees with trunk
489 diameter sensors. *Irrigation Sci.* , 23: 11-19.

490 Goldhamer, D., Fereres, E., Mata, M., Girona, J., Cohen, M. 1999. Sensitivity of
491 continuous and discrete plant and soil water status monitoring in peach trees
492 subjected to deficit irrigation. *J. Am. Soc. Hortic. Sci.* 124, 437-444.

493 Herzog K.M., Hüsler R., Thum R. 1995. Diurnal changes in the radius of a subalpine
494 Norway spruce stem: their relation to the sap flow and their use to estimate
495 transpiration. *Trees* 10, 94-101.

496 Intrigliolo, D.S., Castel, J.R. 2006. Usefulness of diurnal trunk shrinkage as a water
497 stress indicator in plum trees. *Tree Physiol.*, 26 303–311

498 Intrigliolo, D.S., Castel, J.R. 2007. Crop load affects maximum daily trunk shrinkage of
499 plum trees. *Tree Physiol.*, 27: 89-96.

500 Katerji,N., Tardieu,F.,Bethenod,O., Quetin, P. 1994. Behavior of Maize stem diameter
501 during drying cycles:comparison of two methods for detecting water stress. *Crop*
502 *Sci.* 34, 165-169.

503 Klepper, B., Browing, V.D., Taylor, H.M. 1971. Stem diameter in relation to plant water
504 status. *Plant Physiol.* 48, 683-685.

505 Marsal, J., Girona, J. 1997. Leaf water relations parameters in almond compared to
506 hazelnut trees during a deficit irrigation period. *J. Amer. Soc. Hort. Sci.* 122,
507 582-587.

508 Martín-Vertedor, A.I., Pérez-Rodríguez, J.M., Prieto-Losada, H., Fereres, E. 2011.
509 Interactive responses to water deficits and crop load in olive (*Olea europaea* L.,
510 cv. Morisca) I. - Growth and water relations. *Agr. Water Manage.* 98, 941-949.

511 Moreno, F., Conejero, W., Martin-Palomo, M. J., Giron, I. F., Torrecillas, A., 2006.
512 Maximum daily trunk shrinkage reference values for irrigation scheduling in
513 olive trees. *Agr. Water Manage.* 84, 290-294.

514 Moriana, A., Fereres, E. 2002. Plant indicators for scheduling irrigation of young olive
515 trees. *Irrig. Sci.* 21, 83-90.

516 Moriana, A., Fereres, E. 2004. Establishing references values of trunk diameter
517 fluctuations and stem water potential for irrigation scheduling of olive trees.
518 *Acta Hortic.* 664, 407-412.

519 Moriana, A., Fereres, E., Orgaz, F., Castro, J., Humanes, M.D., Pastor, M. 2000. The
520 relations between trunk diameter fluctuations and tree water status in olive tree
521 (*Olea europaea* L). *Acta Hortic.* 537, 293-297.

522 Moriana, A., Orgaz, F., Fereres, E., Pastor, M. 2003. Yield responses of a mature olive
523 orchard to water deficits. *J. Am. Soc. Hortic. Sci.* 128, 425-431.

524 Moriana, A., Giron, I.F., Martin-Palomo, M.J., Conejero, W., Ortuno, M.F., Torrecillas,
525 A., Moreno, F. 2010. New approach for olive trees irrigation scheduling using
526 trunk diameter sensors. *Agric. Water Manage.* 97, 1822-1828.

527 Moriana, A., Moreno, F., Giron, I.F., Conejero, W. Ortuño, M.F., Morales, D., Corell,
528 M., Torrecillas, A. 2011. Seasonal changes of maximum daily shrinkage
529 reference equations for irrigation scheduling in olive trees: Influence of fruit
530 load. *Agric. Water Manage.* 99, 121-127.

531 Naor, A. 2006. Irrigation scheduling and evaluation of tree water status in deciduous
532 orchard. *Hort. Rev.* 32, 111-165.

533 Naor, A., Gal, Y., Peres, M. 2006. The inherent variability of water stress indicators in
534 apple, nectarine and pear orchards and the validity of a leaf selection procedure
535 for water potential measurements. *Irrig. Sci.* 24,129-135.

536 Ortuño, M. F., Garcia-Orellana, Y., Conejero, W. 2009. Assessment of maximum daily
537 trunk shrinkage signal intensity threshold values for deficit irrigation in lemon
538 trees. *Agric. Water Manage.* 96, 80-86.

539 Ortuño, M.F., Conejero, W., Moreno, F., Moriana, A., Intrigliolo, D.S., Biel, C.,
540 Mellisho, C.D., Pérez-Pastor, A., Domingo, R., Ruiz-Sánchez, M.C., Casadesus,
541 J., Bonany, J., Torrecillas, A. 2010. Could trunk diameter sensors be used in
542 woody crops for irrigation scheduling? A review of current knowledge and
543 future perspectives. *Agric. Water Manage.* 97, 1–11.

544 Pagán, E., Pérez-Pastor, A., Domingo, R., Conesa, M.R., Robles, J.M., Botía, P.,
545 García-Oller, I., Caro, M. 2008. Feasibility study of the maximum daily trunk
546 shrinkage for scheduling mandarin trees irrigation. *Ital. J. Agron.* 3, 691-692.

547 Pérez-López, D., Moriana, A., Rapoport, H., Olmedilla, N., Ribas, F. 2008. New
548 approach for using trunk growth rate and endocarp development in the irrigation
549 scheduling of young olive orchard. *Sci. Hort.* 115, 244-251.

550 Scholander, P.F., Hammel, H.T., Bradstreet, E.A., Hemmingsen, E.A. 1965. Sap
551 pressure in vascular plant. *Sci.* 148, 339-346.

552 Simonneau, T., Habib, R., Goutouly, J.P., Huguet, J.G. 1993. Diurnal changes in stem
553 diameter depend upon variations in water content: direct evidence in peach trees.
554 *J. Exp. Bot.* 44, 615-621.

555

556

557

558

559

560

561 Table 1. Dimensions and distance in the orchard of the experiment. There were no
562 significant differences between orchard 1 and orchard 2 in crown volume or trunk
563 diameter.

	Distance (mxm)	Ground cover	Crown Volume (m ³ tree ⁻¹)	Trunk Diameter (m)
Orchard 1	7x5	40%	31.3±4	0.24±0.01
Orchard 2	7x7	24%	23.6±2	0.23±0.01

564

565

566

567

568

569

570

571

572

573

574

575

576

Table 2. Crop evapotranspiration, irrigation applied and yield components: yield (Yield (kg tree⁻¹) and harvest (t ha⁻¹)).

	ETc (mm)	Irrigation applied (mm)	Yield (kg tree ⁻¹)	Yield (t ha ⁻¹)
Orchard 1	299.7	285.4	8.9 ± 6.0	2.5 ± 1.7
Orchard 2	214.0	196.7	2.9 ± 1.4	0.6 ± 0.3

580 **Figure captions**

581 Fig. 1. Daily mean (T_m , solid line) and maximum (T_{max} , dotted line) air temperature (A),
582 reference evapotranspiration (ET_o, solid line) and daily rainfall (vertical bars) (B) and
583 volumetric soil water content (θ_v) down to 1m depth (Orchard 1: open symbols,
584 Orchard 2 solid symbols) values during the experimental period (C). Horizontal lines
585 (C) represent volumetric soil water content at permanent wilting point (WP), at field
586 capacity (FC) and at saturation (S), respectively.

587 Fig. 2. Midday stem water potential (Ψ_{stem}) (A), maximum daily trunk shrinkage (MDS)
588 (B) and maximum daily diameter during experimental period (C) (Orchard 1: open
589 symbols and Orchard 2 solid symbols). Each point is the average of 6 measurements at
590 Orchard 1 and 5 measurements at orchard 2.

591 Fig. 3. Relationship between maximum daily trunk shrinkage (MDS) and maximum
592 temperature at orchard 1 (\square) and orchard 2 (\triangle). In the figure different reference
593 baselines are presented. In all of them, the best fit for each orchard is presented. Orchard
594 1, $MDS = -667 + 34T_{max}$ (bold solid line, $R^2 = 0.60^{***}$, $RMSE = 107 \mu m$, $n = 132$). Orchard
595 2, $MDS = -757 + 37T_{max}$ (bold dash line, $R^2 = 0.67^{***}$, $RMSE = 84 \mu m$, $n = 91$) (A)
596 Comparison between the best fit and the multi-seasons approach (solid line, $MDS = -$
597 $640 + 36 T_{max}$, Moriana et al, 2011). (B) Comparison between the best fit and the early
598 approach in orchard 1 (gray line, $MDS = -137 + 13T_{max}$, $R^2 = 0.52^{***}$, $RMSE = 52 \mu m$,
599 $n = 30$) and orchard 2 (gray dash line, $MDS = -478 + 26T_{max}$, $R^2 = 0.58^{***}$, $RMSE = 64 \mu m$;
600 $n = 21$). (C) Comparison between the best fit and the early-y approach in orchard 1 (gray
601 line, $MDS = -792 + 36T_{max}$) and orchard 2 (gray dash line, $MDS = -760 + 36T_{max}$).

602 Fig. 4. Relationship between measured and estimated maximum daily trunk shrinkage
603 (MDS) data from the beginning of the massive pit hardening. In all the figures the 1:1
604 line is represented. The graph compares the results in orchard 1 (\square) and orchard 2 (\triangle)
605 using the multi-seasons (A), early (B) and early-y (C) approaches. (A) Orchard 1(gray
606 line, $Y=265+0.6X$, $R^2=0.65^{***}$; RMSE=70 μm , $n=101$) and orchard 2(gray dash line,
607 $Y=272+0.6X$, $R^2=0.61^{***}$, RMSE=67 μm , $n=70$). (B) Orchard 1 (gray line,
608 $Y=190+0.2X$, $R^2=0.65^{***}$, RMSE=25 μm , $n=101$) and orchard 2 (gray dash line,
609 $Y=181+0.43X$, $R^2=0.61^{***}$. RMSE=48 μm , $n=70$). (C) Orchard 1(gray line,
610 $Y=114+0.6X$, $R^2=0.65^{***}$, RMSE=70 μm , $n=101$) and Orchard 2 (gray dash line,
611 $Y=152+0.6X$, $R^2=0.61^{***}$, RMSE=67 μm , $n=70$).

612

613

614

615

616

617

618

619

620

621

622

623