

Splash singularity for water waves

Angel Castro^a, Diego Córdoba^b, Charles L. Fefferman^{c,1}, Francisco Gancedo^d, and Javier Gómez-Serrano^e

^aInstituto de Ciencias Matemáticas, Consejo Superior de Investigaciones Científicas, C/ Nicolás Cabrera, 13-15, Campus Cantoblanco, 28049 Madrid, Spain; ^bInstituto de Ciencias Matemáticas, Consejo Superior de Investigaciones Científicas, C/ Nicolás Cabrera, 13-15, Campus Cantoblanco, 28049 Madrid, Spain; ^cDepartment of Mathematics, Princeton University, 1102 Fine Hall, Washington Road, Princeton, NJ 08544; ^dUniversity of Chicago, 5734 University Avenue, Chicago, IL 60637; and ^eInstituto de Ciencias Matemáticas, Consejo Superior de Investigaciones Científicas, C/ Nicolás Cabrera, 13-15, Campus Cantoblanco, 28049 Madrid, Spain

Contributed by Charles L. Fefferman, October 6, 2011 (sent for review July 4, 2011)

We exhibit smooth initial data for the two-dimensional (2D) water-wave equation for which we prove that smoothness of the interface breaks down in finite time. Moreover, we show a stability result together with numerical evidence that there exist solutions of the 2D water-wave equation that start from a graph, turn over, and collapse in a splash singularity (self-intersecting curve in one point) in finite time.

blow-up | Euler | incompressible | free boundary

We consider the two-dimensional (2D) water-wave equation, which governs the motion of the interface between a 2D inviscid incompressible irrotational fluid and a vacuum, taking gravity into account but neglecting surface tension. We prove that an initially smooth interface may in finite time become singular by the mechanism illustrated in Fig. 1. We call such a singularity a “splash.” We also present numerical evidence for a scenario in which the interface starts out as a smooth graph, then “turns over” after finite time, and finally produces a splash, as in Fig. S1.

The equations of motion in \mathbb{R}^2 for the density $\rho = \rho(x, t)$, $x \in \mathbb{R}^2$, $t \geq 0$, the velocity $v = (v^1, v^2)$ and the pressure $p = p(x, t)$ are:

$$\begin{cases} \rho(v_t + v \cdot \nabla v) = -\nabla p - (0, \rho), \\ \rho_t + v \cdot \nabla \rho = 0, \\ \nabla \cdot v = 0. \end{cases} \quad [1]$$

Above, the acceleration due to gravity is taken equal to one for the sake of simplicity.

The free boundary is parameterized by

$$\partial\Omega^j(t) = \{z(\alpha, t) = (z_1(\alpha, t), z_2(\alpha, t)) : \alpha \in \mathbb{R}\},$$

where the regions $\Omega^j(t)$ are defined by

$$\rho(x_1, x_2, t) = \begin{cases} 0, & x \in \Omega^1(t) \\ 1, & x \in \Omega^2(t) = \mathbb{R}^2 - \Omega^1(t). \end{cases}$$

We assume that the fluid is irrotational; i.e., the vorticity $\nabla^\perp \cdot v = 0$, in the interior of each domain Ω^j ($j = 1, 2$). The vorticity will be supported on the free boundary curve $z(\alpha, t)$ and it has the form

$$\nabla^\perp \cdot v(x, t) = \omega(\alpha, t) \delta(x - z(\alpha, t)),$$

i.e., the vorticity is a Dirac measure defined by

$$\langle \nabla^\perp \cdot v, \eta \rangle = \int_{\mathbb{R}} \omega(\alpha, t) \eta(z(\alpha, t)) d\alpha,$$

with $\eta(x)$ a test function. We present results for the following geometries: open curves asymptotic to the horizontal at infinity, periodic curves in the horizontal variable and closed contours.

However, in this paper we will only deal with the periodic case as the others are similar. One method to derive the equations for the evolution of v, ρ is to write the velocity as the orthogonal gradient of the stream function, take the curl and recover the velocity

by inverting the Laplacian; i.e., we apply the Biot-Savart law. Here we use the fact that the vorticity is concentrated on the interface;

$$v(x, t) = \nabla^\perp \Delta^{-1}(\nabla^\perp \cdot v)(x, t) = \frac{1}{2\pi} \int_{\mathbb{R}} \frac{(x - z(\alpha, t))^\perp}{|x - z(\alpha, t)|^2} \omega(\alpha, t) d\alpha,$$

with $x \neq z(\alpha, t)$.

Taking limits of the above equation, by approaching the boundary in the normal direction, we obtain the velocity of the interface, to which we can add any term c in the tangential direction z_α without modifying the geometry of the interface. Thus the interface satisfies

$$z_t(\alpha, t) = BR(z, \omega)(\alpha, t) + c(\alpha, t)z_\alpha(\alpha, t), \quad [2]$$

where the Birkhoff-Rott integral is defined by

$$BR(z, \omega) = \frac{1}{2\pi} P.V. \int_{\mathbb{R}} \frac{(z(\alpha, t) - z(\beta, t))^\perp}{|z(\alpha, t) - z(\beta, t)|^2} \omega(\alpha, t) d\alpha.$$

The system is closed by using Euler equations;

$$\begin{aligned} \omega_t(\alpha, t) &= -2\partial_t BR(z, \omega)(\alpha, t) \cdot \partial_\alpha z(\alpha, t) \\ &\quad - \partial_\alpha \left(\frac{|\omega|^2}{4|\partial_\alpha z|^2} \right)(\alpha, t) + \partial_\alpha (c\omega)(\alpha, t) \\ &\quad + 2c(\alpha, t) \partial_\alpha BR(z, \omega)(\alpha, t) \cdot \partial_\alpha z(\alpha, t) - 2\partial_\alpha z_2(\alpha, t). \end{aligned} \quad [3]$$

Then the dynamic equations, for the interface $z = z(\alpha, t)$ and the vorticity are the system given by references [2] and [3] and are known as the water-wave equations.

Taking the divergence of the Euler Eq. 1 and recalling that the flow is irrotational in the interior of the regions $\Omega^j(t)$, we find that

$$-\Delta p = |\nabla v|^2 \geq 0$$

which, together with the fact that the pressure is zero on the interface implies by Hopf's lemma in $\Omega^2(t)$ that

$$\sigma(\alpha, t) \equiv -|z_\alpha^\perp(\alpha, t)| \partial_n p(z(\alpha, t), t) > 0,$$

where ∂_n denotes the normal derivative. This inequality is known as the Rayleigh-Taylor condition (see refs. 1 and 2) which was first proved by Wu in refs. 3 and 4.

The first results concerning the Cauchy problem for the linearized version of water waves and small data in Sobolev spaces are due to Craig (5), Nalimov (6), Beale, et al. (7) and Yoshida

Author contributions: A.C., D.C., C.L.F., F.G., and J.G.-S. designed research; performed research; and wrote the paper.

The authors declare no conflict of interest.

¹To whom correspondence should be addressed. E-mail: cf@math.princeton.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1115948108/-DCSupplemental.

(8). The well posedness in Sobolev spaces for the water-wave problem was proven by Wu in ref. 3 with the assumption that the initial free interface is non-self-intersecting (satisfies the arc-chord condition). For recent work on local existence see Wu (4), Christodoulou-Lindblad (9), Lindblad (10), Coutand-Shkoller (11), Shatah-Zeng (12), Zhang-Zhang (13), Córdoba et al. (14), Lannes (15, 16), Alazard-Metivier (17), and Ambrose-Masmoudi (18). The issue of long time existence has been treated in Alvarez-Lannes (19) where well posedness over large time scales is shown and different asymptotic regimes are justified. Wu proved in ref. 20 exponential time of existence in two dimensions for small initial data, and Germain, et al. in ref. 21 and Wu in ref. 22 global existence for small data in the three dimensional case (2D interface). In refs. 23 and 24, Castro, et al. showed that there exist large initial data parameterized as a graph for which in finite time the interface reaches a regime in which it is no longer a graph. For previous numerical simulations showing this phenomenon see Beale, et al. (25).

The outline of the paper is the following: in *The Equations in the Tilde Domain* we will describe the equations in a transformed domain which will circumvent the problem of having a singularity where the arc-chord condition fails as the curve self-intersects; i. e., a splash singularity forms. In *Local Existence at the Splash* we will outline the proof of a local existence theorem for the equations in the new domain, both for the analytic case and for Sobolev spaces. *Structural Stability* will be devoted to a stability theorem, whereas *Numerical Results* will comment on the numerical results obtained towards the splash singularity starting from a graph. Finally in *Further Research* we describe the ideas that we hope will lead to a computer-assisted proof of the existence of a solution that starts as a graph and ends in a splash.

The Equations in the Tilde Domain

In this section we will rewrite the equations by applying a transformation from the original coordinates to new ones which we will denote with a tilde. The purpose of this transformation is to be able to deal with the failure of the arc-chord condition. We start by reformulating the set of equations, in the nontilde domain, for the case of a periodic contour in terms of the velocity potential. From [1] and because v is irrotational in $\Omega^2(t)$ we have that:

$$\begin{aligned} \Delta\psi(x,y,t) &= 0 \quad \text{in } \Omega^2(t) & \partial_n\psi|_{z(\alpha,t)} &= -\frac{\Phi_\alpha(\alpha,t)}{|z_\alpha(\alpha,t)|} \\ \psi(x+2\pi,y,t) &= \psi(x,y,t) \quad \text{in } \Omega^2(t) & \psi(x,y) & \text{ is } O(1) \text{ as } y \rightarrow -\infty \\ v &\equiv \nabla^\perp\psi \quad \text{in } \Omega^2(t) & z_t(\alpha,t) &= u(\alpha,t) + c(\alpha,t)z_\alpha(\alpha,t) \\ \phi(x,y,t) & \text{ is the harmonic conjugate of } \psi(x,y,t) \quad \text{in } \Omega^2(t) \\ \Phi_t(\alpha,t) &= \frac{1}{2}|u(\alpha,t)|^2 + c(\alpha,t)u(\alpha,t) \cdot z_\alpha(\alpha,t) - z_2(\alpha,t) + p^*(t) \\ z(\alpha,0) &= z^0(\alpha) & \Phi_\alpha(\alpha,0) &= \Phi_\alpha^0(\alpha) \end{aligned} \quad [4]$$

where ϕ is the velocity potential, $\Phi(\alpha,t)$ is its limit at the interface coming from the fluid region, $\nabla\phi = v$, p^* is a function of t alone, c is a free quantity which represents the reparameterization freedom, $u(\alpha,t)$ is the limit of the velocity at the interface coming from the fluid region, ψ is the stream function, and $\Phi_\alpha^0(\alpha)$ has zero mean.

Although we may take as an initial condition the tangential component of the velocity multiplied by the modulus of the tangent vector; i.e., Φ_{α_t} , we can also solve the system [4] by taking as an initial condition the normal component of the initial velocity multiplied by the modulus of the normal vector; i.e., Ψ_α , ($\Psi(\alpha,t) = \psi(z(\alpha,t),t)$), as we can transform one into the other.

Let us consider $\tilde{z}(\alpha,t) = (\tilde{z}_1(\alpha,t), \tilde{z}_2(\alpha,t)) \equiv P(z(\alpha,t))$ where P is a conformal map defined in the water region that will be given as:

$$P(w) = \left(\tan\left(\frac{w}{2}\right) \right)^{1/2}, \quad w \in \mathbb{C},$$

for a branch of the square root that separates the self-intersecting points of the interface. Here $P(w)$ will refer to a 2D vector whose components are the real and imaginary parts of $P(w_1 + iw_2)$. In this setting, $P^{-1}(z)$ will be well defined modulo multiples of 2π .

The water-wave equations are invariant under time reversal. To obtain a solution that ends in a splash, we can therefore take our initial condition to be a splash, and show that there is a smooth solution for small times $t > 0$. As initial data we are interested in considering a curve that intersects itself at one point, as in Fig. S1. More precisely, we will use as initial data *splash curves* which are defined as follows:

Definition 2.1: We say that $z(\alpha)$ is a *splash curve* if

- $z_1(\alpha) - \alpha, z_2(\alpha)$ are smooth functions and 2π -periodic.
- $z(\alpha)$ satisfies the arc-chord condition at every point except at α_1 and α_2 , with $\alpha_1 < \alpha_2$ where $z(\alpha_1) = z(\alpha_2)$ and $|z_\alpha(\alpha_1)|, |z_\alpha(\alpha_2)| > 0$.
- The curve $z(\alpha)$ separates the complex plane into two regions; a connected water region and a vacuum region. The water region contains each point $x + iy$ for which y is large negative. We choose the parametrization such that the normal vector $n = \frac{(-\partial_\alpha z^2(\alpha), \partial_\alpha z^1(\alpha))}{|\partial_\alpha z(\alpha)|}$ points to the vacuum region.
- We can choose a branch of the function P on the water region such that the curve $\tilde{z}(\alpha) = P(z(\alpha))$ satisfies:

$\tilde{z}_1(\alpha)$ and $\tilde{z}_2(\alpha)$ are smooth and 2π -periodic.

\tilde{z} is a closed contour.

\tilde{z} satisfies the arc-chord condition.

We will choose the branch of the root that produces that

$$\lim_{y \rightarrow -\infty} P(x + iy) = -e^{-i\pi/4}$$

independently of x .

- $P(w)$ is analytic in w and $\frac{dP}{dw}(w) \neq 0$ if w belongs to the water region.
- $\tilde{z}(\alpha) \neq q^l$ for $l = 0, \dots, 4$, where

$$q^0 = (0,0), \quad q^l = \left(\pm \frac{1}{\sqrt{2}}, \pm \frac{1}{\sqrt{2}} \right), \quad l = 1,2,3,4. \quad [5]$$

From now on, we will always work with splash curves as initial data. Condition 6 will be used in the local existence theorems and can be proved to hold for short enough time as long as the initial condition satisfies it. We will also need that the interface passes below the points $(\pm\pi, 0)$ (or, equivalently, that those points belong to the vacuum region) in order for the tilde region to be a closed curve and the vacuum region to lie on the outer part. For a splash curve this is trivial from the definition. For more information about the transformation of both regions, check Figs. S1 and S2 and notice that we rule out the scenarios in Fig. S3.

We will now write [4] in the new tilde coordinates. We define the following quantities:

$$\tilde{\psi}(x,y,t) \equiv \psi(P^{-1}(x,y),t),$$

$$\tilde{\phi}(x,y,t) \equiv \phi(P^{-1}(x,y),t),$$

$$\tilde{v}(x,y,t) \equiv \nabla\tilde{\phi}(x,y,t).$$

Let us note that as ψ and ϕ are 2π periodic, the resulting $\tilde{\psi}$ and $\tilde{\phi}$ are well defined. We do not have problems with the harmonicity of $\tilde{\psi}$ or $\tilde{\phi}$ at the point which is mapped from minus infinity (which belongs to the water region) by P as ϕ and ψ are well defined at infinity. Also, the periodicity of ϕ and ψ causes $\tilde{\phi}$ and $\tilde{\psi}$ to be continuous (and harmonic) at the interior of $P(\Omega^2(t))$.




Fig. 1. Splash singularity. The interface collapses at one point.

Let us assume that there exists a solution of [4] and that we take $u_n = \frac{\Psi_n}{|z_n|}$ such that $u_n(\alpha_1), u_n(\alpha_2) < 0$ for all $0 < t < T, T$ small enough, thus $z(\alpha, t)$ satisfies the arc-chord condition and does not touch the removed branch from $P(w)$.

The water waves system in the new coordinates reads

$$\tilde{z}_t(\alpha, t) = Q^2(\alpha, t)BR(\tilde{z}, \tilde{\omega})(\alpha, t) + \tilde{c}(\alpha, t)\tilde{z}_\alpha(\alpha, t) \quad [6]$$

and the evolution equation for $\tilde{\omega}$

$$\begin{aligned} \tilde{\omega}_t(\alpha, t) = & -2\partial_t BR(\tilde{z}, \tilde{\omega})(\alpha, t) \cdot \tilde{z}_\alpha(\alpha, t) - |BR(\tilde{z}, \tilde{\omega})|^2 \partial_\alpha Q^2(\alpha, t) \\ & - \partial_\alpha \left(\frac{Q^2(\alpha, t)}{4} \frac{\tilde{\omega}(\alpha, t)^2}{|\tilde{z}_\alpha(\alpha, t)|^2} \right) + 2\tilde{c}(\alpha, t)\partial_\alpha BR(\tilde{z}, \tilde{\omega}) \cdot \tilde{z}_\alpha(\alpha, t) \\ & + \partial_\alpha (\tilde{c}(\alpha, t)\tilde{\omega}(\alpha, t)) - 2\partial_\alpha (P_2^{-1}(\tilde{z}(\alpha, t))), \end{aligned} \quad [7]$$

where $Q^2(\alpha, t) = \left| \frac{dP}{dw}(z(\alpha, t)) \right|^2$.

Remark 2.2: Eq. 7 is analogous to [3]. In fact, if we set $Q \equiv 1$ in [7] we recover [3].

Note that for the tilde domain, the Rayleigh-Taylor condition is the same as in the first domain; i.e:

$$\nabla p(\alpha, t) \cdot z_\alpha^\perp(\alpha, t) = \nabla \tilde{p}(\alpha, t) \cdot \tilde{z}_\alpha^\perp(\alpha, t),$$

where $\tilde{p} = p \circ P^{-1}$.

Our strategy will be the following: we will consider the evolution of the solutions in the tilde domain and then see that everything works fine in the original domain.

We will have to obtain the normal velocity once given the tangential velocity, and vice versa:

$$\tilde{\Phi}_\alpha(\alpha, t) = \tilde{u}(\alpha, t) \cdot \tilde{z}_\alpha(\alpha, t) = BR(\tilde{z}, \tilde{\omega}) \cdot \tilde{z}_\alpha(\alpha, t) + \frac{\tilde{\omega}(\alpha, t)}{2}.$$

From the previous formula, we can invert the equation and get $\tilde{\omega}$, plug it into the following expression for $\tilde{\psi}$:

$$\tilde{\psi}(\tilde{x}) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \log(|\tilde{x} - \tilde{z}(\alpha)|) \tilde{\omega}(\alpha) d\alpha$$

and restrict ourselves to the interface to get $\tilde{\Psi}(\alpha, t)$. Taking a derivative in α we can recover the normal component of the velocity. An analogous reasoning can be done to get the tangential velocity from the normal by solving the complementary Neumann problem for ϕ .

We now note that a solution of the system [6]–[7] in the tilde domain gives rise to a solution of the system [4] in the nontilde domain, by inverting the map P . This fact will be the implication used in Theorem 3.1 (finding a solution in the tilde domain, and therefore in the nontilde).

Remark 2.3: A similar argument works for the other two settings (closed contour and asymptotic to horizontal) by choosing an appropriate $P(w)$ that separates the singularity.

Local Existence at the Splash

The main result in this section is a local existence proof for the splash singularity. To avoid the arc-chord condition failure, we will prove the local existence in the tilde domain in two different settings, namely in the space of analytic functions and the Sobolev space H^s .

Theorem 3.1. Let $z^0(\alpha)$ be a splash curve such that $z^0(\alpha) - \alpha, z^0_2(\alpha) \in H^4(\mathbb{T})$. Let $u^0(\alpha) \cdot (z^0_\alpha)^\perp(\alpha) \in H^4(\mathbb{T})$ satisfying:

- $\left(u^0 \cdot \frac{(z^0_\alpha)^\perp}{|z^0_\alpha|} \right)(\alpha_1) < 0, \quad \left(u^0 \cdot \frac{(z^0_\alpha)^\perp}{|z^0_\alpha|} \right)(\alpha_2) < 0,$
- $\int_{\partial\Omega} u^0 \cdot \frac{(z^0_\alpha)^\perp}{|z^0_\alpha|} ds = \int_{\mathbb{T}} u^0(\alpha) \cdot (z^0_\alpha)^\perp d\alpha = 0.$

Then there exist a finite time $T > 0$, a time-varying curve $z(\alpha, t) \in C([0, T]; H^4(\mathbb{T}))$ satisfying:

- $z_1(\alpha, t) - \alpha, z_2(\alpha, t)$ are 2π -periodic,
- $z(\alpha, t)$ satisfies the arc-chord condition for all $t \in (0, T]$,

and $u(\alpha, t) \in C([0, T]; H^3(\mathbb{T}))$ which provides a solution of the water-wave Eq. 4 with $z(\alpha, 0) = z^0(\alpha)$ and $u(\alpha, 0) \cdot (z^0_\alpha)^\perp(\alpha, 0) = u^0(\alpha) \cdot (z^0_\alpha)^\perp(\alpha)$.

Sketch of the proof: Using the fact that there is local existence to the initial data in the tilde domain and applying P^{-1} to the solution obtained there, we can get a curve $z(\alpha, t)$ that solves the water waves equation in the nontilde domain, which leads to the proof of Theorem 3.1. Details on the local existence in the tilde domain are shown below.

Local Existence for Analytic Initial Data in the Tilde Domain. In this subsection, we will work on the tilde domain, and all tildes will be dropped for the sake of simplicity.

We will work with $c = 0$. We have the following system:

$$\begin{cases} z_t = \left| \frac{dP}{dw}(P^{-1}(z)) \right|^2 u \\ \Phi_t = \frac{1}{2} \left| \frac{dP}{dw}(P^{-1}(z)) \right|^2 |u|^2 - P_2^{-1}(z) \\ u = BR(z, \omega) + \frac{\omega}{2|z_\alpha|^2} z_\alpha \\ \Phi_\alpha = \frac{\omega}{2} + BR(z, \omega) \cdot z_\alpha \\ \left| \frac{dP}{dw}(P^{-1}(z(\alpha, t))) \right|^2 = \frac{1}{16} \left| \frac{1+(z_1(\alpha, t)+iz_2(\alpha, t))^4}{z_1(\alpha, t)+iz_2(\alpha, t)} \right|^2 \\ P_2^{-1}(z(\alpha, t)) = \ln \left| \frac{i+(z_1(\alpha, t)+iz_2(\alpha, t))^2}{i-(z_1(\alpha, t)+iz_2(\alpha, t))^2} \right| \end{cases} \quad [8]$$

We demand that $z^0(\alpha) \neq (0, 0)$ to find the function $\frac{dP}{dw}(P^{-1}(z(\alpha, t)))$ well defined. This condition is going to remain true for short time. We also consider $z^0(\alpha) \neq q^l, l = 1, \dots, 4$ in [5] to get $P_2^{-1}(z(\alpha, t))$ well defined which is going to remain true for short time.

We consider the space

$$\begin{aligned} H^3(\partial S_r) = & \left\{ f \text{ analytic in } S_r = \{\alpha + i\eta, |\eta| < r\}: \|f\|_r^2 < \infty \right. \\ & \|f\|_r^2 = \|f\|_{L^2(\partial S_r)}^2 + \|\partial_\alpha^3 f\|_{L^2(\partial S_r)}^2 \quad \text{where} \\ & \left. \|f\|_{L^2(\partial S_r)}^2 = \sum_{\pm} \int_{-\pi}^{\pi} |f(\alpha \pm ir)|^2 d\alpha, f \text{ } 2\pi\text{-periodic} \right\} \end{aligned}$$

and $(z, \Phi) \in (H^3(\partial S_r))^3 \equiv X_r$.

We have the following theorem:

Theorem 3.2. Let $z^0(\alpha)$ be a splash curve and let $u^0 \cdot \frac{z_\alpha^0}{|z_\alpha^0|}(\alpha) = \frac{\Phi_\alpha^0}{|z_\alpha^0|}(\alpha)$ be the initial tangential velocity such that

$$(z^0(\alpha) - \alpha, z_\alpha^0(\alpha), \Phi^0(\alpha)) \in X_{r_0},$$

for some $r_0 > 0$, and satisfying:

1. $\left(u^0 \cdot \frac{(z_\alpha^0)^\perp}{|z_\alpha^0|}\right)(\alpha_1) < 0, \quad \left(u^0 \cdot \frac{(z_\alpha^0)^\perp}{|z_\alpha^0|}\right)(\alpha_2) < 0,$
2. $\int_{\partial\Omega} u^0 \cdot \frac{(z_\alpha^0)^\perp}{|z_\alpha^0|} ds = \int_{\mathbb{T}} u^0(\alpha) \cdot (z_\alpha^0)^\perp d\alpha = 0.$

Then there exist a finite time $T > 0$, $0 < r < r_0$, a time-varying curve $\tilde{z}(\alpha, t)$ satisfying:

1. $P^{-1}(\tilde{z}_1(\alpha, t)) - \alpha, P^{-1}(\tilde{z}_2(\alpha, t))$ are 2π -periodic,
 2. $P^{-1}(\tilde{z}(\alpha, t))$ satisfies the arc-chord condition for all $t \in (0, T]$,
- and $\tilde{u}(\alpha, t)$ with

$$(\tilde{z}_1(\alpha, t), \tilde{z}_2(\alpha, t), \tilde{\Phi}(\alpha, t)) \in C([0, T], X_r)$$

which provides a solution of the water waves Eq. 8 with $\tilde{z}^0(\alpha) = P(z^0(\alpha))$ and $\tilde{u}(\alpha, 0) \cdot (\tilde{z}_\alpha)^{\perp}(\alpha, 0) = \tilde{u}^0(\alpha) \cdot (z^0)^\perp_\alpha(\alpha)$.

The main tool in the proof is to use an abstract Cauchy-Ko-walewski theorem from refs. 26 and 27 as in for example ref. 23.

Local Existence for Initial Data in Sobolev Spaces in the Tilde Domain. We will take the following $\tilde{c}(\alpha, t)$:

$$\begin{aligned} \tilde{c}(\alpha, t) &= \frac{\alpha + \pi}{2\pi} \int_{-\pi}^{\pi} (Q^2 BR(\tilde{z}, \tilde{\omega}))_\beta(\beta, t) \cdot \frac{\tilde{z}_\beta(\beta, t)}{|\tilde{z}_\beta(\beta, t)|^2} d\beta \\ &\quad - \int_{-\pi}^{\alpha} (Q^2 BR(\tilde{z}, \tilde{\omega}))_\beta(\beta, t) \cdot \frac{\tilde{z}_\beta(\beta, t)}{|\tilde{z}_\beta(\beta, t)|^2} d\beta. \end{aligned}$$

We will also define an auxiliary function $\tilde{\varphi}(\alpha, t)$ analogous to the one introduced in ref. 7 (for the linear case) and ref. 18 (non-linear case) which helps us to bound several of the terms that appear:

$$\tilde{\varphi}(\alpha, t) = \frac{Q^2(\alpha, t) \tilde{\omega}(\alpha, t)}{2|\tilde{z}_\alpha(\alpha, t)|} - \tilde{c}(\alpha, t) |\tilde{z}_\alpha(\alpha, t)|. \quad [9]$$

Theorem 3.3. Let $z^0(\alpha)$ be a splash curve such that $z^0_1(\alpha) - \alpha, z^0_2(\alpha) \in H^4(\mathbb{T})$. Let $u^0(\alpha) \cdot (z_\alpha^0)^\perp(\alpha) \in H^4(\mathbb{T})$ satisfying:

1. $\left(u^0 \cdot \frac{(z_\alpha^0)^\perp}{|z_\alpha^0|}\right)(\alpha_1) < 0, \quad \left(u^0 \cdot \frac{(z_\alpha^0)^\perp}{|z_\alpha^0|}\right)(\alpha_2) < 0,$
2. $\int_{\partial\Omega} u^0 \cdot \frac{(z_\alpha^0)^\perp}{|z_\alpha^0|} ds = \int_{\mathbb{T}} u^0(\alpha) \cdot (z_\alpha^0)^\perp d\alpha = 0.$

Then there exist a finite time $T > 0$, a time-varying curve $\tilde{z}(\alpha, t) \in C([0, T]; H^4)$ satisfying:

1. $P^{-1}(\tilde{z}_1(\alpha, t)) - \alpha, P^{-1}(\tilde{z}_2(\alpha, t))$ are 2π -periodic,
 2. $P^{-1}(\tilde{z}(\alpha, t))$ satisfies the arc-chord condition for all $t \in (0, T]$,
- and $\tilde{u}(\alpha, t) \in C([0, T]; H^3(\mathbb{T}))$ which provides a solution of the water waves Eqs. 6-7 with $\tilde{z}^0(\alpha) = P(z^0(\alpha))$ and $\tilde{u}(\alpha, 0) \cdot (\tilde{z}_\alpha)^\perp(\alpha, 0) = \tilde{u}^0(\alpha) \cdot (z^0)^\perp_\alpha(\alpha)$.

Sketch of the Proof: In the proof, for the sake of simplicity, we will drop the tildes from the notation.

The proof will use the properties of $c(\alpha, t)$ and $\varphi(\alpha, t)$ to get an extra cancellation to help us derive energy estimates. Moreover, this choice of c will ensure that the length of the tangent vector of $z(\alpha, t)$ depends only on time.

Here we define the energy $E(t)$ by

$$\begin{aligned} E(t) &= \|z\|_{H^3}^2(t) + \int_{\mathbb{T}} \frac{Q^2 \sigma_z}{|z_\alpha|^2} |\partial_\alpha^4 z|^2 d\alpha + \|F(z)\|_{L^\infty}^2(t) + \|\omega\|_{H^2}^2(t) \\ &\quad + \|\varphi\|_{H^{3+\frac{1}{2}}}^2(t) + \frac{|z_\alpha|^2}{m(Q^2 \sigma_z)(t)} + \sum_{l=0}^4 \frac{1}{m(q^l)(t)}, \end{aligned}$$

where the L^∞ norm of the function

$$F(z) \equiv \frac{|\beta|}{|z(\alpha, t) - z(\alpha - \beta, t)|}, \quad \alpha, \beta \in \mathbb{T}$$

measures the arc-chord condition,

$$\begin{aligned} \sigma_z &\equiv \left(BR_l(z, \omega) + \frac{\varphi}{|z_\alpha|} BR_\alpha(z, \omega) \right) \cdot z_\alpha^\perp + \frac{\omega}{2|z_\alpha|^2} \left(z_\alpha + \frac{\varphi}{|z_\alpha|} z_\alpha \right) \\ &\quad \cdot z_\alpha^\perp + Q \left| BR(z, \omega) + \frac{\omega}{2|z_\alpha|^2} z_\alpha \right|^2 (\nabla Q)(z) \cdot z_\alpha^\perp + (\nabla P_2^{-1})(z) \cdot z_\alpha^\perp \end{aligned}$$

[10]

is the Rayleigh-Taylor function,

$$m(Q^2 \sigma_z)(t) \equiv \min_{\alpha \in \mathbb{T}} Q^2(\alpha, t) \sigma_z(\alpha, t),$$

and finally

$$m(q^l)(t) \equiv \min_{\alpha \in \mathbb{T}} |z(\alpha, t) - q^l|$$

for $l = 0, \dots, 4$. We proceed as in ref. 14: The bound for the operator $(I + J)^{-1}$, where $J\omega = 2BR(z, \omega) \cdot z_\alpha$, and some rather routine estimates allow us to find

$$\begin{aligned} \frac{d}{dt} \left(\|z\|_{H^3}^2(t) + \|F(z)\|_{L^\infty}^2(t) + \|\omega\|_{H^2}^2(t) + \|\varphi\|_{L^2}^2(t) \right. \\ \left. + \frac{|z_\alpha|^2}{m(Q^2 \sigma_z)(t)} + \sum_{l=0}^4 \frac{1}{m(q^l)(t)} \right) \leq CE^k(t), \end{aligned}$$

for C and k universal constants. Above we use that

$$\begin{aligned} \|\partial_\alpha^4 z\|_{L^2}^2(t) &= \int_{\mathbb{T}} \frac{Q^2 \sigma_z |z_\alpha|^2}{Q^2 \sigma_z |z_\alpha|^2} |\partial_\alpha^4 z|^2 d\alpha \\ &\leq \frac{|z_\alpha|^2}{m(Q^2 \sigma_z)} \int_{\mathbb{T}} \frac{Q^2 \sigma_z}{|z_\alpha|^2} |\partial_\alpha^4 z|^2 d\alpha \leq E^2(t). \end{aligned}$$

Further we obtain

$$\frac{d}{dt} \left(\int_{\mathbb{T}} \frac{Q^2 \sigma_z}{|z_\alpha|^2} |\partial_\alpha^4 z|^2 d\alpha \right) \leq CE^k(t) + S(t),$$

where

$$S(t) = 2 \int_{\mathbb{T}} \frac{Q^2 \sigma_z}{|z_\alpha|^2} \partial_\alpha^4 z \cdot \frac{z_\alpha^\perp}{|z_\alpha|} H(\partial_\alpha^4 \varphi) d\alpha.$$

We use [9], [7], and [10] to get

$$\partial_\alpha^3 \varphi_t = -\frac{Q^2 \sigma_z}{|z_\alpha|^2} \partial_\alpha^4 z \cdot \frac{z_\alpha^\perp}{|z_\alpha|} + \text{“control”},$$

where “control” is given by lower order terms and unbounded terms (such as $\partial_\alpha^4 \varphi$) that can be estimated with energy methods in terms of $E(t)$. Therefore, it allows us to get

$$\frac{d}{dt} \|\Lambda^{1/2} \partial_\alpha^3 \varphi\|_{L^2}^2(t) \leq CE^k(t) - S(t)$$

which together with above inequalities yields

$$\left| \frac{d}{dt} E(t) \right| \leq CE^k(t).$$

Local existence follows using standard arguments with the apriori energy estimate.

Structural Stability

Again, in this section, we will omit the tildes from the notation. This section is devoted to establish a stability result. This result will allow us to conclude the following: if (x, γ) approximately satisfies Eq. 11, then near to (x, γ) there exists an exact solution (z, ω) . Below is the theorem.

Theorem 4.1. *Let*

$$D(\alpha, t) \equiv z(\alpha, t) - x(\alpha, t),$$

$$d(\alpha, t) \equiv \omega(\alpha, t) - \gamma(\alpha, t),$$

$$\mathcal{D}(\alpha, t) \equiv \varphi(\alpha, t) - \zeta(\alpha, t),$$

where (x, γ, ζ) are the solutions of

$$\begin{cases} x_t = Q^2(x)BR(x, \gamma) + bx_\alpha + f \\ b = \underbrace{\frac{\alpha + \pi}{2\pi} \int_{-\pi}^{\pi} (Q^2 BR(x, \gamma))_\alpha \frac{x_\alpha}{|x_\alpha|^2} d\alpha}_{b_s} - \underbrace{\int_{-\pi}^{\alpha} (Q^2 BR(x, \gamma))_\beta \frac{x_\alpha}{|x_\alpha|^2} d\beta + \frac{\alpha + \pi}{2\pi} \int_{-\pi}^{\pi} f_\alpha \frac{x_\alpha}{|x_\alpha|^2} d\alpha - \int_{-\pi}^{\alpha} f_\beta \frac{x_\beta}{|x_\beta|^2} d\beta}_{b_e} \\ \gamma_t + 2BR_t(x, \gamma) \cdot x_\alpha = -(Q^2(x))_\alpha |BR(x, \gamma)|^2 + 2bBR_\alpha(x, \gamma) \cdot x_\alpha + (b\gamma)_\alpha - \left(\frac{Q^2(x)\gamma^2}{4|x_\alpha|^2} \right)_\alpha - 2(P_2^{-1}(x))_\alpha + g \\ \zeta(\alpha, t) = \frac{Q_\alpha^2(x, \gamma)}{2|x_\alpha(\alpha, t)|} - b_s(\alpha, t)|x_\alpha(\alpha, t)|, \end{cases} \quad [11]$$

(z, ω, φ) are the solutions of [11] with $f \equiv g \equiv 0$ and \mathcal{E} the following norm for the difference

$$\mathcal{E}(t) \equiv \left(\|D\|_{H^3}^2 + \int_{-\pi}^{\pi} \frac{Q^2 \sigma_z}{|z_\alpha|^2} |\partial_\alpha^4 D|^2 d\alpha + \|d\|_{H^2}^2 + \|\mathcal{D}\|_{H^{3+1/2}}^2 \right).$$

Then we have that

$$\left| \frac{d}{dt} \mathcal{E}(t) \right| \leq \mathcal{E}(t)(\mathcal{E}(t) + \delta(t)),$$

where

$$\mathcal{E}(t) = \mathcal{E}(E(t), \|x\|_{H^{5+1/2}}(t), \|\gamma\|_{H^{4+1/2}}(t), \|\zeta\|_{H^{4+1/2}}(t), \|F(x)\|_{L^\infty}(t))$$

and

$$\delta(t) = (\|f\|_{H^{5+1/2}}(t) + \|g\|_{H^{3+1/2}}(t))^k + (\|f\|_{H^{5+1/2}}(t) + \|g\|_{H^{3+1/2}}(t))^2,$$

with k big enough. Here $E(t)$ is defined in the proof of Theorem 3.3.

Sketch of the Proof: The equation for (x, γ, ζ) is the same as the one for (z, ω, φ) but for f and g . The function b is chosen in such a way that $|x_\alpha|$ only depends on time which allows us to get the following estimates:

$$\frac{d}{dt} (\|D\|_{H^3}^2 + \|d\|_{H^2}^2 + \|\mathcal{D}\|_{L^2}^2) \leq \mathcal{E}(t)(\mathcal{E}(t) + \delta(t)).$$

Further we obtain

$$\frac{d}{dt} \left(\int_{\mathbb{T}} \frac{Q^2 \sigma_z}{|z_\alpha|^2} |\partial_\alpha^4 D|^2 d\alpha \right) \leq \mathcal{E}(t)(\mathcal{E}(t) + \delta(t)) + \mathcal{S}(t)$$

with

$$\mathcal{S}(t) = 2 \int_{\mathbb{T}} \frac{Q^2 \sigma_z}{|z_\alpha|^2} \partial_\alpha^4 D \cdot \frac{z_\alpha^\perp}{|z_\alpha|} H(\partial_\alpha^4 \mathcal{D}) d\alpha.$$

For \mathcal{D} one finds that

$$\partial_\alpha^3 \mathcal{D}_t = - \frac{Q^2 \sigma_z}{|z_\alpha|^2} \partial_\alpha^4 D \cdot \frac{z_\alpha^\perp}{|z_\alpha|} + \text{“control”},$$

where “control” denotes terms which can be estimated by $\mathcal{E}(t)(\mathcal{E}(t) + \delta(t))$, which yields

$$\frac{d}{dt} \|\Lambda^{1/2} \partial_\alpha^3 \mathcal{D}\|_{L^2}^2(t) \leq \mathcal{E}(t)(\mathcal{E}(t) + \delta(t)) - \mathcal{S}(t).$$

Then the desired estimate follows.

Numerical Results

In order to illustrate the splash singularity, several numerical simulations were performed. The simulations were done following the scheme proposed by Beale, Hou, and Lowengrub (25) adapted to the equations on the tilde domain (i.e., taking into account the impact of Q on the equation). Instead of having an evolution equation for ω , a velocity potential ϕ is introduced and its evolution through time subject to the constraint imposed by being a potential is studied. The initial data on the nontilde domain was given by:

$$\begin{aligned} z_1^0(\alpha) &= \alpha + \frac{1}{4} \left(-\frac{3\pi}{2} - 1.9 \right) \sin(\alpha) + \frac{1}{2} \sin(2\alpha) \\ &\quad + \frac{1}{4} \left(\frac{\pi}{2} - 1.9 \right) \sin(3\alpha) \\ z_2^0(\alpha) &= \frac{1}{10} \cos(\alpha) - \frac{3}{10} \cos(2\alpha) + \frac{1}{10} \cos(3\alpha). \end{aligned}$$

Note that $z(\frac{\pi}{2}) = z(-\frac{\pi}{2})$ (splash). Instead of prescribing an initial condition for ω , we prescribed the normal component of the velocity to ensure a more controlled direction of the fluid. From that we got the initial $\omega(\alpha, 0)$ using the following relations. Let ψ be such that $\nabla^\perp \psi = v$ and $\Psi(\alpha)$ its restriction to the interface. Recall that we can transform the initial condition on the normal component of the velocity into an initial condition on the tangential component by applying the transformations described in *The Equations in the Tilde Domain*. The initial normal velocity is then prescribed by setting

$$u_n^0(\alpha)|z_\alpha(\alpha)| = \Psi_\alpha(\alpha) = 3 \cdot \cos(\alpha) - 3.4 \\ \cdot \cos(2\alpha) + \cos(3\alpha) + 0.2 \cdot \cos(4\alpha).$$

The simulations were done using a spatial mesh of $N = 2,048$ nodes and a time step $\Delta t = 10^{-7}$. The time direction was set to run backwards (from the splash to the graph) and the graph was obtained at approximately $T_g = 6.5 \cdot 10^{-3}$. Note that the normal component of the velocity $u_n^0(\pm \frac{\pi}{2}) > 0$ at the splash, which satisfies the hypotheses of Theorem 3.1 as we are running time backwards. Getting the potential of the initial condition from ω_0 and z_0 and the transformation of all the initial data to the tilde domain is a trivial computation, taking care to choose the appropriate branch of the square root. See Figs. S1 and S2.

Further Research

We would like to exhibit a water-wave solution whose interface starts as an H^4 -smooth graph at time zero, and ends in a splash at time T . We sketch a few ideas that may lead to a rigorous computer-assisted proof of the existence of such a solution. We will work in the tilde domain.

A simulation as in *Numerical Results* leads to an approximate solution $x(\alpha, t)$, $\gamma(\alpha, t)$ with the desired properties. Thus $x(\cdot, t)$ describes a graph when $t = 0$ and a splash when $t = T$. Moreover, we believe that equations similar to [11] hold, with very small f and g .

We may suppose that x and γ are known piecewise-polynomial functions on $[0, 2\pi] \times [0, T]$. Using interval arithmetic (28), one can compute rigorous upper bounds for appropriate Sobolev norms of f and g . We hope that these upper bounds will be very small.

Next, we solve the water-wave Eqs. 2, 3 for z, ω , starting at time T , and proceeding backwards in time. We take our initial (z, ω) at time T to be a splash, very close to $(x(\cdot, T), \gamma(\cdot, T))$ in a high Sobolev norm $\|\cdot\|_{H^s}$.

We want to compare the exact solution (z, ω) with the approximate solution (x, γ) , using the quantity $\mathcal{E}(t)$ as in Theorem 4.1.

Because (z, ω) and (x, γ) are very close at time T , we will be able to show easily that

$$\mathcal{E}(T) < \varepsilon_1, \quad \text{for a very small, computable constant } \varepsilon_1. \quad [12]$$

Moreover, the functions x and γ are known; and we also know upper bounds for Sobolev norms of f and g . Therefore, the ideas in the proof of Theorem 4.1, together with interval arithmetic, should lead to a rigorous proof of the differential inequality

$$\left| \frac{d}{dt} \mathcal{E}(t) \right| \leq C_1 \mathcal{E}(t) + \varepsilon_2, \quad [13]$$

where C_1 and ε_2 are computable constants.

We hope that ε_2 is very small, because f and g have small Sobolev norms; and we hope that C_1 won't be too big.

Once we establish [12] and [13], we will then know that our water-wave solution (z, ω) exists for all time $t \in [0, T]$, and that $\mathcal{E}(0) < \varepsilon_3$ for a (hopefully small) computable constant ε_3 .

From the definition of $\mathcal{E}(t)$, we will then easily deduce that $z(\cdot, 0) - x(\cdot, 0)$ has norm at most ε_4 in $H^4(\mathbb{R}/2\pi\mathbb{Z})$, for a computable constant ε_4 .

If ε_4 is small enough, this in turn implies that the interface $z(\cdot, 0)$ is an H^4 -smooth graph. Thus (z, ω) is an exact solution of the water-wave equation, whose interface is an H^4 -smooth graph at time 0, and a splash at time T .

We hope that a proof along these lines can be made to work.

ACKNOWLEDGMENTS. A.C., D.C., F.G., and J.G.S. were partially supported by the Grant MTM2008-03754 of the Ministerio de Ciencia e Innovación (MCINN) (Spain) and the Grant StG-203138CDSIF of the European Research Council (ERC). C.F. was partially supported by National Science Foundation (NSF) Grant DMS-0901040. F.G. was partially supported by NSF grant DMS-0901810.

*Because x and γ are piecewise polynomials, we should not assume that $|\partial_x x(\alpha, t)|$ is a function of t alone. Hence, we cannot take $(z, \omega) = (x, \gamma)$ at time T .

1. Rayleigh L (1878) On the instability of jets. *P Lond Math Soc* s1-10:4-13.
2. Taylor G (1950) The instability of liquid surfaces when accelerated in a direction perpendicular to their planes. I. *P Roy Soc Lond A Mat* 201:192-196.
3. Wu S (1997) Well-posedness in Sobolev spaces of the full water wave problem in 2-D. *Invent Math* 130:39-72.
4. Wu S (1999) Well-posedness in Sobolev spaces of the full water wave problem in 3-D. *J Am Math Soc* 12:445-495.
5. Craig W (1985) An existence theory for water waves and the Boussinesq and Korteweg-de Vries scaling limits. *Commun Part Diff Equations* 10:787-1003.
6. Nalimov VI (1974) *The Cauchy-Poisson problem* (Dinamika Sploshn, Sredy), Vyp. 18 Dinamika Zidkost. so Svobod. Granicami, pp 104-210.
7. Beale JT, Hou TY, Lowengrub JS (1993) Growth rates for the linearized motion of fluid interfaces away from equilibrium. *Communications on Pure and Applied Mathematics Comptes Rendus Mathematique* 46:1269-1301.
8. Yoshida H (1982) Gravity waves on the free surface of an incompressible perfect fluid of finite depth. *Publ Res I Math Sci* 18:49-96.
9. Christodoulou D, Lindblad H (2000) On the motion of the free surface of a liquid. *Comm Pure Appl Math* 53:1536-1602.
10. Lindblad H (2005) Well-posedness for the motion of an incompressible liquid with free surface boundary. *Ann of Math (2)* 162:109-194.
11. Coutand D, Shkoller S (2007) Well-posedness of the free-surface incompressible Euler equations with or without surface tension. *J Am Math Soc* 20:829-930.
12. Shatah J, Zeng C (2008) Geometry and a priori estimates for free boundary problems of the Euler equation. *Comm Pure Appl Math* 61:698-744.
13. Zhang P, Zhang Z (2008) On the free boundary problem of three-dimensional incompressible Euler equations. *Comm Pure Appl Math* 61:877-940.
14. Córdoba A, Córdoba D, Gancedo F (2010) Interface evolution: water waves in 2-D. *Adv Math* 223:120-173.
15. Lannes D (2005) Well-posedness of the water-waves equations. *J Am Math Soc* 18:605-654.
16. Lannes D (2010) A stability criterion for two-fluid interfaces and applications., Arxiv preprint arXiv:1005.4565.
17. Alazard T, Métivier G (2009) Paralinearization of the Dirichlet to Neumann operator, and regularity of three-dimensional water waves. *Commun Part Diff Equations* 34:1632-1704.
18. Ambrose DM, Masmoudi N (2009) The zero surface tension limit of three-dimensional water waves. *Indiana U Math J* 58:479-521.
19. Alvarez-Samaniego B, Lannes D (2008) Large time existence for 3D water-waves and asymptotics. *Invent Math* 171:485-541.
20. Wu S (2009) Almost global wellposedness of the 2-D full water wave problem. *Invent Math* 177:45-135.
21. Germain P, Masmoudi N, Shatah J (2009) Global solutions for the gravity water waves equation in dimension 3. *C R Math Acad Sci Paris* 347:897-902.
22. Wu S (2011) Global wellposedness of the 3-D full water wave problem. *Invent Math* 184:125-220.
23. Castro A, Córdoba D, Fefferman C, Gancedo F, López-Fernández M (2011) Rayleigh-Taylor breakdown for the Muskat problem with applications to water waves. *Ann of Math*, Arxiv preprint arXiv:1102.1902..
24. Castro A, Córdoba D, Fefferman C, Gancedo F, López-Fernández M (2011) Turning waves and breakdown for incompressible flows. *Proc Nat'l Acad Sci USA* 108:4754-4759.
25. Beale JT, Hou TY, Lowengrub J (1996) Convergence of a boundary integral method for water waves. *SIAM J Numer Anal* 33:1797-1843.
26. Nirenberg L (1972) An abstract form of the nonlinear Cauchy-Kowalewski theorem. *J Differ Geom* 6:561-576.
27. Nishida T (1977) A note on a theorem of Nirenberg. *J Differ Geom* 12:629-633.
28. Moore R, Bierbaum F (1979) *Methods and applications of interval analysis*, (Society for Industrial & Applied Mathematics (SIAM), PA), ISBN: 0-89871-161-4, volume 2, pp xi-190.