## LOW-PRESSURE GRANULITE-FACIES METAMORPHISM IN THE PRYDZ BAY REGION, EAST ANTARCTICA (ABSTRACT)

Yoichi Motoyoshi<sup>1</sup>, D. E. Thost<sup>2</sup> and B. J. Hensen<sup>2</sup>

<sup>1</sup>National Institute of Polar Research, 9–10, Kaga 1-chome, Itabashi-ku, Tokyo 173

<sup>2</sup>Department of Applied Geology, University of New South Wales,

Kensington, New South Wales 2033, Australia

The terrain along the coastal line of Prydz Bay is underlain by granulite-facies rocks of various kinds including pelitic to psammitic gneisses, basic to intermediate gneisses and subordinate amounts of calc-silicate rocks and ultramafic rocks. Extensive partial melting has been observed in pelitic to psammitic gneisses. The age of metamorphism has been dated to be Late Proterozoic based on a Rb-Sr whole rock isochron (SHERATON and COLLERSON; BMR J. Aust. Geol. Geophys., 8, 119, 1983).

The rocks have been subjected to low-pressure granulite-facies metamorphism as evidenced by the following petrographical observations.

- 1) Pelitic to psammitic gneisses: Cordierite occurs regionally in pelitic gneisses. Andalusite and sillimanite coexist in a cordierite gneiss, in which andalusite occurs as a retrograde mineral being associated with chlorite.
- 2) Basic to intermediate gneisses: Orthopyroxene occurs regionally in basic to intermediate gneisses, being occasionally associated with clinopyroxene, amphibole and biotite. Garnet is rare, but where it occurs, coronas composed of plagioclase+clinopyroxene+magnetite symplectite are observed around garnet.
- 3) Ultramafic rocks: Olivene+plagioclase intergrowth is observed in ultramafic rocks. Mineral equilibria and geothermo-barometries yield initial P-T conditions around 750°C at  $5\pm1$  kbar.

Prior to the low-pressure granulite-facies metamorphism (possibly in Late Proterozoic), the rocks had been recrystallized at much higher P-T conditions around 950–1050°C at  $\simeq$  10 kbar deduced from local occurrence of pyrope-rich garnet (up to  $X_{Mg}$  70–72) in basic granulites.

(Received February 28, 1990; Revised manuscript received April 4, 1990)