

EFFECTS OF TEMPERATURE ON PHOTOSYNTHETIC RATES IN
DIATOMS ISOLATED FROM THE SOUTHERN OCEAN
(ABSTRACT)

Shigenobu TAKEDA¹, Isamu SAKAGUCHI¹, Kyoji SHINSHIMA¹, Michiyasu KIYONO¹,
Kentaro WATANABE² and Hideki NUMANAMI³

¹Biology Department, Central Research Institute of Electric Power Industry,
1646, Abiko, Abiko 270-11

²National Institute of Polar Research, 9-10, Kaga 1-chome, Itabashi-ku, Tokyo 173

³Tokyo University of Fisheries, 5-7, Kounan 4-chome, Minato-ku, Tokyo 108

The rates of photosynthesis of Antarctic diatoms were determined as a function of temperature. Clonal cultures of diatoms were isolated from open surface waters (0 m, -0.7~2.2°C) in the Indian sector of the Southern Ocean (59~68°S) during the austral summer. Stock cultures were grown in f/2 medium at 1~4°C under 16:8 hr light:dark cycle at an intensity of 100~200 $\mu\text{Einst m}^{-2} \text{s}^{-1}$ for 7~11 months. For photosynthetic rate measurements in the land laboratory, the Antarctic diatoms were incubated in 100 ml polycarbonate bottles (three light and two dark) for 4 hr under a saturating intensity of 200 $\mu\text{Einst m}^{-2} \text{s}^{-1}$ at different temperatures of -2.5°C, 0°C, 3°C, 5°C, 7°C, 10°C, and 15°C. Photosynthetic rates, based on the stable ¹³C isotope method, increased by a factor of 2.3~5.9 with temperature from -2.5°C (0.16~0.64 $\mu\text{gC } \mu\text{gChl } a^{-1} \text{ h}^{-1}$) to 7°C (0.58~1.89 $\mu\text{gC } \mu\text{gChl } a^{-1} \text{ h}^{-1}$). At higher temperatures the photosynthetic rates of *Chaetoceros* sp. and *Nitzschia* sp.1 were decreased rapidly (0.20 and 1.04 $\mu\text{gC } \mu\text{gChl } a^{-1} \text{ h}^{-1}$). On the other hand, photosynthetic rates of *Nitzschia* sp.2, *Nitzschia* sp.3, and *Nitzschia* sp.4 were either not decreased or slightly increased up to 15°C (0.76, 1.37, and 2.40 $\mu\text{gC } \mu\text{gChl } a^{-1} \text{ h}^{-1}$). The temperatures of maximum photosynthetic rates in these species were clearly higher than those in situ (-0.7~2.2°C) and in stock culture (1~4°C). Nevertheless, these five species all stopped growing at 15°C within 24 hr, and *Nitzschia* sp.1 showed little growth even at 7°C. It thus seems that high photosynthetic rates observed above the natural ambient temperature (-1.8~5°C) may remain for the restricted period that can be endured by the cells.

(Received March 31, 1993; Revised manuscript received June 28, 1993)