

## 東南極、昭和基地で記録された脈動と微気圧変動 からみた大気-海洋-固体地球の相互作用

金尾政紀<sup>1</sup>、石原吉明<sup>2</sup>、山本真行<sup>3</sup>、豊国源知<sup>1</sup>、Alessia Maggi<sup>4</sup>

<sup>1</sup> 国立極地研究所

<sup>2</sup> 国立天文台

<sup>3</sup> 高知工科大学

<sup>4</sup> ストラスブルク地球物理研究所

## Atmosphere - Ocean - Solid Earth Interaction from Microseisms & Microbaroms recorded at Syowa Station, East Antarctica

Kanao, M.<sup>1</sup>, Ishihara, Y.<sup>2</sup>, Yamamoto, M.-Y.<sup>3</sup>, Toyokuni, G.<sup>1</sup>, and Maggi, A.<sup>4</sup>

*1 National Institute of Polar Research*

*2 National Astronomical Observatory*

*3 Kochi University of Technology*

*4 Institut de Physique du Globe de Strasbourg*

Microseisms and microbaroms originated from the Southern Ocean are clearly recorded by both the broadband seismograph and infrasound sensor deployed at Syowa Station (39E, 69S), East Antarctica. A continuous images are achieved for the double-frequency microseism / microbaroms (DFM) with peaks between 4 and 10 s during a whole season. The peak amplitudes of DFM reflect the large influence of winter extratropical cyclonic storms (brizzard) in the Southern Ocean. The DFM have relatively lower amplitudes during austral winters, caused by the larger amount of sea ice extent around the Lützow-Holm Bay with decreasing the ocean wave loading effects. On the contrary, single-frequency microseism (SFM, with periods between 12 and 30 s) can be observable only by seismograph under excellent storm conditions particularly in local winter. On the infrasound data, moreover, long stand signals are identified with harmonic over tones at a few Hz to lower most human audible band. It probably related to the ice vibrations in the vicinity of the Station. Microseism measurements are a useful proxy for characterizing ocean wave climate and global storm intensity, complementing other estimates by ocean buoys or satellite measurements. A continuous monitoring both by broadband seismograph and infrasound observations firmly contribute to the Federation of Digital Seismographic Network (FDSN) and the Comprehensive Nuclear-Test-Ban Treaty (CTBTO) in southern high latitude, together with the Pan-Antarctic Observations System (PANTOS) under the Scientific Committee on Antarctic Research (SCAR).