

東南極セール・ロンダーネ山地に分布する原生代後期の未成熟弧に形成した 変トータル岩とアダカイト質花崗岩

亀井淳志¹, 堀江憲路², 大和田正明³, 柚原雅樹⁴, 中野伸彦⁵, 小山内康人⁵,
足立達朗⁵, 原 有希¹, 寺尾まどか¹, 手打晋二郎¹, 志村俊昭⁶, 東田和弘⁷,
外田智千², 岩田智加¹, 白石和行², 石塚英男⁸, 高橋裕平⁹

¹ 島根大学

² 国立極地研究所

³ 山口大学

⁴ 福岡大学

⁵ 九州大学

⁶ 新潟大学

⁷ 名古屋大学博物館

⁸ 高知大学

⁹ 産業技術総合研究所

Late Proterozoic juvenile arc metatonalite and adakitic granitoids in the Sør Rondane Mountains, East Antarctica

Atsushi Kamei¹, Kenji Horie², Masaaki Owada³, Masaki Yuhara⁴, Nobuhiko Nakano⁵, Yasuhito Osana⁵,
Tatsuro Adachi⁵, Yuki Hara¹, Madoka Terao¹, Shinjiro Teuchi¹, Toshiaki Shimura⁶, Kazuhiro Tsukada⁷,
Tomokazu Hokada², Chika Iwata¹, Kazuyuki Shiraishi², Hideo Ishizuka⁸, Yuhei Takahashi⁹

¹ Shimane University

² National Institute of Polar Research

³ Yamaguchi University

⁴ Fukuoka University

⁵ Kyushu University

⁶ Niigata University

⁷ Nagoya University Museum

⁸ Kochi University

⁹ Geological Survey of Japan, AIST

Late Middle Proterozoic to early Late Proterozoic granitoids are scattered throughout the Dronning Maud Land, which lies between 20° W and 45° E in East Antarctica. Recently, the geochronological similarities between the granitoids in western and central Dronning Maud Land have been clarified, with age data being concentrated in the range 1200–1050 Ma (e.g., Grantham et al., 1995, 2011; Groenewald et al., 1995; Jacobs et al., 1996; Board et al., 2005). Moreover, on the basis of the data from Mikhalsky and Jacobs (2004) and Grantham et al. (2011), the rocks can be categorized primarily as high- to medium-K series calc-alkaline granitoids. However, these characteristics of granitoids may not continue into the eastern part because a huge granitic pluton, which composed of metatonalite, in the Sør Rondane Mountains at the eastern Dronning Maud Land exhibits a younger solidification age of 960–920 Ma (Takahashi et al., 1990; Shiraishi et al., 2008) and geochemical features similar to those of low-K Archean tonalite–trondjemite–granodiorite (TTG) associations (Ikeda and Shiraishi, 1998). The existence of this granitic discontinuity has been postulated by a previous study (e.g., Jacobs et al., 2008); however, the cause of the discontinuity has not been clarified to date. In this study, we explore this discontinuity on the basis of a detailed examination of the metatonalite in the Sør Rondane Mountains.

We obtained significant data on lithological and geochemical investigations and some U–Pb SHRIMP zircon dating throughout the area of the metatonalite. Metatonalite is dominant over roughly 100 × 20 km² area in the southwestern end of the mountain range and is classified into five lithologies: gneissose Bt–Hbl metatonalite, weak gneissose Hbl–Bt metatonalite, Hbl metagabbro, Hbl–Bt tonalitic gneiss, and Bt metatonalite. The gneissose Bt–Hbl metatonalite is the main lithotype widely distributed over this area, which is geochemically categorized as low-K tholeiitic granitoid. Petrological studies suggest that the tholeiitic magma was derived from low-K basalt melting at the crustal depth, and the most plausible tectonic setting is a juvenile oceanic arc. The other four metaplutonic rocks are scattered as stocks or small intrusions in this area. They are geochemically regarded as calc-alkaline adakites related to oceanic slab melting. U–Pb SHRIMP zircon ages of the tholeiitic

metatonalite are concentrated at 998–995 Ma, whereas the calc-alkaline adakitic rocks are younger and divided between ages 945–920 Ma and 772 Ma. We believe that the tholeiitic metatonalite was formed first as a juvenile arc component between 998–995 Ma, followed by adakitic magmatism and oceanic slab melting at 945–920 Ma and 772 Ma. This Late Proterozoic tholeiitic granitoid is not distributed in the western to central Dronning Maud Land. Moreover, the exposure of an adakitic equivalent continues at the eastern side of the Sør Rondane Mountains. This suggests that the tectonic framework of the Sør Rondane Mountains, eastern Dronning Maud Land, is different from the western to central Dronning Maud Land. Our results would provide a constraint on the geological history of Dronning Maud Land during the late Middle Proterozoic to early Late Proterozoic.

References

- Board, W.S., Frimmel, H.E., and Armstrong, R.A., Pan-African tectonism in the western Maud-Belt: P–T–t path for high-grade gneisses in the H.U. Sverdrupfjella, East Antarctica. *Journal of Petrology* 46, 671–699, 2005.
- Grantham, G.H., Jackson, C., Moyes, A.B., Groenewald, P.B., Harris, P.D., Ferrar, G., and Krynauw, J.R., The tectonothermal evolution of the Kirwanveggen–H.U. Sverdrupfjella areas, Dronning Maud Land, Antarctica. *Precambrian Research* 75, 209–230, 1995.
- Grantham, G.H., Manhica, A.D.S.T., Armstrong, R.A., Kruger, F.J., and Loubser, M., New SHRIMP, Rb/Sr and Sm/Nd isotope and whole rock chemical data from central Mozambique and western Dronning Maud Land, Antarctica: Implications for the nature of the eastern margin of the Kalahari Craton and the amalgamation of Gondwana. *Journal of Asian Earth Science* 59, 74–100, 2011.
- Groenewald, P.B., Moyes, A.B., Grantham, G.H., and Krynauw, J.R., East Antarctic crustal evolution: geological constraints and modeling in western Dronning Maud Land. *Precambrian Research* 75, 231–251, 1995.
- Ikeda, Y., and Shiraishi, K., Petrogenesis of tonalitic rocks from the Sør Rondane Mountains, East Antarctica. *Polar Geoscience* 11, 143–153, 1998.
- Jacobs, J., Bauer, W., Spaeth, G., Thomas, R.J., and Weber, K., Lithology and structure of the Grenville-aged (~1.1Ga) basement of Heimefrontfjella (East Antarctica). *Geologische Rundschau* 85, 800–822, 1996.
- Jacobs, J., Pisarevsky, S., Thomas, R.J., and Becker, T., The Kalahari Craton during the assembly and dispersal of Rodinia. *Precambrian Research* 160, 142–158, 2008.
- Mikhalsky E.V., and Jacobs J., Orthogneisses in central Dronning Maud Land, East Antarctica: their origin and tectonic setting. International GeoMaud expedition of the BGR to central Dronning Maud Land in 1995/96 - Geological results. In: Paech H.-J. (Ed.), *Geologisches Jahrbuch*. 96 B, 49–76, 2004.
- Shiraishi, K., Dunkley, D.J., Hokada, T., Fanning, C.M., Kagami, H., and Hamamoto, T., Geochronological constraints on the Late Proterozoic to Cambrian crustal evolution of eastern Dronning Maud Land, East Antarctica: a synthesis of SHRIMP U–Pb age and Nd model age data. *Geodynamic Evolution of East Antarctica: A Key to the East–West Gondwana Connection*. In: Satish, K.M., Motoyoshi, Y., Osanai, Y., Hiroi, Y., Shiraishi, K. (Eds.), *Geological Society of London Special Publications* 308, 21–67, 2008.
- Takahasi, Y., Arakawa, Y., Sakiyama, T., Osanai, Y., and Makimoto, H., Rb–Sr and K–Ar whole rock ages of the plutonic bodies from the Sør Rondane Mountains, East Antarctica. *Proceeding of NIPR Symposium on Antarctic Geoscience* 4, 1–8, 1990.