

# アラスカの山岳氷河アイスコアから復元された降水量変動と太平洋十年規模変動との関係

的場澄人<sup>1</sup>、對馬あかね<sup>1,2</sup>、白岩孝行<sup>1</sup>

<sup>1</sup>北海道大学 低温科学研究所

<sup>2</sup>北海道大学大学院 環境科学院

## Relation between annual accumulation reconstructed from ice cores at Alaskan alpine glaciers and Pacific Climate Shift

Sumito Matoba<sup>1</sup>, Akane Tsushima<sup>1,2</sup>, and Takayuki Shiraiwa<sup>1</sup>

<sup>1</sup>Institute of Low Temperature Science, Hokkaido University

<sup>2</sup>Graduate School for Environmental Science, Hokkaido University

To reconstruct centurial climate and environmental changes at the northern North Pacific region, we analyze the ice cores obtained from alpine glaciers at Alaska and Kamchatka. In this contribution, we present data extracted from two ice cores obtained from summit glacier of Mount Wrangell in Wrangell-St Elias Mountains and from the ice field near Aurora Peak in the Alaska Range in Alaska. Profiles of  $\delta D$  of both ice cores shows remarkable seasonal variations, and we estimated yearly variations of annual accumulation for 50 years from the seasonal variations of  $\delta D$  in the ice cores. Annual accumulation at Aurora Peak showed significant increase from 1970s to present at the rate of 28mm/yr. This significant increase trend has not been observed at any meteorological stations at ground level. Annual accumulation at Mount Wrangell showed high values before 1977, low values from 1977 to 1990, and high values after 1990. The shift of accumulation in 1977 is likely associated with the Pacific climate shift called Regime shift or Pacific Decadal Oscillation. The annual accumulation in the Wrangell ice core after 1990 anti-correlated with surface level pressure over the Gulf of Alaska.




Figure 1. Location of drilling sites




Figure 2. Estimated annual accumulation of Aurora Peak and Mount Wrangell