## 南極域の風送バイオエアロゾル実相調査

小林 史尚  $^1$ 、牧 輝弥  $^1$ 、柿川 真紀子  $^2$ 、山田 丸  $^3$ 、松木 篤  $^4$ 、長沼 毅  $^6$ 、東 朋美  $^5$ 、岩坂 泰信  $^4$ 

<sup>1</sup> 金沢大学,理工研究域

<sup>2</sup> 金沢大学,環日本海域環境研究センター

<sup>3</sup> 金沢大学,イノベーション創成センター

<sup>4</sup> 金沢大学,フロンティアサイエンス機構

<sup>5</sup> 広島大学大学院,生物圏科学研究科

<sup>6</sup> 金沢大学,医薬保健研究域

## The research of atmosphere bioaerosol over Antarctica

Fumihisa Kobayashi<sup>1</sup>, Teruya Maki<sup>1</sup>, Makiko Kakikawa<sup>2</sup>, Maromu Yamada<sup>3</sup>, Atsushi Matsuki<sup>4</sup>, Takeshi Naganuma<sup>5</sup>,

Tomomi Higashi<sup>6</sup>, Yasunobu Iwasaka<sup>4</sup>

<sup>1</sup>College of Science and Engineering, Kanazawa University

<sup>2</sup>Institute of Nature and Environmental Technology, Kanazawa University

<sup>3</sup>Center for Innovation, Kanazawa University

<sup>4</sup>Frontier Science Organization, Kanazawa University

<sup>5</sup>Graduate School of Biosphere Science, Hiroshima University

<sup>6</sup>College of Medical, Pharmaceutical and Health Sciences, Kanazawa University

We carried out the direct sampling and bio-analysis of bioaerosols in yellow dust (KOSA) using bioaerosol sampler which was developed on our own terms, a tethered balloon, and an airplane over Japan and China<sup>1-3)</sup>. More than sixty strains were isolated as atmospheric bioaerosol over Dunhuang City in China and Suzu City in Japan using separated rich-culture method, metagenomic method, and separated culture method of halotolerant bacteria by a tethered balloon or an airplane. The study of atmospheric bioaerosol over Antarctica will be focused on because it is attracting attention to find the microorganism in Antarctic ice cores<sup>4)</sup>, investigate the long-range transport of atmospheric bioaerosol<sup>5)</sup>, and be starting the worldwide bioaerosol observations<sup>6)</sup>. To apply our previous methods in the study on KOSA bioaerosol, the direct sampling and bio-analysis of atmospheric bioaerosol over Antarctica will be tried. The results and finding of atmospheric bioaerosol over Antarctica will be establish the strong and important impacts to not only biogeography, ecology, history of the Earth, theory of evolution, influence of health, ice-forming nucleus, and global long-range transports of atmospheric bioaerosols, but also investigation on ecosystem in the sky.

## References

- 1) Kobayashi, F., M. Kakikawa, *et al.*, Study on atmospheric diffusion of bioaerosols in a KOSA source region, *Earozoru Kenkyu*, **22**, 218—227, 2007.
- 2) Maki, T., S. Susuki, *et al.*, Phylogenetic analysis of atmospheric halotolerant bacterial communities at high altitude in an Asian dust (KOSA) arrival region, Suzu City, *Science of the Total Environment*, **408**, 4556-4562, 2010.
- 3) Kakikawa, M., F. Kobayashi, *et al.*, Dustborne microorganisms in the atmosphere over an Asian dust source region, Dunhuang, *Air Quality, Atmosphere, and Health*, **1**, 195-202, 2008.
- 4) Segawa, T., K. Ushida, *et al.*, Bacterial communities in two Antarctic ice cores analyzed by 16S rRNA gene sequencing analysis, *Polar Science*, **4**, 215-227, 2010.
- 5) Iwasaka, Y., G.-Y. Shi, *et al.*, Mixture of Kosa (Asian dust) and bioaerosols detected in the atmosphere over the Kosa particles source regions with balloon-bore measurements: possibility of long-range transport, *Air Quality, Atmosphere, and Health*, **2**, 29-38, 2009.
- 6) Kellogg, C.A., D.W. Griffin, *et al.*, Characterization of aerosolized bacteria and fungi from desert dust events in Mali, West Africa, *Aerobiologia*, **20**, 99-110, 2004.