

## 隆起海浜地形・地質からみたグリーンランド氷床融解史の再評価

三浦英樹<sup>1</sup>、前杵英明<sup>2</sup>、奥野淳一<sup>1</sup>

<sup>1</sup> 国立極地研究所

<sup>2</sup> 広島大学

### Reappraisal for the melting history of the Greenland Ice sheet from geological analysis on the raised beach deposits

Hideki Miura<sup>1</sup>, Hideaki Maemoku<sup>2</sup>, Jun'ichi Okuno<sup>1</sup>

<sup>1</sup>National Institute of Polar Research

<sup>2</sup>Hiroshima University

The past melting history of the Greenland ice sheet is the great key for elucidating the future sea-level rising. The GIA (Glacial isostatic adjustment) model using the spatial and temporal variations of relative sea-level changes is one of useful techniques for the reconstruction of ice melting history. Although many radiocarbon dating ages for the reconstruction of the sea-level history have been obtained from the coastal area of Greenland (ex. Kelly, 1973; Ten Brink, 1974, 1975; Weidick, 1968, 1972 in West Greenland), the detailed geomorphological and Quaternary stratigraphical investigations have not been enough. For example, though Reeh (1989) and Henriksen (2008) compiled the map showing the amount of uplift during the Holocene along the coastal area of Greenland, some isolated high coastal existences of the place of the amount of upheaval suggest that the field confirmation of Holocene marine limits is questionable. Actually, Fleming and Lambeck (2004) have tried to make the GIA models of Greenland since the Last Glacial Maximum (LGM) to reproduce the sea level history with extraordinary upheaval point and obtained the result that changes in ice thickness since the LGM were > 500 m along the present-day outer coast and > 1500 m along some parts of the present-day ice margin. In this presentation, we introduce the report of the reappraisal for the melting history of the Greenland Ice sheet from detailed geological analysis on the raised beach deposits and GIA modelling.

#### References

- Fleming, K. and Lambeck, K., Constraints on the Greenland Ice Sheet since the Last Glacial Maximum from sea-level observations and glacial-rebound models. *Quaternary Science Reviews*, 23, 1053–1077, 2004.
- Henriksen, N., *Geological History of Greenland: Four billion years of Earth evolution*, Geological Survey of Denmark and Greenland (GEUS), 2008.
- Kelly, M. Radiocarbon dated shell samples from Nordre Strømfjord, West Greenland. Rapport, 59, *Grønlands Geologiske Undersøgelse*, Copenhagen, 1973.
- Reeh, N., Part 3. Quaternary Geology of Greenland. Fluton, R.J. (ed.) *Quaternary Geology of Canada and Greenland: 739–822*, Canada, 1989.
- Rinterknecht, V., Gorokhovich, Y., Schaefer, J. and Caffee, M., Preliminary 10Be chronology for the last deglaciation of the western margin of the Greenland Ice Sheet. *Journal of Quaternary Science*, 24, 270–278, 2009.
- Ten Brink, N.W., Glacio-isostasy: new data from West Greenland and geophysical implications. *Geological Society of America Bulletin*, 85, 219–228, 1974.
- Ten Brink, N.W., Holocene history of the Greenland ice sheet based on radiocarbon-dated moraines in West Greenland. *Meddelelser om Grønland*, 201, 1–44, 1975.
- Weidick, A., Observations on some Holocene glacier fluctuations in West Greenland. *Meddelelsser om Grønland*, 165, 1–203, 1968.
- Weidick, A., Holocene shore-lines and glacial stages in Greenland: an attempt at correlation. Rapport, 41, *Grønlands Geologiske Undersøgelse*, Copenhagen, 1972.
- Weidick, A., Oerter, H., Reeh, N., Thomsen, H.H. and Thoring, L., The recession of Inland Ice margin during the Holocene climatic optimum in the JaconsHAVN Isfjord area of West Greenland. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 82, 389–399, 1990.