

4-4-2022

## Protein kinase C (Pkc)- $\delta$ mediates arginine-induced glucagon secretion in pancreatic $\alpha$ -cells

Norikiyo Honzawa  
*Jikei University School of Medicine*

Kei Fujimoto  
*Gunma University*

Masaki Kobayashi  
*Jikei University School of Medicine*

Daisuke Kohno  
*Gunma University*

Osamu Kikuchi  
*Gunma University*

*See next page for additional authors*

Follow this and additional works at: [https://digitalcommons.wustl.edu/open\\_access\\_pubs](https://digitalcommons.wustl.edu/open_access_pubs)

---

### Recommended Citation

Honzawa, Norikiyo; Fujimoto, Kei; Kobayashi, Masaki; Kohno, Daisuke; Kikuchi, Osamu; Yokota-Hashimoto, Hiromi; Wada, Eri; Ikeuchi, Yuichi; Tabei, Yoko; Dorn, Gerald W II; Utsunomiya, Kazunori; Nishimura, Rimei; and Kitamura, Tadahiro, "Protein kinase C (Pkc)- $\delta$  mediates arginine-induced glucagon secretion in pancreatic  $\alpha$ -cells." *International journal of molecular science*. 23,7. . (2022).  
[https://digitalcommons.wustl.edu/open\\_access\\_pubs/11627](https://digitalcommons.wustl.edu/open_access_pubs/11627)

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact [vanam@wustl.edu](mailto:vanam@wustl.edu).

---

## Authors

Norikiyo Honzawa, Kei Fujimoto, Masaki Kobayashi, Daisuke Kohno, Osamu Kikuchi, Hiromi Yokota-Hashimoto, Eri Wada, Yuichi Ikeuchi, Yoko Tabei, Gerald W Dorn II, Kazunori Utsunomiya, Rimei Nishimura, and Tadahiro Kitamura



Article

# Protein Kinase C (Pkc)- $\delta$ Mediates Arginine-Induced Glucagon Secretion in Pancreatic $\alpha$ -Cells

Norikiyo Honzawa <sup>1,2</sup>, Kei Fujimoto <sup>3,\*</sup> , Masaki Kobayashi <sup>2</sup>, Daisuke Kohno <sup>2</sup>, Osamu Kikuchi <sup>2</sup> , Hiromi Yokota-Hashimoto <sup>2</sup>, Eri Wada <sup>2</sup>, Yuichi Ikeuchi <sup>2</sup>, Yoko Tabei <sup>2</sup>, Gerald W. Dorn II <sup>4</sup>, Kazunori Utsunomiya <sup>1</sup>, Rimei Nishimura <sup>1</sup> and Tadahiro Kitamura <sup>2,\*</sup>

- <sup>1</sup> Division of Diabetes, Metabolism and Endocrinology, Department of Internal Medicine, Jikei University School of Medicine, 3-25-8 Nishishinbashi, Minato-ku, Tokyo 105-8461, Japan; mamnoon.am.m.m.h.m@gmail.com (N.H.); kazu-utsunomiya@jikei.ac.jp (K.U.); rimei@jikei.ac.jp (R.N.)
- <sup>2</sup> Metabolic Signal Research Center, Institute for Molecular and Cellular Regulation, Gunma University, 3-39-15 Showa-machi, Maebashi 371-8512, Japan; masakikobayashi@gunma-u.ac.jp (M.K.); daisuke.kohno@gunma-u.ac.jp (D.K.); m10702010@gunma-u.ac.jp (O.K.); hiyokota@gunma-u.ac.jp (H.Y.-H.); wada@riem.nagoya-u.ac.jp (E.W.); m1920045@gunma-u.ac.jp (Y.I.); tabei-y@gunma-u.ac.jp (Y.T.)
- <sup>3</sup> Division of Diabetes, Metabolism and Endocrinology, Department of Internal Medicine, Jikei University Daisan Hospital, 4-11-1, Izumihoncho, Komae-shi, Tokyo 201-8601, Japan
- <sup>4</sup> Center for Pharmacogenomics, Division of Cardiology, Department of Internal Medicine, Washington University School of Medicine, St. Louis, MO 63110, USA; gdorn@dom.wustl.edu
- \* Correspondence: fuji@jikei.ac.jp (K.F.); kitamura@gunma-u.ac.jp (T.K.); Tel.: +81-3-3480-1151 (K.F.); +81-27-220-8849 (T.K.)



**Citation:** Honzawa, N.; Fujimoto, K.; Kobayashi, M.; Kohno, D.; Kikuchi, O.; Yokota-Hashimoto, H.; Wada, E.; Ikeuchi, Y.; Tabei, Y.; Dorn, G.W., II; et al. Protein Kinase C (Pkc)- $\delta$  Mediates Arginine-Induced Glucagon Secretion in Pancreatic  $\alpha$ -Cells. *Int. J. Mol. Sci.* **2022**, *23*, 4003. <https://doi.org/10.3390/ijms23074003>

Academic Editors: Katsuya Tanabe and Shunichiro Asahara

Received: 15 March 2022

Accepted: 1 April 2022

Published: 4 April 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

**Abstract:** The pathophysiology of type 2 diabetes involves insulin and glucagon. Protein kinase C (Pkc)- $\delta$ , a serine–threonine kinase, is ubiquitously expressed and involved in regulating cell death and proliferation. However, the role of Pkc $\delta$  in regulating glucagon secretion in pancreatic  $\alpha$ -cells remains unclear. Therefore, this study aimed to elucidate the physiological role of Pkc $\delta$  in glucagon secretion from pancreatic  $\alpha$ -cells. Glucagon secretions were investigated in Pkc $\delta$ -knockdown InR1G9 cells and pancreatic  $\alpha$ -cell-specific Pkc $\delta$ -knockout ( $\alpha$ Pkc $\delta$ KO) mice. Knockdown of Pkc $\delta$  in the glucagon-secreting cell line InR1G9 cells reduced glucagon secretion. The basic amino acid arginine enhances glucagon secretion via voltage-dependent calcium channels (VDCC). Furthermore, we showed that arginine increased Pkc $\delta$  phosphorylation at Thr<sup>505</sup>, which is critical for Pkc $\delta$  activation. Interestingly, the knockdown of Pkc $\delta$  in InR1G9 cells reduced arginine-induced glucagon secretion. Moreover, arginine-induced glucagon secretions were decreased in  $\alpha$ Pkc $\delta$ KO mice and islets from  $\alpha$ Pkc $\delta$ KO mice. Pkc $\delta$  is essential for arginine-induced glucagon secretion in pancreatic  $\alpha$ -cells. Therefore, this study may contribute to the elucidation of the molecular mechanism of amino acid-induced glucagon secretion and the development of novel antidiabetic drugs targeting Pkc $\delta$  and glucagon.

**Keywords:** pancreatic  $\alpha$ -cell; protein kinase C- $\delta$ ; glucagon; arginine

## 1. Introduction

The onset and progression of type 2 diabetes mellitus (T2DM) is due to relative insulin deficiency, but the contribution of impaired glucagon to T2DM is unclear. Surprisingly, it has been reported that the glucose tolerance of mice lacking pancreatic  $\alpha$ -cells or glucagon receptors does not worsen when pancreatic  $\beta$ -cells are destroyed with streptozotocin [1–3]. This indicates that the presence of glucagon is more critical for glucose intolerance than insulin deficiency [4]. Therefore, the importance of glucagon is now being reaffirmed.

Although still controversial, glucagon secretion from pancreatic  $\alpha$ -cells is regulated by various intracellular signals, including a rise in Ca<sup>2+</sup> influx in  $\alpha$ -cells [5,6]. However, it is unclear whether glucose per se enhances Ca<sup>2+</sup> influx in pancreatic  $\alpha$ -cells [7–10]. Alternatively, it was also reported that Ca<sup>2+</sup> influx is not necessarily involved, but cyclic adenosine monophosphate (cAMP) is involved in glucagon secretion from pancreatic  $\alpha$ -cells [11].

Additionally, reports show that insulin affects the cAMP-PKA pathway downstream of somatostatin receptor (SSTR) 2 and regulates glucagon secretion in pancreatic  $\alpha$ -cells [12,13]. Arginine is a basic non-essential amino acid and enhances glucagon secretion via voltage-dependent calcium channels (VDCC) [14]. It promotes the secretion of growth hormone and insulin. It is used clinically as a secretory provocation test for these hormones, but its detailed mechanism is unknown.


Protein kinase C (Pkc), a serine–threonine kinase, is ubiquitously expressed and involved in the regulation of cellular development, differentiation, death, and proliferation [15–17]. Pkc is divided into three sub-families, conventional, novel, and atypical, according to the difference in the activation manner. The conventional-type Pkc is activated by diacylglycerol (DAG) and  $\text{Ca}^{2+}$ , while the novel-type Pkc is activated by DAG. How atypical-type Pkc is activated has not been clarified. The primary substrate for Pkc is myristoylated alanine-rich C-kinase substrate (MARCKS), a cytoskeleton component, also known as an indicator of Pkc activation [18]. MARCKS translocates from the cell membrane to cytoplasm in response to Pkc activation and regulates exocytosis [18]. Pkc $\delta$ , a novel-type Pkc, comprises the regulatory domain consisting of conserved (C) 1 and C2-like regions and the catalytic domain consisting of C3 and C4 regions. DAG binding to the Pkc $\delta$  C1 region triggers Pkc $\delta$  activation and cleavage of Pkc $\delta$  by caspase-3, which produces the Pkc $\delta$ -catalytic fragment (CF). Phosphorylation at Thr<sup>505</sup> in the Pkc $\delta$  C4 region and Pkc $\delta$  translocation from the cytoplasm to the cell membrane contribute to its stabilization.

For diabetes, hyperglycemia enhances glycolysis followed by DAG production from glyceraldehyde 3-phosphate (de novo DAG production pathway) [19]. Additionally, diabetes is often comorbid with dyslipidemia, such as free fatty acid (FFA) elevation. FFA enters cells through fatty acid transporters and is metabolized to DAG via Acyl-CoA. Therefore, Pkc $\delta$  is activated due to increased DAG levels in diabetes. Importantly, systemic Pkc $\delta$ -knockout mice had improved retinopathy by suppressing retinal vascular proliferation, as well as improved nephropathy by suppressing glomerular podocyte loss, even under diabetic conditions [20,21]. Pancreatic  $\beta$ -cell-specific Pkc $\delta$  kinase-negative transgenic mice had improved glucose tolerance via increased insulin secretion [22]. Therefore, Pkc $\delta$  is a therapeutic target for diabetes and diabetic complications. Alternatively, Pkc $\delta$  is expressed in pancreatic  $\alpha$ -cells and regulates glucagon secretion [23]. Pan-Pkc activator enhanced, while Pkc $\delta$  inhibitors suppressed, glucagon secretion from isolated islets [24,25]. However, the physiological role of Pkc $\delta$  in pancreatic  $\alpha$ -cells is unclear. Therefore, we elucidated the physiological roles of Pkc $\delta$  in pancreatic  $\alpha$ -cells using InR1G9 cells and pancreatic  $\alpha$ -cell-specific Pkc $\delta$ -knockout ( $\alpha$ Pkc $\delta$ KO) mice.

## 2. Results

### 2.1. Knockdown of Pkc $\delta$ Decreased Glucagon Secretion in InR1G9 Cells


Pharmaceutical studies on isolated islets suggested that Pkc $\delta$  is involved in regulating glucagon secretion [25]. To confirm the physiological roles of Pkc $\delta$  in  $\alpha$ -cells, we investigated glucagon secretion in InR1G9 cells [26,27]. Findings revealed that Pkc $\delta$  inhibitor, rottlerin, significantly decreased glucagon secretion in InR1G9 cells (Figure 1a), which is consistent with previous a report [25]. However, because rottlerin is not a specific inhibitor for Pkc $\delta$  [28], the knockdown of Pkc $\delta$  in InR1G9 cells using short interfering ribonucleic acid (siRNA) was conducted. The knockdown efficiencies by *Prkcd* #1 and #2 siRNA at the protein level were 36.0% and 46.8%, respectively (Figure 1b,c). Additionally, the expression levels of Pkc $\delta$ -CF, which is a catalytic fragment of Pkc $\delta$  and indicates its activation, were significantly reduced by these *Prkcd* siRNAs (Figure 1b,d). Because of the better knockdown efficiency, we used *Prkcd* #2 siRNA in this experiment. The knockdown efficiency by *Prkcd* #2 siRNA at the mRNA level was 49.9% in InR1G9 cells (Figure 1e). Importantly, we found that glucagon secretion was significantly reduced by Pkc $\delta$  knockdown, indicating that Pkc $\delta$  is essential for glucagon secretion in InR1G9 cells (Figure 1f).



**Figure 1.** Knockdown of Pkc $\delta$  decreased glucagon secretion from InR1G9 cells. (a) Glucagon secretion was significantly reduced by 10- $\mu$ M Pkc $\delta$  inhibitor rottlerin in InR1G9 cells. (b–d) InR1G9 cells were transfected with scrambled siRNA, *Prkcd* #1 siRNA, or *Prkcd* #2 siRNA for 48 h. Cell lysates were examined for Pkc $\delta$  and Pkc $\delta$ -CF expression by Western blotting.  $\alpha$ -tubulin was employed as an internal control. Pkc $\delta$  expressions were reduced by around 40–50% *Prkcd* #1 siRNA and *Prkcd* #2 siRNA compared to the scrambled siRNA in InR1G9 cells. Pkc $\delta$ -CF expressions were also reduced by *Prkcd* #1 siRNA and *Prkcd* #2 siRNA in InR1G9 cells. (e) Pkc $\delta$  mRNA expression was significantly reduced by *Prkcd* #2 siRNA in InR1G9 cells. (f) Glucagon secretion was significantly reduced by *Prkcd* #2 siRNA in InR1G9 cells. We performed triplicate and three independent experiments. \*  $p < 0.05$ , \*\*  $p < 0.01$ ;  $t$ -test (a,e,f) or ANOVA (Bonferroni adjustment) (c,d). Data are expressed as means  $\pm$  SEM.

## 2.2. Arginine-Induced Glucagon Secretion Was Decreased by Pkc $\delta$ Knockdown in InR1G9 Cells

Arginine (via activating VDCC) and 3-Isobutyl 1-methylxanthine (IBMX) (via inhibiting phosphodiesterase and increasing cAMP) are known as glucagon secretagogues. As shown in Figure 2a, both arginine and IBMX significantly increased glucagon secretion in InR1G9 cells. Arginine significantly enhanced Pkc $\delta$  phosphorylation at Thr<sup>505</sup> in InR1G9 cells (Figure 2b,c), indicating that Pkc $\delta$  is activated by arginine. In these results, decreased PKC $\delta$  protein levels might be caused by enhanced PKC $\delta$  cleavage, which is also a marker for PKC $\delta$  activation. Furthermore, we investigated the localization of Pkc substrate MARCKS using green fluorescent protein (GFP)-labeled MARCKS (MARCKS-GFP). Under the basal condition, MARCKS-GFP localizing on the cell membrane was translocated to the cytoplasm by the Pkc activator, phorbol 12-myristate 13-acetate (PMA) (Figure 2d). Arginine also significantly induced cytoplasmic localization of MARCKS-GFP, which was canceled by the Pkc $\delta$  inhibitor, rottlerin (Figure 2d). These results suggested that Pkc $\delta$  is activated by arginine in InR1G9 cells. We investigated whether Pkc $\delta$  mediated arginine-induced glucagon secretion. Arginine increased glucagon secretion in InR1G9 cells, which was significantly suppressed by Pkc $\delta$  knockdown (Figure 2e). Thus, Pkc $\delta$  mediates arginine-induced glucagon secretion in InR1G9 cells.




**Figure 2.** Arginine-induced glucagon secretion was decreased by *Pkcδ* knockdown in InR1G9 cells. (a) Glucagon secretion was significantly increased by 15-mM arginine and 100-mM IBMX in InR1G9 cells. (b–c) InR1G9 cells were treated with or without 15-mM arginine for 30 min; then cell lysates were examined for *Pkcδ* and p*Pkcδ* (Thr505) by Western blotting.  $\beta$ -actin was employed as an internal control. Arginine significantly increased *Pkcδ* (Thr505)/*Pkcδ* relative ratio in InR1G9 cells. (d) InR1G9 cells were transfected with plasmid encoding MARCKS conjugated with GFP for 48 h followed by 100-mM PMA, 15-mM arginine, or 15-mM arginine plus 10- $\mu$ M rottlerin administration. MARCKS localized on the cell membrane was translocated to the cytoplasm by PMA. Arginine also translocated MARCKS to the cytoplasm, which was canceled by rottlerin. Representative photomicrographs are shown. Scale bars represent 20  $\mu$ m. (e) Arginine-induced glucagon secretion was significantly reduced by *Prkcd* #2 siRNA in InR1G9 cells. We performed triplicate and three independent experiments. \*  $p < 0.05$ ;  $t$ -test (c) or ANOVA (Bonferroni adjustment) (a,e). Data are expressed as means  $\pm$  SEM.

### 2.3. Establishment of $\alpha$ -Cell-Specific *Pkcδ*-Knockout Mice

To investigate the physiological roles of *Pkcδ* in vivo, we generated pancreatic  $\alpha$ -cell-specific *Pkcδ*-knockout mice using the Cre/LoxP system. We first crossed *Prkcd*<sup>flxed</sup> [29] with *Gcg*<sup>Cre</sup> mice [30]. However, because Cre recombinase expression levels in the  $\alpha$ -cells of *Gcg*<sup>Cre</sup> mice were approximately 50% (data not shown), as previously reported [31], we crossed *Prkcd*<sup>flxed</sup> with *Gcg*<sup>CreERT2</sup> mice, which reportedly had higher Cre expression efficiency in  $\alpha$ -cells [32]. Because commercially available *Pkcδ* antibodies were not good enough for immunohistochemistry, we employed a Cre reporter mouse (*Rosa26*<sup>tdTomato</sup> mouse) to test the Cre expression efficiency [33]. Native fluorescence of tdTomato allowed for direct visualization of Cre expression. Glucagon-positive and tdTomato-positive cells were mostly merged in *Gcg*<sup>CreERT2</sup>: *Rosa26*<sup>tdTomato</sup> mice (Figure 3a). Cre expression efficiency in  $\alpha$ -cells was 96.6% (1285 tdTomato-positive cells/1330 glucagon-positive cells) in these mice; therefore, we expect that most *Pkcδ* should be deleted in  $\alpha$ -cells of *Gcg*<sup>CreERT2</sup>: *Prkcd*<sup>flxed</sup> mice. Another advantage of *Gcg*<sup>CreERT2</sup> mice was that we excluded the influence


of *pkcδ* deletion during the embryonic stage by treating the mice with tamoxifen at the adult stage. Here, we used our homemade N-terminal glucagon antibody (rat monoclonal IgG antibody) for staining  $\alpha$ -cells. To check the specificity of this antibody for glucagon, we performed glucagon staining together with GFP in the pancreatic sections of *Gcg<sup>gfp/+</sup>* and *Gcg<sup>gfp/gfp</sup>* mice [34]. Because the region including exon 2 and 3 of the glucagon gene is replaced by the *gfp* sequence, glucagon is not expressed in *Gcg<sup>gfp/gfp</sup>* mice. Though glucagon was stained in GFP-positive  $\alpha$ -cells of *Gcg<sup>gfp/+</sup>* mice, glucagon was not detected in GFP-positive  $\alpha$ -cells of *Gcg<sup>gfp/gfp</sup>* mice (Supplementary Figure S1), indicating high specificity of the glucagon antibody used in this study. We also confirmed the establishment of  $\alpha$ Pkc $\delta$ KO mice at the DNA level. In this system, the recombinant *Prkcd* allele (predicted band) should be detected only in Pkc $\delta$ -deleted cells. As shown in Figure 3b, the predicted and floxed bands were observed in the genomic DNA extracted from the islets of *Gcg<sup>CreERT2</sup>; Prkcd<sup>floxed</sup>* mice but not in the genomic DNA extracted from the tails of these mice, showing the establishment of  $\alpha$ -cell-specific Pkc $\delta$ -knockout mice.



**Figure 3.** Establishment of  $\alpha$ -cell-specific Pkc $\delta$ -knockout mice. (a) Pancreatic sections of *Gcg<sup>CreERT2</sup>; Rosa26<sup>tdTomato</sup>* mice were subjected to immunohistochemistry with rat monoclonal anti-glucagon antibody. Native fluorescence of tdTomato is shown in red, glucagon is shown in green, and DAPI is shown in blue. Merge (yellow) indicates the colocalization of tdTomato with glucagon. Glucagon-positive and tdTomato-positive cells were mostly merged in *Gcg<sup>CreERT2</sup>; Rosa26<sup>tdTomato</sup>* mice.  $n = 4$ , each group. Scale bars represent 200  $\mu$ m (upper panels) and 50  $\mu$ m (lower panels). These studies were performed under blinded analysis. (b) Genomic DNA was extracted from the tails and the pancreatic islets of *Gcg<sup>CreERT2/CreERT2</sup>; Prkcd<sup>floxed/floxed</sup>*, *Gcg<sup>CreERT2/+</sup>; Prkcd<sup>floxed/floxed</sup>*, and *Gcg<sup>+/+</sup>; Prkcd<sup>floxed/floxed</sup>* mice. Because the region between loxP sites is excised in the tissues where Cre recombinase is expressed, the smaller PCR products (predicted bands) were detected in the islets but not in the tails of *Gcg<sup>CreERT2/CreERT2</sup>; Prkcd<sup>floxed/floxed</sup>*; and *Gcg<sup>CreERT2/+</sup>; Prkcd<sup>floxed/floxed</sup>* mice. This result indicated that Pkc $\delta$  was deleted in the islets of  $\alpha$ Pkc $\delta$ KO mice.

#### 2.4. Arginine-Induced Glucagon Secretion Decreased in $\alpha$ Pkc $\delta$ KO Mice and the Islets from $\alpha$ Pkc $\delta$ KO Mice

We first examined the body weight, glucose tolerance, insulin sensitivity, and area of pancreatic  $\alpha$ - and  $\beta$ -cells in  $\alpha$ Pkc $\delta$ KO mice. There were no significant differences in body weight (Supplementary Figure S2a), fasting and random-fed blood glucose levels (Figure 4a,b), intraperitoneal glucose tolerance test (IPGTT) (Supplementary Figure S2b), and insulin tolerance test (ITT) (Supplementary Figure S2c) in  $\alpha$ Pkc $\delta$ KO compared to the control mice. Moreover, the area of  $\alpha$ - and  $\beta$ -cells and the percentage of the  $\alpha$ -cell area against the  $\beta$ -cell area were comparable between the  $\alpha$ Pkc $\delta$ KO and control mice (Supplementary Figure S2d–g). Furthermore, fasting and random-fed glucagon levels were similar in  $\alpha$ Pkc $\delta$ KO and control mice (Figure 4c,d). However, glucagon secretion 15 min after 3 g/kg arginine intraperitoneal administration was significantly decreased in  $\alpha$ Pkc $\delta$ KO mice compared to the control mice (Figure 4e). Additionally, fasting, random-fed, and arginine-induced plasma insulin levels were comparable between  $\alpha$ Pkc $\delta$ KO and control mice (Supplementary Figure S3a–c). Furthermore, arginine-induced glucagon secretion was significantly lower in the islets isolated from  $\alpha$ Pkc $\delta$ KO mice than the islets isolated from the control mice (Figure 4f). Altogether, although neither fasting, random-fed glucagon levels, glucose tolerance, nor insulin tolerance were unchanged, arginine-induced glucagon secretion significantly decreased in  $\alpha$ Pkc $\delta$ KO mice and islets isolated from  $\alpha$ Pkc $\delta$ KO mice.



**Figure 4.** Arginine-induced glucagon secretion decreased in  $\alpha$ Pkc $\delta$ KO mice and the islets from  $\alpha$ Pkc $\delta$ KO mice. (a–d) Unaltered fasting (a) and random-fed (b) blood glucose levels and fasting (c) and random-fed (d) plasma glucagon levels in  $\alpha$ Pkc $\delta$ KO mice. (e) Plasma glucagon levels 15 min after 3 g/kg arginine intraperitoneal administration was significantly lower in  $\alpha$ Pkc $\delta$ KO than control mice.  $n = 10$ , each group. \*  $p < 0.05$ ;  $t$ -test. Data are expressed as a box plot.  $\alpha$ Pkc $\delta$ KO indicates  $Gcg^{CreERT2/+}; Prkcd^{floxed/floxed}$  mice and Cont indicates  $Gcg^{+/+}; Prkcd^{floxed/floxed}$  mice. (f) 15-mM arginine-induced glucagon secretion was significantly lower in islets isolated from  $\alpha$ Pkc $\delta$ KO mice than those from control mice. We used the islets isolated from six mice for each group ( $n = 6$ ). We performed triplicate experiments. \*  $p < 0.05$ ; ANOVA (Bonferroni adjustment). Data are expressed as means  $\pm$  SEM.  $\alpha$ Pkc $\delta$ KO indicates  $Gcg^{CreERT2/+}; Prkcd^{floxed/floxed}$  mice and Cont indicates  $Gcg^{+/+}; Prkcd^{floxed/floxed}$  mice.



### 3. Discussion

Studies have suggested that Pkc $\delta$  regulates glucagon secretion; however, these studies were mainly performed using nonspecific Pkc activators and inhibitors in isolated islets. Therefore, the specific and physiological roles of Pkc $\delta$  on glucose metabolism in vivo remain unclear. This study showed that arginine activated Pkc $\delta$  and enhanced glucagon secretion in InR1G9 cells. It also showed that arginine-induced glucagon secretion was decreased by Pkc $\delta$  knockdown in InR1G9 cells and islets isolated from  $\alpha$ Pkc $\delta$ KO mice. Although arginine-induced glucagon secretion was also reduced in  $\alpha$ Pkc $\delta$ KO mice, glucose and insulin tolerance remained unchanged, which is probably due to compensation mechanisms for controlling glucose metabolism. To our knowledge, this is the first report showing the physiological roles of Pkc $\delta$  in pancreatic  $\alpha$ -cells on glucagon secretion and glucose metabolism in vivo.

Arginine enhances glucagon secretion in pancreatic  $\alpha$ -cells, but the detailed mechanism is unclear. Arginine-induced glucagon secretion is directly inhibited by gliclazide, which suppresses adenosine triphosphate-sensitive potassium ( $K_{ATP}$ ) channels in  $\alpha$ -cells [14]. Here, deletion of Pkc $\delta$  suppressed arginine-induced glucagon secretion in InR1G9 cells,  $\alpha$ Pkc $\delta$ KO mice, and islets from  $\alpha$ Pkc $\delta$ KO mice (Figures 2e and 4e,f). Pkc $\delta$  was also reported to enhance  $Ca^{2+}$ -dependent exocytosis and is related to soluble *n*-ethylmaleimide sensitive factor attachment protein receptor (SNARE) proteins' phosphorylation [23,35]. Therefore, Pkc $\delta$  might be involved in the  $K_{ATP}$  channel–VDCC pathway and exocytosis in  $\alpha$ -cells. In this study, although Pkc $\delta$  knockdown reduced glucagon secretion in InR1G9 cells (Figure 1f), knockout of Pkc $\delta$  did not affect glucagon secretion in  $\alpha$ Pkc $\delta$ KO mice and islets from  $\alpha$ Pkc $\delta$ KO mice under basal conditions (Figure 4c,d,f). This discrepancy could be caused by the paracrine effects of other islet hormones, such as insulin and somatostatin, though fasting and random-fed plasma insulin levels were unchanged in  $\alpha$ Pkc $\delta$ KO (Supplementary Figure S3a,b). Indeed, insulin and somatostatin regulate glucagon secretion in  $\alpha$ -cells by lowering cAMP [12,13].  $\alpha$ -cell-specific insulin receptor KO mice showed increased glucagon secretion under fasting and fed conditions [36]. Altogether, Pkc $\delta$  might be involved in the downstream of arginine signaling in  $\alpha$ -cells, and more importantly, in arginine-induced glucagon secretion.

Because hyperglycemia increases DAG production and activates Pkc $\delta$  in various cell types, Pkc $\delta$  could be pathologically involved in diabetes progression [19]. Indeed, Pkc $\delta$  inhibition reportedly protected against the progression of diabetic nephropathy and diabetic retinopathy [20,21]. We previously reported that T2DM patients have higher plasma glucagon levels than healthy subjects, and proposed that glucagon could become a diagnostic marker and a new therapeutic target for T2DM [37,38]. This study revealed that Pkc $\delta$  is essential for arginine-induced glucagon secretion (Figures 2e and 4e,f), suggesting that Pkc $\delta$  might be related to postprandial hypersecretion of glucagon in T2DM. Alternatively, Pkc $\delta$  is involved in cell proliferation and apoptosis. Furthermore, increased  $\beta$ -cell number and insulin content were observed in  $\beta$ -cell-specific Pkc $\delta$  kinase-negative transgenic mice [22]. However, in this study, the area of  $\alpha$ -cells in  $\alpha$ Pkc $\delta$ KO mice was unaltered (Supplementary Figure S2d,e), suggesting that Pkc $\delta$  deletion in  $\alpha$ -cells does not affect  $\alpha$ -cell proliferation and apoptosis. Thus, Pkc $\delta$  inhibition in  $\alpha$ -cells could become a novel therapeutic target for diabetes without affecting  $\alpha$ -cell proliferation and cell death.

This study has several limitations. First is the usage of tumorigenic cell line InR1G9. Because tumorigenic cells are oncogenic, their character may differ from native  $\alpha$ -cells. Additionally, although InR1G9 cells secrete glucagon, they are derived from insulinoma [26]. Because the technique for isolating  $\alpha$ -cells from islets has only been recently developed [39–41], we will need to reconfirm the results in InR1G9 cells using isolated  $\alpha$ -cells in the future. The other limitation is the character of  $Gcg^{CreERT2}$  mice. Because  $Gcg^{CreERT2}$  mice possess the CreERT2-coding gene in exon 2 of the *preproglucagon* gene, they heterozygously lack the *preproglucagon* gene. Although it was reported that no significant difference was observed in glucagon secretion between  $Gcg^{CreERT2/+}$  and wild-type mice, the effect of heterozygous *preproglucagon* gene deletion cannot be excluded entirely [32].

In conclusion, we elucidated that Pkc $\delta$  is essential for arginine-induced glucagon secretion in pancreatic  $\alpha$ -cells using InR1G9 cells,  $\alpha$ Pkc $\delta$ KO mice, and islets from  $\alpha$ Pkc $\delta$ KO mice. Thus, this study may contribute to the elucidation of the mechanism of amino acid-induced glucagon secretion and the development of novel antidiabetic drugs targeting Pkc $\delta$  and glucagon.

## 4. Materials and Methods

### 4.1. Cell Culture

Glucagon-secreting InR1G9 cells were cultured in RPMI1640 (#189-02025, Wako, Osaka, Japan) supplemented with 10% FBS (#10270-106, Life Technologies, Carlsbad, CA, USA) and 1% penicillin/streptomycin (#268-23191, Wako, Osaka, Japan). All cells were grown at 37 °C in humidified air containing 5% (*v/v*) CO<sub>2</sub>.

### 4.2. Glucagon Secretion Assay

InR1G9 cells seeded on six-well dishes or six isolated islets seeded on 24-well dishes were pre-incubated with KRB (0.13-M NaCl, 4.7-mM KCL, 1.2-mM KH<sub>2</sub>PO<sub>4</sub>, 1.2-mM MgSO<sub>4</sub>·7H<sub>2</sub>O, 10-mM HEPES, 2- $\mu$ M CaCl<sub>2</sub>, pH 7.3) for 0.5 h. InR1G cells were incubated for 0.5 h with 15-mM arginine (#015-04613, Wako, Osaka, Japan), 100-mM IBMX (#095-03413, Wako, Osaka, Japan), or 10- $\mu$ M rottlerin (#12006, CAY, Funakoshi, Japan). Isolated islets were incubated for 2 h with 15 mM arginine. Supernatants were subjected to glucagon assay using glucagon sandwich ELISA (#10-1271-01, Mercodia, Sweden). Experiments were repeated at least three times. Glucagon concentrations were normalized by total protein content using the Pierce BCA assay kit (#23227, Life Technologies, Carlsbad, CA, USA).

### 4.3. Protein Extraction and Western Blotting

Whole-cell lysates were extracted using a lysis buffer (50-mM Tris (pH 8.0), 150-mM NaCl, 10% Glycerol, 1% Triton-X100, 0.1% SDS, 5-mM EDTA, 1-mM DTT, 10- mM NaF, 1-mM Na<sub>3</sub>VO<sub>4</sub>). Equal amounts of protein samples were loaded onto SDS-PAGE gels, electrophoresed, and transferred onto nitrocellulose membranes. After blocking with non-fat milk, membranes were incubated with primary antibodies followed by the corresponding secondary horseradish peroxidase-conjugated antibodies. The signal intensity was measured using a LAS-4010 mini-Luminescent Image Analyzer (FUJIFILM, Tokyo, Japan). Pkc $\delta$  (#ab182126, abcam, Japan), phospho-Pkc $\delta$  (Thr505) (#9374, Cell Signaling Technology Japan, Tokyo, Japan),  $\beta$ -actin (#sc-47778, Santa Cruz Biotechnology, Dallas, TX, USA), and  $\alpha$ -tubulin antibodies (#sc-5286, Santa Cruz Biotechnology, Dallas, TX, USA) were used.

### 4.4. RNA Extraction and Quantitative PCR

Total RNA was generated from InR1G9 cells using the RNAiso Plus kit (#9108, Takara Bio, Shiga, Japan). Additionally, RNA was reverse-transcribed into cDNA using the ImProm II Reverse Transcription System (#A3800, Promega, Madison, WI). To evaluate the mRNA expressions of Pkc $\delta$  and  $\beta$ -actin, quantitative real-time PCR analysis using Applied Biosystems ViiA<sup>7</sup> Real-Time PCR System (Life Technologies, Carlsbad, CA, USA) was conducted with Power SYBR Green Master Mix (#A25742, Life Technologies, Carlsbad, CA, USA). All reactions were normalized using  $\beta$ -actin. Primers for Pkc $\delta$  were purchased from Takara bio (#MA167545), and those for  $\beta$ -actin were as follows: 5'-AGCCTTCCTTCTTGGGTA-3' (forward) and 5'-GAGCAATGATCTTGATCTTC-3' (reverse).

### 4.5. siRNA Transfection

InR1G9 cells were transfected with scramble control (#4390843, Ambion, Life Technologies, Carlsbad, CA, USA), *Prkcd* Silencer Select siRNA #1 (#4390771, s71696, Ambion, Life Technologies, Carlsbad, CA, USA) or siRNA #2 (#4390771, s71698, Ambion, Life Technologies, Carlsbad, CA, USA) for 48 h. Additionally, transfections were performed with Lipofectamine RNAiMAX (#13778, Life Technologies, Carlsbad, CA, USA). The knock-

down efficiency was evaluated 48 h after transfection, followed by Western blotting or quantitative RT-PCR analyses.

#### 4.6. MARCKS Transfection and Immunocytochemistry

InR1G9 cells seeded on four-well chamber slides (# 154526, Life Technologies, Carlsbad, CA, USA) were transfected with a plasmid encoding MARCKS-GFP for 48 h using Lipofectamine 2000 transfection (#11668027, Life Technologies, Carlsbad, CA, USA), as previously described [18]. Then, InR1G9 cells were treated with 100-nM PMA (#AG-CN2-0010, Adipogen Life Sciences, San Diego, CA, USA), 15-mM arginine, or 15-mM arginine plus 10- $\mu$ M rottlerin for 0.5 h. Afterwards, the cells were fixed in 4% paraformaldehyde (PFA) and the localization of MARCK-GFP was observed by a confocal laser-scanning microscope (FV1000, Olympus Life Science, Tokyo, Japan).

#### 4.7. Animals and Physiological Experiments

All experimental procedures were performed in accordance with the Guide for the Care and Use of Laboratory Animals of the Science Council of Japan and approved by the Animal Experiment Committee of Gunma University.  $Gcg^{CreERT2}$  and  $Pkcd^{floxed}$  mice were mated to generate  $\alpha Pkc\delta KO$  mice. To induce Cre-mediated recombination,  $\alpha Pkc\delta KO$  mice were subcutaneously injected with 6 mg tamoxifen (2 mg, every other day) at four-weeks old and were characterized at 24-weeks old. Animal studies were performed following a normal chow diet (CE-2, CLEA Japan, Tokyo, Japan) using male mice. In this study,  $Gcg^{CreERT2/+}$ ;  $Prkcd^{floxed/floxed}$  and  $Gcg^{+/+}$ ;  $Prkcd^{floxed/floxed}$  mice served as  $\alpha Pkc\delta KO$  mice and controls, respectively. Arginine (pH 7.4, 3 g/kg weight) was injected intraperitoneally, and 15 min later, plasma glucagon levels were measured in  $\alpha Pkc\delta KO$  and control mice. Insulin (0.75 U/kg weight, Eli Lilly Japan KK, Hyogo, Japan) and D(+)-glucose (#049-31165, Wako, Osaka, Japan) were injected intraperitoneally. Blood glucose was measured using Glutest ai (#GT-1840, SANWA KAGAKU KENKYUSHO, Aichi, Japan). Plasma glucagon levels were measured by glucagon sandwich ELISA (#10-1281-01, Mercodia, Sweden). Arginine or glucose injection was performed after overnight fasting.

#### 4.8. Frozen Section Preparation and Immunohistochemistry

For immunohistochemistry,  $\alpha Pkc\delta KO$  mice (24-week-old males) were anesthetized and perfused transcardially with ice-cold 0.05-M phosphate-buffered saline (pH 7.4) followed by 4% PFA for fixation. The pancreas was dissected and fixed in 4% PFA. Samples were frozen in OCT compound and sectioned in 5  $\mu$ m-thick slices. Samples were permeabilized and blocked with 0.1% Triton-X-100/PBS/5% normal donkey serum (NDS) for 1 h. Rat monoclonal anti-glucagon (1:200, #52A1A, Immuno-Biological Laboratories, Gunma, Japan) and guinea pig polyclonal anti-insulin antibodies (1:500, #A0564, Dako, USA) were diluted in 0.1% Triton-X-100/PBS/5% NDS, then added to the sections and incubated overnight at 4 °C. After three washes with PBS, secondary antibodies (1:200, Jackson ImmunoResearch, Philadelphia, PA, USA) diluted in 0.1% Triton-X-100/PBS/5% NDS were added to the sections and incubated for 1 h at room temperature. Images were acquired using a confocal laser-scanning microscope (FV1000, Olympus Life Science, Japan).

#### 4.9. PCR Detecting the Recombined Allele

Genomic DNA was extracted from the tails and the pancreatic islets of  $Gcg^{CreERT2/CreERT2}$ ;  $Prkcd^{floxed/floxed}$ ,  $Gcg^{CreERT2/+}$ ;  $Prkcd^{floxed/floxed}$ , and  $Gcg^{+/+}$ ;  $Prkcd^{floxed/floxed}$  mice using REDxtract-N-Amp kit (#254-457-8, Sigma-Aldrich, St. Louis, MO, USA). PCR was performed using these DNAs as templates and the following primers: GT-1: 5'-ACCAGCGAT TTGAGA AGAAGC-3', GT-5: 5'-TCATCTGTACCTTCCACACCA-3', GT-6: 5'-AGAACCTC CATCACGAAGAAC-3'. The recombined PCR products were loaded onto 2% agarose gel, electrophoresed, and detected using PrintgraphClassic (#WSE-5400, ATTO CORPORATION, Tokyo, Japan).

#### 4.10. Pancreatic Islet Isolation

Mice were anesthetized with isoflurane and laparotomy was performed. First, the duodenum and small intestine were moved to the left side of the mouse, after which the common bile duct and papilla of Vater were checked and the proximal part (liver side) of the common bile duct was clamped. Then, a partial incision was made in the duodenum and a 27 G winged needle with a rounded tip was inserted into the common bile duct (CBD) from the duodenum. Collagenase (#C7657, Sigma-Aldrich, Japan) dissolved in medium 199 (#M0393, Sigma-Aldrich, Japan) was injected into the CBD. After confirming the swelling of the pancreas, the pancreas was excised from the rectum, stomach, duodenum, and small intestine. After that, the pancreas was added into a 50 mL tube and incubated in a water bath at 37 °C for 17 min. Then, medium 199 containing 10% FBS was added to a 50 mL tube to stop the collagenase reaction, and the mixture was shaken for 10 s. After centrifuging at 1000 rpm and 4 °C for 3 min, the supernatants were removed, then fresh medium 199 was added, and the pancreatic tissue was suspended again. The suspension was transferred to a 6 cm dish, then islets were picked up using a stereomicroscope and used in the experiment.

#### 4.11. Statistical Analyses

Data were presented as mean  $\pm$  SEM. Significant differences between the two groups were assessed using Student's or Welch's *t*-test. For multiple comparisons, we used the repeated measure one-way ANOVA with Bonferroni adjustment. A *p*-value less than 0.05 was considered significant. Statistical analyses were performed using IBM SPSS Statistics version 27 software.

**Supplementary Materials:** The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/ijms23074003/s1>.

**Author Contributions:** N.H. collected, analyzed, interpreted the data, and drafted the article. K.F. designed, conceived, and reviewed the experiments. M.K., D.K., O.K., H.Y.-H., E.W., Y.I., Y.T., G.W.D.II, K.U. and R.N. reviewed the data. T.K. conceived the experiments, reviewed the data, edited the article, and approved the version to be published. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was supported by The Jikei University Research Fund for Graduate Students (to N.H.) and Grant-in-Aid for Scientific Research (C) (18K08491) from the Ministry of Education, Culture, Sports, Science, and Technology of Japan (to K.F.).

**Institutional Review Board Statement:** This study was approved by the Animal Care and Use Committee (Permission number: 18-027) and Safety Committee for Recombinant DNA Experiments (Permission number: 18-063) in Gunma University.

**Acknowledgments:** We thank Y. Fujitani (Laboratory of Developmental Biology & Metabolism, Institute for Molecular and Cellular Regulation, Gunma University) and members of his laboratory for providing *Gcg<sup>CreERT2</sup>* mice and critical discussions of the data.

**Conflicts of Interest:** K.F. received honorarium from Sanofi K.K, Merck Sharpe & Dohme, Taisho Pharmaceutical, Eli Lilly Japan KK, TERUMO Corporation, ARKRAY, Astellas Pharma, AstraZeneca, Nippon Boehringer Ingelheim, Mitsubishi Tanabe Pharma Corporation, Ono Pharmaceutical, Novo Nordisk Pharma, Kissei Pharmaceutical, Sumitomo Dainippon Pharma, Kowa Pharmaceutical, Takeda Pharmaceutical, Daiichi Sankyo, Chugai Pharmaceutical, Abbot Japan, Otsuka Pharmaceutical and Kyowa Hakko Kirin. R.N. received honorarium from Sanofi K.K, Nippon Boehringer Ingelheim, Takeda Pharmaceutical, Kissei Pharmaceutical, Novartis Pharma, Eli Lilly Japan KK, Novo Nordisk Pharma, MSD, Astellas Pharma, Taisho Pharmaceutical, Ono Pharmaceutical.

## Abbreviations

cAMP	cyclic adenosine monophosphate
CBD	common bile duct
CF	catalytic fragment
DAG	diacylglycerol
FFA	free fatty acid
Gcg	glucagon
GFP	green fluorescent protein
IBMX	3-Isobutyl 1-methylxanthine
ITT	insulin tolerance test
IPGTT	intraperitoneal glucose tolerance test
K <sub>ATP</sub>	adenosine triphosphate-sensitive potassium
KO	knockout
KRB	krebs ringer buffer
MARCKS	myristoylated alanine-rich C-kinase substrate
NDS	normal donkey serum
PFA	paraformaldehyde
Pkc	protein kinase C
PMA	phorbol 12-myristate 13-acetate
siRNA	small interfering ribonucleic acid
SNARE	soluble <i>n</i> -ethylmaleimide sensitive factor attachment protein receptor
SSTR	somatostatin receptor
T2DM	type 2 diabetes
VDCC	voltage-dependent calcium channels
αPkcδKO	α-cell-specific Pkcδ knockout

## References

- Hancock, A.S.; Du, A.; Liu, J.; Miller, M.; May, C.L. Glucagon deficiency reduces hepatic glucose production and improves glucose tolerance in adult mice. *Mol. Endocrinol.* **2010**, *24*, 1605–1614. [[CrossRef](#)] [[PubMed](#)]
- Lee, Y.; Wang, M.Y.; Du, X.Q.; Charron, M.J.; Unger, R.H. Glucagon receptor knockout prevents insulin-deficient type 1 diabetes in mice. *Diabetes* **2011**, *60*, 391–397. [[CrossRef](#)] [[PubMed](#)]
- Lee, Y.; Berglund, E.D.; Wang, M.Y.; Fu, X.; Yu, X.; Charron, M.J.; Burgess, S.C.; Unger, R.H. Metabolic manifestations of insulin deficiency do not occur without glucagon action. *Proc. Natl. Acad. Sci. USA* **2012**, *109*, 14972–14976. [[CrossRef](#)] [[PubMed](#)]
- Unger, R.H.; Cherrington, A.D. Glucagonocentric restructuring of diabetes: A pathophysiologic and therapeutic makeover. *J. Clin. Investig.* **2012**, *122*, 4–12. [[CrossRef](#)] [[PubMed](#)]
- Barg, S.; Galvanovskis, J.; Göpel, S.O.; Rorsman, P.; Eliasson, L. Tight coupling between electrical activity and exocytosis in mouse glucagon-secreting α-cells. *Diabetes* **2000**, *49*, 1500–1510. [[CrossRef](#)] [[PubMed](#)]
- Gilon, P. The role of alpha-cells in islet function and glucose homeostasis in health and type 2 diabetes. *J. Mol. Biol.* **2020**, *432*, 1367–1394. [[CrossRef](#)]
- Le Marchand, S.J.; Piston, D.W. Glucose suppression of glucagon secretion: Metabolic and calcium responses from α-cells in intact mouse pancreatic islets. *J. Biol. Chem.* **2010**, *285*, 14389–14398. [[CrossRef](#)]
- Le Marchand, S.J.; Piston, D.W. Glucose decouples intracellular Ca<sup>2+</sup> activity from glucagon secretion in mouse pancreatic islet α-cells. *PLoS ONE* **2012**, *7*, e47084. [[CrossRef](#)]
- Quoix, N.; Cheng-Xue, R.; Mattart, L.; Zeinoun, Z.; Guiot, Y.; Beauvois, M.C.; Henquin, J.C.; Gilon, P. Glucose and pharmacological modulators of ATP-sensitive K<sup>+</sup> channels control [Ca<sup>2+</sup>]<sub>i</sub> by different mechanisms in isolated mouse α-cells. *Diabetes* **2009**, *58*, 412–421. [[CrossRef](#)]
- Dickerson, M.T.; Dadi, P.K.K.; Altman, M.K.; Verlage, K.R.; Thorson, A.S.; Jordan, K.L.; Vierra, N.C.; Amarnath, G.; Jacobson, D.A. Glucose-mediated inhibition of calcium-activated potassium channels limits α-cell calcium influx and glucagon secretion. *Am. J. Physiol. Endocrinol. Metab.* **2019**, *316*, E646–E659. [[CrossRef](#)]
- Yu, Q.; Shuai, H.; Ahooghalandari, P.; Gylfe, E.; Tengholm, A. Glucose controls glucagon secretion by directly modulating cAMP in α cells. *Diabetologia* **2019**, *62*, 1212–1224. [[CrossRef](#)]
- Elliott, A.D.; Ustione, A.; Piston, D.W. Somatostatin and insulin mediate glucose-inhibited glucagon secretion in the pancreatic α-cell by lowering cAMP. *Am. J. Physiol. Endocrinol. Metab.* **2015**, *308*, E130–E143. [[CrossRef](#)] [[PubMed](#)]
- Briant, L.J.B.; Reinbothe, T.M.; Spiliotis, I.; Miranda, C.; Rodriguez, B.; Rorsman, P. δ-cells, and β-cells are electrically coupled and regulate α-cell activity via somatostatin. *J. Physiol.* **2018**, *596*, 197–215. [[CrossRef](#)] [[PubMed](#)]
- Cejvan, K.; Coy, D.H.; Holst, J.J.; Cerasi, E.; Efendic, S. Gliclazide directly inhibits arginine-induced glucagon release. *Diabetes* **2002**, *51*, S381–S384. [[CrossRef](#)] [[PubMed](#)]

15. Kikkawa, U.; Matsuzaki, H.; Yamamoto, T. Protein kinase C8 (PKC8): Activation mechanisms and functions. *J. Buxhem.* **2002**, *132*, 831–839.
16. Reyland, M.E. Protein kinase C isoforms: Multi-functional regulators of cell life and death. *Front. Biosci.* **2009**, *14*, 2386–2399. [[CrossRef](#)]
17. Salzer, E.; Santos-Valente, E.; Keller, B.; Warnatz, K.; Boztug, K. Protein kinase C  $\delta$ : A gatekeeper of immune homeostasis. *J. Clin. Immunol.* **2016**, *36*, 631–640. [[CrossRef](#)]
18. Suzuki, Y.; Zhang, H.; Saito, N.; Kojima, I.; Urano, T.; Mogami, H. Glucagon-like peptide 1 activates protein kinase C through  $\text{Ca}^{2+}$ -dependent activation of phospholipase C in insulin-secreting cells. *J. Biol. Chem.* **2006**, *281*, 28499–28507. [[CrossRef](#)]
19. Xia, P.; Inoguchi, T.; Kern, T.S.; Engerman, R.L.; Oates, P.J.; King, G.L. Characterization of the mechanism for the chronic activation of diacylglycerol-protein kinase C pathway in diabetes and hypergalactosemia. *Diabetes* **1994**, *43*, 1122–1129. [[CrossRef](#)]
20. Geraldes, P.; Hiraoka-Yamamoto, J.; Matsumoto, M.; Clermont, A.; Leitges, M.; Marette, A.; Aiello, L.P.; Kern, T.S.; King, G.L. Activation of PKC- $\delta$  and SHP-1 by hyperglycemia causes vascular cell apoptosis and diabetic retinopathy. *Nat. Med.* **2009**, *15*, 1298–1306. [[CrossRef](#)]
21. Mima, A.; Kitada, M.; Geraldes, P.; Li, Q.; Matsumoto, M.; Mizutani, K.; Qi, W.; Li, C.; Leitges, M.; Rask-Madsen, C.; et al. Glomerular VEGF resistance induced by PKCdelta/SHP-1 activation and contribution to diabetic nephropathy. *FASEB J.* **2012**, *26*, 2963–2974. [[CrossRef](#)] [[PubMed](#)]
22. Hennige, A.M.; Häring, H.U.; Ranta, F.; Heinzlmann, I.; Düfer, M.; Lutz, S.Z.; Lammers, R.; Drews, G.; Ullrich, A.S. Overexpression of kinase-negative protein kinase C $\delta$  in pancreatic  $\beta$ -cells protects mice from diet-induced glucose intolerance and  $\beta$ -cell dysfunction. *Diabetes* **2010**, *59*, 119–127. [[CrossRef](#)] [[PubMed](#)]
23. De Marinis, Y.Z.; Zhang, E.; Amisten, S.; Taneera, J.; Renstrom, E.; Rorsman, P.; Eliasson, L. Enhancement of glucagon secretion in mouse and human pancreatic alpha cells by protein kinase C (PKC) involves intracellular trafficking of PKC $\alpha$  and PKC $\delta$ . *Diabetologia* **2010**, *53*, 717–729. [[CrossRef](#)] [[PubMed](#)]
24. Hii, C.S.; Stutchfield, J.; Howell, S.L. Enhancement of glucagon secretion from isolated rat islets of Langerhans by phorbol 12-myristate 13-acetate. *Biochem. J.* **1986**, *233*, 287–289. [[CrossRef](#)]
25. Yamamoto, K.; Mizuguchi, H.; Tokashiki, N.; Kobayashi, M.; Tamaki, M.; Sato, Y.; Fukui, H.; Yamauchi, A. Protein kinase C-signaling regulates glucagon secretion from pancreatic islets. *J. Med. Investig.* **2017**, *64*, 122–128. [[CrossRef](#)]
26. Takaki, R.; Ono, J.; Nakamura, M.; Yokogawa, Y.; Kumae, S.; Hiraoka, T.; Yamaguchi, K.; Hamaguchi, K.; Uchida, S. Isolation of glucagon-secreting cell lines by cloning insulinoma cells. *In Vitro Cell. Dev. Biol.* **1986**, *22*, 120–126. [[CrossRef](#)]
27. Ono, J.; Yamaguchi, K.; Okeda, T.; Asano, T.; Takaki, R. Characterization of secretory responses of a glucagon-producing In-R1-G9 cell line. *Duthetes Res. Clin. Pract.* **1988**, *4*, 203–207. [[CrossRef](#)]
28. Soltoff, S.P. Rottlerin: An inappropriate and ineffective inhibitor of PKC $\delta$ . *Trends Pharmacol. Sci.* **2007**, *28*, 453–458. [[CrossRef](#)]
29. Song, M.; Matkovich, S.J.; Zhang, Y.; Hammer, D.J.; Dorn, G.W., 2nd. Combined cardiomyocyte PKC $\delta$  and PKC $\epsilon$  gene deletion uncover their central role in restraining developmental and reactive heart growth. *Sci. Signal.* **2015**, *8*, ra39. [[CrossRef](#)]
30. Herrera, P.L. Adult insulin- and glucagon-producing cells differentiate from two independent cell lineages. *Development* **2000**, *127*, 2317–2322. [[CrossRef](#)]
31. Tuduri, E.; Denroche, H.C.; Kara, J.A.; Asadi, A.; Fox, J.K.; Kieffer, T.J. Partial ablation of leptin signaling in mouse pancreatic  $\alpha$ -cells does not alter either glucose or lipid homeostasis. *Am. J. Physiol. Endocrinol. Metab.* **2014**, *306*, E748–E755. [[CrossRef](#)] [[PubMed](#)]
32. Shiota, C.; Prasadani, K.; Guo, P.; Fusco, J.; Xiao, X.; Gittes, G.K. Gcg (CreERT2) knockin mice as a tool for genetic manipulation in pancreatic alpha cells. *Diabetologia* **2017**, *60*, 2399–2408. [[CrossRef](#)] [[PubMed](#)]
33. Madisen, L.; Zwingman, T.A.; Sunkin, S.M.; Oh, S.W.; Zariwala, H.A.; Gu, H.; Ng, L.L.; Palmiter, R.D.; Hawrylycz, M.J.; Jones, A.R.; et al. A robust and high-throughput Cre reporting and characterization system for the whole mouse brain. *Nat. Neurosci.* **2010**, *13*, 133–140. [[CrossRef](#)] [[PubMed](#)]
34. Hayashi, Y.; Yamamoto, M.; Mizoguchi, H.; Watanabe, C.; Ito, R.; Yamamoto, S.; Sun, X.Y.; Murata, Y. Mice deficient for glucagon gene-derived peptides display normoglycemia and hyperplasia of islet  $\alpha$ -cells but not of intestinal L-cells. *Mol. Endocrinol.* **2009**, *23*, 1990–1999. [[CrossRef](#)] [[PubMed](#)]
35. Andersson, S.A.; Pedersen, M.G.; Vikman, J.; Eliasson, L. Glucose-dependent docking and SNARE protein-mediated exocytosis in mouse pancreatic alpha-cell. *Pflug. Arch.* **2011**, *462*, 443–454. [[CrossRef](#)]
36. Kawamori, D.; Kurpad, A.J.; Hu, J.; Liew, C.W.; Shih, J.L.; Ford, E.L.; Herrera, P.L.; Polonsky, K.S.; McGuinness, O.P.; Kulkarni, R.N. Insulin signaling in  $\alpha$  cells modulates glucagon secretion In Vivo. *Cell Metab.* **2009**, *9*, 350–361. [[CrossRef](#)]
37. Ichikawa, R.; Takano, K.; Fujimoto, K.; Motomiya, T.; Kobayashi, M.; Kitamura, T.; Shichiri, M. Basal glucagon hypersecretion and response to oral glucose load in prediabetes and mild type 2 diabetes. *Endocr. J.* **2019**, *66*, 663–675. [[CrossRef](#)]
38. Kobayashi, M.; Satoh, H.; Matsuo, T.; Kusunoki, Y.; Tokushima, M.; Watada, H.; Namba, M.; Kitamura, T. Plasma glucagon levels measured by sandwich ELISA are correlated with impaired glucose tolerance in type 2 diabetes. *Endocr. J.* **2020**, *63*, 903–922. [[CrossRef](#)]
39. Dorrell, C.; Grompe, M.T.; Pan, F.C.; Zhong, Y.; Canaday, P.S.; Shultz, L.D.; Greiner, D.L.; Wright, C.V.; Streeter, P.R.; Grompe, M. Isolation of mouse pancreatic alpha, beta, duct and acinar populations with cell surface markers. *Mol. Cell. Endocrinol.* **2011**, *339*, 144–150. [[CrossRef](#)]

40. Muraro, M.J.; Dharmadhikari, G.; Grun, D.; Groen, N.; Dielen, T.; Jansen, E.; van Gurp, L.; Engelse, M.A.; Carlotti, F.; de Koning, E.J. A single-cell transcriptome atlas of the human pancreas. *Cell Syst.* **2016**, *3*, 385–394.e3. [[CrossRef](#)]
41. Zadeh, E.H.G.; Huang, Z.; Xia, J.; Li, D.; Davidson, H.W.; Li, W.H. ZIGIR, a granule-specific Zn<sup>2+</sup> indicator, reveals human islet alpha cell heterogeneity. *Cell Rep.* **2020**, *32*, 107904. [[CrossRef](#)] [[PubMed](#)]