



Projective and Reedy Model Category Structures for (Infinitesimal) Bimodules over an Operad

Julien Ducoulombier¹ · Benoit Fresse² · Victor Turchin³

Received: 13 January 2021 / Accepted: 27 January 2022 / Published online: 8 April 2022
© The Author(s) 2022

Abstract

We construct and study projective and Reedy model category structures for bimodules and infinitesimal bimodules over topological operads. Both model structures produce the same homotopy categories. For the model categories in question, we build explicit cofibrant and fibrant replacements. We show that these categories are right proper and under some conditions left proper. We also study the extension/restriction adjunctions.

Keywords Model category · Operad · Homotopy theory · Reedy category · Cofibrant/fibrant replacements · Algebras and modules over operads · Infinitesimal bimodules and bimodules · Left/right properness · Extension/restriction adjunctions

Contents

Introduction	826
1 Model Category Structures for Operads	830
1.1 The Projective Model Categories of G -Spaces and of Σ -Sequences	831

Communicated by Michael Batanin.

J.D. was partially supported by the NCCR SwissMAP funded by the Swiss National Science Foundation and the ERC starting Grant GRAPHCPX and the Max Planck Institute for Mathematics in Bonn. B.F. acknowledges support from the Labex CEMPI (ANR-11-LABX-0007-01) and from the FNS-ANR project OCHoTop (ANR-18CE93-0002-01). V.T. has benefited from a visiting position of the Labex CEMPI (ANR-11-LABX-0007-01) at the Université de Lille and a visiting position at the Max Planck Institute for Mathematics in Bonn for the achievement of this work. He was also partially supported by the Simons Foundation Grant, award ID: 519474. J.D. and V.T. acknowledge the University of Lille for hospitality.

✉ Julien Ducoulombier
julien@mpim-bonn.mpg.de
Benoit Fresse
benoit.fresse@univ-lille.fr
Victor Turchin
turchin@ksu.edu

¹ Max Planck Institute for Mathematics, Vivatsgasse 7, 53111 Bonn, Germany
² Univ. Lille, CNRS, UMR 8524 - Laboratoire Paul Painlevé, F-59000 Lille, France
³ Department of Mathematics, Kansas State University, 138 Cardwell Hall, Manhattan, KS 66506, USA

1.2	The Reedy Model Categories of Λ - and $\Lambda_{>0}$ -Sequences	834
1.3	A Fibrant Replacement Functor for Λ - and $\Lambda_{>0}$ -Sequences	835
1.4	The Projective/Reedy Model Category of Operads	840
1.5	Properties of Cofibrant Operads	841
2	The Projective Model Category of $(P-Q)$ -Bimodules	847
2.1	Properties of the Category of Bimodules	848
2.1.1	Bimodules as Algebras over a Colored Operad	848
2.1.2	The Free Bimodule Functor	854
2.1.3	Combinatorial Description of the Pushout	857
2.2	The Model Category Structure	860
2.2.1	Relative Left Properness of the Projective Model Category	861
2.2.2	Extension/Restriction Adjunction for the Projective Model Category of Bimodules	869
3	The Reedy Model Category of $(P-Q)$ -Bimodules	871
3.1	Properties of the Reedy Model Category of Bimodules	874
3.1.1	A Reedy Fibrant Replacement Functor for Bimodules	874
3.1.2	Characterization of Reedy Cofibrations for Bimodules	881
3.1.3	Left Properness and Extension/Restriction Functors for the Reedy Model Structure	885
3.2	The Connection Between the Model Category Structures on Bimodules	886
3.2.1	Quillen Adjunction Between the Model Category Structures	887
3.2.2	Boardman–Vogt Type Resolution in the Projective/Reedy Model Category	888
3.2.3	A Functorial Cofibrant Replacement in the Projective/Reedy Model Category	891
3.3	The Reedy Fibrant Replacement as an Internal Hom	891
3.3.1	The Right Closed Monoidal Category of Symmetric Sequences	892
3.3.2	The Tensor-Hom Adjunction	892
3.3.3	The Reedy Fibrant Replacement	893
3.4	The Subcategory of Reduced Bimodules	894
3.4.1	Main Properties	894
3.4.2	Quillen Adjunction Between the Reedy Model Categories	896
4	The Projective Model Category of O Infinitesimal Bimodules	897
4.1	Properties of the Category of Infinitesimal Bimodules	898
4.1.1	Infinitesimal Bimodules as Algebras over a Colored Operad	898
4.1.2	The Free Bimodule Functor	899
4.1.3	Combinatorial Description of the Pushout	901
4.2	Model Category Structure	902
4.2.1	Left Properness of the Projective Model Category	902
4.2.2	Extension/Restriction Adjunction for the Projective Model Category of Infinitesimal Bimodules	902
5	The Reedy Model Category of O Infinitesimal Bimodules	904
5.1	Properties of the Reedy Model Category of Infinitesimal Bimodules	904
5.1.1	Fibrant Replacement Functor for Infinitesimal Bimodules	905
5.1.2	Characterization of Cofibrations/Left properness/Extension-restriction adjunction	906
5.2	The Connection Between the Model Category Structures on Infinitesimal Bimodules	908
5.2.1	Quillen Adjunction Between the Model Category Structures	908
5.2.2	Cofibrant Resolution in the Projective/Reedy Model Category	909
A	Equivariant Homotopy Theory	912
A.1	Projective Cofibrations for Monoidal Action	912
A.2	Cellularly Equivariant Cofibrations	917
	References	919

Introduction

In this paper, we set up a homotopy theory for the categories of bimodules and of infinitesimal bimodules over topological operads. More precisely, we study two model structures, the *projective* model structure and the Reedy model structure, which we define for both bimodule categories. The Reedy and the projective model structures have the same class of weak equivalences and, therefore, produce equivalent homotopy categories. Both model categories find important applications in the manifold functor calculus, specifically in the problems of

delooping the functor calculus towers [14–16]. It is well known that the arity zero elements essentially complicate the homotopy theory of such objects. However, in practical examples the arity zero component of the studied objects is often reduced to a point. Such objects are called *reduced*. Motivated by the homotopy theory of the little 2-discs operad, the second author developed the Reedy model structure for reduced operads [18, 19]. We adapt this theory to the setting of bimodules and of infinitesimal bimodules. One of the advantages of the Reedy model structure in comparison to the projective one is that the cofibrant resolutions are smaller as they do not take into account the arity zero component. This makes the constructions of delooping in [14–16] simpler. By contrast, while all the objects are fibrant for the projective model categories, there is no obvious Reedy fibrant coresolution. Consequently, both model structures have their advantages and it can be convenient to be able to switch from one structure to another.

The starting idea of the Reedy model structure for reduced operads is to encode the operadic composition operations with the unique point in arity zero in an extension of the diagram structure which underlies our objects. In the usual category of symmetric operads, the diagram structure of the objects is governed by the category $\Sigma = \coprod_n \Sigma_n$ which is defined by taking the disjoint union of the symmetric groups Σ_n . In what follows, we use the expression ‘ Σ -sequence’ for the objects of the category of diagrams over Σ . We use the notation ΣSeq for this category of diagrams and the notation $\Sigma Operad$ for the category of symmetric operads. To formalize the construction of the Reedy model structure, we consider the category Λ , which has the finite sets $[n] = \{1, \dots, n\}$ as objects and all injective maps of finite sets $u : \{1, \dots, m\} \hookrightarrow \{1, \dots, n\}$ as morphisms. We use the expression ‘ Λ -sequence’ for the objects of the category of contravariant diagrams over Λ , and we use the notation ΛSeq for the category of Λ -sequences. The composition operations with the arity zero term $P(0) = *$ in a reduced operad P are equivalent to restriction operators $u^* : P(n) \rightarrow P(m)$, which can be associated to the injective maps of finite sets $u : \{1, \dots, m\} \hookrightarrow \{1, \dots, n\}$ and hence to the morphisms in the category Λ . This observation implies that the category of reduced operads is identified with a category $\Lambda_* Operad$, whose objects are operads shaped on this category of finite sets and injections Λ instead of the category of permutations Σ .

The category of Λ -sequences inherits a Reedy model structure, in which the fibrations are defined by using a natural notion of matching object. The Reedy model structure of reduced operads is precisely defined by transferring this Reedy model structure on the category of Λ -sequences ΛSeq to our category of operads $\Lambda_* Operad$, while the projective model structure of symmetric operads is defined by transferring the projective model structure on the category of Σ -sequences ΣSeq to $\Sigma Operad$.

Throughout this paper, we work in the category of topological spaces, and we therefore deal with operads in topological spaces. In that context, the projective model category of symmetric operads is known to be left proper relative to Σ -cofibrant operads (i.e. operads that are cofibrant as Σ -sequences) and right proper [27] making the homotopy colimits and limits easier to identify in this category. Furthermore, all operads are fibrant in the projective model category of symmetric operads in topological spaces. In the Reedy model category of reduced operads, the objects are not necessarily fibrant. We use a notion of matching object to define the class of fibrations (and unfortunately, we have no explicit definition of a fibrant coresolution functor at the time), but the class of cofibrations is larger. In fact, a morphism of reduced operads is a cofibration with respect to the Reedy model structure if and only if this morphism defines a projective cofibration of operads after forgetting the arity zero components [19, Theorem 8.4.12] (thus if and only if this morphism defines a cofibration in the projective model category of operads with a void component in arity zero). This result implies that the Reedy model category of reduced operads is also left proper

relative to Σ -cofibrant reduced operads. Together with Willwacher, the second and third authors [21] showed that, for any reduced operads P and Q , there is a weak equivalence of derived mapping spaces

$$\Sigma\text{Operad}^h(P; Q) \simeq \Lambda_*\text{Operad}^h(P; Q).$$

Hence we can use the Reedy model category to compute mapping spaces in the usual category of topological operads.

The main purpose of this work is to extend the results of these operadic homotopy theories to the setting of bimodules and of infinitesimal bimodules. First of all, we define our counterparts of the projective and Reedy model structures for bimodules and infinitesimal bimodules in topological spaces. We also address the definition of these model structures for truncated bimodules.

For the definition of the projective model structure, we work out difficulties that occur in the context of topological spaces, notably regarding the application of the small object argument (see the discussion of [29]). This question is independent from other works on the projective model categories of modules over operads carried out in the litterature. In fact, the projective model structure of bimodules was defined in [35] for bimodules in simplicial sets and for bimodules in a category of simplicial bimodules over a ring. The paper [24] gives the definition of an analogous model structure for left modules over non-symmetric operads and for left modules over symmetric operads when every Σ -sequence is projectively cofibrant in the base category (for instance, when the base category is a category of chain complexes over a characteristic zero field). The book [20], by the second author, provides a general study of the homotopy theory of modules and bimodules over operads, but deals with semi-model structures (with a restriction of the application of the axioms to maps with a cofibrant source) to get results that are valid in any base monoidal model category. In the paper, we prove that, when we work in the category of topological spaces, we have a full validity of the definition of the projective model category of bimodules over a pair of operads, and this result holds without any assumption on our operads. We get the same result for the definition of the projective model category of infinitesimal bimodules.

For the definition of the Reedy model categories of bimodules and of infinitesimal bimodules, we rely on a preliminary definition of a fibrant coresolution functor and we apply a transfer argument, using an adjunction between (infinitesimal) bimodules and Λ -sequences. We just need a mild assumption on our operads to ensure the validity of the definition of the Reedy model structures (technically, we just need to consider well-pointed operads, in which the inclusion of the operadic unit in arity one defines a cofibration of spaces).

We use the notation $\Sigma\text{Bimod}_{P; Q}$ for the category of bimodules associated to a pair of operads (P, Q) , while we adopt the notation ΣIbimod_O for the category of infinitesimal bimodules over an operad O . To distinguish the Reedy model structure from the projective model structure, we adopt the convention to keep these notations $\Sigma\text{Bimod}_{P; Q}$ and ΣIbimod_O when we equip these bimodule categories with the projective model structure, and we pass to the notations $\Lambda\text{Bimod}_{P; Q}$ and ΛIbimod_O when we consider the Reedy model structure. We prove that our model categories have the following features.

- ▶ Sections 3.1.1 and 5.1.1: All the objects in $\Sigma\text{Bimod}_{P; Q}$ and ΣIbimod_O are fibrant. Furthermore, we give explicit fibrant coresolutions in the Reedy model categories $\Lambda\text{Bimod}_{P; Q}$ and ΛIbimod_O .
- ▶ Sections 2.2.1, 3.1.3, 4.2.1 and 5.1.2: The categories $\Sigma\text{Bimod}_{P; Q}$, $\Lambda\text{Bimod}_{P; Q}$, ΣIbimod_O and ΛIbimod_O are right proper. Moreover, if P is either projectively or Reedy cofibrant, and Q, O are componentwise cofibrant, then the categories $\Sigma\text{Bimod}_{P; Q}$

and $\Lambda\text{Bimod}_{P;Q}$ are left proper relative to componentwise cofibrant objects while ΣIbimod_O and ΛIbimod_O are left proper.

- ▶ Sections 3.1.2 and 5.1.2: Let $Q_{>0}$ and $O_{>0}$ be the sub-operads obtained from Q and O , respectively, by removing the arity zero components. Any map in $\Lambda\text{Bimod}_{P;Q}$ and ΛIbimod_O is a cofibration if and only if the corresponding map in $\Sigma\text{Bimod}_{P;Q_{>0}}$ and $\Sigma\text{Ibimod}_{O_{>0}}$, respectively, is a cofibration.
- ▶ Sections 3.2.1 and 5.2.1: If M and N are $(P-Q)$ -bimodules, while M' and N' are O infinitesimal bimodules, then one has weak equivalences between derived mapping spaces

$$\begin{aligned} \Sigma\text{Bimod}_{P;Q}^h(M;N) &\simeq \Lambda\text{Bimod}_{P;Q}^h(M;N) && \text{and} \\ \Sigma\text{Ibimod}_O^h(M';N') &\simeq \Lambda\text{Ibimod}_O^h(M';N'). \end{aligned}$$

- ▶ Sections 2.2.2, 3.1.3, 4.2.2 and 5.1.2: Let $\phi_1 : P \rightarrow P'$, $\phi_2 : Q \rightarrow Q'$ and $\phi : O \rightarrow O'$ be weak equivalences between Σ -cofibrant operads P, P' and componentwise cofibrant operads Q, Q', O, O' , then the extension and restriction functors form Quillen equivalences

$$\begin{aligned} \phi_1 : \Sigma\text{Bimod}_{P;Q} &\rightleftarrows \Sigma\text{Bimod}_{P';Q'} : \phi^*, & \phi_1 : \Sigma\text{Ibimod}_O &\rightleftarrows \Sigma\text{Ibimod}_{O'} : \phi^*, \\ \phi_1 : \Lambda\text{Bimod}_{P;Q} &\rightleftarrows \Lambda\text{Bimod}_{P';Q'} : \phi^*, & \phi_1 : \Lambda\text{Ibimod}_O &\rightleftarrows \Lambda\text{Ibimod}_{O'} : \phi^*. \end{aligned}$$

(Also, see Sects. 2.2.2 and 3.4.1 for a refinement of this result when we forget about the arity zero components of bimodules or, respectively, if such components are reduced to a point.)

Organization of the paper: We review the background of our constructions in the first section of the paper. We review the definition of the projective model category of Σ -sequences and the definition of the Reedy model category of Λ -sequences. We also build an explicit fibrant coresolution in the category of Λ -sequences. Then we recall the definition of the projective model category of operads and the Reedy model category of reduced operads together with their properties. Most of the new results in this section appear at the very end in Sect. 1.5, where we study some natural properties of cofibrant operads.

In the second section, we define the projective model category of $(P-Q)$ -bimodules. First, we show that this category is equivalent to the category of algebras over a colored operad. Then we give combinatorial descriptions of the free bimodule functor and of pushouts. After that, we define the projective model category structure for bimodules and we prove that this model structure is relatively left proper and that the extension/restriction adjunctions along weak equivalences of operads form Quillen equivalences.

In the third section, we study the Reedy model category of bimodules. We check that (almost all) the constructions introduced in the second section can be extended to the Reedy model category. Furthermore, we give an explicit Reedy fibrant coresolution as well as a characterization of cofibrations. As a consequence of this characterization, we show that this model category is relatively left proper and that the extension/restriction adjunctions along weak equivalences of reduced operads form Quillen equivalences. Both model structures having the same set of weak equivalences, they produce the same homotopy category. Then we construct a functorial cofibrant resolution for bimodules in both (projective and Reedy) model structures. As an application, we explain how our Reedy fibrant coresolution can be expressed in terms of internal hom in the category of Σ -sequences. In the last subsection, assuming that both operads P and Q are reduced, we study the subcategory $\Lambda_*\text{Bimod}_{P;Q}$ of reduced bimodules equipped with the Reedy model category structure. This subcategory

enjoys slightly better properties as we compare the Reedy model structures of bimodules and of reduced bimodules.

In the fourth section we adapt the results from the second section to the context of infinitesimal bimodules. In that case, the proofs are easier since pushouts of infinitesimal bimodules coincide with pushouts taken componentwise. Similarly, we show that this category is equivalent to the category of algebras over a colored operad. After that, we introduce the projective model category structure and we prove that the extension/restriction adjunctions along weak equivalences of operads form Quillen equivalences.

In the fifth section we adapt the results from the third section to the context of infinitesimal bimodules. In the same way, we build an explicit fibrant coresolution. We give a characterization of cofibrations and, as a consequence of it, we show that the extension/restriction adjunctions along weak equivalences of reduced operads form Quillen equivalences. We then compare the projective and Reedy model structures on infinitesimal bimodules. At the end we exhibit explicit cofibrant resolutions.

The last sixth section is an “Appendix” where several technical lemmas that we use from the equivariant homotopy theory are formulated and proved.

Notation: In [18, 19], the notation $\Lambda_*\text{Operad}$ actually refers to a category of Λ -operads, which is defined by dropping the arity zero component of reduced operads. But we do not use this convention in this paper. We therefore forget about the refined structure of a Λ -operad and we use the notation $\Lambda_*\text{Operad}$ for the category of reduced operads. We keep the letter Λ in order to emphasize the underlying Λ -diagram structure of our objects, but we forget about further reductions in the definition of our structures. We write Λ_* instead of Λ to remind that the operads in this category are reduced.

In the paper we use many different sets of rooted trees. As a general rule, we use letter \mathbb{P} for sets of planar trees and letter \mathbb{T} for non-planar trees. Usually the set $\ell(T)$ of leaves of a planar tree T is labelled by a permutation in $\Sigma_{|T|}$, where $|T|$ is the number of leaves in the tree. Internal vertices in these trees are usually allowed to have any arity unless we use the superscript ≥ 1 or ≥ 2 , like in $\mathbb{P}^{\geq 1}$ or $\mathbb{T}^{\geq 2}$ meaning that the arities of vertices are ≥ 1 or ≥ 2 , respectively. We use different terms to deal with trees in the context of operads, such as set of leaves $\ell(T)$, set of vertices $V(T)$, set of edges $E(T)$, the arity $|v|$ of a vertex v , etc from [6, Section 5.8]. Even though we formally do not define these terms, we show them on figures, so that the reader can easily guess the meaning of those words without referring to loc. cit.

For a subspace of a space X , we sometimes use notation ∂X . By $\partial \prod_{i \in I} X_i$ we understand a subspace in $\prod_{i \in I} X_i$ consisting of points with at least one coordinate in ∂X_i . A point in the product space $\prod_{i \in I} X_i$ is usually denoted by $\{x_i\}_{i \in I}$ or just $\{x_i\}$ if there is no ambiguity about the set I .

1 Model Category Structures for Operads

In this section, we introduce the categories ΣSeq and ΛSeq as well as their model category structures called *projective* and *Reedy* model category structures. These are categories whose objects are sequences of topological spaces with some extra structures. We also define the categories of operads ΣOperad and reduced operads $\Lambda_*\text{Operad}$. Both categories inherit model category structures from the following adjunctions in which the functors \mathcal{F}^Σ and \mathcal{F}^Λ

are the left adjoints to the forgetful functors:

$$\mathcal{F}^\Sigma : \Sigma Seq \rightleftarrows \Sigma Operad : \mathcal{U}^\Sigma \quad \text{and} \quad \mathcal{F}^\Lambda : \Lambda_{>0} Seq \rightleftarrows \Lambda_* Operad : \mathcal{U}^\Lambda.$$

Here $\Lambda_{>0} Seq$ can be interpreted as the full model subcategory of ΛSeq composed of objects whose arity zero component is the one point topological space (see Sect. 1.2). Both model categories $\Sigma Operad$ and $\Lambda_* Operad$ have intensively been studied by Berger-Moerdijk [6] and the second author [18, 19]. We list their properties in Sect. 1.4. Usually, in order to define a model category structure from an adjunction, we use the following statement also called the *transfer principle*:

Theorem 1.1 [6, Section 2.5] *Let \mathcal{D} be a cofibrantly generated model category with a set of generating cofibrations \mathcal{D}_c and a set of generating acyclic cofibrations \mathcal{D}_{ac} . Let $L : \mathcal{D} \rightleftarrows \mathcal{C} : R$ be an adjunction with left adjoint L and right adjoint R . Assume that \mathcal{C} is bicomplete. Define a map f in \mathcal{C} to be a weak equivalence (respectively, a fibration) if $R(f)$ is a weak equivalence (respectively, a fibration) in \mathcal{D} . If the following conditions are satisfied:*

- (i) *both sets $L(\mathcal{D}_c)$ and $L(\mathcal{D}_{ac})$ permit the small object argument;*
- (ii) *\mathcal{C} has a fibrant replacement functor for objects;*
- (iii) *\mathcal{C} has a functorial path object for fibrant objects, i.e. for any fibrant object X there is a functorial factorization of the diagonal map into a weak equivalence followed by a fibration*

$$X \xrightarrow{\simeq} Path(X) \twoheadrightarrow X \times X;$$

then we have a cofibrantly generated model category structure on \mathcal{C} in which the set of generating cofibrations (respectively acyclic cofibrations) is given by $L(\mathcal{D}_c)$ (respectively, $L(\mathcal{D}_{ac})$). Furthermore, this model category structure makes the adjunction $(L; R)$ into a Quillen adjunction.

As explained in the following subsections, all objects in the category ΣSeq are fibrant and the identity functor produces a functorial fibrant replacement in the category $\Sigma Operad$. So, the transfer principle can easily be applied to the adjunction $(\mathcal{F}^\Sigma; \mathcal{U}^\Sigma)$. Unfortunately, the objects in $\Lambda_{>0} Seq$ are not necessarily fibrant and the second author proves in [19] the existence of the model category structure for reduced operads without the transfer principle. In the present work, we build an explicit functorial fibrant replacement in both categories ΛSeq and $\Lambda_{>0} Seq$. This resolution will be enhanced in the next sections in order to define Reedy model category structures for (infinitesimal) bimodules using the transfer principle.

Both categories ΣSeq and ΛSeq are obtained as categories of functors from Σ and Λ to topological spaces. So the following model category structures are particular cases of model categories of diagrams. We refer to [5, 22, 25] for a comprehensive study of projective model categories of diagrams over a discrete category (and of dual injective model categories of diagrams), to [11, 33] for a study of projective model categories of diagrams in the enriched setting. For Reedy model structures and applications to simplicial homotopy theory, we refer the reader to [23, 34] and to [1, 4, 9, 36, 37] for generalizations to enriched categories or extended Reedy categories.

1.1 The Projective Model Categories of G-Spaces and of Σ -Sequences

- *The model category of spaces.* In what follows, by spaces we mean compactly generated, but not necessarily Hausdorff, topological spaces. Such spaces are often called k -spaces

[28]. One has a natural *kellyfication functor* from the category of all topological spaces to k -spaces. The topology of mapping spaces, products, subspaces and more generally limits in this category are defined by taking kellyfication of their usual compact-open, product and subspace topologies. The coproducts, quotients and more generally any colimits of k -spaces are automatically k -spaces and kellyfication is not necessary. The category of k -spaces has the advantage of being cartesian closed [31, Theorem 5.5]. This statement implies that the cartesian products distribute over colimits, which is a prerequisite for the theory of operads and operadic objects. Moreover, it implies that the product of two quotient maps (in particular, of a quotient map and an identity one) is again a quotient map in the category of k -spaces [10, Corollary 5.9.10]. We use this observation in our constructions of free objects and pushouts in the category of operads and in the categories of bimodules over operads.

The category of spaces, fixed in the previous paragraph (thus, the category of k -spaces), is denoted by Top and is equipped with the Quillen model category structure (see [28, Theorem 2.4.23]) in which a continuous map is a weak equivalence (respectively, a fibration) if it is a weak homotopy equivalence (respectively, a Serre fibration). According to this definition, all spaces are fibrant and the model category Top is cofibrantly generated. The set of generating cofibrations S_c and the set of generating acyclic cofibrations S_{ac} are the following ones, where S^{-1} denotes the empty set:

$$S_c = \{ S^{n-1} \hookrightarrow D^n, n \geq 0 \} \quad \text{and} \quad S_{ac} = \{ D^n \times \{0\} \hookrightarrow D^n \times [0, 1], n \geq 0 \}.$$

- *Projective model category of G -spaces.* Let G be a topological monoid. The category $G\text{-}Top$ of G -spaces consists of spaces equipped with a right action of G . There is an adjunction $G[-] : Top \rightleftarrows G\text{-}Top : \mathcal{U}$, where \mathcal{U} is the forgetful functor and $G[-]$ is the functor sending a space X to the G -space $G[X] = X \times G$. As a consequence of Theorem 1.1, the category $G\text{-}Top$ inherits a cofibrantly generated model category structure whose sets of generating cofibrations and acyclic cofibrations are $G[S_c]$ and $G[S_{ac}]$, respectively. Indeed, the identity functor provides a fibrant replacement functor while, for any G -space X , the functorial path object is given by the mapping space

$$Path(X) = Map([0, 1], X).$$

Cofibrations and fibrations in this category will be called *G -cofibrations* and *G -fibrations*, respectively. We will be mostly using spaces with a right action of a monoid (or a group). At few occasions we will need to deal with spaces endowed with a left action. Such spaces will be called *left G -spaces* and the category of such will be denoted by $G^{op}\text{-}Top$.

- *Projective model category of Σ -sequences.* Let Σ be the category whose objects are finite sets $[n] = \{1, \dots, n\}$, with $n \geq 0$, and morphisms are bijections between them. By a Σ -sequence, we mean a contravariant functor from Σ to the category of spaces. In practice, a Σ -sequence is given by a family of spaces $X(0), X(1), \dots$ together with an action of the symmetric group: for each permutation $\sigma \in \Sigma_n$, there is a map

$$\begin{aligned} \sigma^* : X(n) &\longrightarrow X(n); \\ x &\longmapsto x \cdot \sigma, \end{aligned} \tag{1}$$

satisfying the relations $(x \cdot \sigma) \cdot \tau = x \cdot (\sigma\tau)$, with $\tau \in \Sigma_n$, and $x \cdot e = x$. A morphism between Σ -sequences is a family of continuous maps that should preserve the right action of the symmetric groups. We denote by ΣSeq the category of Σ -sequences and by $\Sigma_{>0} Seq$ its subcategory composed of Σ -sequences whose arity 0 component is empty.

Given an integer $r \geq 0$, we also consider the category of r -truncated Σ -sequences $T_r \Sigma Seq$ which we define as follows. Let $T_r \Sigma$ be the category with objects $[n] = \{1, \dots, n\}, 0 \leq n \leq r$,

and bijections between them. An r -truncated Σ -sequence is a contravariant functor from $T_r \Sigma$ to the category of spaces. In practice, an r -truncated Σ -sequence is given by a family of spaces $X(0), \dots, X(r)$ together with an action of the corresponding symmetric group Σ_n for each $n \leq r$.

A (possibly truncated) Σ -sequence is said to be *pointed* if there is a distinguished element $*_1 \in X(1)$ called *unit*. It is said *well-pointed* if the inclusion $*_1 \rightarrow X(1)$ is a cofibration. Recall that we deal with the Quillen model structure on topological spaces. Thus, we consider the cofibrations of the Quillen model structure in this definition, not the Hurewicz cofibrations of the usual notion of well-pointed space. Note that a space T is well-pointed in this sense as soon as it is cofibrant (with respect to the Quillen model structure). Indeed, in this case, T occurs as a retract of a (generalized) CW-complex and we may just observe that the inclusion of any choice of base point in a (generalized) CW-complex is a cofibration to conclude that this is the same for T . From this observation, we deduce that a (truncated) Σ -sequence X is well-pointed in the sense of our definition if and only if it is pointed and its component $X(1)$ is cofibrant.

There is an obvious functor called *truncation functor*

$$T_r(-) : \Sigma Seq \longrightarrow T_r \Sigma Seq.$$

One has

$$\Sigma Seq = \prod_{n \geq 0} \Sigma_n\text{-Top}, \quad \Sigma_{>0} Seq = \prod_{n \geq 1} \Sigma_n\text{-Top} \quad \text{and} \quad T_r \Sigma Seq = \prod_{0 \leq n \leq r} \Sigma_n\text{-Top}.$$

Since $\Sigma_n\text{-Top}$ is a cofibrantly generated model category for any $n \geq 0$, the categories ΣSeq , $\Sigma_{>0} Seq$ and $T_r \Sigma Seq$ are endowed with a cofibrantly generated model category structure, called *the projective model category structure*, in which all objects are fibrant. More precisely, a map between (possibly truncated) Σ -sequences is a weak equivalence (respectively, a fibration) if the map is degreewise a weak homotopy equivalence (respectively, a Serre fibration). The sets of generating cofibrations S_c and acyclic cofibrations S_{ac} of ΣSeq (respectively, $\Sigma_{>0} Seq$ and $T_r \Sigma Seq$) are given by

$$S_c = \bigcup_{\substack{n \geq 0 \\ (\text{resp. } n > 0 \text{ and} \\ 0 \leq n \leq r)}} \left(S_c^n \times \prod_{m \neq n} 1_m \right) \quad \text{and} \quad S_{ac} = \bigcup_{\substack{n \geq 0 \\ (\text{resp. } n > 0 \text{ and} \\ 0 \leq n \leq r)}} \left(S_{ac}^n \times \prod_{m \neq n} 1_m \right)$$

where S_c^n and S_{ac}^n are the sets of generating cofibrations and acyclic cofibrations, respectively, of $\Sigma_n\text{-Top}$ while $1_m : \emptyset \rightarrow \emptyset$ is the identity map of the initial object of $\Sigma_m\text{-Top}$ (see [26, Proposition 11.1.10]).

• *Notation for cofibrations.* Let \mathcal{C} be a category together with a functor \mathcal{U} from \mathcal{C} to the category ΣSeq (respectively the categories $\Sigma_{>0} Seq$ and $T_r \Sigma Seq$). In the rest of the paper, an object C in the category \mathcal{C} is said to be Σ -*cofibrant* if the underlying Σ -sequence $\mathcal{U}(C)$ is cofibrant in the projective model category ΣSeq (respectively, $\Sigma_{>0} Seq$ and $T_r \Sigma Seq$). It is called *componentwise cofibrant* if every component $U(C)(n)$ is cofibrant in Top . In case the forgetful functor U factors through the category of pointed (truncated) Σ -sequences, the object C is called *well-pointed* if $U(C)(1)$ is cofibrant. In the following the category \mathcal{C} will be the category of operads ΣOperad , of reduced operads $\Lambda_*\text{Operad}$, of bimodules $\Sigma\text{Bimod}_{P; Q} = \Lambda\text{Bimod}_{P; Q}$, of reduced bimodules $\Lambda_*\text{Bimod}_{P; Q}$, or of infinitesimal bimodules $\Sigma\text{Ibimod}_O = \Lambda\text{Ibimod}_O$.

1.2 The Reedy Model Categories of Λ - and $\Lambda_{>0}$ -Sequences

• *The category of Λ -sequences.* We refer the reader to [18, 19] for a detailed account on the categories introduced in this subsection. Let Λ be the category whose objects are finite sets $[n] = \{1, \dots, n\}$, with $n \geq 0$, and morphisms are injective maps between them. (Hence, Σ is the subcategory of isomorphisms of Λ .) By a Λ -sequence, we understand a contravariant functor from Λ to spaces and we denote the corresponding category by ΛSeq . In practice, such an object is given by a Σ -sequence $X(0), X(1), \dots$ together with maps generated by applications of the form

$$s_i^* : X(n) \longrightarrow X(n - 1), \quad \text{with } 1 \leq i \leq n, \tag{2}$$

associated to the injective maps

$$s_i : [n - 1] \longrightarrow [n] ; \ell \longmapsto \begin{cases} \ell & \text{if } \ell < i, \\ \ell + 1 & \text{if } \ell \geq i. \end{cases}$$

Given an integer $r \geq 0$, we also consider the full subcategory $T_r \Lambda$ whose objects are families of finite sets $[n] = \{1, \dots, n\}$, with $0 \leq n \leq r$. An r -truncated Λ -sequence is a contravariant functor from $T_r \Lambda$ to spaces and we denote by $T_r \Lambda Seq$ the associated category. We will also be using the categories $\Lambda_{>0}$ and $T_r \Lambda_{>0}$ which are full subcategories of non-empty objects of Λ and $T_r \Lambda$. The categories $\Lambda_{>0} Seq$ and $T_r \Lambda_{>0} Seq$ are similarly defined. There exist obvious truncation functors

$$T_r(-) : \Lambda Seq \longrightarrow T_r \Lambda Seq \quad \text{and} \quad T_r(-) : \Lambda_{>0} Seq \longrightarrow T_r \Lambda_{>0} Seq.$$

• *Useful adjunctions.* The inclusions of categories $\Sigma \subset \Lambda$ and $T_r \Sigma \subset T_r \Lambda$ induce adjunctions between the categories of (possibly truncated) Σ -sequences and Λ -sequences

$$\Lambda[-] : \Sigma Seq \rightleftarrows \Lambda Seq : \mathcal{U} \quad \text{and} \quad \Lambda_r[-] : T_r \Sigma Seq \rightleftarrows T_r \Lambda Seq : \mathcal{U}$$

where \mathcal{U} is the obvious forgetful functor, which forgets about the operations generated by (2). The functor $\Lambda[-]$ sends a Σ -sequence X to the Λ -sequence $\Lambda[X]$ given by

$$\Lambda[X](n) := \coprod_{\substack{\Lambda_+([n];[m]) \\ m \geq n}} X(m), \quad \text{for all } n \geq 0,$$

where Λ_+ is the subcategory of order preserving injective maps. A point in $\Lambda[X](n)$ is denoted by $(h; x)$ with $h : [n] \rightarrow [m]$ an order preserving injective map and $x \in X(m)$. For any permutation $\sigma \in \Sigma_n$ and any order preserving injective map $h : [n] \rightarrow [m]$, we denote by $\sigma_h \in \Sigma_m$ the permutation

$$\sigma_h(i) := \begin{cases} h(\sigma(j)), & \text{if } h(j) = i, \\ i, & \text{otherwise.} \end{cases}$$

According to this notation, the Λ -structure on $\Lambda[X]$ is given by the following formulas:

$$\begin{aligned} \sigma^* : \Lambda[X](n) &\longrightarrow \Lambda[X](n) && ; (h; x) \longmapsto (h; x \cdot \sigma_h), \\ s_i^* : \Lambda[X](n) &\longrightarrow \Lambda[X](n - 1) && ; (h; x) \longmapsto (h \circ s_i; x). \end{aligned}$$

• *The matching object.* For a (possibly truncated) Λ -sequence X , the *matching object* of X , denoted by $\mathcal{M}(X)$, is the (possibly truncated) Σ -sequence defined as follows:

$$\mathcal{M}(X)(n) = \lim_{\substack{h \in \Lambda_+([n];[l]) \\ \ell < n}} X(\ell). \tag{3}$$

Let $\sigma \in \Sigma_n$ be a permutation and $h : [\ell] \rightarrow [n]$ be an order preserving inclusion. We denote by $h \cdot \sigma : [\ell] \rightarrow [n]$ the unique order preserving inclusion whose image is $\text{Im}(\sigma \circ h)$ and $\sigma[h] \in \Sigma_\ell$ is the unique permutation satisfying $\sigma[h](i) < \sigma[h](j)$ if and only if $\sigma(h(i)) < \sigma(h(j))$. According to this notation, the action of the symmetric group on the matching object is the following one:

$$\begin{aligned} \sigma^* : \mathcal{M}(X)(n) &\longrightarrow \mathcal{M}(X)(n); \\ x = \{x_h\}_h &\longmapsto \sigma^*(x) = \{x_{h \cdot \sigma} \cdot \sigma[h]\}_h. \end{aligned}$$

• *The Reedy model category of Λ -sequences.* According to [19, Theorem 8.3.19], the categories ΛSeq , $\Lambda_{>0} \text{Seq}$, $T_r \Lambda \text{Seq}$ and $T_r \Lambda_{>0} \text{Seq}$ are endowed with cofibrantly generated model category structures in which weak equivalences are objectwise weak homotopy equivalences. A morphism $f : X \rightarrow Y$ is a fibration if the corresponding maps

$$X(n) \longrightarrow \mathcal{M}(X)(n) \times_{\mathcal{M}(Y)(n)} Y(n),$$

whenever defined, are Serre fibrations. The set of generating cofibrations (respectively the set of generating acyclic cofibrations) consists of maps of the form

$$(\Lambda[f], \iota) : \Lambda[X] \coprod_{\partial \Lambda[X]} \partial \Lambda[Y] \longrightarrow \Lambda[Y],$$

where $f : X \rightarrow Y$ is a generating cofibration (respectively a generating acyclic cofibration) in the projective model category of Σ -sequences. The map $\iota : \partial \Lambda[Y] \rightarrow \Lambda[Y]$ is the inclusion where $\partial \Lambda[-]$ is the functor from Σ -sequences to Λ -sequences left adjoint to the matching object functor \mathcal{M} and expressed by the formula

$$\partial \Lambda[Y](n) = \text{colim}_{\substack{h \in \Lambda_+([n]; [\ell]) \\ \ell > n}} \Lambda[Y](\ell) = \coprod_{\substack{\Lambda_+([n]; [\ell]) \\ \ell > n}} Y(\ell).$$

Remark 1.2 The truncation functors

$$T_r : \Lambda \text{Seq} \rightarrow T_r \Lambda \text{Seq}; \quad T_r : \Lambda_{>0} \text{Seq} \rightarrow T_r \Lambda_{>0} \text{Seq}; \quad T_r : T_{r'} \Lambda \text{Seq} \rightarrow T_r \Lambda \text{Seq}, \quad r' > r,$$

preserve fibrations, cofibrations and weak equivalences.

For fibrations and weak equivalences, the statement follows from definition. For cofibrations we recall [19, Theorem 8.3.20] that a morphism in the Reedy model structure is a cofibration if and only if it is a projective cofibration in the corresponding category of (truncated) Σ -sequences.

1.3 A Fibrant Replacement Functor for Λ - and $\Lambda_{>0}$ -Sequences

As explained previously, in order to apply the transfer principle, described in Theorem 1.1, we need a functorial fibrant replacement. Unfortunately, the objects in ΛSeq and $\Lambda_{>0} \text{Seq}$ are not necessarily fibrant and therefore the identity functors can not be regarded as a fibrant replacement one. To solve this problem, we build explicit and functorial fibrant coresolutions

$$(-)^f : \Lambda \text{Seq} \longrightarrow \Lambda \text{Seq} \quad \text{and} \quad (-)^f : \Lambda_{>0} \text{Seq} \longrightarrow \Lambda_{>0} \text{Seq}.$$

For this purpose, we need some notation. For any map $h \in \Lambda_+([\ell]; [n])$, we denote by $h_c \in \Lambda_+([n - \ell]; [n])$ its complementary map which is the unique order preserving inclusion so that $\text{Im}(h) \cap \text{Im}(h_c) = \emptyset$. Then, for any pair of order preserving inclusions of the form

$s_i : [\ell] \rightarrow [\ell + 1]$ and $h : [\ell + 1] \rightarrow [n]$, we denote by $\varepsilon_{h;i} \in [n - \ell]$ the unique index such that the following diagram commute:

$$\begin{array}{ccc}
 [n - \ell - 1] & \xrightarrow{s_{\varepsilon_{h;i}}} & [n - \ell] \\
 & \searrow h_c & \swarrow (h \circ s_i)_c \\
 & & [n]
 \end{array}$$

Let X be a Λ -sequence. The space $X^f(n)$ is the subspace

$$X^f(n) \subset \prod_{\substack{\Lambda_+([\ell], [n]) \\ \ell \leq n}} \text{Map}([0, 1]^{n-\ell}; X(\ell)),$$

consisting of families of maps $\{f_h\}_{h \in \Lambda_+([\ell], [n])}$ such that

$$\begin{array}{ccc}
 [0, 1]^{n-\ell-1} & \xrightarrow{\tau_1[\varepsilon_{h;i}]} & [0, 1]^{n-\ell} \\
 \downarrow f_h & & \downarrow f_{h \circ s_i} \\
 X(\ell + 1) & \xrightarrow{s_i^*} & X(\ell)
 \end{array} \tag{4}$$

where $\tau_t[k] : [0, 1]^{n-\ell-1} \rightarrow [0, 1]^{n-\ell}$, with $t \in [0, 1]$ and $k \in [n - \ell - 1]$, inserts t at the k -th position:

$$\tau_t[k](t_1, \dots, t_{n-\ell-1}) = (t'_1, \dots, t'_{n-\ell}) \quad \text{with} \quad t'_j = \begin{cases} t_j & \text{if } j < k, \\ t & \text{if } j = k, \\ t_{j-1} & \text{if } j > k. \end{cases}$$

• *The Λ -structure on X^f .* In order to describe the Λ -structure we consider the following notation. For any order preserving inclusions $s_i : [n - 1] \rightarrow [n]$ and $h : [\ell] \rightarrow [n - 1]$, we denote by $\vartheta_{h;i}$ the unique index such that the following diagram commute:

$$\begin{array}{ccc}
 [n - \ell - 1] & \xrightarrow{h_c} & [n - 1] \\
 \downarrow s_{\vartheta_{h;i}} & & \downarrow s_i \\
 [n - \ell] & \xrightarrow{(s_i \circ h)_c} & [n]
 \end{array}$$

According to this notation, the Λ -structure operations are given by

$$\begin{aligned}
 s_i^* : X^f(n) &\longrightarrow X^f(n - 1) ; \{f_h\}_{h \in \Lambda_+([\ell], [n])} \longmapsto \{(s_i^* \circ f)_h\}_{h \in \Lambda_+([\ell], [n-1])}, \\
 \sigma^* : X^f(n) &\longrightarrow X^f(n) \quad ; \{f_h\}_{h \in \Lambda_+([\ell], [n])} \longmapsto \{(f \cdot \sigma)_h\}_{h \in \Lambda_+([\ell], [n])},
 \end{aligned}$$

where the continuous maps $(s_i^* \circ f)_h$ and $(f \cdot \sigma)_h$ are the following ones:

$$\begin{aligned}
 (s_i^* \circ f)_h : [0, 1]^{n-\ell-1} &\longrightarrow X(\ell) ; (t_1, \dots, t_{n-\ell-1}) \longmapsto f_{s_i \circ h}(\tau_0[\vartheta_{h;i}](t_1, \dots, t_{n-\ell-1})), \\
 (f \cdot \sigma)_h : [0, 1]^{n-\ell} &\longrightarrow X(\ell) ; (t_1, \dots, t_{n-\ell}) \longmapsto f_{h \cdot \sigma}(\sigma_{[h_c]^{-1}(1)}, \dots, \sigma_{[h_c]^{-1}(n-\ell)}) \cdot \sigma[h].
 \end{aligned} \tag{5}$$

• *The fibrant replacement functor:* The Λ -sequence X^f is obviously functorial along the Λ -sequence X . Furthermore, there is a map $\varphi : X \rightarrow X^f$ sending a point $x \in X(n)$ to the family of constant maps $\{\varphi(x)_h\}_{h \in \Lambda_+([\ell], [n])}$ obtained using the Λ -structure of X :

$$\begin{aligned} \varphi(x)_h : \quad [0, 1]^{n-\ell} &\longrightarrow X(\ell); \\ (t_1, \dots, t_{n-\ell}) &\longmapsto h^*(x). \end{aligned}$$

The map is well defined and preserves the Λ -structures. Indeed, one has the following equalities:

$$\begin{aligned} s_i^*(\varphi(x)) &= s_i^* \left(\left\{ \begin{array}{l} \varphi(x)_h : [0, 1]^{n-\ell} \longrightarrow X(\ell); \\ (t_1, \dots, t_{n-\ell}) \longmapsto h^*(x). \end{array} \right\}_{h \in \Lambda_+([\ell], [n])} \right) \\ &= \left\{ \begin{array}{l} s_i^* \circ \varphi(x)_h : [0, 1]^{n-\ell-1} \longrightarrow X(\ell); \\ (t_1, \dots, t_{n-\ell-1}) \longmapsto \varphi(x)_{s_i \circ h}(t_1, \dots, t_{\partial_{h:i}-1}, 0, t_{\partial_{h:i}}, \dots, t_{n-\ell-1}). \end{array} \right\}_{h \in \Lambda_+([\ell], [n-1])} \\ \varphi(s_i^*(x)) &= \left\{ \begin{array}{l} \varphi(x)_h : [0, 1]^{n-\ell-1} \longrightarrow X(\ell); \\ (t_1, \dots, t_{n-\ell-1}) \longmapsto h^*(s_i^*(x)). \end{array} \right\}_{h \in \Lambda_+([\ell], [n-1])} \end{aligned}$$

Proposition 1.3 *The map $\phi : X \rightarrow X^f$ is a weak equivalence of Λ -sequences.*

Proof More precisely, we show that the map of Λ -sequences $\varphi_n : X(n) \rightarrow X^f(n)$ is a homotopy equivalence of Σ -sequences. For this purpose, we introduce a map of Σ -sequences (which is not a map of Λ -sequences) $\psi : X^f \rightarrow X$ given by

$$\begin{aligned} \psi_n : \quad X^f(n) &\longrightarrow X(n); \\ \{f_h\}_{h \in \Lambda_+([\ell], [n])} &\longmapsto f_{[n] \rightarrow [n]}(*), \end{aligned}$$

which makes φ into a deformation retract. The homotopy consists in bringing the parameters to 1:

$$\begin{aligned} H_n : [0, 1] \times X^f(n) &\longrightarrow X^f(n); \\ t ; \{f_h\} &\longmapsto \{H_n(t ; f_h)\}, \end{aligned}$$

with $H_n(t ; f_h)(t_1, \dots, t_{n-\ell}) = f_h((1-t)t_1 + t, \dots, (1-t)t_{n-\ell} + t)$. □

Proposition 1.4 *The Λ -sequence X^f is Reedy fibrant.*

Lemma 1.5 (Fiber product version of Reedy’s patching lemma, see Lemma 1.3 in [35]) *For any commutative diagram of spaces of the form*

$$\begin{array}{ccccc} A & \xrightarrow{f} & B & \xleftarrow{g} & C \\ v_A \downarrow & & v_B \downarrow & & v_C \downarrow \\ A' & \xrightarrow{f'} & B' & \xleftarrow{g'} & C' \end{array}$$

the induced map between the limits of the horizontal diagrams

$$v : \lim(A \rightarrow B \leftarrow C) \longrightarrow \lim(A' \rightarrow B' \leftarrow C')$$

is a fibration if the map v_C as well as the map

$$(v_A; f) : A \longrightarrow B \times_{B'} A' \tag{6}$$

are fibrations. Furthermore, the map v is an acyclic fibration if the map v_C and (6) are acyclic fibrations.

Proof of Proposition 1.4 We show that the map from $X^f(n)$ to the matching object $\mathcal{M}(X^f)(n)$ is a Serre fibration. Let us remark that the spaces $X^f(n)$ and $M(X^f)(n)$ can be expressed in terms of pullback diagrams. More precisely, one has

$$X^f(n) = \lim \left(A \longrightarrow \prod_{s_i: [n-1] \rightarrow [n]} X(n-1) \longleftarrow X(n) \right), \quad \text{where}$$

$$A \subset \prod_{\substack{\Lambda_+([\ell], [n]) \\ \ell < n}} \text{Map}([0, 1]^{n-\ell}; X(\ell))$$

is the subspace satisfying the condition (4). The map from A to the product $\prod_{s_i} X(n-1)$ sends a family of maps $\{f_h\}$ to the family of points $\{f_{s_i}(1)\}$ by taking the evaluation at the point 1. Furthermore, one has the commutative diagram

$$\begin{array}{ccc} A & \longrightarrow & \prod_{s_i: [n-1] \rightarrow [n]} X(n-1) \longleftarrow X(n) \\ \downarrow & & \downarrow \qquad \qquad \downarrow \\ \mathcal{M}(X^f)(n) & \longrightarrow & * \longleftarrow * \end{array}$$

According to Lemma 1.5 and since $X(n)$ is fibrant, we only need to check that the map

$$A \longrightarrow A' = \mathcal{M}(X^f)(n) \times \prod_{s_i: [n-1] \rightarrow [n]} X(n-1). \tag{7}$$

is a Serre fibration. In other words, if we denote by $\partial'[0, 1]^{n-\ell}$ the subspace of $[0, 1]^{n-\ell}$ composed of the point $(1, \dots, 1)$ and the points having at least one coordinate equal to 0, then A' is the following subspace satisfying the relation (4):

$$A' \subset \prod_{\substack{\Lambda_+([\ell], [n]) \\ \ell < n}} \text{Map}(\partial'[0, 1]^{n-\ell}; X(\ell)).$$

Let us notice that the inclusion from $\partial'[0, 1]^{n-\ell}$ into $[0, 1]^{n-\ell}$ is a cofibration as an inclusion of CW-complexes. Unfortunately, we can not deduce directly the result due to condition (4). To solve this problem, we introduce a cofiltration of the map (7) according to the dimension of the cubes. Let us consider the following subspaces:

$$A_k \subset \prod_{\substack{\Lambda_+([\ell], [n]) \\ n-k \leq \ell < n}} \text{Map}([0, 1]^{n-\ell}; X(\ell)) \quad \text{and}$$

$$A'_k \subset \prod_{\substack{\Lambda_+([\ell], [n]) \\ n-k \leq \ell < n}} \text{Map}(\partial'[0, 1]^{n-\ell}; X(\ell))$$

satisfying condition (4). In particular, one has $A_n = A$ and $A'_n = A'$. Furthermore, the spaces A_k and A'_k can be obtained from A_{k-1} and A'_{k-1} , respectively, using the following pullback diagrams:

$$\begin{array}{ccc}
 A_k \longrightarrow \prod_{[n-k] \rightarrow [n]} \text{Map}([0, 1]^k; X(n-k)) & & A'_k \longrightarrow \prod_{[n-k] \rightarrow [n]} \text{Map}(\partial'[0, 1]^k; X(n-k)) \\
 \downarrow & & \downarrow \\
 A_{k-1} \rightrightarrows \prod_{[n-k] \rightarrow [n-k+1] \rightarrow [n]} \text{Map}([0, 1]^{k-1}; X(n-k)) & & A'_{k-1} \rightrightarrows \prod_{[n-k] \rightarrow [n-k+1] \rightarrow [n]} \text{Map}(\partial'[0, 1]^{k-1}; X(n-k))
 \end{array}$$

We prove by induction that the maps $A_k \rightarrow A'_k$ are Serre fibrations. First, the map

$$A_1 = \prod_{[n-1] \rightarrow [n]} \text{Map}([0, 1]; X(n)) \longrightarrow \prod_{[n-1] \rightarrow [n]} \text{Map}(\partial[0, 1]; X(n)) = A'_1$$

is obviously a Serre fibration since the inclusion from $\partial[0, 1] = \{0, 1\}$ into the interval $[0, 1]$ is a cofibration. From now on, we assume that the map $A_{k-1} \rightarrow A'_{k-1}$ is a Serre fibration. Then we consider the commutative diagram

$$\begin{array}{ccccc}
 \prod_{[n-k] \rightarrow [n]} \text{Map}([0, 1]^k; X(n-k)) & \longrightarrow & \prod_{[n-k] \rightarrow [n-k+1] \rightarrow [n]} \text{Map}([0, 1]^{k-1}; X(n-k)) & \longleftarrow & A_{k-1} \\
 \downarrow & & \downarrow & & \downarrow \\
 \prod_{[n-k] \rightarrow [n]} \text{Map}(\partial'[0, 1]^k; X(n-k)) & \longrightarrow & \prod_{[n-k] \rightarrow [n-k+1] \rightarrow [n]} \text{Map}(\partial'[0, 1]^{k-1}; X(n-k)) & \longleftarrow & A'_{k-1}
 \end{array}$$

According to Lemma 1.5, one has to check that the map from the space

$$\prod_{[n-k] \rightarrow [n]} \text{Map}([0, 1]^k; X(n-k))$$

to the limit of the diagram

$$\begin{array}{ccc}
 & & \prod_{[n-k] \rightarrow [n-k+1] \rightarrow [n]} \text{Map}([0, 1]^{k-1}; X(n-k)) \\
 & & \downarrow \\
 \prod_{[n-k] \rightarrow [n]} \text{Map}(\partial'[0, 1]^k; X(n-k)) & \longrightarrow & \prod_{[n-k] \rightarrow [n-k+1] \rightarrow [n]} \text{Map}(\partial'[0, 1]^{k-1}; X(n-k))
 \end{array}$$

is a Serre fibration. The limit corresponds to the space

$$\prod_{[n-k] \rightarrow [n]} \text{Map}(\partial[0, 1]^k; X(n-k))$$

and the map

$$\prod_{[n-k] \rightarrow [n]} \text{Map}([0, 1]^k; X(n-k)) \longrightarrow \prod_{[n-k] \rightarrow [n]} \text{Map}(\partial[0, 1]^k; X(n-k))$$

is obviously a Serre fibration since the inclusion from $\partial[0, 1]^k$ to $[0, 1]^k$ is a cofibration. \square

Remark 1.6 The same strategy can be used in order to get a fibrant replacement functor for r -truncated Λ -sequences. In that case, we only need to restrict our construction to order preserving inclusions $h : [\ell] \rightarrow [n]$ with $n \leq r$. Similarly, we get fibrant replacement functors for the categories $\Lambda_{>0}Seq$ and $T_r \Lambda_{>0}Seq$.

1.4 The Projective/Reedy Model Category of Operads

An operad is a pointed Σ -sequence O together with operations called *operadic compositions*

$$\circ_i : O(n) \times O(m) \longrightarrow O(n + m - 1), \quad \text{with } 1 \leq i \leq n. \tag{8}$$

These operations satisfy associativity and unit axioms as well as compatibility relations with the symmetric group action. More precisely, for any integers $i \in \{1, \dots, n\}$, $j \in \{i + 1, \dots, n\}$, $k \in \{1, \dots, m\}$ and any permutations $\sigma \in \Sigma_n$ and $\tau \in \Sigma_m$, one has the following commutative diagrams:

$$\begin{array}{ccc} O(n) \times O(m) \times O(\ell) & \xrightarrow{\circ_i \times id} & O(n + m - 1) \times O(\ell) \\ id \times \circ_k \downarrow & & \downarrow \circ_{i+k-1} \\ O(n) \times O(m + \ell - 1) & \xrightarrow{\circ_i} & O(n + m + \ell - 2) \end{array}$$

Linear associativity axiom

$$\begin{array}{ccc} O(n) \times O(m) \times O(\ell) & \xrightarrow{\circ_i \times id} & O(n + m - 1) \times O(\ell) \\ \circ_j \times id \downarrow & & \downarrow \circ_{i+m-1} \\ O(n + \ell - 1) \times O(m) & \xrightarrow{\circ_i} & O(n + m + \ell - 2) \end{array}$$

Ramified associativity axiom

$$\begin{array}{ccc} O(n) \times O(1) & \xleftarrow{id \times *1} & O(n) & \xrightarrow{*1 \times id} & O(1) \times O(n) \\ & \searrow \circ_i & \parallel & \swarrow \circ_1 & \\ & & O(n) & & \end{array}$$

Unit axiom

$$\begin{array}{ccc} O(n) \times O(m) & \xrightarrow{\circ_i} & O(n + m - 1) \\ \sigma^* \times \tau^* \downarrow & & \downarrow (\sigma \circ_{\sigma(i)} \tau)^* \\ O(n) \times O(m) & \xrightarrow{\circ_{\sigma(i)}} & O(n + m - 1) \end{array}$$

Compatibility with the symmetric group action

where the permutation $\sigma \circ_{\sigma(i)} \tau$ is obtained from the well known operadic compositions on the symmetric groups (see [18, Proposition 1.1.9]).

A map between operads should preserve the operadic compositions. We denote by ΣOperad the category of topological operads. The category of operads is obviously endowed with a forgetful functor to the category of Σ -sequences by forgetting the operadic composition (8):

$$U^\Sigma : \Sigma\text{Operad} \longrightarrow \Sigma Seq. \tag{9}$$

• *The category of reduced operads and their underlying Λ -structures.* An operad O is said to be *reduced* if $O(0)$ is the one point topological space. This point is denoted by $*_0$. We denote by $\Lambda_*\text{Operad}$ the category of reduced operads. This category is equipped with a forgetful functor to the category of $\Lambda_{>0}$ -sequences, which consists in forgetting the arity zero component and the operadic compositions (8) for $m \geq 1$:

$$U^\Lambda : \Lambda_*\text{Operad} \longrightarrow \Lambda_{>0}Seq. \tag{10}$$

(Note also that the category of $\Lambda_{>0}$ -sequences is equivalent to the category of reduced Λ -sequences, i.e Λ -sequences X so that $X(0) = *$.) Indeed, if O is a reduced operad, then the $\Lambda_{>0}$ -structure on $U^\Lambda(O)$ is generated by the operations of the form

$$\begin{aligned} s_i^* : U^\Lambda(O)(n) = O(n) &\longrightarrow U^\Lambda(O)(n - 1) = O(n - 1); \\ \theta &\longmapsto \theta \circ_i *_0. \end{aligned}$$

- *The model category of algebras over an operad.* An algebra over a (possibly reduced) operad O , or O -algebra, is a topological space X together with operations of the form

$$\alpha_n : O(n) \times X^{\times n} \longrightarrow X, \quad \text{with } n \geq 0,$$

compatible with the operadic structure (see [18, Section 1.1.13 and Figure 1.9]). The category of O -algebras is denoted by Alg_O . It has been proved in [8, Theorem 2.1] that the category of algebras over any operad O in Top inherits a cofibrantly generated model category structure by using the transfer principle applied to the adjunction $\mathcal{F}_O : Top \rightleftarrows Alg_O : \mathcal{U}$ where the free algebra functor \mathcal{F}_O is the left adjoint to the forgetful functor \mathcal{U} .

- *The projective and Reedy model categories of operads and reduced operads.* Both forgetful functors (9) and (10) have left adjoints, which we respectively denote by \mathcal{F}^Σ and \mathcal{F}^Λ , and which, given a Σ - or a $\Lambda_{>0}$ -sequence X , produce the free operad generated by X . Explicitly, elements of $\mathcal{F}^\Sigma(X)$ and $\mathcal{F}^\Lambda(X)$ are described as rooted trees (without univalent vertices for reduced operads) with internal vertices labelled by elements of the sequence. We refer the reader to [18] for a detailed account on these adjunctions:

$$\mathcal{F}^\Sigma : \Sigma Seq \rightleftarrows \Sigma Operad : \mathcal{U}^\Sigma \quad \text{and} \quad \mathcal{F}^\Lambda : \Lambda_{>0} Seq \rightleftarrows \Lambda_* Operad : \mathcal{U}^\Lambda.$$

As a consequence of the transfer principle 1.1, the category $\Sigma Operad$ inherits a cofibrantly generated model category structure which we call the *projective* model structure (we refer the reader to [6] for more details). Similarly to the usual category of Λ -sequences, the category $\Lambda_{>0} Seq$ is also endowed with a (cofibrantly generated) Reedy model category structure. The second author in [19] proves that $\Lambda_* Operad$ has a cofibrantly generated model category structure called the *Reedy* model structure. In both cases, a map of (possibly reduced) operads $f : P \rightarrow Q$ is a weak equivalence (respectively, a fibration) if the map $\mathcal{U}^\Sigma(f)$ or $\mathcal{U}^\Lambda(f)$ is a weak equivalence (respectively, a fibration) in the appropriate category.

Theorem 1.7 *The projective and the Reedy model structures have the following properties:*

- ▶ [6, Section 2.5]: All operads are fibrant in $\Sigma Operad$.
- ▶ [27, Theorem 3.1.10]: The category of operads (respectively of reduced operads) is left proper (see Sect. 2.2.1) relative to the class of Σ -cofibrant operads (respectively of reduced Σ -cofibrant operads).
- ▶ [19, Theorem 8.4.12]: A map of reduced operads $\phi : P \rightarrow Q$ is a cofibration in $\Lambda_* Operad$ if and only if the corresponding map $\phi_{>0} : P_{>0} \rightarrow Q_{>0}$ is a cofibration in $\Sigma Operad$ where $P_{>0}$ and $Q_{>0}$ are the sub-operads obtained from P and Q , respectively, by redefining the arity zero components to be empty.
- ▶ [21, Theorem 1]: If P and Q are reduced operads, then one has a weak equivalence between the derived mapping spaces

$$\Sigma Operad^h(P; Q) \simeq \Lambda_* Operad^h(P; Q).$$

- ▶ [6, Theorem 4.4], [20, Theorem 15.A]: If $\phi : P \rightarrow Q$ is a weak equivalence between Σ -cofibrant operads, then the extension $\phi_!$ and restriction ϕ^* functors (see Sect. 2.2.2) form a Quillen equivalence

$$\phi_! : Alg_P \rightleftarrows Alg_Q : \phi^*.$$

1.5 Properties of Cofibrant Operads

The results of this section are used in Sect. 2.2.1 in order to prove that the category of bimodules is relatively left proper.

Proposition 1.8 *If P is a cofibrant operad, then its suboperad $P_{>0}$, obtained by forgetting its arity zero component, is also cofibrant.*

Proof Since an operad is cofibrant if and only if it is a retract of a cellular one, we can assume that P is cellular: $P = \text{colim}_{\alpha < \lambda} P_\alpha$, where P_α is obtained from $P_{<\alpha} := \text{colim}_{\beta < \alpha} P_\beta$ using the pushout

$$\begin{array}{ccc}
 \mathcal{F}(\partial X_\alpha) & \longrightarrow & \mathcal{F}(X_\alpha) \\
 \downarrow & & \downarrow \\
 P_{<\alpha} & \longrightarrow & P_\alpha,
 \end{array}
 \tag{11}$$

where $\mathcal{F}(-)$ is the free operadic functor (see [6] for a combinatorial description of \mathcal{F} in terms of trees) and each $\partial X_\alpha \rightarrow X_\alpha$ is a generating Σ -cofibration. We need to show that $P_{>0}$ is also cellular. Note that as an operad in sets, P is a free operad generated by the Σ -sequence $X = \coprod_{\alpha \in \lambda} X_\alpha \setminus \partial X_\alpha$. We claim that $P_{>0}$ is also free as an operad in sets being generated by its Σ -subsequence represented by trees, whose vertices are labelled by X , with the property that only their root vertex can have leaves and in fact must have at least one leaf attached, see Fig. 1.

This set of generating trees splits into cells of $P_{>0}$, namely by the way from which X_α 's, $\alpha \in \lambda$, the labels come from. So the set of cells can be described as the set of trees as above with vertices labelled by elements α 's from λ . One can define a total ordering of this set that prescribes in which order the new cells are attached. Given two such trees, we compare first the maximal elements from λ they have as labels (including their arity zero vertices). If their maximal elements are the same, we compare which one has more such maximal labels. If the numbers of such labels are the same, we compare their next to maximal labels. And so on. If they have exactly the same sets of labels, we put any random order between them, or we put them together in one bigger cell. \square

Let O be an operad. By $\Sigma_k \wr O(1)$, we understand the monoid that acts on $O(k)$ and is given by the following extension

$$1 \longrightarrow O(1)^{\times k} \longrightarrow \Sigma_k \wr O(1) \longrightarrow \Sigma_k \longrightarrow 1.$$




Fig. 1 An element in a free operad and its decomposition in the new positive arity generators

Proposition 1.9 *If O is a cofibrant operad, then each space $O(n)$ is $\Sigma_n \wr O(1)$ -cofibrant.*

For each $1 < k < n$, we denote by $O(n; k)$ the subspace of $O(n)$ which consists of points of the form

$$(\theta_1 \circ_1 \theta_2) \cdot \sigma,$$

with $\theta_2 \in O(i), 2 \leq i \leq k, \theta_1 \in O(n - i + 1)$ while σ preserves the position of $k + 1, \dots, n$ and shuffles $\{1, \dots, i\}$ with $\{i + 1, \dots, k\}$. Both spaces $O(n)$ and $O(n; k)$ inherit an action of the monoid $(\Sigma_k \times \Sigma_{n-k}) \wr O(1) := (\Sigma_k \wr O(1)) \times (\Sigma_{n-k} \wr O(1))$ and the inclusion from $O(n; k)$ into $O(n)$ is a $(\Sigma_k \times \Sigma_{n-k}) \wr O(1)$ -equivariant map.

Proposition 1.10 *If O is a cofibrant operad, then $O(n; k)$ is $(\Sigma_k \times \Sigma_{n-k}) \wr O(1)$ -cofibrant and the map*

$$O(n; k) \rightarrow O(n) \tag{12}$$

is a $(\Sigma_k \times \Sigma_{n-k}) \wr O(1)$ -cofibration. As a consequence, $O(n; k)$ is $(\Sigma_k \wr O(1)) \times \Sigma_{n-k}$ -cofibrant and the map (12) is a $(\Sigma_k \wr O(1)) \times \Sigma_{n-k}$ -cofibration.

These propositions are proved by similar arguments which are both adaptation of Berger-Moerdijk’s proof of [6, Proposition 4.3] stating that cofibrant operads are always Σ -cofibrant. The latter result is obtained by iteratively using [6, Lemma 5.10]. Its slightly stronger version [7, Lemma 2.5.3] is Lemma A.1, both being equivariant pushout-product type statements with respect to a discrete group action. In the “Appendix” we formulate and prove their analogue – Lemma A.5 for topological monoids that applies to our case of action by monoids $\Sigma_n \wr O(1), (\Sigma_k \times \Sigma_{n-k}) \wr O(1), (\Sigma_k \wr O(1)) \times \Sigma_{n-k}$ or alike.

Proof of Proposition 1.9 Let O be a cofibrant operad. By Proposition 1.8, without loss of generality, we can assume that $O(0) = \emptyset$. Since O is cofibrant, this operad is a retract of a cellular operad P . In what follows, we denote by P_1 the sub-operad of P obtained by taking the restriction to the arity 1:

$$P_1(n) = \begin{cases} P(1) & \text{if } n = 1, \\ \emptyset & \text{otherwise.} \end{cases}$$

In the absence of arity zero operations, the cellular attachments can be reordered so that first we attach arity one cells, then arity two cells and so on. In particular, the map of operads $P_1 \rightarrow P$ can also be seen as a cellular extension:

$$P_1 \longrightarrow P_2 \longrightarrow \dots \longrightarrow P_\alpha \longrightarrow P_{\alpha+1} \longrightarrow \dots \longrightarrow P, \tag{13}$$

where $P_\alpha, \alpha > 1$, is obtained from $P_{<\alpha} := \text{colim}_{\beta < \alpha} P_\beta$ using a pushout of the form (11), where each $\partial X_\alpha \rightarrow X_\alpha$ is a generating Σ -cofibration concentrated in arity $ar_\alpha \geq 2$.

For $n \geq 2$, one has $P_1(n) = \emptyset$, which is $\Sigma_n \wr P(1)$ -cofibrant. In what follows, we will show that each map $P_{<\alpha}(n) \rightarrow P_\alpha(n)$ is a $\Sigma_n \wr P(1)$ -cofibration. For this purpose, we need a combinatorial description of the pushout (11) using the language of trees. Let $\mathbb{P}_n^{\geq 1}$ be the set of planar rooted trees having exactly n leaves indexed by an element of the symmetric group Σ_n with internal vertices of arity ≥ 1 . According to this notation, $P_\alpha(n)$ is obtained from the set of trees $\mathbb{P}_n^{\geq 1}$ by indexing the vertices by points in $P_{<\alpha}(n)$ and X_α . More precisely, one has

$$P_\alpha(n) = \coprod_{T \in \mathbb{P}_n^{\geq 1}} \prod_{v \in V(T)} \left[P_{<\alpha}(|v|) \prod_{\partial X_\alpha(|v|)} X_\alpha(|v|) \right] / \sim, \tag{14}$$

where the equivalence relation is generated by the relation contracting two consecutive vertices indexed by points in $P_{<\alpha}(n)$ using its operadic structure, the compatibility with the symmetric group action and the removal of vertices indexed by the unit $*_1 \in P_{<\alpha}(1) = P(1)$. Let us remark that the arity one vertices are necessarily indexed by points in $P(1)$.

We equip $P_\alpha(n)$ with the filtration (17) given by the number of vertices indexed by X_α . Let $\mathbb{T}_n^{\geq 2}[m]$, $n \geq 2$ and $m \geq 1$, be the set of non-planar rooted trees with internal vertices of arity ≥ 2 , having n leaves and two kinds of vertices called auxiliary and primary, respectively. The corresponding sets of vertices of a tree T are denoted by $V_{aux}(T)$ and $V_{pri}(T)$. Furthermore, we assume that there are no consecutive primary vertices and each tree $T \in \mathbb{T}_n^{\geq 2}[m]$ has exactly m auxiliary vertices. For any $T \in \mathbb{T}_n^{\geq 2}[m]$, we denote by $E_1(T)$ the subset of the set $E(T)$ of edges which consists of: the root edge of T if it is adjacent to an auxiliary vertex; the leaf edges of T connected to auxiliary vertices; the inner edges of T connecting two auxiliary vertices. According to this notation, we set

$$\bar{X}_\alpha(T) = \prod_{v \in V_{pri}(T)} P_{<\alpha}(n)(|v|) \times \prod_{v \in V_{aux}(T)} X_\alpha(|v|) \times \prod_{e \in E_1(T)} P(1). \tag{15}$$

The subspace $\partial \bar{X}_\alpha(T) \subset \bar{X}_\alpha(T)$ is defined as the one composed of elements having at least one auxiliary vertex indexed by ∂X_α . The automorphism group $Aut(T)$ of the tree T acts on these spaces $\partial \bar{X}_\alpha(T)$ and $\bar{X}_\alpha(T)$ by permuting the incoming edges of the vertices. We choose a bijection of the set $\ell(T)$ of leaves of T with the set $[n]$. This allows us to consider the group $Aut(T)$ as a subgroup of the symmetric group Σ_n observing how it permutes the n leaves of T . Denote by $Aut(T) \wr P(1)$ the submonoid of $\Sigma_n \wr P(1)$ defined as the pullback

$$\begin{array}{ccc} Aut(T) \wr P(1) & \longrightarrow & Aut(T) \\ \downarrow & & \downarrow \\ \Sigma_n \wr P(1) & \longrightarrow & \Sigma_n. \end{array} \tag{16}$$

One has the filtration by the number m of vertices labelled by X :

$$P_{<\alpha}(n) = P_\alpha(n)_0 \longrightarrow \dots \longrightarrow P_\alpha(n)_{m-1} \longrightarrow P_\alpha(n)_m \longrightarrow \dots \longrightarrow P_\alpha(n). \tag{17}$$

The inclusion $P_\alpha(n)_{m-1} \rightarrow P_\alpha(n)_m$ fits in the following pushout diagram of $\Sigma_n \wr P(1)$ -spaces:

$$\begin{array}{ccc} \prod_{T \in \mathbb{T}_n^{\geq 2}[m]} \left(\partial \bar{X}_\alpha(T) \times_{Aut(T) \wr P(1)} \Sigma_n \wr P(1) \right) & \longrightarrow & \prod_{T \in \mathbb{T}_n^{\geq 2}[m]} \left(\bar{X}_\alpha(T) \times_{Aut(T) \wr P(1)} \Sigma_n \wr P(1) \right) \\ \downarrow & & \downarrow \\ P_\alpha(n)_{m-1} & \longrightarrow & P_\alpha(n)_m. \end{array} \tag{18}$$

The map $P_\alpha(n)_{m-1} \rightarrow P_\alpha(n)_m$ is a $\Sigma_n \wr P(1)$ -cofibration if the upper horizontal arrow in the above diagram is one. According to Lemma A.3, the extension functor preserves cofibrations and we are only left to showing that every inclusion $\partial \bar{X}_\alpha(T) \rightarrow \bar{X}_\alpha(T)$ is an $Aut(T) \wr P(1)$ -cofibration.

The map $\partial \bar{X}_\alpha(T) \rightarrow \bar{X}_\alpha(T)$ is an $Aut(T) \wr P(1)$ -cofibration. We induct over the number of vertices in T . The base of induction is when T has 0 vertices for which the statement is vacuously true. Now assume that T has ≥ 1 vertices and the statement holds for any other rooted tree with less vertices. We also assume that for any i , $P_{<\alpha}(i)$ is $\Sigma_i \wr P(1)$ -cofibrant as our argument follows a double induction. Let r denote the root vertex. Let us assume that its first ℓ incoming edges are grafted to trees T_1, \dots, T_ℓ , for some $1 \leq \ell \leq |r|$, and the remaining $\ell - |r|$ edges connect to leaves. Consider the following short exact sequence of monoids:

$$1 \rightarrow \prod_{i=1}^{\ell} (Aut(T_i) \wr P(1)) \rightarrow Aut(T) \wr P(1) \rightarrow Aut(r) \times (\Sigma_{|r|-\ell} \wr P(1)) \rightarrow 1,$$

where $Aut(r) \subset \Sigma_\ell$ is the subgroup of permutation of the first ℓ incoming edges of r as a result of $Aut(T)$ -action. It is easy to see that this sequence is split-surjective (see Definition A.4) and thus Lemma A.5 can be applied. There are two cases to consider: the vertex r is auxiliary or it is primary. In both cases, $G_2 = Aut(r) \times \Sigma_{|r|-\ell}$ and we explain how to apply Lemma A.5.

In the first case we take $A \rightarrow B$ to be $P(1) \times \partial X_\alpha(|r|) \times P(1)^{\times(|r|-\ell)} \rightarrow P(1) \times X_\alpha(|r|) \times P(1)^{\times(|r|-\ell)}$, where the first factor $P(1)$ corresponds to the root edge, while the other factors $P(1)$ correspond to the leaf edges connected to r . It is an $Aut(r) \times (\Sigma_{|r|-\ell} \wr P(1))$ -cofibration by restriction (Lemma A.3) and also Lemma A.5. For $X \rightarrow Y$ we take $\partial \prod_{i=1}^{\ell} \bar{X}_\alpha(T_i) \rightarrow \prod_{i=1}^{\ell} \bar{X}_\alpha(T_i)$. It is a $\prod_{i=1}^{\ell} (Aut(T_i) \wr P(1))$ -cofibration according to Example A.6. Denote in this case by ${}_{P(1)\setminus} \bar{X}_\alpha(T)$ the product (15) with one factor $P(1)$ missing, which corresponds to the root edge. We also denote by ${}_{P(1)\setminus} \partial \bar{X}_\alpha(T)$ the image of $\partial \bar{X}_\alpha(T)$ under the projection $\bar{X}_\alpha(T) \rightarrow {}_{P(1)\setminus} \bar{X}_\alpha(T)$. Note that the induction step also allows us to conclude that the inclusion ${}_{P(1)\setminus} \partial \bar{X}_\alpha(T) \rightarrow {}_{P(1)\setminus} \bar{X}_\alpha(T)$ is an $Aut(T) \wr P(1)$ -cofibration.

In the second case, if r is primary, then we take for $A \rightarrow B$ the inclusion $\emptyset \rightarrow P_{<\alpha}(|r|)$. It is an $Aut(r) \times (\Sigma_{|r|-\ell} \wr P(1))$ -cofibration by restricting from $\Sigma_{|r|} \wr P(1)$ and applying Lemma A.3. For $X \rightarrow Y$ we take the map $\partial \prod_{i=1}^{\ell} ({}_{P(1)\setminus} \bar{X}_\alpha(T_i)) \rightarrow \prod_{i=1}^{\ell} ({}_{P(1)\setminus} \bar{X}_\alpha(T_i))$.

The space $O(n)$ is $\Sigma_n \wr O(1)$ -cofibrant. Since the operad O is a retract of the operad P , the monoid $\Sigma_n \wr O(1)$ is a retract of the monoid $\Sigma_n \wr P(1)$. Let $f : \Sigma_n \wr O(1) \rightarrow \Sigma_n \wr P(1)$ and $g : \Sigma_n \wr P(1) \rightarrow \Sigma_n \wr O(1)$ be these maps of monoids such that $g \circ f = id$. These maps give rise to Quillen adjunctions:

$$\Sigma_n \wr O(1)\text{-}Top \xrightleftharpoons[i_2]{i_1} \Sigma_n \wr P(1)\text{-}Top \xrightleftharpoons[j_2]{j_1} \Sigma_n \wr O(1)\text{-}Top$$

For any $A \in \Sigma_n \wr O(1)\text{-}Top$ and $B \in \Sigma_n \wr P(1)\text{-}Top$, the objects $j_2(A)$ and $i_2(B)$ are defined by $j_2(A) = A$ and $i_2(B) = B$ as spaces, with the structure operations such that:

$$j_2(A) \times \Sigma_n \wr P(1) \xrightarrow{id \times g} A \times \Sigma_n \wr O(1) \longrightarrow A,$$

$$i_2(B) \times \Sigma_n \wr O(1) \xrightarrow{id \times f} B \times \Sigma_n \wr P(1) \longrightarrow B.$$

In order to show that $O(n)$ is $\Sigma_n \wr O(1)$ -cofibrant, we consider the lifting problem

$$\begin{array}{ccc} \emptyset & \longrightarrow & X \\ \downarrow & & \downarrow \simeq \\ O(n) & \longrightarrow & Y, \end{array} \tag{19}$$

where $X \rightarrow Y$ is an acyclic fibration in $\Sigma_n \wr O(1)\text{-Top}$. Then we apply the functor j_2 to Diagram (19). The map $j_2(X) \rightarrow j_2(Y)$ is still an acyclic fibration in $\Sigma_n \wr P(1)\text{-Top}$ because the restriction functor j_2 creates fibrations and weak equivalences (in the sense that a map β in $\Sigma_n \wr O(1)\text{-Top}$ is a fibration or a weak equivalence precisely if $j_2(\beta)$ is so in $\Sigma_n \wr P(1)\text{-Top}$). Furthermore, O being a retract of P , the space $j_2(O(n))$ is a retract of $P(n)$ in $\Sigma_n \wr P(1)\text{-Top}$. Since the latter is a $\Sigma_n \wr P(1)$ -cofibration, the space $j_2(O(n))$ is a $\Sigma_n \wr P(1)$ -cofibrant object and there is a map $h : j_2(O(n)) \rightarrow j_2(X)$ solution of the lifting problem (19) in $\Sigma_n \wr P(1)\text{-Top}$. Finally, the map $i_2(h)$ provides a solution to the lifting problem (1.10) due to the relation $g \circ f = id$. \square

Proof of Proposition 1.10 In what follows, we overuse the notation introduced in the proof of Proposition 1.9. By Proposition 1.8, without loss of generality we can assume that $O(0) = \emptyset$. Arguing in the same way as at the end of the proof of Proposition 1.9, we can assume that O is a cellular operad. Indeed, O being a retract of a cellular operad P , the inclusion $O(n; k) \rightarrow O(n)$ is also a retract of $P(n; k) \rightarrow P(n)$. For simplicity and to agree with the notation Proposition 1.9, we assume that $O = P$.

Recall filtration (13) in $P(n)$. We consider a similar filtration in $P(n; k)$ by taking $P_\alpha(n; k) := P(n; k) \cap P_\alpha(n)$. Similarly, we filter the inclusion $P(n; k) \rightarrow P(n)$ taking $P'_\alpha(n) := P(n; k) \cup P_\alpha(n)$. Denote by $P_{<\alpha}(n; k) := \text{colim}_{\beta < \alpha} P_\beta(n; k)$ and $P'_{<\alpha}(n) := \text{colim}_{\beta < \alpha} P'_\beta(n)$. In order to prove the proposition, one has to show that the inclusions $P_{<\alpha}(n; k) \rightarrow P_\alpha(n; k)$ and $P'_{<\alpha}(n) \rightarrow P'_\alpha(n)$ are $(\Sigma_k \times \Sigma_{n-k}) \wr P(1)$ -cofibrations.

This is done by filtering these inclusions similarly to (17). Namely, we define $P_\alpha(n; k)_m := P(n; k) \cap P_\alpha(n)_m$ and $P'_\alpha(n)_m := P(n; k) \cup P_\alpha(n)_m$. We are left to showing that the inclusions $P_\alpha(n; k)_{m-1} \rightarrow P_\alpha(n; k)_m$ and $P'_\alpha(n)_{m-1} \rightarrow P'_\alpha(n)_m$ are $(\Sigma_k \times \Sigma_{n-k}) \wr P(1)$ -cofibrations.

Let $\mathbb{T}_{n,k}^{\geq 2}[m]$, $1 < k < n$, denote the set of exactly the same trees as in $\mathbb{T}_n^{\geq 2}[m]$ in which in addition k (out of n) leaves are marked as special. For $T \in \mathbb{T}_{n,k}^{\geq 2}[m]$, we denote by $U(T) \in \mathbb{T}_n^{\geq 2}[m]$ the tree obtained by forgetting which leaves are special. Let $Aut(T)$ denote the group of automorphisms of T . For each tree $T \in \mathbb{T}_{n,k}^{\geq 2}[m]$ we choose an ordering of its leaves so that we count special leaves first. This ordering gives us an inclusion $Aut(T) \rightarrow \Sigma_n$ (as well as an inclusion $Aut(U(T)) \rightarrow \Sigma_n$). One obviously has $Aut(T) = Aut(U(T)) \cap (\Sigma_k \times \Sigma_{n-k})$. We similarly define

$$Aut(T) \wr P(1) := (Aut(U(T)) \wr P(1)) \cap ((\Sigma_k \times \Sigma_{n-k}) \wr P(1)).$$

Consider the subset $\mathbb{T}_{n,k}^{\geq 2}[m]_I \subset \mathbb{T}_{n,k}^{\geq 2}[m]$ composed of trees that contain a vertex whose all incoming edges are special leaf edges. Its complement is denoted by $\mathbb{T}_{n,k}^{\geq 2}[m]_{II} := \mathbb{T}_{n,k}^{\geq 2}[m] \setminus \mathbb{T}_{n,k}^{\geq 2}[m]_I$.

One has similar to (18) pushout diagrams of $(\Sigma_k \wr P(1) \times \Sigma_{n-k} \wr P(1))$ -spaces.

$$\begin{CD} \prod_{T \in \mathbb{T}_{n,k}^{\geq 2}[m]_I} \partial \bar{X}_\alpha(T) @>>> \prod_{T \in \mathbb{T}_{n,k}^{\geq 2}[m]_{II}} \bar{X}_\alpha(T) \\ @V \downarrow VV @VV \downarrow V \\ P_\alpha(n; k)_{m-1} @>>> P_\alpha(n; k)_m \end{CD}$$

(20)

$$\begin{array}{ccc}
 \coprod_{T \in \mathbb{T}_{n,k}^{\geq 2}[m]_{II}} \partial \overline{X}_\alpha(T) \times_{Aut(T) \wr P(1)} ((\Sigma_k \times \Sigma_{n-k}) \wr P(1)) & \longrightarrow & \coprod_{T \in \mathbb{T}_{n,k}^{\geq 2}[m]_{II}} \overline{X}_\alpha(T) \times_{Aut(T) \wr P(1)} ((\Sigma_k \times \Sigma_{n-k}) \wr P(1)) \\
 \downarrow & & \downarrow \\
 P'_\alpha(n)_{m-1} & \longrightarrow & P'_\alpha(n)_m.
 \end{array} \tag{21}$$

In the above $\overline{X}_\alpha(T)$ is defined by (15).

The upper horizontal arrows in diagrams (20) and (21) are $(\Sigma_k \wr P(1) \times \Sigma_{n-k} \wr P(1))$ -cofibrations. Indeed, from the proof of Proposition 1.9 we know that each inclusion $\partial \overline{X}_\alpha(T) \rightarrow \overline{X}_\alpha(T)$ is an $Aut(U(T)) \wr P(1)$ -cofibration. Applying Lemma A.3 to the restriction along $Aut(T) \wr P(1) \rightarrow Aut(U(T)) \wr P(1)$, we get that such inclusion is an $Aut(T) \wr P(1)$ -cofibration. Applying again this lemma to the induction along $Aut(T) \wr P(1) \rightarrow (\Sigma_k \times \Sigma_{n-k}) \wr P(1)$ and using the fact that the diagrams (20) and (21) are pushout squares, we conclude that $P_\alpha(n; k)_{m-1} \rightarrow P_\alpha(n; k)_m$ and $P'_\alpha(n)_{m-1} \rightarrow P'_\alpha(n)_m$ are $(\Sigma_k \times \Sigma_{n-k}) \wr P(1)$ -cofibrations. Applying again Lemma A.3 to the restriction along $(\Sigma_k \wr P(1)) \times \Sigma_{n-k} \rightarrow (\Sigma_k \times \Sigma_{n-k}) \wr P(1)$, these maps are also $(\Sigma_k \wr P(1)) \times \Sigma_{n-k}$ -cofibrations. \square

2 The Projective Model Category of (P-Q)-Bimodules

Let P and Q be two operads. A $(P-Q)$ -bimodule is a Σ -sequence $M \in \Sigma Seq$ together with operations

$$\begin{aligned}
 \gamma_r : M(n) \times \prod_{1 \leq i \leq n} Q(m_i) &\longrightarrow M(\sum_i m_i), \text{ right operations,} \\
 \gamma_\ell : P(n) \times \prod_{1 \leq i \leq n} M(m_i) &\longrightarrow M(\sum_i m_i), \text{ left operations,}
 \end{aligned} \tag{22}$$

satisfying the following relations, with $\sigma \in \Sigma_n$ and $\tau_i \in \Sigma_{m_i}$:

$$\begin{array}{ccc}
 M(n) \times \prod_{1 \leq i \leq n} Q(m_i) \times \prod_{1 \leq j \leq m_i} Q(k_{i,j}) & \longrightarrow & M(n) \times \prod_{1 \leq i \leq n} Q(\sum_j k_{i,j}) \\
 \downarrow & & \downarrow \\
 M(\sum_i m_i) \times \prod_{1 \leq j \leq m_i} Q(k_{i,j}) & \longrightarrow & M(\sum_{i,j} k_{i,j}) \\
 \text{Associativity for the right operations} & & \text{Associativity for the left operations}
 \end{array}$$

$$\begin{array}{ccc}
 P(n) \times \prod_{1 \leq i \leq n} P(m_i) \times \prod_{1 \leq j \leq m_i} M(k_{i,j}) & \longrightarrow & P(n) \times \prod_{1 \leq i \leq n} M(\sum_j k_{i,j}) \\
 \downarrow & & \downarrow \\
 P(\sum_i m_i) \times \prod_{1 \leq j \leq m_i} M(k_{i,j}) & \longrightarrow & M(\sum_{i,j} k_{i,j}) \\
 \text{Associativity for the left operations} & & \text{Associativity for the right operations}
 \end{array}$$

$$\begin{array}{ccc}
 P(n) \times \prod_{1 \leq i \leq n} M(m_i) \times \prod_{1 \leq j \leq m_i} Q(k_{i,j}) & \longrightarrow & P(n) \times \prod_{1 \leq i \leq n} M(\sum_j k_{i,j}) \\
 \downarrow & & \downarrow \\
 M(\sum_i m_i) \times \prod_{1 \leq j \leq m_i} Q(k_{i,j}) & \longrightarrow & M(\sum_{i,j} k_{i,j}) \\
 \text{Compatibility between the left and right operations} & & \text{Compatibility with the unit of the operad}
 \end{array}$$

$$\begin{array}{ccc}
 M(n) \times Q(1)^{\times n} & \longleftarrow & M(n) & \longrightarrow & P(1) \times M(n) \\
 & \searrow \gamma_r & \parallel & \swarrow \gamma_\ell & \\
 & & M(n) & &
 \end{array}$$

$$\begin{array}{ccc}
 M(n) \times \prod_{1 \leq i \leq n} Q(m_i) & \longrightarrow & M(m_1 + \dots + m_n) \\
 \sigma^* \times \prod_i (\tau_i)^* \downarrow & & \downarrow (\sigma(\tau_1, \dots, \tau_n))^* \\
 M(n) \times \prod_{1 \leq i \leq n} Q(m_{\sigma(i)}) & \longrightarrow & M(m_1 + \dots + m_n) \\
 \text{Right compatibility with the symmetric group action} & &
 \end{array}
 \qquad
 \begin{array}{ccc}
 P(n) \times \prod_{1 \leq i \leq n} M(m_i) & \longrightarrow & M(m_1 + \dots + m_n) \\
 \sigma^* \times \prod_i (\tau_i)^* \downarrow & & \downarrow (\sigma(\tau_1, \dots, \tau_n))^* \\
 P(n) \times \prod_{1 \leq i \leq n} M(m_{\sigma(i)}) & \longrightarrow & M(m_1 + \dots + m_n) \\
 \text{Left compatibility with the symmetric group action} & &
 \end{array}$$

where the permutation $\sigma(\tau_1, \dots, \tau_n)$ is obtained from the well known operadic compositions on the symmetric groups (see [18, Proposition 1.1.9]).

As part of the left operations, there is a map $\gamma_0 : P(0) \rightarrow M(0)$ in arity 0. A map between $(P-Q)$ -bimodules should preserve these operations. We denote by $\Sigma\text{Bimod}_{P; Q}$ the category of $(P-Q)$ -bimodules. Thanks to the unit in $Q(1)$, the right operations γ_r can equivalently be defined as a family of continuous maps

$$\circ^i : M(n) \times Q(m) \longrightarrow M(n + m - 1), \quad \text{with } 1 \leq i \leq n.$$

Given an integer $r \geq 0$, we also consider the category of r -truncated bimodules $T_r \Sigma\text{Bimod}_{P; Q}$. An object is an r -truncated Σ -sequence endowed with left and right operations (22) under the conditions $n \leq r$ and $\sum m_i \leq r$ for γ_r and the condition $\sum m_i \leq r$ for γ_ℓ . One has an obvious truncation functor

$$T_r(-) : \Sigma\text{Bimod}_{P; Q} \longrightarrow T_r \Sigma\text{Bimod}_{P; Q}.$$

In the rest of the paper, we use the notation

$$\begin{aligned}
 x \circ^i q &= \circ^i(x; q), \text{ for } x \in M(n) \text{ and } q \in Q(m), \\
 p(x_1, \dots, x_n) &= \gamma_\ell(p, x_1, \dots, x_n), \text{ for } p \in P(n) \text{ and } x_i \in M(m_i).
 \end{aligned}$$

Example 2.1 If $\eta : P \rightarrow Q$ is a map of operads, then η is also a map of P -bimodules. Indeed, any operad is a bimodule over itself while the P -bimodule structure on Q is given by the following formulas:

$$\begin{aligned}
 \circ^i : Q(n) \times P(m) &\longrightarrow Q(m + n - 1); \\
 (q; p) &\longmapsto q \circ_i \eta(p), \\
 \gamma_\ell : P(n) \times Q(m_1) \times \dots \times Q(m_n) &\longrightarrow Q(m_1 + \dots + m_n); \\
 (p, q_1, \dots, q_n) &\longmapsto (\dots ((\eta(p) \circ_n q_n) \circ_{n-1} q_{n-1}) \dots) \circ_1 q_1.
 \end{aligned}$$

2.1 Properties of the Category of Bimodules

In this subsection we introduce some basic properties related to the category of $(P-Q)$ -bimodules where P and Q are two fixed operads. First, we show that the category of $(P-Q)$ -bimodules is equivalent to the category of algebras over an explicit colored operad denoted by $P+Q$. Thereafter, we build the free bimodule functor using the language of trees. Using this explicit construction of the free functor, we are able to give a combinatorial description of the pushout for bimodules.

2.1.1 Bimodules as Algebras over a Colored Operad

Given two operads P and Q , we build a colored operad $P+Q$ such that the category of $(P-Q)$ -bimodules is equivalent to the category of $(P+Q)$ -algebras. By a colored operad C , with set of colors S , we understand a family of spaces $C = \{C(s_1, \dots, s_n; s_{n+1}), n \geq$

$0, s_i \in S\}$, equipped with an action of permutations such that $\sigma^* : C(s_1, \dots, s_n; s_{n+1}) \rightarrow C(s_{\sigma(1)}, \dots, s_{\sigma(n)}; s_{n+1})$, for $\sigma \in \Sigma_n$, with units $*_s \in C(s; s)$ for $s \in S$, and with operadic compositions

$$\begin{aligned} \circ_i &: C(s_1, \dots, s_n; s_{n+1}) \times C(s'_1, \dots, s'_m; s_i) \\ &\longrightarrow C(s_1, \dots, s_{i-1}, s'_1, \dots, s'_m, s_{i+1}, \dots, s_n; s_{n+1}), \end{aligned}$$

that satisfy associativity, unit and equivariance relations which are similar to the ones introduced in Sect. 1.4. An algebra over C is a family of spaces $\{X_s, s \in S\}$ together with operations of the form

$$\alpha[s_1, \dots, s_n; s_{n+1}] : C(s_1, \dots, s_n; s_{n+1}) \times X_{s_1} \times \dots \times X_{s_n} \longrightarrow X_{s_{n+1}},$$

compatible with the operadic structure.

The operad $P + Q$ has the set of non-negative integers $S = \mathbb{N}$ as colors so that an element $\theta \in (P + Q)(n_1, \dots, n_k; m)$ governs an operation of the form $\theta : M(n_1) \times \dots \times M(n_k) \rightarrow M(m)$ on a $(P-Q)$ -bimodule M . We define this operad $P + Q$ by a presentation by generators and relations. We take two kinds of generating operations, which respectively encode the left P -action and the right Q -action of a $(P-Q)$ -bimodule structure. We actually integrate the first kind of generators in a colored operad P_1 (encoding the left operations) and the second kind of generators in another colored operad Q_1 (encoding the right operations), which is concentrated in arity one. We explain the definition of these operads P_1 and Q_1 and their action on a $(P-Q)$ -bimodule M in the next paragraph. We shape the composite elements of the operad $P + Q$ on trees equipped with two sets of vertices, left vertices, represented by diamonds \blacklozenge , which correspond to the P_1 -factors, and right vertices, represented by circles \bullet , which correspond to the Q_1 -factors. We describe the structure of these trees in a second step and we explain afterwards the definition of the operad $P + Q$ with elements shaped on such trees, moded out by relations which reflect the structure relations of left and right actions on bimodules. (We can actually identify $P + Q$ with a coproduct of the colored operads P_1 and Q_1 moded out by extra relations which reflect the commutation of left and right actions on a bimodule.)

Definition 2.2 The colored operads P_1 and Q_1

- To any collection of non-negative integers n_1, \dots, n_k , we associate the space $P_1(n_1, \dots, n_k; n_1 + \dots + n_k)$ such that:

$$P_1(n_1, \dots, n_k; n_1 + \dots + n_k) := P(k).$$

We have an action of permutations $\sigma^* : P_1(n_1, \dots, n_k; n_1 + \dots + n_k) \rightarrow P_1(n_{\sigma(1)}, \dots, n_{\sigma(k)}; n_1 + \dots + n_k)$, induced by the symmetric structure of the operad, unit elements $*_n \in P_1(n, n)$, given by the unit of P , and “colored” composition operations

$$\begin{aligned} \circ_i &: P_1(n_1, \dots, n_k; m) \times P_1(n'_1, \dots, n'_k; n_i) \longrightarrow P_1(n_1, \dots, n_{i-1}, n'_1, \dots, n'_k, n_{i+1}, \dots, n_k; m), \\ p &: p' \longmapsto p \circ_i p', \end{aligned} \tag{23}$$

induced by the composition operations of P , for $n_i = n'_1 + \dots + n'_k$ and $m = n_1 + \dots + n_k$. We immediately see that these structure operations fulfill the unit, associativity and equivariance axioms of colored operads (as a consequence of the operad axioms in P). We also see that the left P -action on a $(P-Q)$ -bimodule M gives an operation

$$P_1(n_1, \dots, n_k; n_1 + \dots + n_k) \times M(n_1) \times \dots \times M(n_k) \rightarrow M(n_1 + \dots + n_k),$$

which makes M and algebra over the colored operad P_1 .

- To any integers n and m , we associate the space $Q_1(n; m)$ such that:

$$Q_1(n; m) := \coprod_{\alpha: [m] \rightarrow [n]} \prod_{i \in [n]} Q(|\alpha^{-1}(i)|).$$

A point in the above space is denoted by $(\alpha; \{q_i\}_{i \in [n]})$. We have a unit element $*_n \in Q_1(n; n)$ given by collection of operadic units $1 \in Q(1)$ in the factor of $Q_1(n; n)$ indexed by the identity map $id : [n] \rightarrow [n]$. We have compositions of the form

$$\begin{aligned} \mu : Q_1(n; m) \times Q_1(m; \ell) &\longrightarrow Q_1(n; \ell); \\ (\alpha; \{q_i\}); (\alpha'; \{q'_j\}) &\longmapsto (\alpha \circ \alpha'; \{z_i\}_{i \in [n]}), \end{aligned} \tag{24}$$

where z_i , with $i \in [n]$, is obtained using the operadic structure of Q :

$$\begin{aligned} z_i := \sigma_i(\alpha, \alpha')^* &\left[\left(\cdots \left((q_i \circ_{l_i} q'_{b_{l_i}}) \circ_{\ell_{i-1}} q'_{b_{l_{i-1}}} \right) \cdots \right) \circ_1 q'_{b_1} \right], \quad \text{with} \\ \alpha^{-1}(i) = \{b_1 < \cdots < b_{\ell_i}\}. \end{aligned}$$

Here $\sigma_i(\alpha, \alpha')^*$ is the inverse of the shuffle permutation of the set $(\alpha \circ \alpha')^{-1}(i)$, reordering this set as $(\alpha')^{-1}(b_1), \dots, (\alpha')^{-1}(b_{\ell_i})$. The operadic axioms imply that these operations are associative and unital, so that Q_1 forms a colored operad (concentrated in arity one). We also see that the right Q -action on a $(P - Q)$ -bimodule M gives an operation

$$\zeta : Q_1(n; m) \times M(n) \rightarrow M(m)$$

by

$$\zeta((\alpha; \{q_i\}), x) = \sigma_\alpha^* \left(\left(\cdots (x \circ^n q_n) \cdots \right) \circ^1 q_1 \right),$$

for any $(\alpha; \{q_i\}) \in Q_1(n; m)$, $x \in M(n)$, where $\sigma_\alpha \in \Sigma_m$ is the inverse of the corresponding shuffle of the blocks of size $|\alpha^{-1}(1)|, \dots, |\alpha^{-1}(n)|$ in $[m]$. We readily deduce from the bimodule axioms that this operation is unital and associative with respect to the composition operation (24), so that M forms an algebra over the colored operad Q_1 .

Definition 2.3 The set of trees $\mathbb{P}[n_1, \dots, n_k; m]$

Let n_1, \dots, n_k and m be non-negative integers. An element in $\mathbb{P}[n_1, \dots, n_k; m]$ is a tuple $T = (T, V_\ell(T), V_r(T), f)$, where T is a planar rooted tree having k leaves indexed by a permutation from Σ_k and having two kinds of vertices called *left* and *right* vertices, respectively. The sets $V_\ell(T)$ and $V_r(T)$ consist of left vertices and right vertices, respectively. In particular, right vertices are necessarily of arity one and are represented by circles \bullet in the tree while the left vertices are represented by diamonds \blacklozenge (see Fig. 2). Left vertices are allowed to be of any arity ≥ 0 .

The assignment $f : E(T) \rightarrow \mathbb{N}$ labels the edges of the planar tree by integers. In particular, the outgoing edge of the tree is labelled by m while the k leaves are labelled by n_1, \dots, n_k , according to the permutation. We denote by $n_1^v, \dots, n_{|v|}^v$ the integers labelling the incoming edges of a vertex v and by n_0^v its output edge according to the orientation toward the root. Furthermore, if v is a left vertex, then one has the relation $n_0^v = n_1^v + \cdots + n_{|v|}^v$.

Construction 2.4 The colored operad $P+Q$, with set of colors \mathbb{N} (non-negative integers), is obtained from the sets of trees $\mathbb{T}[n_1, \dots, n_k; m]$ by indexing the left vertices by points in P_1





Fig. 2 Illustration of an element in $\mathbb{P}[n_1, n_2, n_3, n_4, n_5; m]$

and the right vertices by points in Q_1 . More precisely, one has


$$(P+Q)(n_1, \dots, n_k; m) := \coprod_{T \in \mathbb{T}[n_1, \dots, n_k; m]} \left[\prod_{v \in V_r(T)} Q_1(n_v^v; n_0^v) \times \prod_{v \in V_l(T)} P_1(n_v^v, \dots, n_{|v|}^v; n_0^v) \right] / \sim$$

where the equivalence relation is generated by the following axioms:

- If a vertex is indexed by the unit of the operad P_1 or Q_1 , then we remove it:



- If a left vertex is indexed by a point of the form $p \cdot \sigma$, with $p \in P(k)$ and $\sigma \in \Sigma_k$, then one has the following identity in which $\tau = \sigma(id_{\Sigma_{n_1}}, \dots, id_{\Sigma_{n_k}}) \in \Sigma_{n_1 + \dots + n_k}$ is the element permuting the blocks $\{1, \dots, n_1\}, \dots, \{n_1 + \dots + n_{k-1} + 1, \dots, n_1 + \dots + n_k\}$ in $[n_1 + \dots + n_k]$:








Fig. 3 Illustration of the operadic composition \circ_2

- ▶ If there are two consecutive left vertices or two consecutive right vertices, then we contract the edge connecting them using the operation (24) or (23):



- ▶ If the incoming edges of a left vertex are connected to right vertices, then we can permute them:



Let x and y be two points in $(P+Q)(n_1, \dots, n_k; m)$ and $(P+Q)(n'_1, \dots, n'_k; n_i)$, respectively. The operadic composition $x \circ_i y$ consists in grafting the tree indexing y into the i -th leaf of the tree indexing x and keeping the labels of each tree in order to decorate the new one.

Remark 2.5 According to the relations introduced in the previous construction, for any integers n_1, \dots, n_k and m , each point in the component $(P+Q)(n_1, \dots, n_k; m)$ has a representative element having exactly one left vertex and one right vertex such that the right vertex is the root of the tree, see Fig. 4. Thanks to the second relation of Construction 2.4, one can order the leaves from 1 to k . Therefore, as a space,

$$(P+Q)(n_1, \dots, n_k; m) = P(k) \times \left[\coprod_{\alpha: [m] \rightarrow [n_1 + \dots + n_k]} \prod_{1 \leq i \leq n_1 + \dots + n_k} Q(|\alpha^{-1}(i)|) \right]. \tag{25}$$

Proposition 2.6 *The category of $(P-Q)$ -bimodules is equivalent to the category of $(P+Q)$ -algebras.*

Proof In one direction, we already explained that a $(P-Q)$ -bimodule M inherits an action of the operads P_1 and Q_1 that generate $P+Q$. We just check that these actions are compatible with the relations of Construction 2.4 when we compose them to get an action of $P+Q$ on M (the compatibility with the first three relations follows from the unit, associativity and




Fig. 4 Illustration of a representative element

equivariance of the action of the operads P_1 and Q_1 on M , the compatibility with the fourth relation follows from the commutation of the left and right actions).

In the converse direction, we show that a $(P+Q)$ -algebra M' inherits a bimodule structure. First, M' is a Σ -sequence thanks to the action of the left vertices indexed by $(\sigma; \{*_1\})$, where σ is a permutation from some Σ_k and $\{*_1\}$ is a collection of k identity elements from $Q(1)$. Then we just inverse the constructions of Definition 2.2 in order to retrieve left and right operations on M' . To be explicit, in order to define the left operations γ_ℓ , we consider the inclusion $\iota_\ell : P(k) \rightarrow (P+Q)(n_1, \dots, n_k; n_1 + \dots + n_k)$, for any integers n_1, \dots, n_k , sending a point $p \in P(k)$ to the k -corolla whose root is a left vertex indexed by p . The map γ_ℓ is defined as the following composite:

$$\begin{array}{ccc}
 P(k) \times M'(n_1) \times \dots \times M'(n_k) & \xrightarrow{\gamma_\ell} & M'(n_1 + \dots + n_k) \\
 \downarrow \iota_\ell \times id \times \dots \times id & \nearrow & \\
 (P+Q)(n_1, \dots, n_k; n_1 + \dots + n_k) \times M'(n_1) \times \dots \times M'(n_k) & &
 \end{array}$$

Similarly, for any pair of integers $(n; m)$ and $1 \leq i \leq n$, there is a map $\iota_r : Q(m) \rightarrow (P+Q)(n; n+m-1)$ sending a point $q \in Q(m)$ to the element $(\alpha; \{q_j\})$ where α is given by

$$\alpha : [n+m-1] \longrightarrow [n]; j \mapsto \begin{cases} j & \text{if } j \leq i, \\ i & \text{if } i < j < i+m, \\ j-m+1 & \text{if } j \geq i+m. \end{cases}$$

One has $q_i = q$ and $q_j = *_1$ for any $j \neq i$. The right operation o^i , with $1 \leq i \leq n$, is defined as the following composite map:

$$\begin{array}{ccc}
 M'(n) \times Q(m) & \xrightarrow{o^i} & M'(n+m-1) \\
 \downarrow id \times \iota_r & & \uparrow \\
 M'(n) \times (P+Q)(n; n+m-1) & \xrightarrow{\cong} & (P+Q)(n; n+m-1) \times M'(n)
 \end{array}$$

The relations introduced in Construction 2.4 readily imply that these operations satisfy the bimodule axioms. □



Fig. 5 Examples of a tree with section $T_1 \in s\mathbb{P}_5$ and a reduced tree with section $T_2 \in rs\mathbb{P}_6$

2.1.2 The Free Bimodule Functor

We denote by ΣSeq_P and $T_r \Sigma Seq_P$ the categories of Σ -sequences and r -truncated Σ -sequences M equipped with a map $\gamma_0 : P(0) \rightarrow M(0)$. In other words, if P_0 is the Σ -sequence given by $P_0(0) = P(0)$ and the empty set otherwise, then one has the following identities:

$$\Sigma Seq_P := P_0 \downarrow \Sigma Seq \quad \text{and} \quad T_r \Sigma Seq_P := P_0 \downarrow T_r \Sigma Seq. \tag{26}$$

Furthermore, there is a forgetful functor from the category of (possibly truncated) bimodules to the category of Σ -sequences endowed with a map from P_0 :

$$\mathcal{U}^\Sigma : \Sigma \text{Bimod}_{P; Q} \longrightarrow \Sigma Seq_P : \quad \text{and} \quad \mathcal{U}^{T_r \Sigma} : T_r \Sigma \text{Bimod}_{P; Q} \longrightarrow T_r \Sigma Seq_P. \tag{27}$$

Their left adjoints denoted $\mathcal{F}_{P; Q}^\Sigma$ and $\mathcal{F}_{P; Q}^{T_r \Sigma}$, respectively, are some versions of free functors. As usual in the operadic theory, the free functor can be described as a coproduct indexed by a particular set of trees. In that case, we use the set of *trees with section* which are pairs $T = (T; V^p(T))$, where T is a planar rooted tree (whose leaves are labelled by some permutation) and $V^p(T)$ is a subset of vertices, called *pearls*, satisfying the following condition: each path from a leaf or a univalent vertex to the root passes through a unique pearl.

The set of pearls forms a section cutting the tree into two parts. Note that the arity zero vertices can be either above or on the section, but never below. We denote by $V^u(T)$ (respectively, $V^d(T)$) the vertices above the section (respectively below the section). A tree with section is said to be *reduced* if each non-pearl vertex is connected to a pearl by an inner edge. We denote by $s\mathbb{P}_n$ and $rs\mathbb{P}_n$ the sets of trees with section and reduced trees with section, respectively, having exactly n leaves labelled by a permutation of $\{1 \dots n\}$.

Construction 2.7 Let $M = \{M(n)\}$ be a Σ -sequence equipped with a map $\gamma_0 : P(0) \rightarrow M(0)$. The space $\mathcal{F}_{P; Q}^\Sigma(M)(n)$ is obtained from the set of reduced trees with section by indexing the pearls by points in M whereas the vertices above the section (respectively below the section) are indexed by points in the operad Q (respectively the operad P). More precisely, one has

$$\mathcal{F}_{P; Q}^\Sigma(M)(n) = \left(\coprod_{T \in rs\mathbb{P}_n} \prod_{p \in V^p(T)} M(|p|) \times \prod_{v \in V^d(T)} P(|v|) \times \prod_{v \in V^u(T)} Q(|v|) \right) / \sim. \tag{28}$$

A point in $\mathcal{F}_{P; Q}^\Sigma(M)$ is denoted by $[T; \{m_p\}; \{p_v\}; \{q_v\}]$ where T is a reduced tree with section while $\{m_p\}_{p \in V^p(T)}$, $\{p_v\}_{v \in V^d(T)}$ and $\{q_v\}_{v \in V^u(T)}$ are points in M , P and Q , respectively. The equivalence relation is generated by the following relations:




Fig. 6 Illustration of the compatibility with the symmetric group





Fig. 7 Illustration of the γ_0 -relation

(i) *The unit relation:* if a vertex is indexed by the unit of the operad P or Q , then we can remove it:



(ii) *The compatibility with the symmetric group action:* if a vertex is labelled by $x \cdot \sigma$, with x a point in $P(n)$, $Q(n)$ or $M(n)$ and $\sigma \in \Sigma_n$, then we can remove σ by permuting the incoming edges (see Fig. 6).

(iii) *The γ_0 -relation:* if a pearl is indexed by a point of the form $\gamma_0(x)$, with $\gamma_0 : P(0) \rightarrow M(0)$ and $x \in P(0)$, then we contract its output edge using the operadic structures of P (see Fig. 7). In particular, if the vertex below the section indexed by $p \in P(n)$ is connected only to univalent pearls indexed by $\gamma_0(p_1), \dots, \gamma_0(p_n)$, respectively, then we can contract all the incoming edges. The new vertex so obtained is a pearl indexed by $\gamma_0((\dots((p \circ_n p_n) \circ_{n-1} p_{n-1}) \dots) \circ_1 p_1)$.

The right operation \circ^i with an element $q \in Q(m)$ consists in grafting the m -corolla indexed by q into the i -th leaf of the reduced tree with section T . If the so obtained element contains an inner edge joining two consecutive vertices other than a pearl, then we contract it using the operadic structure of Q (see Fig. 8).

The left operation between an element $p \in P(n)$ and a family of points $[T_i ; \{m_i^p\} ; \{p_i^v\} ; \{q_i^v\}] \in \mathcal{F}_{P; Q}^\Sigma(M)$, with $1 \leq i \leq n$, is defined as follows: each tree T_i , with $1 \leq i \leq n$, is grafted from left to right to a leaf of the n -corolla whose vertex is indexed by p . If the so obtained element contains inner edges joining two consecutive vertices other than pearls, then we contract them using operadic structure of P .

Finally, one has the following map sending $p \in P(0)$ to the element $[T ; \{\gamma_0(p)\} ; \emptyset ; \emptyset]$ where T is the perled 0-corolla whose root is indexed by $\gamma_0(p)$:

$$\gamma'_0 : P(0) \longrightarrow \mathcal{F}_{P; Q}^\Sigma(M)(0).$$

Similarly, the free r -truncated bimodule functor $\mathcal{F}_{P; Q}^{T_r, \Sigma}$ is obtained from the formula (28) by taking the restriction of the coproduct to the reduced trees with section having at most r




Fig. 8 Illustration of the right operation $\circ^1 : \mathcal{F}_{P; Q}^\Sigma(M)(5) \times Q(3) \rightarrow \mathcal{F}_{P; Q}^\Sigma(M)(7)$

leaves and such that each pearl has at most r incoming edges. The equivalence relation, the left and right operations and the map γ'_0 are defined in the same way. Finally, one has two functors:

$$\mathcal{F}_{P; Q}^\Sigma : \Sigma Seq_P \longrightarrow \Sigma \text{Bimod}_{P; Q} \quad \text{and} \quad \mathcal{F}_{P; Q}^{T_r \Sigma} : T_r \Sigma Seq_P \longrightarrow T_r \Sigma \text{Bimod}_{P; Q}.$$

Theorem 2.8 *One has the following adjunctions:*

$$\begin{aligned} \mathcal{F}_{P; Q}^\Sigma : \Sigma Seq_P \rightleftarrows \Sigma \text{Bimod}_{P; Q} : \mathcal{U}^\Sigma \quad \text{and} \\ \mathcal{F}_{P; Q}^{T_r \Sigma} : T_r \Sigma Seq_P \rightleftarrows T_r \Sigma \text{Bimod}_{P; Q} : \mathcal{U}^{T_r \Sigma}. \end{aligned} \tag{29}$$

Proof Let M' be a $(P-Q)$ -bimodule and $f : M \rightarrow M'$ be a morphism in the category ΣSeq_P . One has to prove that there exists a unique map of $(P-Q)$ -bimodules $\tilde{f} : \mathcal{F}_{P; Q}^\Sigma(M) \rightarrow M'$ such that the following diagram commutes:

$$\begin{array}{ccc} M & \xrightarrow{f} & M' \\ \downarrow i & \dashrightarrow \exists! \tilde{f} & \\ \mathcal{F}_{P; Q}^\Sigma(M) & & \end{array} \tag{30}$$

We build the map \tilde{f} by induction on the number of vertices in the set $np(T) = V(T) \setminus V^p(T)$. Let $[(T; \sigma); \{m_p\}; \{p_v\}; \{q_v\}]$ be a point in $\mathcal{F}_{P; Q}^\Sigma(M)$ such that $|nb(T)| = 0$ and σ is the permutation indexing the leaves of T . By construction, T is necessarily a pearl corolla with only one vertex labelled by $m_r \in M$. Due to the commutativity of Diagram (30), the following equality has to be satisfied:

$$\tilde{f}([(T; \sigma); \{m_p\}; \{p_v\}; \{q_v\}]) = f(m_r) \cdot \sigma.$$

Let $[(T; \sigma); \{m_p\}; \{p_v\}; \{q_v\}]$ be a point in $\mathcal{F}_{P; Q}^\Sigma(M)$ where T has only one vertex v which is not a pearl. There are two cases to consider. If v is the root of the tree T , then the root is labelled by a point $p_v \in P$ and $[(T; \sigma); \{m_p\}; \{p_v\}; \{q_v\}]$ has a decomposition of the form

$$p_v([(T_1; id); \{m_1\}; \emptyset; \emptyset], \dots, [(T_{|v|}; id); \{m_{|v|}\}; \emptyset; \emptyset]) \cdot \sigma,$$

where T_i is a pearl corolla labelled by $m_i \in M$. Since \tilde{f} has to be a $(P-Q)$ -bimodule map, one has the equality

$$\tilde{f}([(T; \sigma); \{m_p\}; \{p_v\}; \{q_v\}]) = p_v(f(m_1), \dots, f(m_{|v|})) \cdot \sigma.$$

If the root is a pearl, then there exists a unique inner edge e such that $s(e) = v$ and $t(e) = r$. So, the point $[(T; \sigma); \{m_p\}; \{p_v\}; \{q_v\}]$ has a decomposition on the form

$([(T_1; id); \{m_p\}; \emptyset; \emptyset] \circ^i q_{s(e)}) \cdot \sigma$ with $q_{s(e)} \in Q$ and $m_{t(e)} \in M$. Since \tilde{f} has to be an $(P-Q)$ -bimodule map, there is the equality

$$\tilde{f}([(T; \sigma); \{m_p\}; \{p_v\}; \{q_v\}]) = (f(m_{t(e)}) \circ^i q_{s(e)}) \cdot \sigma.$$

Assume \tilde{f} has been defined for $|np(T)| \leq n$. Let $[(T; \sigma); \{m_p\}; \{p_v\}; \{q_v\}]$ be a point in $\mathcal{F}_{P; Q}^\Sigma(M)$ such that $|np(T)| = n + 1$. By definition, there is an inner edge e whose target vertex is a pearl. So, the point $[(T; \sigma); \{m_p\}; \{p_v\}; \{q_v\}]$ has a decomposition of the form $([(T_1; id); \{m_p\}; \{p_v\}; \{q_v\} \setminus \{q_{s(e)}\}] \circ^i q_{s(e)}) \cdot \sigma$ where T_1 is a planar tree with section such that $|np(T_1)| = n$. Since \tilde{f} has to be a $(P-Q)$ -bimodule map, there is the equality

$$\tilde{f}([(T; \sigma); \{m_p\}; \{p_v\}; \{q_v\}]) = (\tilde{f}([(T_1; id); \{m_p\}; \{p_v\}; \{q_v\} \setminus \{q_{s(e)}\}]) \circ^i q_{s(e)}) \cdot \sigma.$$

Due to the $(P-Q)$ -bimodule axioms, \tilde{f} does not depend on the choice of the decomposition and \tilde{f} is a $(P-Q)$ -bimodule map. The uniqueness follows from the construction. Similarly, we can prove that the functor $\mathcal{F}_{P; Q}^{T, \Sigma}$ is the left adjoint to the forgetful functor. \square

2.1.3 Combinatorial Description of the Pushout

Let P and Q be two topological operads. In the following, we use the notation introduced for the free bimodule functor in order to give an explicit description of the pushout in the category of $(P-Q)$ -bimodules. This description will be used in the next subsections. We fix the following diagram in the category of $(P-Q)$ -bimodules

$$\begin{array}{ccc} A & \xrightarrow{f_1} & C \\ f_2 \downarrow & & \\ B & & \end{array} \tag{31}$$

Then we consider the Σ -sequence D obtained from A, B, C and the sets $s\mathbb{P}_n$ of trees with section (see Sect. 2.1.2) by indexing the pearls of such trees by points in B or C whereas the other vertices below the section (respectively above the section) are indexed by points in the operad P (respectively the operad Q). More precisely, one has

$$D(n) = \left(\prod_{T \in s\mathbb{P}_n} \prod_{p \in V^P(T)} \left(B(|p|) \sqcup_{A(|p|)} C(|p|) \right) \times \prod_{v \in V^d(T)} P(|v|) \times \prod_{v \in V^u(T)} Q(|v|) \right) / \sim. \tag{32}$$

By abuse of notation, we denote by $[T; \{m_p\}; \{p_v\}; \{q_v\}]$ a point in $D(n)$. The equivalence relation is generated by relations (i), (ii), (iii) in Construction 2.7 (unit relation, compatibility with the symmetric group action and γ_0 -relation). Furthermore, one has also the following relations:

- (iv) *Pushout relation 1*: each inner edge, which is not connected to a pearl, is contracted using the operadic structures of P and Q (see Fig. 9).
- (v) *Pushout relation 2*: every inner edge above the section connected to a pearl is contracted using the right Q -module structures of B and C .
- (vi) *Pushout relation 3*: if a vertex v below the section is connected to pearls indexed by points in B (respectively, C), then we contract the incoming edges of v using the P -module structure of B (respectively, the left P -module structure of C) (see Fig. 9).




Fig. 9 Illustration of the pushout relations (iv) and (vi)

The Σ -sequence D inherits a $(P-Q)$ -bimodule structure from B and C . The right operations are defined using the right module structures of B and C . Similarly, the left operation between an element $p \in P(n)$ and a family of points $[T_i; \{m_p^i\}; \{p_v^i\}; \{q_v^i\}] \in D$, with $1 \leq i \leq n$, is defined as follows: each tree T_i , with $1 \leq i \leq n$, is grafted from left to right to a leaf of the n -corolla whose vertex is indexed by p . Moreover, there is a map

$$\gamma'_0 : P(0) \longrightarrow D(0)$$

sending a point $p \in P(0)$ to the 0-corolla labelled by $\gamma_0(p) \in A(0)$.

The reader can check that the so obtained bimodule is well defined and that this construction works in the context of truncated bimodules. If A_r, B_r, C_r are r -truncated bimodules and f_1, f_2 are r -truncated bimodule maps, then the pushout in the category of r -truncated bimodules D_r is obtained from the formula (32) by taking the restriction of the coproduct to the trees with section having at most r leaves and such that each pearl has at most r incoming edges. The equivalence relation, the left and right operations and the map γ'_0 are defined in the same way.

Proposition 2.9 *One has the following identities:*

$$D = \operatorname{colim}_{\Sigma \text{Bimod}_{P,Q}} (B \leftarrow A \rightarrow C) \quad \text{and}$$

$$D_r = \operatorname{colim}_{T_r \Sigma \text{Bimod}_{P,Q}} (B_r \leftarrow A_r \rightarrow C_r).$$

Proof We need to check the universal property of the pushout in the category of $(P-Q)$ -bimodules. Let D' be a $(P-Q)$ -bimodule together with $(P-Q)$ -bimodule maps $g_1 : B \rightarrow D'$ and $g_2 : C \rightarrow D'$ such that $g_1 \circ f_1 = g_2 \circ f_2$. One has to show that there is a unique $(P-Q)$ -bimodule map $\delta : D \rightarrow D'$ such that the following diagram commutes:

$$\begin{array}{ccc}
 A & \xrightarrow{f_2} & C \\
 f_1 \downarrow & & \downarrow \\
 B & \longrightarrow & D \\
 & \searrow \delta & \swarrow g_2 \\
 & & D'
 \end{array}
 \quad (33)$$

Let $[(T; \sigma); \{m_p\}; \{p_v\}; \{q_v\}]$ be a point D where σ is the permutation labelling the leaves. Due to the pushout relations, we can assume that T is a reduced tree with section without vertices above the section. If the tree with section T has only one vertex (which is necessarily a pearl) indexed by m_r in B or C , then, due to the commutative diagram (33), δ must be defined as follows:

$$\delta([(T; \sigma); \{m_p\}; \{p_v\}; \{q_v\}]) = \begin{cases} g_1(m_r) \cdot \sigma & \text{if } m_r \in B, \\ g_2(m_r) \cdot \sigma & \text{if } m_r \in C. \end{cases}$$

If the tree with section T has more than 2 vertices, then the root of T is indexed by a point p_r in the operad P and the point $[(T; \sigma); \{m_p\}; \{p_v\}; \{q_v\}]$ has a decomposition of the form

$$(p_r([(T_1; id); \{m_1\}; \emptyset; \emptyset], \dots, [(T_{|v|}; id); \{m_{|v|}\}; \emptyset; \emptyset])) \cdot \sigma$$

where $T_1, \dots, T_{|v|}$ are corollas. Since δ is a bimodule map, one has

$$\begin{aligned} &\delta([(T; \sigma); \{m_p\}; \{p_v\}; \{q_v\}]) \\ &= (p_r(\delta([(T_1; id); \{m_1\}; \emptyset; \emptyset]), \dots, \delta([(T_{|v|}; id); \{m_{|v|}\}; \emptyset; \emptyset]))) \cdot \sigma. \end{aligned}$$

Thanks to the $(P-Q)$ -bimodule axioms, δ does not depend on the choice of the representative element and δ is a $(P-Q)$ -bimodule map. The uniqueness follows from the construction. The same arguments work for the truncated case. \square

Lemma 2.10 *Let $\partial X \rightarrow X$ be a morphism of ΣSeq_P which defines a closed inclusion of topological spaces objectwise. For every pushout diagram of the form*

$$\begin{CD} \mathcal{F}_{P;Q}^\Sigma(\partial X) @>>> \mathcal{F}_{P;Q}^\Sigma(X) \\ @VVV @VVV \\ B @>>> D, \end{CD} \tag{34}$$

the $(P-Q)$ -bimodule map $B \rightarrow D$ is also a closed inclusion of topological spaces objectwise.

Proof Let $s\mathbb{P}'_n$ denote the set of planar trees with section and with n leaves labelled by a permutation from Σ_n , whose set V^P of pearls is partitioned into two subsets $V^P = V^{P_{pri}} \sqcup V^{P_{aux}}$ of primary and auxiliary pearls, respectively. One has

$$\begin{aligned} D(n) &= \left(\coprod_{T \in s\mathbb{P}'_n} X(T) \right) / \sim, \text{ where} \\ X(T) &= \prod_{p \in V^{P_{pri}}(T)} B(|p|) \times \prod_{p \in V^{P_{aux}}(T)} X(|p|) \times \prod_{v \in V^d(T)} P(|v|) \times \prod_{v \in V^u(T)} Q(|v|). \end{aligned}$$

The relations are (i)-(vi), see above. We denote by π_T the map $\pi_T : X(T) \rightarrow D(n)$.

A closed injective map is always a topological inclusion. We prove first that each map $i : B(n) \rightarrow D(n)$, $n \geq 0$, is injective and then that it is closed.

As a set, $D(n)$ does not depend on the topology of X . So, we can choose a topology such that each $X(i) \setminus \partial X(i)$ is discrete and open in $X(i)$. In the latter case, one has $D = B \coprod \mathcal{F}_{P;Q}^\Sigma(P_0 \sqcup (X \setminus \partial X))$ and the inclusion of B in this coproduct is obviously injective. Indeed, a tree with a vertex labelled by $x \in X \setminus \partial X$ in the coproduct can never loose such vertex by means of relations (i)-(vi) of the coproduct, and therefore can not produce a relation in B .

Now we check that for any closed $C \subset B(n)$, the set $i(C)$ is closed in $D(n)$. Consider

$$X^\partial(T) := \prod_{p \in V^{P_{pri}}(T)} B(|p|) \times \prod_{p \in V^{P_{aux}}(T)} \partial X(|p|) \times \prod_{v \in V^d(T)} P(|v|) \times \prod_{v \in V^u(T)} Q(|v|).$$

Since each composition map $\pi_T^\partial : X^\partial(T) \rightarrow B(n)$ is continuous, the set $(\pi_T^\partial)^{-1}(C)$ is closed in $X^\partial(T)$. On the other hand, since each map $\partial X(|p|) \rightarrow X(|p|)$ is a closed inclusion, the map $X^\partial(T) \rightarrow X(T)$ is also one. (Here, we use the fact that kellyfication preserves closed inclusions.) Thus $(\pi_T^\partial)^{-1}(C) = (\pi_T)^{-1}(i(C))$ is closed in $X(T)$ for every T and we conclude that so is $i(C)$ in $D(n)$. \square

2.2 The Model Category Structure

By using the identifications (26), the categories ΣSeq_P and $T_r \Sigma Seq_P$ inherit model category structures from the categories of Σ -sequences and truncated Σ -sequences, respectively. More precisely, a map is a weak equivalence, a fibration or a cofibration if the corresponding map is a weak equivalence, a fibration or a cofibration in the category of (truncated) Σ -sequences. In particular, ΣSeq_P and $T_r \Sigma Seq_P$ are cofibrantly generated and all their objects are fibrant. Their (acyclic) generating cofibrations are $\{P_0 \sqcup \partial X \rightarrow P_0 \sqcup X\}$, where $\{\partial X \rightarrow X\}$ is the set of (acyclic) generating cofibrations of ΣSeq or $T_r \Sigma Seq$, respectively. By applying the transfer principle 1.1 to the adjunctions

$$\begin{aligned} \mathcal{F}_{P;Q}^\Sigma : \Sigma Seq_P &\rightleftarrows \Sigma \text{Bimod}_{P;Q} : \mathcal{U}^\Sigma & \text{and} \\ \mathcal{F}_{P;Q}^{T_r \Sigma} : T_r \Sigma Seq_P &\rightleftarrows T_r \Sigma \text{Bimod}_{P;Q} : \mathcal{U}^{T_r \Sigma}, \end{aligned} \tag{35}$$

we get the following statement:

Theorem 2.11 *For any pair (P, Q) of topological operads, the category of (truncated) (P, Q) -bimodules $\Sigma \text{Bimod}_{P;Q}$ (respectively, $T_r \Sigma \text{Bimod}_{P;Q}$, $r \geq 0$) inherits a cofibrantly generated model category structure, called the projective model category structure, in which all objects are fibrant. The model structure in question makes the adjunctions (35) into Quillen adjunctions. More precisely, a bimodule map f is a weak equivalence (respectively, a fibration) if and only if the induced map $\mathcal{U}^\Sigma(f)$ or $\mathcal{U}^{T_r \Sigma}(f)$ is a weak equivalence (respectively, a fibration) in the category of (possibly truncated) Σ -sequences.*

Proof According to the transfer principle 1.1, we have to check the small object argument as well as the existence of a functorial fibrant replacement and a functorial factorization of the diagonal map in the category $\Sigma \text{Bimod}_{P;Q}$.

We check first the small object argument. Let $\mathcal{F}_{P;Q}^\Sigma(X)$ be a domain of an element in the set of generating (acyclic) cofibrations in $\Sigma \text{Bimod}_{P;Q}$. Let $\lambda = \aleph_1$ be the first uncountable ordinal. Assume that $\{M_\alpha\}_{\alpha < \lambda}$ is a λ -sequence of (P, Q) -bimodules, such that each map $M_{<\alpha} := \text{colim}_{\beta < \alpha} M_\beta \rightarrow M_\alpha$ fits into a pushout square

$$\begin{array}{ccc} \mathcal{F}_{P;Q}^\Sigma(\partial Y) & \longrightarrow & \mathcal{F}_{P;Q}^\Sigma(Y) \\ \downarrow & & \downarrow \\ M_{<\alpha} & \longrightarrow & M_\alpha, \end{array} \tag{36}$$

where $\partial Y \rightarrow Y$ is a possibly infinite coproduct in ΣSeq_P of generating (acyclic) cofibrations. Due to the adjunction (35), one has the identity

$$\Sigma \text{Bimod}_{P;Q}(\mathcal{F}_{P;Q}^\Sigma(X); \text{colim}_{\alpha < \lambda} M_\alpha) \cong \Sigma Seq_P(X; \mathcal{U}(\text{colim}_{\alpha < \lambda} M_\alpha)).$$

Since the forgetful functor is monadic, it preserves filtered colimits. So, one has

$$\Sigma Seq_P(X; \mathcal{U}(\text{colim}_{\alpha < \lambda} M_\alpha)) \cong \Sigma Seq_P(X; \text{colim}_{\alpha < \lambda} \mathcal{U}(M_\alpha)).$$

It follows from Lemma 2.10 that $M_{<\alpha} \rightarrow M_\alpha$ is an objectwise closed inclusion. The sequence $X \setminus P_0$ is concentrated in only one arity where it is a finite union of spheres (or discs), thus a separable space. Therefore, X is \aleph_1 -small relative to componentwise closed inclusions¹ and

¹ In fact X is \aleph_0 -small relative to componentwise closed inclusions of T_1 -spaces, as X being compact can not contain a discrete countable closed subspace. Note, however, that in our category Top non- T_1 spaces are allowed.

one has the identities

$$\begin{aligned} &\Sigma Seq_P(X; \operatorname{colim}_{\alpha < \lambda} \mathcal{U}(M_\alpha)) \\ &\cong \operatorname{colim}_{\alpha < \lambda} \Sigma Seq_P(X; \mathcal{U}(M_\alpha)) \cong \operatorname{colim}_{\alpha < \lambda} \Sigma \operatorname{Bimod}_{P; Q}(\mathcal{F}_{P; Q}^\Sigma(X); M_\alpha). \end{aligned}$$

This proves the small object argument.

Since all objects are fibrant in the category of Σ -sequences, the identity functor provides a functorial fibrant replacement. For any $M \in \Sigma \operatorname{Bimod}_{P; Q}$, one needs to prove the existence of an element $Path(M) \in \Sigma \operatorname{Bimod}_{P; Q}$ inducing a factorization of the diagonal map

$$\Delta : M \xrightarrow[f_1]{\simeq} Path(M) \xrightarrow[f_2]{\twoheadrightarrow} M \times M,$$

where f_1 is a weak equivalence and f_2 is a Serre fibration. Let us consider

$$Path(M)(n) = \operatorname{Map}([0, 1]; M(n)).$$

This Σ -sequence inherits a bimodule structure from M . The map from M to $Path(M)$, sending a point to the constant path, is clearly a homotopy equivalence. Furthermore, the map

$$f_2 : \operatorname{Map}([0, 1]; M(n)) \longrightarrow \operatorname{Map}(\partial[0, 1]; M(n)) = (M \times M)(n)$$

induced by the inclusion $i : \partial[0, 1] \rightarrow [0, 1]$ is a Serre fibration since the map i is a cofibration. □

Alternative proof As explained in Sect. 2.1.1, the category of $(P-Q)$ -bimodules is equivalent to the category of algebras over a colored operad $P+Q$. According to the general result of [8, Theorem 2.1], the category of algebras over any topological operad has a projective model structure. To be precise, let $Seq = Top^{\mathbb{N}}$ be the projective model category of sequences of topological spaces. The adjunction between the forgetful functor and the free $(P+Q)$ -algebra functor

$$\mathcal{F}_{P+Q} : Seq \rightleftarrows \operatorname{Alg}_{P+Q} : \mathcal{U},$$

induces a cofibrantly generated model category structure on the category of $(P+Q)$ -algebras. In this model structure, a map of $(P+Q)$ -algebras f is a weak equivalence (respectively, a fibration) if the corresponding map $\mathcal{U}(f)$ is a weak equivalence (respectively, a fibration) in the category of sequences. This model structure coincides with the model structure described in the theorem because the projective model structure on ΣSeq is itself transferred from Seq . □

2.2.1 Relative Left Properness of the Projective Model Category

• *Definitions of relative left and right properness.* First, we recall the definition in a general setting. Let \mathcal{C} be a model category and let \mathcal{S} be a class of objects of \mathcal{C} . The model category \mathcal{C} is said to be left proper (respectively right proper) relative to \mathcal{S} if for each pushout diagram (respectively pullback diagram) of the form

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ \downarrow g & \simeq & \downarrow j \\ C & \xrightarrow{i} & D \end{array} \quad (\text{respectively, } \begin{array}{ccc} A & \xrightarrow{i} & B \\ \downarrow j & & \downarrow g \\ C & \xrightarrow[f]{\simeq} & D \end{array})$$

with g a cofibration (respectively, a fibration) and f a weak equivalence between objects in \mathcal{S} , the morphism i is also a weak equivalence. In particular, the category \mathcal{C} is said to be left proper (respectively right proper) if \mathcal{C} is left proper (respectively right proper) relative to all the objects. Furthermore, a model category \mathcal{C} is said to be proper (relative to \mathcal{S}) if the category is both left and right proper (relative to \mathcal{S}). The advantage of such a category is that we have a criterion allowing to identify homotopy invariant colimits or/and limits.

Proposition 2.12 [32, Proposition A.2.4.4] *Let \mathcal{C} be a model category which is left proper relative to a class of objects \mathcal{S} . For any commutative diagram in the category \mathcal{C} of the form*

$$\begin{array}{ccccc}
 A & \xleftarrow{f} & B & \xrightarrow{g} & C \\
 v_A \downarrow & & v_B \downarrow & & v_C \downarrow \\
 A' & \xleftarrow{f'} & B' & \xrightarrow{g'} & C'
 \end{array}$$

the induced map between the colimits of the horizontal diagrams

$$\text{colim}(A \leftarrow B \rightarrow C) \longrightarrow \text{colim}(A' \leftarrow B' \rightarrow C')$$

is a weak equivalence if the vertical morphisms are weak equivalences; one of the pairs (f, C) or (g, A) consists of a cofibration and an object in \mathcal{S} ; one of the pairs (f', C') or (g', A') consists of a cofibration and an object in \mathcal{S} .

Dually, let \mathcal{C} be a model category which is right proper relative to a class \mathcal{S} . For any commutative diagram in the category \mathcal{C} of the form

$$\begin{array}{ccccc}
 A & \xrightarrow{f} & B & \xleftarrow{g} & C \\
 v_A \downarrow & & v_B \downarrow & & v_C \downarrow \\
 A' & \xrightarrow{f'} & B' & \xleftarrow{g'} & C'
 \end{array}$$

the induced map between the limits of the horizontal diagrams

$$\text{lim}(A \rightarrow B \leftarrow C) \longrightarrow \text{lim}(A' \rightarrow B' \leftarrow C')$$

is a weak equivalence if the vertical morphisms are weak equivalences; one of the pairs (f, C) or (g, A) consists of a fibration and an object in \mathcal{S} ; one of the pairs (f', C') or (g', A') consists of a fibration and an object in \mathcal{S} .

Remark 2.13 Even when the category \mathcal{C} is not left or right proper, the statement of Proposition 2.12 still holds provided one of the pairs $(f; C)$ or $(g; A)$ consists of a (co)fibration and a (co)fibrant object whereas one of the pairs $(f'; C')$ or $(g'; A')$ consists of a (co)fibration and a (co)fibrant object.

• *Application to the projective model category of bimodules.* Let P and Q be two topological operads. From now on, we focus our attention on the projective model category of $(P\text{-}Q)$ -bimodules. More precisely, we show that this category is right proper relative to all the objects and is relatively left proper.

Theorem 2.14 *The projective model category $\Sigma\text{Bimod}_{P; Q}$ is right proper.*

Proof We consider the following pullback diagram in which f is a weak equivalence and g is a fibration:

$$A = \lim (C \xrightarrow[\simeq]{f} D \xleftarrow{g} B).$$

The map $i : A \rightarrow B$ is equivalent to the following map between pullback diagrams:

$$\begin{array}{ccc} A & \xlongequal{\quad} & \lim (C \xrightarrow[\simeq]{} D \xleftarrow{\quad} B) \\ \downarrow i & & \downarrow \simeq \\ B & \xlongequal{\quad} & \lim (D \xlongequal{\quad} D \xleftarrow{\quad} B) \end{array}$$

According to Remark 2.13 and since all the objects in the projective model category of bimodules are fibrant, the map $i : A \rightarrow B$ is also a weak equivalence. Consequently, the category $\Sigma\text{Bimod}_{P; Q}$ is right proper relative to all the objects. \square

Lemma 2.15 *Let $C \leftarrow A \rightarrow B$ be a diagram in a cocomplete category \mathcal{C} . If $A \rightarrow C$ is a retract of $A' \rightarrow C'$, then the morphism $C \rightarrow \text{colim}(C \leftarrow A \rightarrow B)$ is a retract of $C' \rightarrow \text{colim}(C' \leftarrow A' \rightarrow A \rightarrow B)$.*

Proof Since $A \rightarrow C$ is a retract of $A' \rightarrow C'$, one has a commutative diagram

$$\begin{array}{ccccc} C & \xrightarrow{g_1} & C' & \xrightarrow{g_2} & C \\ \uparrow & & \uparrow & & \uparrow \\ A & \xrightarrow{h_1} & A' & \xrightarrow{h_2} & A \\ \downarrow & & \downarrow & & \downarrow \\ B & \xlongequal{\quad} & B & \xlongequal{\quad} & B \end{array} \tag{37}$$

such that $g_2 \circ g_1 = id$ and $h_2 \circ h_1 = id$. By taking the pushout of the vertical diagrams in (37), we get the commutative diagram

$$\begin{array}{ccccc} C & \xrightarrow{g_1} & C' & \xrightarrow{g_2} & C \\ \downarrow & & \downarrow & & \downarrow \\ C \sqcup_A B & \xrightarrow[k_1=g_1 \sqcup_{h_1} id]{} & C' \sqcup_{A'} B & \xrightarrow[k_2=g_2 \sqcup_{h_2} id]{} & C \sqcup_A B \end{array}$$

in which $g_2 \circ g_1 = id$ and $k_2 \circ k_1 = id$. \square

Theorem 2.16 *If P and Q are operads, such that $P(0) \in Top$ is cofibrant, $P_{>0} \in \Sigma\text{Operad}$ is a cofibrant operad, and Q is a componentwise cofibrant operad, then the projective model category $\Sigma\text{Bimod}_{P; Q}$ is left proper relative to the class \mathcal{S} of componentwise cofibrant bimodules M for which the arity zero left action map $\gamma_0 : P(0) \rightarrow M(0)$ is a cofibration.*

Corollary 2.17 *If Q is a componentwise cofibrant operad and P is either a Reedy cofibrant or projectively cofibrant operad, then the projective model category $\Sigma\text{Bimod}_{P; Q}$ is relatively left proper.*

Proof An operad is Reedy cofibrant if and only if it is reduced and its positive arity part is cofibrant. This proves the first case. The second case follows from Proposition 1.8. \square

Proof of Theorem 2.16 We first prove this theorem assuming $P(0) = \emptyset$ and at the end we explain how the argument has to be adjusted to the general case $P(0) \neq \emptyset$.

In order to prove that the category is left proper relative to \mathcal{S} , we consider the following pushout diagram in the category of $(P\text{-}Q)$ -bimodules in which the map f is a weak equivalence between componentwise cofibrant objects and the map g is a cofibration:

$$D = \operatorname{colim} (C \xleftarrow{g} \circlearrowleft A \xrightarrow{\simeq} B).$$

Since any cofibration is a retract of a cellular extension, we can assume that g is a cellular extension due to Lemma 2.15. On the other hand, any cellular extension is a possibly transfinite sequence of cell attachments. By Theorem 2.18a, cofibrations with domain in \mathcal{S} are componentwise cofibrations. Thus, without loss of generality, we can assume that g is a cellular attachment and we restrict our study to diagrams of the form

$$\begin{array}{ccccc} \mathcal{F}_{P;Q}^\Sigma(\partial X) & \longrightarrow & A & \longrightarrow & B \\ \downarrow & & \downarrow & & \downarrow \\ \mathcal{F}_{P;Q}^\Sigma(X) & \longrightarrow & C & \longrightarrow & D, \end{array}$$

where $\partial X \rightarrow X$ is a generating cofibration in the category of Σ -sequences and both squares are pushout diagrams. The strategy is to use the explicit description of the pushout from Sect. 2.1.3 and to introduce a filtration in Σ -sequences C and D according to the number of vertices in the trees indexed by X :

$$\begin{array}{ccccccccccccccc} A = C_0 & \longrightarrow & C_1 & \longrightarrow & \cdots & \longrightarrow & C_{i-1} & \longrightarrow & C_i & \longrightarrow & \cdots & \longrightarrow & C & (38) \\ \downarrow & & \downarrow & & & & \downarrow & & \downarrow & & & & \downarrow & \\ B = D_0 & \longrightarrow & D_1 & \longrightarrow & \cdots & \longrightarrow & D_{i-1} & \longrightarrow & D_i & \longrightarrow & \cdots & \longrightarrow & D. \end{array}$$

We will prove that in (38), each horizontal map is a componentwise cofibration and each vertical map is a weak equivalence.

Let $rs\mathbb{P}_n[i]$ denote the set of reduced planar trees with section with n leaves labelled by a permutation in Σ_n . In addition we assume that it has two types of pearls: i auxiliary ones and some number of primary ones, primary ones coming first. Moreover, it is required that all incoming edges of any primary pearl are leaf edges, while all incoming edges of any auxiliary pearl connects it to an internal vertex.

It follows from Construction 2.7 of a free bimodule and the combinatorial description of a pushout (Sect. 2.1.3) that any element in $C(n)$ or $D(n)$ can be obtained as a tree $T \in rs\mathbb{P}_n[i]$ (for some $i \geq 0$), whose root, primary pearls, auxiliary pearls, and vertices above the section are labelled by P , A (or B), X , and Q , respectively. The filtration (38) is by the number i of auxiliary pearls in such elements. To show that maps $C_{i-1}(n) \rightarrow C_i(n)$ and $D_{i-1}(n) \rightarrow D_i(n)$ are cofibrations, we additionally filter this inclusion by the total number ℓ of pearls:

$$\begin{array}{ccccccccccccccc} C_{i-1}(n) =: C_i(n)_{i-1} & \longrightarrow & C_i(n)_i & \longrightarrow & \cdots & \longrightarrow & C_i(n)_{\ell-1} & \longrightarrow & C_i(n)_\ell & \longrightarrow & \cdots & \longrightarrow & C_i(n) \\ \downarrow & & \downarrow & & & & \downarrow & & \downarrow & & & & \downarrow \\ D_{i-1}(n) =: D_i(n)_{i-1} & \longrightarrow & D_i(n)_i & \longrightarrow & \cdots & \longrightarrow & D_i(n)_{\ell-1} & \longrightarrow & D_i(n)_\ell & \longrightarrow & \cdots & \longrightarrow & D_i(n) \end{array} \quad (39)$$

We will show that each horizontal inclusion is a cofibration and each vertical map is an equivalence, which would prove the theorem.

The maps $C_i(n)_{\ell-1} \rightarrow C_i(n)_\ell$ and $D_i(n)_{\ell-1} \rightarrow D_i(n)_\ell$ are cofibrations. For $\ell \geq i$, let $rs\mathbb{P}_n[i; \ell] \subset rs\mathbb{P}_n[i]$ be the subset of planar trees with section with exactly ℓ pearls (see Fig. 10). Let $rs\mathbb{T}_n[i; \ell] := rs\mathbb{P}_n[i; \ell]/\sim$ be the set of equivalence classes, where two trees are equivalent if they are isomorphic as non-planar trees by an isomorphism that forgets the labels of leaves and that sends primary pearls to primary ones and same with auxiliary ones.

For $T \in rs\mathbb{T}_n[i; \ell]$, denote by $Aut(T)$ the group of automorphisms of T that does not mix the auxiliary pearls with the primary ones (only arity zero ones can potentially be permuted, but we do not allow it). A choice of representative of T in $rs\mathbb{P}_n[i; \ell]$ gives a labeling of leaves by the set $[n]$ and therefore induces a homomorphism $Aut(T) \rightarrow \Sigma_n$. Denote by $Aut_0(T)$ the kernel of this homomorphism which is the subgroup of $Aut(T)$ permuting the univalent vertices and the auxiliary pearls that have only univalent vertices above them. Let p_1, \dots, p_ℓ be the pearls of the chosen planar representative of T labelled in the planar order they appear.

Then we consider the spaces

$$M_1(T) = \prod_{\ell-i+1 \leq j \leq \ell} X(|p_j|) \times \prod_{v \in V^u(T)} Q(|v|),$$

where $V^u(T)$ is the set of vertices of T above the section, and


$$M_2(T; A) = P(\ell) \times_{P(1)^{\ell-i}} \prod_{1 \leq j \leq \ell-i} A(|p_j|) \\ = \text{coeq} \left(P(\ell) \times P(1)^{\ell-i} \times \prod_{1 \leq j \leq \ell-i} A(|p_j|) \rightrightarrows P(\ell) \times \prod_{1 \leq j \leq \ell-i} A(|p_j|) \right).$$

The two arrows in the coequalizer correspond respectively to the action of $P(1)^{\ell-i}$ on the first $(\ell - i)$ inputs of $P(\ell)$ (by operadic composition) and on $\prod_{1 \leq j \leq \ell-i} A(|p_j|)$ (by the left P -action on A).

In other words, $M_1(T)$ is the space of indexations of the i last pearls by points in X and the vertices above the section by points in the operad Q . Similarly, $M_2(T; A)$ is the space of indexations of the other pearls by points in A and the root by a point in $P(\ell)$. This presentation is not unique, that is why we take the coequalizer.

Moreover, we denote by $M_1^-(T)$ the subspace of $M_1(T)$ formed by points having at least one pearl indexed by a point in ∂X . The space $M_2^-(T; A)$ consists of points in $M_2(T; A)$ for which the root is indexed by a point $p \in P(\ell; \ell - i) \subset P(\ell)$ (see notation before

Fig. 10 Illustration of a tree in $rs\mathbb{P}_9[4; 7] \subset rs\mathbb{P}_9[4]$ labelled by P, A, X and Q



Proposition 1.10). We define

$$M(T; A) := M_1(T) \times M_2(T; A) \quad \text{and}$$

$$M^-(T; A) := M_1^-(T) \times M_2^-(T; A) \quad \coprod_{M_1^-(T) \times M_2^-(T; A)} M_1^-(T) \times M_2(T; A).$$

The spaces $M_2(T; B)$, $M_2^-(T; B)$, $M(T; B)$, $M^-(T; B)$ are defined similarly. One has the following pushout squares

$$\begin{array}{ccc}
 \coprod_{T \in rs\mathbb{T}_n[i; \ell]} M^-(T; A) \times_{Aut(T)} \Sigma_n & \longrightarrow & C_i(n)_{\ell-1} \\
 \downarrow & & \downarrow \\
 \coprod_{T \in rs\mathbb{T}_n[i; \ell]} M(T; A) \times_{Aut(T)} \Sigma_n & \longrightarrow & C_i(n)_\ell
 \end{array}
 \qquad
 \begin{array}{ccc}
 \coprod_{T \in rs\mathbb{T}_n[i; \ell]} M^-(T; B) \times_{Aut(T)} \Sigma_n & \longrightarrow & D_i(n)_{\ell-1} \\
 \downarrow & & \downarrow \\
 \coprod_{T \in rs\mathbb{T}_n[i; \ell]} M(T; B) \times_{Aut(T)} \Sigma_n & \longrightarrow & D_i(n)_\ell
 \end{array}
 \tag{40}$$

To prove that the right vertical maps are cofibrations one needs to show that the left ones are such. Recall that $Aut_0(T)$ denotes the kernel of $Aut(T) \rightarrow \Sigma_n$. Equivalently, for every $T \in rs\mathbb{T}_n[i; \ell]$ we need to show that the maps

$$\begin{aligned}
 M^-(T; A)/Aut_0(T) &\rightarrow M(T; A)/Aut_0(T) \quad \text{and} \\
 M^-(T; B)/Aut_0(T) &\rightarrow M(T; B)/Aut_0(T)
 \end{aligned}
 \tag{41}$$

are cofibrations. We also need later that the sources of the maps are cofibrant.

Let G_1 be the subgroup of $Aut_0(T)$ that fixes all the pearls. It permutes the arity zero vertices above the section. The subgroup G_1 is normal and the quotient group $G_2 = Aut_0(T)/G_1$ is responsible for the permutations of the arity zero primary pearls and the auxiliary pearls that have only arity zero vertices above them (in particular they can be of arity zero themselves). Note that $G_2 \subset \Sigma_{\ell-i} \times \Sigma_i$. By construction, the spaces $M_1^-(T)$, $M_1(T)$, $M_2^-(T; A)$ and $M_2(T; A)$ are equipped with an action of $Aut(T)$ and of $Aut_0(T)$ by taking restriction. Moreover, for the last two, the $Aut_0(T)$ -action factors through G_2 .

The inclusion $M_1^-(T) \rightarrow M_1(T)$ is an $Aut_0(T)$ -equivariant G_1 -cofibration. Indeed, $G_1 = \prod_{j=\ell-i+1}^{\ell} \Sigma_{d_j}$, where d_j is the number of univalent vertices above the j -th pearl. Applying iteratively Lemma A.1, we get that the map

$$\partial \prod_{\ell-i+1 \leq j \leq \ell} X(\lfloor p_j \rfloor) \rightarrow \prod_{\ell-i+1 \leq j \leq \ell} X(\lfloor p_j \rfloor)
 \tag{42}$$

is a G_1 -cofibration. On the other hand, since Q is componentwise cofibrant, the space

$$\prod_{v \in V^u(T)} Q(\lfloor v \rfloor)$$

is a cofibrant (in *Top*) G_1 -space. Applying again Lemma A.1, we get that $M_1^-(T) \rightarrow M_1(T)$ is a G_1 -cofibration.

The source of the map (42) is also G_1 -cofibrant as it can be obtained as a sequence of G_1 -cofibrations

$$Y_1 \rightarrow Y_2 \rightarrow \dots \rightarrow Y_i,$$

where

$$Y_m = \partial \left(\prod_{\ell-i+1 \leq j \leq \ell-i+m} X(|p_j|) \right) \times \prod_{\ell-i+m+1 \leq j \leq \ell} \partial X(|p_j|).$$

Note that $Y_1 = \prod_{\ell-i+1 \leq j \leq \ell} \partial X(|p_j|)$ is G_1 -cofibrant being a product of Σ_{d_j} -cofibrant spaces (see Example A.6). We finally conclude that $M_1^-(T)$ is $Aut_0(T)$ -cofibrant.

Besides, we claim that the inclusion from $M_2^-(T; A)$ into $M_2(T; A)$ is a G_2 -cofibrant map between G_2 -cofibrant spaces. The assignment

$$R \mapsto R \times_{P(1)^{\ell-i}} \prod_{1 \leq j \leq \ell-i} A(|p_j|)$$

can be viewed as a functor from $(\Sigma_{\ell-i} \wr P(1)) \times \Sigma_i\text{-Top}$ to $G_2\text{-Top}$. This functor preserves cofibrations. Indeed, it preserves colimits and sends any generating cofibration $S^{k-1} \times (\Sigma_{\ell-i} \wr P(1)) \times \Sigma_i \rightarrow D^k \times (\Sigma_{\ell-i} \wr P(1)) \times \Sigma_i$ to a G_2 -cofibration:

$$\left(S^{k-1} \times \Sigma_{\ell-i} \times \prod_{1 \leq j \leq \ell-i} A(|p_j|) \right) \times \Sigma_i \rightarrow \left(D^k \times \Sigma_{\ell-i} \times \prod_{1 \leq j \leq \ell-i} A(|p_j|) \right) \times \Sigma_i.$$

The map above is a G_2 -cofibration by Lemmas A.1 and A.3 (recall that $G_2 \subset \Sigma_{\ell-i} \times \Sigma_i$). On the other hand, since P is a cofibrant operad, by Proposition 1.10, the inclusion $P(\ell; \ell-i) \rightarrow P(\ell)$ is a $(\Sigma_{\ell-i} \wr P(1)) \times \Sigma_i$ -cofibration with a $(\Sigma_{\ell-i} \wr P(1)) \times \Sigma_i$ -cofibrant source. We conclude that $M_2^-(T; A) \rightarrow M_2(T; A)$ is a G_2 -cofibrant map between G_2 -cofibrant spaces.

Now combining that $M_1^-(T) \rightarrow M_1(T)$ is an $Aut_0(T)$ -equivariant G_1 -cofibration and that $M_2^-(T; A) \rightarrow M_2(T; A)$ is a G_2 -cofibration and applying Lemma A.1, we get that $M^-(T; A) \rightarrow M(T; A)$ is an $Aut_0(T)$ -cofibration. (We get a similar statement for B as well by replacing A with B in all formulas.)

We can similarly conclude that $M_1^-(T) \times M_2^-(T; A)$ is $Aut_0(T)$ -cofibrant and the inclusions

$$M_1^-(T) \times M_2^-(T; A) \rightarrow M_1^-(T) \times M_2(T; A) \rightarrow M^-(T; A)$$

are $Aut_0(T)$ -cofibrations. As a result, $M^-(T; A)$ and (arguing similarly) $M^-(T; B)$ are $Aut_0(T)$ -cofibrant.

The map $C_i(n)_\ell \rightarrow D_i(n)_\ell$ is a weak equivalence. By induction we assume that the map $C_i(n)_{\ell-1} \rightarrow D_i(n)_{\ell-1}$ is a weak equivalence of cofibrant spaces. The spaces $C_i(n)_\ell$ and $D_i(n)_\ell$ are the colimits of the first and second lines, respectively, in the following diagram.

$$\begin{array}{ccccc} \coprod_{T \in rs\mathbb{T}_n[i; \ell]} M(T; A) \times_{Aut(T)} \Sigma_n & \longleftarrow & \coprod_{T \in rs\mathbb{T}_n[i; \ell]} M^-(T; A) \times_{Aut(T)} \Sigma_n & \longrightarrow & C_i(n)_{\ell-1} \\ \downarrow & & \downarrow & & \downarrow \\ \coprod_{T \in rs\mathbb{T}_n[i; \ell]} M(T; B) \times_{Aut(T)} \Sigma_n & \longleftarrow & \coprod_{T \in rs\mathbb{T}_n[i; \ell]} M^-(T; B) \times_{Aut(T)} \Sigma_n & \longrightarrow & D_i(n)_{\ell-1} \end{array}$$

As it follows from Proposition 2.12, the induced map of colimits is an equivalence provided all vertical maps are equivalences. We are left to showing that the left two arrows are such.

In other words, it only remains to prove that for every tree $T \in rs\mathbb{T}_n[i; \ell]$, the maps

$$M^-(T; A) \rightarrow M^-(T; B) \quad \text{and} \quad M(T; A) \rightarrow M(T; B)$$

are weak equivalences (as we already proved that all the four spaces above are $Aut_0(T)$ -cofibrant). The latter fact is deduced from the fact that the induced maps

$$M_2^-(T; A) \rightarrow M_2^-(T; B) \quad \text{and} \quad M_2(T; A) \rightarrow M_2(T; B)$$

are equivalences of cofibrant spaces. The latter is proved by Lemma A.7 for which we take $\Gamma = P(1)^{\ell-i}$, $Z = P(\ell)$ or $P(\ell; \ell - i)$, and the corresponding Γ^{op} -equivariant equivalence between cofibrant (in Top) objects is the map $\prod_{j=1}^{\ell-i} A(|p_j|) \rightarrow \prod_{j=1}^{\ell-i} B(|p_j|)$.

Proving the case $P(0) \neq \emptyset$. The only difference with the case $P(0) = \emptyset$ is that we need to take into account the γ_0 -relation contracting the univalent pearls indexed by points coming from the image of $P(0)$. For this purpose, we change slightly the definition of the spaces $M_2^-(T; A)$ and $M_2^-(T; B)$. Thus one also needs to check the three statements below.

- $M_2^-(T; A)$ and $M_2^-(T; B)$ are G_2 -cofibrant.
- $M_2^-(T; A) \rightarrow M_2(T; A)$ and $M_2^-(T; B) \rightarrow M_2(T; B)$ are G_2 -cofibrations.
- $M_2^-(T; A) \rightarrow M_2^-(T; B)$ is a weak equivalence.

Denote by $\vec{A} := \prod_{j=1}^{\ell-i} A(|p_j|)$ and by $\partial\vec{A} \subset \vec{A}$ the subset consisting of points with at least one coordinate in $\gamma_0(P(0)) \subset A(0)$. One has

$$M_2^-(T; A) = \left(P(\ell) \times_{P(1)^{\ell-i}} \partial\vec{A} \right) \coprod_{P(\ell; \ell-i) \times_{P(1)^{\ell-i}} \partial\vec{A}} \left(P(\ell; \ell - i) \times_{P(1)^{\ell-i}} \vec{A} \right).$$

By the same argument as above, the functor

$$(-) \times_{P(1)^{\ell-i}} \partial\vec{A}: (\Sigma_{\ell-i} \wr P(1)) \times \Sigma_i\text{-Top} \rightarrow G_2\text{-Top}$$

preserves cofibrations. As a consequence $P(\ell; \ell - i) \times_{P(1)^{\ell-i}} \partial\vec{A}$ is G_2 -cofibrant and the inclusion $P(\ell; \ell - i) \times_{P(1)^{\ell-i}} \partial\vec{A} \rightarrow P(\ell) \times_{P(1)^{\ell-i}} \partial\vec{A}$ is a G_2 -cofibration. Moreover, we know that $P(\ell; \ell - i) \times_{P(1)^{\ell-i}} \vec{A}$ is G_2 -cofibrant. We conclude that $M_2^-(T; A)$ is so.

The fact that $M_2^-(T; A) \rightarrow M_2(T; A)$ is a G_2 -cofibration follows from Lemma A.8 applied to $\Gamma = P(1)^{\ell-i}$, $K_1 = 1$, $K = K_2 = G_2$, $A \rightarrow B$ being $P(\ell; \ell - i) \rightarrow P(\ell)$, and $X \rightarrow Y$ being $\partial\vec{A} \rightarrow \vec{A}$.

Finally, to prove that $M_2^-(T; A) \rightarrow M_2^-(T; B)$ is a weak equivalence, we apply Lemma A.7 again to show that

$$\begin{aligned} P(\ell; \ell - i) \times_{P(1)^{\ell-i}} \partial\vec{A} &\rightarrow P(\ell; \ell - i) \times_{P(1)^{\ell-i}} \partial\vec{B}, \\ P(\ell) \times_{P(1)^{\ell-i}} \partial\vec{A} &\rightarrow P(\ell) \times_{P(1)^{\ell-i}} \partial\vec{B}, \\ P(\ell; \ell - i) \times_{P(1)^{\ell-i}} \vec{A} &\rightarrow P(\ell; \ell - i) \times_{P(1)^{\ell-i}} \vec{B}, \end{aligned}$$

are all the three weak equivalences. The statement follows. □

Theorem 2.18 (a) *Let P , Q and S be as in Theorem 2.16. In the category $\Sigma\text{Bimod}_P; Q$, cofibrations with domain in S are componentwise cofibrations. In particular, the class \mathcal{S} of objects is closed under cofibrations and cofibrant $(P-Q)$ -bimodules are always componentwise cofibrant.*

(b) *Assume additionally that Q is Σ -cofibrant. In the category $\Sigma\text{Bimod}_P; Q$, cofibrations with domain in the subclass $S^\Sigma \subset S$ of Σ -cofibrant objects, are Σ -cofibrations. The class S^Σ is closed under cofibrations and cofibrant $(P-Q)$ -bimodules are always Σ -cofibrant.*

Proof Let A be a componentwise cofibrant $(P-Q)$ -bimodule. It is enough to check that any cellular extension $A \rightarrow C$ is a componentwise cofibration (respectively, Σ -cofibration). In the proof of Theorem 2.16 we define a filtration (38) in C in which, as we proved, every inclusion $C_{i-1} \rightarrow C_i$ is a componentwise cofibration.

Additionally assuming that Q and A are Σ -cofibrant, one can show that any such inclusion is a Σ -cofibration. The strategy is the same to filter this inclusion as in (39) and then to show that the left vertical map in the first square (40) is a Σ_n -cofibration. The latter is proved by showing that each inclusion $M^-(T; A) \rightarrow M(T; A)$ is an $Aut(T)$ -cofibration. Let $H_1 \subset Aut(T)$ be the subgroup of elements that fix all the pearls and the leaves above the primary pearls. This group permutes vertices above the section and their leaves. It is normal and the quotient group $H_2 = Aut(T)/H_1$ is responsible for permuting the pearls and leaves above the primary pearls. By essentially the same argument, the inclusion $M_1^-(T) \rightarrow M_1(T)$ is an $Aut(T)$ -equivariant H_1 -cofibration and the inclusion $M_2^-(T; A) \rightarrow M_2(T; A)$ is an H_2 -cofibration (by Lemma A.8 applied for $\Gamma = P(1)^{\ell-i}$, $K = H_2$, $K_1 \subset H_2$ – the subgroup of permutations of leaves above primary pearls). The proof is completed by invoking Lemma A.1. □

2.2.2 Extension/Restriction Adjunction for the Projective Model Category of Bimodules

Let $\phi_1 : P \rightarrow P'$ and $\phi_2 : Q \rightarrow Q'$ be maps of operads. Similarly to the category of algebras (see Theorem 1.7), we show that the projective model categories of $(P-Q)$ -bimodules and of $(P'-Q')$ -bimodules are Quillen equivalent under some conditions on the maps ϕ_1 and ϕ_2 . For this purpose, we recall the constructions of the *restriction* functor ϕ^* and the *extension* functor $\phi_!$ in the context of bimodules:

$$\phi_! : \Sigma\text{Bimod}_{P; Q} \rightleftarrows \Sigma\text{Bimod}_{P'; Q'} : \phi^*.$$

- *The restriction functor.* The restriction functor ϕ^* sends a $(P'-Q')$ -bimodule M to the $(P-Q)$ -bimodule $\phi^*(M) = \{\phi^*(M)(n) = M(n), n \geq 0\}$ in which the $(P-Q)$ -bimodule structure is defined as follows using the $(P'-Q')$ -bimodule structure of M :


$$\begin{aligned} \circ^i : \phi^*(M)(n) \times Q(m) &\longrightarrow \phi^*(M)(n + m - 1); \\ x; q &\longmapsto x \circ^i \phi_2(q), \end{aligned}$$

$$\begin{aligned} \gamma_\ell : P(n) \times \phi^*(M)(m_1) \times \dots \times \phi^*(M)(m_n) &\longrightarrow \phi^*(M)(m_1 + \dots + m_n); \\ p; x_1, \dots, x_n &\longmapsto \phi_1(p)(x_1, \dots, x_n). \end{aligned}$$


- *The extension functor.* The extension functor $\phi_!$ is obtained as a quotient of the free $(P'-Q')$ -bimodule functor introduced in Sect. 2.1.2. More precisely, if M is a $(P-Q)$ -bimodule, then the extension functor is given by the formula

$$\phi_!(M)(n) = \mathcal{F}_{P'; Q'}^\Sigma(\mathcal{U}^\Sigma(M))(n) / \sim$$

where the equivalence relation is generated by the axiom which consists in contracting inner edges having a vertex v below the section (respectively above the section) indexed by a point of the form $\phi_1(p)$ (respectively a point of the form $\phi_2(q)$) using the left P -module structure (respectively the right Q -module structure) of M as illustrated in the following picture:



The $(P'-Q')$ -bimodule structure on the free object is compatible with the equivalence relation and provides a $(P'-Q')$ -bimodule structure on $\phi_!(M)$. Let us remark that the $(P-Q)$ -bimodule map $M \rightarrow \phi^*(\phi_!(M))$, sending a point $x \in M(n)$ to the n -corolla indexed by x , is not necessarily injective. For instance, if there are $q_1 \neq q_2$ in $Q(m)$ and $x \in M(n)$ such that $\phi_2(q_1) = \phi_2(q_2)$ and $x \circ^i q_1 \neq x \circ^i q_2$, then $x \circ^i q_1$ and $x \circ^i q_2$ have the same image in $\phi^*(\phi_!(M))$ as illustrated in the following picture:



Theorem 2.19 Let $\phi_1 : P \rightarrow P'$ be a weak equivalence between Σ -cofibrant operads and $\phi_2 : Q \rightarrow Q'$ be a weak equivalence between componentwise cofibrant operads. The extension and restriction functors, as well as their truncated versions, give rise to Quillen equivalences:

$$\phi_1 : \Sigma\text{Bimod}_{P; Q} \rightleftarrows \Sigma\text{Bimod}_{P'; Q'} : \phi^*, \tag{43}$$

$$\phi_1 : T_r \Sigma\text{Bimod}_{P; Q} \rightleftarrows T_r \Sigma\text{Bimod}_{P'; Q'} : \phi^*. \tag{44}$$

Lemma 2.20 Any pair of operadic maps (ϕ_1, ϕ_2) , with $\phi_1 : P \rightarrow P'$ a weak equivalence between Σ -cofibrant operads and $\phi_2 : Q \rightarrow Q'$ a weak equivalence between componentwise cofibrant operads, induces a weak equivalence $\phi_+ : P+Q \rightarrow P'+Q'$ between Σ -cofibrant colored operads where $P+Q$ and $P'+Q'$ are colored operads obtained from Construction 2.4.

Proof According to Remark 2.5, for any family of integers n_1, \dots, n_k and m , the spaces $(P+Q)(n_1, \dots, n_k; m)$ and $(P'+Q')(n_1, \dots, n_k; m)$ have the following description:

$$\begin{aligned} (P+Q)(n_1, \dots, n_k; m) &\cong P(k) \times Q_1(n_1 + \dots + n_k; m) \cong P(k) \\ &\times \coprod_{\alpha: [m] \rightarrow [n_1 + \dots + n_k]} \prod_{i \in [n_1 + \dots + n_k]} Q(|\alpha^{-1}(i)|), \\ (P'+Q')(n_1, \dots, n_k; m) &\cong P'(k) \times Q'_1(n_1 + \dots + n_k; m) \cong P'(k) \\ &\times \coprod_{\alpha: [m] \rightarrow [n_1 + \dots + n_k]} \prod_{i \in [n_1 + \dots + n_k]} Q'(|\alpha^{-1}(i)|). \end{aligned}$$

Consequently, the weak equivalences $\phi_1 : P \rightarrow P'$ and $\phi_2 : Q \rightarrow Q'$ induce a weak equivalence of colored operads $\phi_+ : P+Q \rightarrow P'+Q'$. Let Σ' be the subgroup Σ_k which can send i to j if and only if $n_i = n_j$. This group acts as a subgroup of Σ_k on the factor $P(k)$ (respectively the factor $P'(k)$) and it acts on the factor $Q_1(n_1 + \dots + n_k; m)$ (respectively, the factor $Q'_1(n_1 + \dots + n_k; m)$) by reordering the summands in the disjoint union labelled by maps $\alpha : [m] \rightarrow [n_1 + \dots + n_k]$. The reordering is induced by permutation of the blocks $\{1, \dots, n_1\}, \dots, \{n_1 + \dots + n_{k-1} + 1, \dots, n_1 + \dots + n_k\}$ in $[n_1 + \dots + n_k]$. Consequently, the colored operads $P+Q$ and $P'+Q'$ are Σ -cofibrant as soon as the operads P and P' are Σ -cofibrant and the components of Q and Q' are cofibrant. \square

Proof of Theorem 2.19 As explained in Sect. 2.1.1, the projective model category of bimodules over an operad is equivalent to the projective model category of algebras over a specific colored operad. If we denote by $P+Q$ and $P'+Q'$ the colored operads obtained from Construction 2.4, then one has

$$\phi_! : \Sigma \text{Bimod}_{P; Q} = \text{Alg}_{P+Q} \rightleftarrows \text{Alg}_{P'+Q'} = \Sigma \text{Bimod}_{P'; Q'} : \phi^*,$$

induced by the extension/restriction adjunction between the categories of algebras. Due to Lemma 2.20, the induced map $\phi_+ : P+Q \rightarrow P'+Q'$ is a weak equivalence between Σ -cofibrant colored operads. Consequently, according to [6, Theorem 4.4] the extension/restriction adjunction between the categories of algebras is a Quillen equivalence. \square

• *Bimodules with the empty set in arity zero.* Consider the case where the acting operads P and Q are trivial in arity zero $P(0) = Q(0) = \emptyset$ and consider the full subcategory $\Sigma_{>0} \text{Bimod}_{P; Q}$ of $(P-Q)$ -bimodules M also satisfying $M(0) = \emptyset$. One can similarly to Construction 2.4 define a colored operad $(P+Q)_{>0}$ that governs this algebraic structure. Its set of colors is the set $\mathbb{N}_{>0}$ of positive integers. Its components can be similarly described:

$$(P+Q)_{>0}(n_1, \dots, n_k; m) = P(k) \times \coprod_{\alpha: [m] \rightarrow [n_1 + \dots + n_k]} \prod_{i \in [n_1 + \dots + n_k]} Q(|\alpha^{-1}(i)|). \tag{45}$$

The crucial difference is that the coproduct is taken over surjective maps $\alpha : [m] \rightarrow [n_1 + \dots + n_k]$.

Proposition 2.21 *Let P and Q be componentwise cofibrant operads satisfying $P(0) = Q(0) = \emptyset$, then the colored operad $(P+Q)_{>0}$ is Σ -cofibrant.*

Proof Consider the component (45) of $(P+Q)_{>0}$. Let Σ' be the subgroup of Σ_k which can send i to j if and only if $n_i = n_j$. One has that $P(k)$ is a cofibrant Σ' -space, while the second factor

$$\coprod_{\alpha: [m] \rightarrow [n_1 + \dots + n_k]} \prod_{i \in [n_1 + \dots + n_k]} Q(|\alpha^{-1}(i)|)$$

is Σ' -cofibrant. Indeed, since all α 's are surjective in the coproduct, the group Σ' acts freely on this disjoint union. Applying Lemma A.1 for $G_1 = 1, G = G_2 = \Sigma'$, we get that the component (45) is Σ' -cofibrant. \square

Theorem 2.22 *Let $\phi_1 : P \rightarrow P'$ and $\phi_2 : Q \rightarrow Q'$ be weak equivalences between componentwise cofibrant operads satisfying $P(0) = P'(0) = Q(0) = Q'(0) = \emptyset$. The extension and restriction functors, as well as their truncated versions, give rise to Quillen equivalences:*

$$\phi_! : \Sigma_{>0} \text{Bimod}_{P; Q} \rightleftarrows \Sigma_{>0} \text{Bimod}_{P'; Q'} : \phi^*, \tag{46}$$

$$\phi_! : T_r \Sigma_{>0} \text{Bimod}_{P; Q} \rightleftarrows T_r \Sigma_{>0} \text{Bimod}_{P'; Q'} : \phi^*. \tag{47}$$

Proof The proof is similar to that of Theorem 2.19. We use Proposition 2.21 that the operads $(P+Q)_{>0}$ and $(P'+Q')_{>0}$ governing these structures are Σ -cofibrant. \square

3 The Reedy Model Category of $(P-Q)$ -Bimodules

Let P be any topological operad and Q be a reduced operad. From now on, we denote by $\Lambda \text{Bimod}_{P; Q}$ and $T_r \Lambda \text{Bimod}_{P; Q}$ the categories of $(P-Q)$ -bimodules and r -truncated




Fig. 11 Illustration of the right action by $*_0$

$(P-Q)$ -bimodules, respectively, equipped with the Reedy model category structures. This structure is transferred from the categories $\Lambda Seq_P := P_0 \downarrow \Lambda Seq$ and $T_r \Lambda Seq_P := P_0 \downarrow T_r \Lambda Seq$, respectively, along the adjunctions

$$\begin{aligned} \mathcal{F}_{P;Q}^\Lambda : \Lambda Seq_P &\rightleftarrows \Lambda \text{Bimod}_{P;Q} : \mathcal{U}^\Lambda, \\ \mathcal{F}_{P;Q}^{T_r \Lambda} : T_r \Lambda Seq_P &\rightleftarrows T_r \Lambda \text{Bimod}_{P;Q} : \mathcal{U}^\Lambda, \end{aligned} \tag{48}$$

where both free functors are obtained from the functors $\mathcal{F}_{P;Q}^\Sigma$ and $\mathcal{F}_{P;Q}^{T_r \Sigma}$ by taking the restriction of the coproduct (28) to the reduced trees with section without univalent vertices other than the pearls. The (acyclic) generating cofibrations in ΛSeq_P and $T_r \Lambda Seq_P$ are $\{P_0 \sqcup \partial X \rightarrow P_0 \sqcup X\}$, where $\{\partial X \rightarrow X\}$ is the set of (acyclic) generating cofibrations of ΛSeq and $T_r \Lambda Seq$, respectively.

If we denote by $Q_{>0}$ the operad obtained from Q by changing the arity 0 component to the empty set (i.e. $Q_{>0}(0) = \emptyset$ and $Q_{>0}(n) = Q(n)$ for $n \geq 1$), then for any (possibly truncated) Λ -sequence M and $n \geq 0$, one has

$$\mathcal{F}_{P;Q}^\Lambda(M)(n) := \mathcal{F}_{P;Q_{>0}}^\Sigma(M)(n) \quad \text{and} \quad \mathcal{F}_{P;Q}^{T_r \Lambda}(M)(n) := \mathcal{F}_{P;Q_{>0}}^{T_r \Sigma}(M)(n). \tag{49}$$

By construction, the above Σ -sequences are equipped with a (possibly truncated) right module structures over $Q_{>0}$. We can extend this structure in order to get a (possibly truncated) right Q -module structures using the operadic structure of Q and the Λ structure of M (see Fig. 11).

Theorem 3.1 *Let P be an operad and Q be a reduced well-pointed operad. The categories $\Lambda \text{Bimod}_{P;Q}$ and $T_r \Lambda \text{Bimod}_{P;Q}$, with $r \geq 0$, admit cofibrantly generated model category structures, called Reedy model category structures, transferred from ΛSeq_P and $T_r \Lambda Seq_P$, respectively, along the adjunctions (48). In particular, these model category structures make the pairs of functors (48) into Quillen adjunctions.*

Proof According to the transfer principle of Theorem 1.1, we have to check the small object argument as well as the existence of a functorial fibrant replacement and a functorial factorization of the diagonal map in the category $\Lambda \text{Bimod}_{P;Q}$.

For the small object argument, let us remark that the pushout in the category of $(P-Q)$ -bimodules is defined exactly as in the category of $(P-Q_{>0})$ -bimodules. More precisely, for any $n \geq 0$, one has:²

$$\text{colim}_{\Lambda \text{Bimod}_{P;Q}} (B \leftarrow A \rightarrow C)(n) = \text{colim}_{\Sigma \text{Bimod}_{P;Q_{>0}}} (B \leftarrow A \rightarrow C)(n). \tag{50}$$

Let $\mathcal{F}_{P;Q}^\Lambda(X)$ be a domain of an element in the set of generating (acyclic) cofibrations in $\Lambda \text{Bimod}_{P;Q}$. Let $\lambda = \aleph_1$ and assume that $\{M_\alpha\}_{\alpha < \lambda}$ is a λ -sequence of $(P-Q)$ -bimodules,

² The forgetful functor from the category of $(P-Q)$ -bimodules to the category of $(P-Q_{>0})$ -bimodules preserves colimits as it admits a right adjoint by Proposition 3.14b for $Q_1 = Q$, $Q_2 = Q_{>0}$, $Y = Q$.

such that each map $M_{<\alpha} := \text{colim}_{\beta < \alpha} M_\beta \rightarrow M_\alpha$ fits into a pushout square

$$\begin{array}{ccc}
 \mathcal{F}_{P;Q}^\Lambda(\partial Y) & \longrightarrow & \mathcal{F}_{P;Q}^\Lambda(Y) \\
 \downarrow & & \downarrow \\
 M_{<\alpha} & \longrightarrow & M_\alpha,
 \end{array} \tag{51}$$

where $\partial Y \rightarrow Y$ is a possibly infinite coproduct in ΛSeq_P of generating (acyclic) cofibrations. It follows from Lemma 2.10 and equations (49) and (50), that each map $M_{<\alpha} \rightarrow M_\alpha$ is an objectwise closed inclusion. Moreover, $(\coprod_{n \geq 0} X(n)) \setminus P(0)$ is a finite union of spheres (or discs), thus separable, and therefore, X is \aleph_1 -small relative to componentwise closed inclusions. So, the same argument used for the proof of the small objects argument in Theorem 2.11 works.

Contrary to the category of $\Sigma \text{Bimod}_{P;Q}$, the objects in the category $\Lambda \text{Bimod}_{P;Q}$ are not necessarily fibrant and the identity functor is not a fibrant replacement functor. The aim of Sect. 3.1.1 is to introduce such a fibrant replacement functor if Q is well-pointed. This fibrant replacement is different from the fibrant coresolution functor for Λ -sequences defined in Sect. 1.3.

From now on, we introduce a functorial path object in the Reedy model category of $(P-Q)$ -bimodules. In other words, for any $M \in \Lambda \text{Bimod}_{P;Q}$ which is fibrant in the category of Λ -sequences, we build an element $\text{Path}(M) \in \Lambda \text{Bimod}_{P;Q}$ such that there is a factorization of the diagonal map

$$\Delta : M \xrightarrow[\simeq]{f_1} \text{Path}(M) \xrightarrow[f_2]{\twoheadrightarrow} M \times M,$$

where f_1 is a weak equivalence and f_2 is a fibration. Let us consider

$$\text{Path}(M)(n) = \text{Map}([0, 1]; M(n)). \tag{52}$$

The object so obtained inherits a bimodule structure from M . The map from M to $\text{Path}(M)$, sending a point to the constant path, is clearly a homotopy equivalence. Furthermore, let us remark that one has the following identities:

$$\begin{aligned}
 \mathcal{M}(\text{Path}(M))(n) &= \text{Map}([0, 1]; \mathcal{M}(M)(n)) \quad \text{and} \\
 \mathcal{M}(M \times M)(n) &= \mathcal{M}(M)(n) \times \mathcal{M}(M)(n).
 \end{aligned}$$

So, the map f_2 is a fibration if the map between the limits induced by the natural transformation

$$\begin{array}{ccccc}
 \text{Path}(M)(n) & \xlongequal{\quad} & \text{Path}(M)(n) & \xlongequal{\quad} & \text{Path}(M)(n) \\
 \downarrow & & \downarrow & & \downarrow \\
 \text{Map}([0, 1]; \mathcal{M}(M)(n)) & \longrightarrow & \mathcal{M}(M)(n) \times \mathcal{M}(M)(n) & \longleftarrow & M(n) \times M(n)
 \end{array}$$

is a Serre fibration. The right vertical map is obviously a Serre fibration because the inclusion from $\partial[0, 1]$ into $[0, 1]$ is a cofibration (see the proof of Theorem 2.11). Moreover, since the inclusion $\partial[0, 1] \rightarrow [0, 1]$ is a cofibration and the map $M(n) \rightarrow \mathcal{M}(M)(n)$ is a fibration, an alternative version of the pushout product lemma (see [26, Section 9.1.5]) implies that the map

$$\text{Map}([0, 1]; M(n)) \longrightarrow \text{Map}([0, 1]; \mathcal{M}(M)(n)) \quad \times \quad \text{Map}(\partial[0, 1]; M(n))$$

$$\qquad \qquad \qquad \text{Map}(\partial[0, 1]; \mathcal{M}(M)(n))$$

is also a Serre fibration. As a consequence of Lemma 1.5, the map f_2 is a Serre fibration. \square

3.1 Properties of the Reedy Model Category of Bimodules

This subsection is divided into three parts. The first one is devoted to the construction of an explicit fibrant coresolution functor for reduced bimodules. In the second part, we characterize (acyclic) cofibrations in the Reedy model category of bimodules as (acyclic) cofibrations in the usual projective model category of bimodules. The last part consists in extending the properties introduced in Sect. 2.2 to the Reedy model category.

3.1.1 A Reedy Fibrant Replacement Functor for Bimodules

Let P be an operad and Q be a reduced operad. The goal of this section is to give an explicit Reedy fibrant replacement in the category of bimodules if the operad Q is well-pointed. A conceptual description of this fibrant coresolution in terms of internal hom is given in Sect. 3.3.

- *The set of trees $\mathbb{P}[n]$.* Let $\mathbb{P}[n]$ be the set of planar trees T whose roots have exactly n incoming edges. We label their leaves with the identity permutation in $\Sigma_{|T|}$, where $|T|$ is the number of leaves in T . We also label the n incoming edges bijectively by the set $[n]$ in the planar order from left to right. Such a tree T is equipped with an orientation towards the root and we say that $v < v'$ if the path joining the vertex v' with the root passes through the vertex v . It makes the set of vertices $V(T)$ into a partially ordered set. Moreover, we consider the operations

$$\delta_{i,m} : \mathbb{P}[n + m - 1] \longrightarrow \mathbb{P}[n], \text{ with } n, m \in \mathbb{N} \text{ and } 1 \leq i \leq n,$$

$$\Gamma_k^m : \mathbb{P}[n] \longrightarrow \mathbb{P}[m], \text{ with } n, m, k \in \mathbb{N} \text{ and } m + k \leq n,$$

The operation $\delta_{i,m}(T)$ is defined as follows. If $m = 0$, then $\delta_{i,m}$ consists in adding an incoming edge to the root of T at the i -th position. The new incoming edge is connected to a univalent vertex. If $m > 0$, then $\delta_{i,m}(T)$ is obtained from T by gluing together the incoming edges $i, i + 1, \dots, i + m - 1$, of the root counted according to the planar order. In both cases, $\delta_{i,m}(T)$ has one additional vertex to those of T . The new vertex has exactly m incoming edges (see Fig. 12).

For any n, m and k such that $m + k \leq n$, the map $\Gamma_k^m : \mathbb{P}[n] \rightarrow \mathbb{P}[m]$ consists in removing $n - m$ incoming edges together with the trees attached to them. The removed edges are those labelled by $[n] \setminus \{k + 1, \dots, k + m\}$ (see Fig. 13).

- *Construction of the fibrant replacement functor.* From now on, we fix a $(P-Q)$ -bimodule M . Let T be an element in $\mathbb{P}[n]$. We consider the spaces $H(T)$ of indexing the vertices of T




Fig. 12 Illustration of the applications $\delta_{2,0}$ and $\delta_{2,3}$




Fig. 13 Illustration of the applications Γ_1^2 and Γ_0^2

other than the root r by real numbers and points in the operad \mathcal{Q} , respectively. More precisely, one has

$$D(T) = \prod_{v \in V(T) \setminus \{r\}} \mathcal{Q}(|v|),$$

$$H(T) = \{ \{t_v\} \in [0, 1]^{|V(T) \setminus \{r\}|} \mid \forall v < v', t_v \leq t_{v'} \}.$$

In other words, $H(T)$ is the space of order preserving maps

$$V(T) \setminus \{r\} \rightarrow [0, 1].$$

Finally, we denote by $M^f(n)$ the subspace

$$M^f(n) \subset \prod_{T \in \mathbb{P}[n]} \text{Map}(H(T) \times D(T); M(|T|)) \tag{53}$$

composed of families of continuous maps $\{f_T\}_{T \in \mathbb{P}[n]}$ satisfying the following conditions:

1. Let T be an element in $\mathbb{P}[n]$ having a bivalent vertex v other than the root. Then one has

$$\begin{array}{ccc} H(T) \times D(T \setminus \{v\}) & \longrightarrow & H(T) \times D(T) \\ \downarrow & & \downarrow f_r \\ H(T \setminus \{v\}) \times D(T \setminus \{v\}) & \xrightarrow{f_{T \setminus \{v\}}} & M(|T \setminus \{v\}|) = M(|T|) \end{array} \tag{54}$$

where $T \setminus \{v\}$ is the tree obtained from T by removing the bivalent vertex v (i.e. by replacing the incoming and outgoing edges of v by a single edge). The upper horizontal map indexes the vertex v by the unit of the operad \mathcal{Q} while the left vertical map is obtained by forgetting the the real number indexing the bivalent vertex v .

2. For any non-root vertex v of T and any permutation σ of the incoming edges of v , one has

$$\begin{array}{ccc} H(T) \times D(T) & \longrightarrow & H(T \cdot \sigma) \times D(T \cdot \sigma) \\ f_r \downarrow & & \downarrow f_{T \cdot \sigma} \\ M(|T|) & \xrightarrow{\sigma_L[T]^*} & M(|T \cdot \sigma|) = M(|T|) \end{array} \tag{55}$$

where $T \cdot \sigma$ is the tree obtained from T by permuting the incoming edges of v according to the permutation σ and $\sigma_L[T] \in \Sigma_{|T|}$ is the induced permutation of the leaves of T . The upper horizontal map sends the decorations of the tree T to the corresponding decorations of $T \cdot \sigma$ and acts on the \mathcal{Q} -decoration of the vertex v using the Σ -structure of the operad \mathcal{Q} .

3. For any inner edge e , which is not connected to the root, one has

$$\begin{array}{ccc}
 H(T/e) \times D(T) & \longrightarrow & H(T) \times D(T) \\
 \downarrow & & \downarrow f_T \\
 H(T/e) \times D(T/e) & \xrightarrow{f_{T/e}} & M(|T/e|) = M(|T|)
 \end{array} \tag{56}$$

where T/e is the tree obtained from T by contracting the edge e . The upper horizontal map indexes the source and the target vertices of e by the real number in $H(T/e)$ corresponding to the vertex resulting from the contraction of e . The left vertical map is defined using the operadic structure of Q .

4. Any tree T has a unique decomposition of the form $T = T_e \circ_{i(e)} T'_e$ along any edge e . Then one has

$$\begin{array}{ccc}
 H(T_e) \times D(T_e) \times \prod_{v \in V(T'_e)} Q(|v|) & \longrightarrow & H(T) \times D(T) \\
 f_{T_e} \times \eta \downarrow & & \downarrow f_T \\
 M(|T_e|) \times Q(|T'_e|) & \xrightarrow{\circ^{i(e)}} & M(|T|)
 \end{array} \tag{57}$$

The upper horizontal map consists in indexing the vertices associated to the tree T'_e by 1. The map $\eta : \prod_{v \in V(T'_e)} Q(|v|) \rightarrow Q(|T'_e|)$ is defined using the operadic structure of Q while the lower horizontal map is obtained using the right Q -module structure of M .

Remark 3.2 Let us notice that, as a special case of the fourth condition, for any univalent vertex v , one has

$$\begin{array}{ccc}
 H(T \setminus \{v\}) \times D(T \setminus \{v\}) & \longrightarrow & H(T) \times D(T) \\
 f_{T \setminus \{v\}} \downarrow & & \downarrow f_T \\
 M(|T \setminus \{v\}|) = M(|T| + 1) & \xrightarrow{\circ^{v,*_0}} & M(|T|)
 \end{array}$$

where $T \setminus \{v\}$ is the tree obtained from T by removing the univalent vertex v (and thus producing one more leaf). The upper horizontal map consists in indexing the vertex v by the real number 1 and the unique point $*_0 \in Q(0)$. The lower horizontal map composes the input of $M(|T \setminus \{v\}|)$, corresponding to the vertex v , with the point $*_0 \in Q(0)$ using the right Q -module structure of M .

• *The Σ -structure on the fibrant coresolution.* The space $M^f(n)$ inherits an action of the permutation group Σ_n . More precisely, for any $\sigma \in \Sigma_n$, we denote by T_σ the tree obtained from $T \in \mathbb{P}[n]$ by permuting the incoming edges associated to the root of T according to the permutation σ . Such a permutation induces the following two bijections:

$$\begin{aligned}
 \sigma_V[T] : V(T \setminus \{r\}) &\longrightarrow V(T_\sigma \setminus \{r\}) \in \Sigma_{|V(T) \setminus \{r\}|}, \\
 \sigma_L[T] : \ell(T) &\longrightarrow \ell(T_\sigma) \in \Sigma_{|T|}.
 \end{aligned} \tag{58}$$

Here, $\ell(T)$ denotes the set $[|T|]$ of leaves. So, the action of the permutation group $\sigma^* : M^f(n) \rightarrow M^f(n)$ sends a family of continuous maps $\{f_T\}_{T \in \mathbb{P}[n]}$ to the family

$\{(f \cdot \sigma)_T\}_{T \in \mathbb{P}[n]}$ given by the formula

$$(f \cdot \sigma)_T : H(T) \times D(T) \longrightarrow M(|T|);$$

$$\{t_v\}, \{q_v\} \longmapsto f_{T_\sigma}(\{t_{\sigma_V[T](v)}\}, \{q_{\sigma_V[T](v)}\}) \cdot \sigma_L[T].$$

• *The bimodule structure on the fibrant coresolution.* Since one has the identity $M^f(0) = M(0)$, one has a map $\gamma_0 : P(0) \rightarrow M^f(0)$ and the Σ -sequence M^f inherits a $(P-Q)$ -bimodule structure whose right operations are given by

$$\circ^i : M^f(n) \times Q(m) \longrightarrow M^f(n + m - 1);$$

$$\{f_T\}_{T \in \mathbb{P}[n]}, q \longmapsto \{(f \circ^i q)_T\}_{T \in \mathbb{P}[n+m-1]}, \tag{59}$$

where $(f \circ^i q)_T$ is the composite map:

$$(f \circ^i q)_T : H(T) \times D(T) \longrightarrow H(\delta_{i,m}(T)) \times D(\delta_{i,m}(T)) \xrightarrow{f_{\delta_{i,m}(T)}} M(|\delta_{i,m}(T)|) = M(|T|).$$

The left hand side map consists in indexing the new vertex by the real number 0 and the point $q \in Q(m)$. Similarly, the left P -module structure on M^f is given by the operations

$$\gamma_\ell : P(k) \times M^f(n_1) \times \dots \times M^f(n_k) \longrightarrow M^f(n_1 + \dots + n_k);$$

$$p, \{f_T^1\}_{T \in \mathbb{P}[n_1]}, \dots, \{f_T^k\}_{T \in \mathbb{P}[n_k]} \longmapsto \{p(f^1, \dots, f^k)_T\}_{T \in \mathbb{P}[n_1 + \dots + n_k]},$$

where $p(f^1, \dots, f^k)_T$ is the composite map

$$\begin{array}{ccc} H(T) \times D(T) & \xrightarrow{p(f^1, \dots, f^k)_T} & M(|T|) \\ \cong \downarrow & & \uparrow \gamma_\ell(p; -, \dots, -) \\ \prod_{1 \leq i \leq k} H(\Gamma_{n_1 + \dots + n_{i-1}}^{n_i}(T)) \times D(\Gamma_{n_1 + \dots + n_{i-1}}^{n_i}(T)) & \xrightarrow{\times_i f_{\Gamma_{n_1 + \dots + n_{i-1}}^{n_i}(T)}} & \prod_{1 \leq i \leq k} M(|\Gamma_{n_1 + \dots + n_{i-1}}^{n_i}(T)|). \end{array}$$

Finally, there is a map of $(P-Q)$ -bimodules $\eta : M \rightarrow M^f$ sending a point $m \in M(n)$ to the family of continuous maps $\{\eta(m)_T\}_{T \in \mathbb{P}[n]}$ given by the formula

$$\eta(m)_T : H(T) \times D(T) \longrightarrow M(|T|);$$

$$\{t_v\}, \{q_v\} \longmapsto m \circ \{q_v\},$$

using the right Q -bimodule structure of M . It means that m is taken for a root and we compose with $\{q_v\}$ using the operadic structure of Q and the right module structure of M .

Proposition 3.3 *The map $\eta : M \rightarrow M^f$ is a weak equivalence.*

Proof The proof is similar to the proof of Proposition 1.3. We show that the map of Λ -sequences $\eta_n : M(n) \rightarrow M^f(n)$ is a homotopy equivalence of Σ -sequences. For this purpose, we introduce a map of Σ -sequences $\psi : M^f \rightarrow M$ given by

$$\psi_n : M^f(n) \longrightarrow M(n);$$

$$\{f_T\}_{T \in \mathbb{P}[n]} \longmapsto f_{C_n}(*),$$

where C_n is the n -corolla whose corresponding space $H(C_n) \times D(C_n)$ is necessarily the one point topological space. The map ψ so obtained makes η into a deformation retract and the homotopy consists in bringing the real numbers indexing the vertices other than the root to 1. □




Fig. 14 Illustration of the construction of $T[h]$

Proposition 3.4 *If the operad \underline{Q} is well-pointed, then the $(P\text{-}Q)$ -bimodule M^f is Reedy fibrant.*

Proof The proof is divided into two parts. First, we identify the matching object $\mathcal{M}(M^f)(n)$ with a space $\mathcal{M}_0(n)$ defined in the same way as $M^f(n)$ by changing slightly the construction of the space $H(T)$. Thereafter, we build a tower of fibrations related to $M^f(n)$ and $\mathcal{M}_0(n)$ according to the number of vertices of the trees in $\mathbb{P}[n]$. In the proof we use equivariant homotopy theory techniques from ‘‘Appendix A.2’’.

Simplification of the matching object: We say that two trees T_1 and T_2 from $\mathbb{P}[n]$ are equivalent if they are isomorphic as non-planar trees and moreover the isomorphism between T_1 and T_2 preserves the order of the incoming edges of the root r . We denote by $\mathbb{T}[n]$ the so obtained set of equivalence classes. For $T \in \mathbb{T}[n]$ we denote by $\text{Aut}(T)$ the set of automorphisms of T which preserve the order of the incoming edges to the root. Let $|T|$ denote the number of leaves of T . For each $T \in \mathbb{T}[n]$ we choose a planar representative and we label the leaves of T in the corresponding planar order they appear. This gives us a homomorphism $\text{Aut}(T) \rightarrow \Sigma_{|T|}$, which is not always injective due to the presence of arity zero vertices.

Because of the relation (55), the space $M^f(n)$ can be equivalently described as the subspace

$$M^f(n) \subset \prod_{T \in \mathbb{T}[n]} \text{Map}_{\text{Aut}(T)}(H(T) \times D(T); M(|T|))$$

composed of families of continuous maps $\{f_T\}_{T \in \mathbb{T}[n]}$ satisfying the conditions (54), (56), (57) properly understood.

For any tree $T \in \mathbb{T}[n]$, we denote by $H_0(T)$ the subspace of $H(T)$ which consists of points having at least one univalent vertex labelled by 0 and connected to the root of T . Let $\mathcal{M}_0(n)$ be the subspace

$$\mathcal{M}_0(n) \subset \prod_{T \in \mathbb{T}[n]} \text{Map}_{\text{Aut}(T)}(H_0(T) \times D(T); M(|T|)),$$

satisfying the relations (54), (56) and (57). In order to show that the space $\mathcal{M}_0(n)$ is homeomorphic to the matching object $\mathcal{M}(M^f)(n)$, we recall that a point in $\mathcal{M}(M^f)(n)$ is a family of maps $\{\phi_{T,h}\}$, indexed by $h \in \Lambda_+([\ell], [n])$, $T \in \mathbb{T}[\ell]$ and $\ell < n$, satisfying some conditions related to the limit (3) as well as the relations (54), (56) and (57).

For each pair (T, h) , with $h \in \Lambda_+([\ell], [n])$ and $T \in \mathbb{T}[\ell]$, we denote by $T[h]$ the tree in $\mathbb{T}[n]$ obtained from T by adding univalent vertices connected to the root according to the order preserving map h (see Fig. 14). So the map

$$\begin{aligned} \alpha : \mathcal{M}_0(n) &\longrightarrow \mathcal{M}(M^f)(n); \\ \phi = \{\phi_T\} &\longmapsto \{(\alpha \circ \phi)_{T,h}\}, \end{aligned}$$

is given by the composite maps

$$(\alpha \circ \phi)_{T,h} : H(T) \times D(T) \longrightarrow H_0(T[h]) \times D(T[h]) \xrightarrow{\phi_{T[h]}} M(|T[h]|) = M(|T|),$$

where the left hand side map consists in indexing the new univalent vertices by the unique point in $Q(0)$ and the real number 0.

Conversely, for any tree $T \in \mathbb{T}[n]$ and $\{t_v\} \in H_0(T)$, we denote by $\ell_{\{t_v\}}$ the number of incoming edges of the root of T which are not connected to univalent vertices indexed by 0. Furthermore, we denote by $h[\{t_v\}] : [\ell_{\{t_v\}}] \rightarrow [n]$ the order preserving map which keeps in mind the position of the incoming edges of the root which are not connected to a univalent vertex indexed by 0. Finally, $T[\{t_v\}] \in \mathbb{T}[\ell_{\{t_v\}}]$ is the tree obtained from T by removing the incoming edges of the root which are connected to a univalent vertex indexed by 0. So, one has the map

$$\begin{aligned} \beta : \mathcal{M}(M^f)(n) &\longrightarrow \mathcal{M}_0(n); \\ \phi = \{\phi_{T,h}\} &\longmapsto \{(\beta \circ \phi)_T\}, \end{aligned}$$

given by

$$\begin{aligned} (\beta \circ \phi)_T : H_0(T) \times D(T) &\longrightarrow M(|T|); \\ \{t_v\}, \{q_v\} &\longmapsto \phi_{T[\{t_v\}],h[\{t_v\}]}(\{\tilde{t}_v\}, \{\tilde{q}_v\}), \end{aligned}$$

where the families $\{\tilde{t}_v\}$ and $\{\tilde{q}_v\}$ are obtained from $\{t_v\}$ and $\{q_v\}$, respectively, by removing the parameters corresponding to the univalent vertices indexed by 0 and connected to the root. The map β is well defined due to the relations induced by the limit and, together with the map α , induces a homeomorphism between $\mathcal{M}_0(n)$ and the matching object $\mathcal{M}(M^f)(n)$. Furthermore, the map from $M^f(n) \rightarrow \mathcal{M}(M^f)(n)$ is equivalent to the restriction map

$$r : M^f(n) \longrightarrow \mathcal{M}_0(n),$$

induced by the inclusion $H_0(T) \rightarrow H(T)$ for any tree $T \in \mathbb{T}[n]$.

Construction of the tower of fibrations: We construct a tower of fibrations according to the number of vertices of the trees in $\mathbb{T}[n]$:

$$\mathcal{M}_0(n) = A_0 \longleftarrow A_1 \longleftarrow \dots \longleftarrow A_{k-1} \longleftarrow A_k \longleftarrow \dots \longleftarrow M^f(n).$$

For this purpose, we introduce the set $\mathbb{T}[n, k]$ of trees in $\mathbb{T}[n]$ having exactly k vertices. In particular, $\mathbb{T}[n, 1]$ has only one element which is the n -corolla C_n . The space A_k is defined as the subspace

$$\begin{aligned} A_k \subset &\left(\prod_{\substack{T \in \mathbb{T}[n,i] \\ i \leq k}} \text{Map}_{\text{Aut}(T)}(H(T) \times D(T); M(|T|)) \right) \\ &\times \left(\prod_{\substack{T \in \mathbb{T}[n,i] \\ i > k}} \text{Map}_{\text{Aut}(T)}(H_0(T) \times D(T); M(|T|)) \right) \end{aligned}$$

composed of families of continuous maps $\{f_T\}_{T \in \mathbb{T}[n]}$ satisfying the conditions (54), (56), (57).

The space A_k is defined by induction from A_{k-1} using the pullback diagram:

$$\begin{array}{ccc}
 A_k & \longrightarrow & \prod_{[T] \in \mathbb{T}[n,k]} \text{Map}_{\text{Aut}(T)}(H(T) \times D(T); M(|T|)) \\
 \downarrow & & \downarrow \\
 A_{k-1} & \longrightarrow & \prod_{[T] \in \mathbb{T}[n,k]} \text{Map}_{\text{Aut}(T)}((H \times D)^-(T); M(|T|)),
 \end{array} \tag{60}$$

In the diagram above, for any tree $T \in \mathbb{T}[n, k]$, $D^-(T)$ is the subspace of $D(T)$ formed by points having at least one bivalent vertex labelled by the unit of the operad \mathcal{Q} . On the other hand, the space $H^-(T) \subset H(T)$ consists of points in $H_0(T)$ or having two consecutive vertices indexed by the same real number or having a vertex indexed by 1. Then we set

$$(H \times D)^-(T) := (H^-(T) \times D(T)) \coprod_{H^-(T) \times D^-(T)} (H(T) \times D^-(T)).$$

To show that the map $A_k \rightarrow A_{k-1}$ is a Serre fibration, one has to prove that the right vertical map in (60) is one. Let t be the number of arity one (non-root) vertices of T . Denote by

$$D_1(T) := \mathcal{Q}(1)^{\times t}, \quad D_{\neq 1}(T) := \prod_{\substack{v \in V(T) \setminus \{r\} \\ |v| \neq 1}} \mathcal{Q}(|v|).$$

One has, $D(T) = D_1(T) \times D_{\neq 1}(T)$. We similarly denote by $D_1^-(T)$ the subset consisting of points having at least one coordinate equal to the unit $*_1 \in \mathcal{Q}(1)$ and define

$$(H \times D_1)^-(T) := (H^-(T) \times D_1(T)) \coprod_{H^-(T) \times D_1^-(T)} (H(T) \times D_1^-(T)).$$

We have

$$\begin{aligned}
 (H \times D)^-(T) &= (H \times D_1)^-(T) \times D_{\neq 1}(T) \quad \text{and} \\
 (H \times D)(T) &= (H \times D_1)(T) \times D_{\neq 1}(T).
 \end{aligned} \tag{61}$$

Recall Definition A.9 of a *cellularly* equivariant cofibration. Let \underline{n} denote the ordered set $\{0, 1, \dots, n + 1\}$. The dual to the simplicial indexing category Δ can be defined as the category having $\{\underline{n}, n \in \mathbb{N}\}$ as the set of objects with morphisms order preserving maps that also preserve both extrema. The space $H(T)$ can be described as the realization of the simplicial set $H_T(\bullet)$ with $H_T(\underline{n})$ being the set of order preserving maps

$$V(T) \setminus \{r\} \rightarrow \underline{n}. \tag{62}$$

Note that $H^-(T)$ is the realization of a simplicial subset $H_T^-(\bullet) \subset H_T(\bullet)$. It consists of order-preserving maps (62) which are either sending an arity zero vertex of T to 0 or non-injective or sending some vertex of T to the maximal element. By Lemma A.11, the inclusion $H^-(T) \rightarrow H(T)$ is a cellularly $\text{Aut}(T)$ -equivariant cofibration. Using the fact that the operad \mathcal{Q} is well-pointed and also applying Lemmas A.12, A.13, A.14, we get that the inclusion $(H \times D_1)^-(T) \rightarrow H(T) \times D_1(T)$ is a cellularly $\text{Aut}(T)$ -equivariant cofibration.

Using the fact that the category Top is cartesian closed, for any $X \in Aut(T)\text{-}Top$, one has a homeomorphism

$$Map_{Aut(T)}(X \times D_{\neq 1}(T); M(|T|)) = Map_{Aut(T)}(X; Map(D_{\neq 1}(T), M(|T|))), \tag{63}$$

where the $Aut(T)$ -action on $Map(D_{\neq 1}(T), M(|T|))$ is defined by $(f \cdot g)(x) = f(x \cdot g^{-1}) \cdot g$, $g \in Aut(T)$. Applying this homeomorphism to (61) and then Lemma A.10 to the right vertical map in (60), we get that the latter is a Serre fibration. \square

Remark 3.5 The same strategy can be used in order to get a fibrant replacement functor for the Reedy model category of r -truncated $(P\text{-}Q)$ -bimodules. The fibrant replacement should be defined as a subspace of the product (53) with an additional restriction $|T| \leq r$. The constraints are the same.

3.1.2 Characterization of Reedy Cofibrations for Bimodules

It has been shown by the second author [19, Theorem 8.3.20] that a map of Λ -sequences is a Reedy cofibration if and only if it is a projective Σ -cofibration. In the context of operads, he also proves [19, Theorem 8.4.12] that a map $\phi : P \rightarrow Q$ between reduced operads is cofibrant in the Reedy model category $\Lambda_*\text{Operad}$ if and only if the corresponding map $\phi_{>0} : P_{>0} \rightarrow Q_{>0}$ is a cofibration in the projective model structure of (not necessarily reduced) operads. In what follows, we prove an analogous version in the context of operadic bimodules.

Theorem 3.6 *Let P and Q be as in Theorem 3.1. A morphism $\phi : M \rightarrow N$ in the category of (possibly truncated) $(P\text{-}Q)$ -bimodules is a Reedy cofibration if and only if the corresponding map ϕ is a cofibration in the projective model category of (possibly truncated) $(P\text{-}Q_{>0})$ -bimodules.*

Proof First, we show that if the induced map is a cofibration in the projective model category of $(P\text{-}Q_{>0})$ -bimodules then the map $\phi : M \rightarrow N$ is a Reedy cofibration in the category of $(P\text{-}Q)$ -bimodules. For this purpose, we consider the following lifting problem in the category of $(P\text{-}Q)$ -bimodules:

$$\begin{array}{ccc}
 M & \xrightarrow{i} & A \\
 \phi \downarrow & \exists \varphi? \nearrow & \downarrow p \\
 N & \xrightarrow{j} & B
 \end{array} \tag{64}$$

where $p : A \rightarrow B$ is an acyclic Reedy fibration. The strategy is to build the map φ by induction using an adjunction between $(P\text{-}Q)$ -bimodules

$$ar_s : \Lambda\text{Bimod}_{P; Q} \rightleftarrows \Lambda\text{Bimod}_{P; Q} : \text{cosk}_s.$$

The functor ar_s , called the arity filtration functor, sends a $(P\text{-}Q)$ -bimodule M to the bimodule $ar_s(M)$ defined as a quotient of the free $(P\text{-}Q_{>0})$ -bimodule generated by the first s components of M where the equivalence relation is determined by the restriction of the bimodule structure on the first s components of M . In other words, if L_s denotes the left adjoint to the truncation functor T_s , then the arity filtration functor is given by $ar_s = L_s \circ T_s$. By construction, $ar_s(M)$ is a $(P\text{-}Q)$ -bimodule and one has the identities

$$ar_{s-1} \circ ar_s = ar_{s-1}, \quad \text{colim}_s ar_s(M) = M \quad \text{and} \quad ar_s(M)(k) = M(k), \text{ for } k \leq s.$$

Let us notice that, even though this functor is called filtration, the natural map $ar_s(M) \rightarrow M$ might not be an inclusion. First, we give an explicit description of the right adjoint of the arity filtration functor.

The Λ -coskeletons associated to a bimodule. According to the notation introduced by the second author [19], the Σ -sequence $cosk_s(M) = \{cosk_s(M)(n), n \geq 0\}$, called the s -th coskeleton associated to the bimodule M , is given by the formula

$$cosk_s(M)(n) = \lim_{\substack{h \in \Lambda_+([i]; [n]) \\ 0 \leq i \leq s}} M(i), \quad \text{with } n \geq 0. \tag{65}$$

As a matter of fact, the right adjoint R_s , to the s -truncation functor T_s , is defined by the same formula. Similarly to the arity filtration functor, the functor $cosk_s$ is also given by

$$cosk_s = R_s \circ T_s.$$

By construction, a point $x \in cosk_s(M)(n)$ is a family of elements $x = \{x_h \in M(i), h \in \Lambda_+([i]; [n]) \text{ and } i \leq s\}$ satisfying the relation of the limit: for any $f \in \Lambda_+([i]; [j])$ and $g \in \Lambda_+([j]; [n])$, one has $f^*(x_g) = x_{g \circ f}$. In particular, $cosk_s(M)(0) = M(0)$ is obviously endowed with a map from $P(0)$. Furthermore, the Σ -sequence $cosk_s(M)$ inherits a Λ -structure. For any $f \in \Lambda_+([n_1]; [n_2])$, one has

$$\begin{aligned} f^* : cosk_s(M)(n_2) &\longrightarrow cosk_s(M)(n_1); \\ \{x_u\}_{h \in \Lambda_+([i]; [n_2])}^{0 \leq i \leq s} &\longmapsto \{x_{f \circ u}\}_{u \in \Lambda_+([i]; [n_1])}^{0 \leq i \leq s}. \end{aligned} \tag{66}$$

The Λ -sequence $cosk_s(M)$ is also a $(P-Q)$ -bimodule. In order to define the right operations, we introduce some notation. Let $n, m > 0, l \in \{1, \dots, n\}$ and $h \in \Lambda_+([i]; [n+m-1])$. If we denote by $\ell_1 \in \Lambda_+([m]; [n+m-1])$ and $\ell_2 \in \Lambda_+([n]; [n+m-1])$ the order preserving maps

$$\begin{aligned} \ell_1 : [m] &\longrightarrow [n+m-1]; \quad \text{and } \ell_2 : [n] \longrightarrow [n+m-1]; \\ \alpha &\longmapsto \alpha + \ell, \quad \alpha \longmapsto \begin{cases} \alpha & \text{if } \alpha \leq \ell, \\ \alpha + m & \text{if } \alpha > \ell, \end{cases} \end{aligned}$$

then there exist unique morphisms h_1 and h_2 making the following diagrams commute:

$$\begin{array}{ccc} [i] & \xrightarrow{h} & [n+m-1] \\ \uparrow & & \uparrow \ell_1 \\ [Im(\ell_1) \cap Im(h)] & \xrightarrow{h_1} & [m] \end{array} \qquad \begin{array}{ccc} [i] & \xrightarrow{h} & [n+m-1] \\ \uparrow & & \uparrow \ell_2 \\ [Im(\ell_2 \setminus \{\ell\}) \cap Im(h)] & \xrightarrow{h_2} & [n] \end{array}$$

Finally, if we denote by $\bar{\ell} = \ell - |\{\alpha \in [i] \mid h(\alpha) < \ell\}|$, then the right operations are defined as follows:

$$\begin{aligned} \circ^i : cosk_s(M)(n) \times Q(m) &\longrightarrow cosk_s(M)(n+m-1); \\ \{x_u\}_{u \in \Lambda_+([i]; [n])}^{0 \leq i \leq s} ; q &\longmapsto \{x_{h_2 \circ \bar{\ell} h_1^*(q)}\}_{h \in \Lambda_+([i]; [n+m-1])}^{0 \leq i \leq s}. \end{aligned} \tag{67}$$

Let $k_1, \dots, k_\ell > 0$ and $h \in \Lambda_+([i]; [k_1 + \dots + k_\ell])$. In order to define the left operation, we introduce the morphism $\ell_i \in \Lambda_+([k_i]; [k_1 + \dots + k_\ell])$ sending α to $\alpha + k_1 + \dots + k_{i-1}$.

Then there exists a unique morphism h_i such that the following diagram commutes:

$$\begin{array}{ccc}
 [i] & \xrightarrow{h} & [k_1 + \dots + k_\ell] \\
 \uparrow & & \uparrow \ell_i \\
 [Im(h) \cap Im(\ell_i)] & \xrightarrow{h_i} & [k_i]
 \end{array}$$

Finally, the left operation is given by the formula

$$\begin{aligned}
 \gamma : P(\ell) \times \text{cosk}_s(M)(k_1) \times \dots \times \text{cosk}_s(M)(k_\ell) &\longrightarrow \text{cosk}_s(M)(k_1 + \dots + k_\ell); \\
 p : \{x_{u_1}^1\}, \dots, \{x_{u_\ell}^\ell\} &\longmapsto \{p(x_{h_1}^1, \dots, x_{h_\ell}^\ell)\}_{h \in \Lambda_+([i]; [k_1 + \dots + k_\ell])}^{0 \leq i \leq s}
 \end{aligned}$$

The arity filtration and the coskeleton functors form an adjunction. For any pair of reduced $(P-Q)$ -bimodules M and N , we define below a homeomorphism between the mapping spaces of $(P-Q)$ -bimodules:

$$F : \Lambda \text{Bimod}_{P; Q}(ar_s(M); N) \cong \Lambda \text{Bimod}_{P; Q}(M; \text{cosk}_s(N)) : G. \tag{68}$$

Let $f : M \rightarrow \text{cosk}_s(N)$ be a bimodule map. The bimodule map $G(f)$ is defined by induction. If $n \leq s$ and $x \in ar_s(M)(n) = M(n)$, then $G(f)(x) = f(x)_{[n] \rightarrow [n]}$, the point indexed by the identity order preserving map. From now on, we assume that $G(f)$ is defined for any element $ar_s(M)$ until the arity $n \geq s$. Let $x \in ar_s(M)(n + 1)$. Then one has

$$G(f)(x) = \begin{cases} G(f)(x') \circ^i y & \text{if } x = x' \circ^i y, \\ y(G(f)(x_1), \dots, G(f)(x_\ell)) & \text{if } x = y(x_1, \dots, x_\ell). \end{cases}$$

Conversely, let $g : ar_s(M) \rightarrow N$ be a bimodule map. Then one has

$$\begin{aligned}
 F(g)_n : M(n) &\longrightarrow \text{cosk}_s(N)(n); \\
 x &\longmapsto \{x_h = g \circ h^*(x)\}_{h \in \Lambda_+([i]; [n])}^{0 \leq i \leq s}.
 \end{aligned}$$

The arity functor ar_s preserves $\Sigma \text{Bimod}_{P, Q>0}$ -cofibrations. The functor ar_s can be defined on the category $\Sigma \text{Bimod}_{P, Q>0}$ and it fits into a Quillen adjunction

$$ar_s : \Sigma \text{Bimod}_{P, Q>0} \rightleftarrows \Sigma \text{Bimod}_{P, Q>0} : tr_s, \tag{69}$$

where tr_s is the truncation functor $tr_s(L)(n) = \begin{cases} L(n), & n \leq s; \\ *, & n > s. \end{cases}$

One has a square of functors

$$\begin{array}{ccc}
 \Lambda \text{Bimod}_{P; Q} & \xrightarrow{ar_s} & \Lambda \text{Bimod}_{P; Q} \\
 \mathcal{U} \downarrow & & \downarrow \mathcal{U} \\
 \Sigma \text{Bimod}_{P, Q>0} & \xrightarrow{ar_s} & \Sigma \text{Bimod}_{P, Q>0}.
 \end{array}$$

The vertical arrows denote the forgetful functor. This square commutes by the same argument as in the case of operads, see [19, Theorem 8.4.12]. One checks it first for free bimodules: $ar_s \mathcal{F}_{P; Q}^\Lambda(L) = \mathcal{F}_{P; Q}^\Lambda(L_{\leq s})$ and $ar_s \mathcal{F}_{P; Q>0}^\Sigma(L) = \mathcal{F}_{P; Q>0}^\Sigma(L_{\leq s})$. Here, $L_{\leq s}$ denotes the sequence

$$\begin{cases} L(n), & n \leq s; \\ \emptyset, & n > s. \end{cases}$$

It follows from (49) that the square commutes for free bimodules. The forgetful functor \mathcal{U} preserves colimits as it has a right adjoint by Proposition 3.14b for $Q_1 = Q, Q_2 = Q_{>0}, Y = Q$. So do the functors ar_s . On the other hand, any bimodule can naturally be seen as a coequalizer of free ones. The adjunction (69) is a Quillen adjunction as tr_s preserves fibrations and equivalences. As a consequence, $ar_s : \Sigma\text{Bimod}_{P, Q_{>0}} \rightarrow \Sigma\text{Bimod}_{P, Q_{>0}}$ preserves cofibrations.

Construction of the lift by induction. We work out our lifting problem (64) by an inductive construction on arity. By definition, one has $ar_0(N) = N_0$ (that is $N(0)$ in arity 0 and the empty set otherwise) and $ar_0(M) = M_0$. Consequently, the bimodule map $\varphi_0 : ar_0(N) \rightarrow A$ is obtained as a lift of the following diagram of P -algebras:

$$\begin{array}{ccc} M(0) & \xrightarrow{i_0} & A(0) \\ \phi_0 \downarrow & & \downarrow p_0 \\ N(0) & \xrightarrow{j_0} & B(0) \end{array}$$

Such a lift exists. Indeed, the left vertical map is a cofibration of P -algebras since the arity functor preserves cofibrations. Moreover, since p is an acyclic Reedy fibration, the map $A(0) \rightarrow B(0) \times_{\mathcal{M}(B)(0)} \mathcal{M}(A)(0) = B(0)$ is an acyclic Serre fibration.

Then we assume that the bimodule map $\varphi_{s-1} : ar_{s-1}(N) \rightarrow A$ is well defined. We consider the following diagram in the category of reduced $(P-Q)$ -bimodules:

$$\begin{array}{ccc} ar_s(M) \coprod_{ar_{s-1}(M)} ar_{s-1}(N) & \xrightarrow{(i\circ\iota; \varphi_{s-1})} & A \\ (ar_s(\phi); \iota) \downarrow & \dashrightarrow \exists \varphi_s? & \downarrow \simeq p \\ ar_s(N) & \xrightarrow{j\circ\iota} & B \end{array} \tag{70}$$

where the upper horizontal map is defined using the map $i \circ \iota : ar_s(M) \rightarrow M \rightarrow A$ on the first summand and the map $\varphi_{s-1} : ar_{s-1}(N) \rightarrow A$ on the second summand. By applying the identifications $ar_{s-1} \circ ar_s = ar_{s-1}$ and $ar_s \circ ar_s = ar_s$ together with the adjunction (68), we get that the lifting problem (70) becomes equivalent to a lifting problem of the following form in the category of reduced $(P-Q)$ -bimodules:

$$\begin{array}{ccc} ar_s(M) & \xrightarrow{F(i\circ\iota)} & \text{cosk}_s(A) \\ \downarrow ar_s(\phi) & \dashrightarrow \exists \tilde{\varphi}_s & \downarrow \simeq \pi \times \text{cosk}_s(p) \\ ar_s(N) & \xrightarrow{F(\varphi_{s-1}) \times F(ar_s(j))} & \text{cosk}_{s-1}(A) \times_{\text{cosk}_{s-1}(B)} \text{cosk}_s(B). \end{array} \tag{71}$$

The left vertical map is a cofibration in $\Sigma\text{Bimod}_{P, Q_{>0}}$ since the arity functor preserves cofibrations. On the other hand, the right vertical map is an acyclic Serre fibration in both the projective model category of Σ -sequences and the Reedy model categories of Λ -sequences, see the proof of [19, Theorem 8.3.20]. In that proof the second author has a similar diagram (loc. cit. equation (6)), but in the category of Λ -sequence. The coskeleton functors are defined on the category ΛSeq by the same formula (65). Thus, the second author shows that for any

Reedy fibration $A \rightarrow B$ of Λ -sequences, the induced map

$$\text{cosk}_s(A) \rightarrow \text{cosk}_{s-1}(A) \times_{\text{cosk}_{s-1}(B)} \text{cosk}_s(B)$$

is an acyclic both Reedy and (therefore) Serre fibration. As a consequence the lift $\tilde{\varphi}_s$ of Diagram (71) exists in the category of $(P\text{-}Q_{>0})$ -bimodules.

The morphism φ is compatible with the Λ -structure. By definition, let us remark that one has the identifications $\text{cosk}_{s-1}(N)(s) = \mathcal{M}(N)(s)$ and $\text{cosk}_{s-1}(A)(s) = \mathcal{M}(A)(s)$. By construction of the map F from (68) applied to $\varphi_{s-1} : ar_{s-1} \circ ar_s(N)(s) \rightarrow A(s)$, on has the factorization

$$F(\varphi_{s-1}) : ar_s(N)(s) = N(s) \longrightarrow \mathcal{M}(N)(s) \xrightarrow{\mathcal{M}(p)} \mathcal{M}(A)(s).$$

Consequently, the commutativity of diagram (71) implies the commutativity of the square:

$$\begin{array}{ccc} ar_s(N)(s) = N(s) & \longrightarrow & \text{cosk}_s(A)(s) = A(s) \\ \downarrow & & \downarrow \pi \\ \mathcal{M}(N)(s) & \longrightarrow & \text{cosk}_{s-1}(A)(s) = \mathcal{M}(A)(s). \end{array}$$

It follows that our morphism $\tilde{\varphi}_s$ intertwines the action of the restriction operators $h : [n_1] \rightarrow [n_2]$. This proves that $\tilde{\varphi}_s$ and, therefore, φ_s are morphisms of reduced bimodules preserving the Λ -structure. Thus so is $\varphi = \lim_s \varphi_s$.

Conversely. The forgetful functor $\mathcal{U} : \Lambda\text{Bimod}_{P;Q} \rightarrow \Sigma\text{Bimod}_{P,Q_{>0}}$ preserves colimits. Therefore, it is enough to check the statement for generating cofibrations in $\Lambda\text{Bimod}_{P;Q}$. The latter are given by applying the free functor $\mathcal{F}_{P;Q}^\Lambda$ to the generating cofibrations of ΛSeq_P . On the other hand, Λ -cofibrations are always Σ -cofibrations and the free functor $\mathcal{F}_{P;Q}^\Lambda$ agrees with $\mathcal{F}_{P;Q}^\Sigma$, see (49). \square

3.1.3 Left Properness and Extension/Restriction Functors for the Reedy Model Structure

As seen in Sect. 2.2, under some conditions on the operads, the projective model category of bimodules is relatively left proper and the extension/restriction adjunctions are Quillen equivalences. In what follows, we show that the Reedy model category inherits the same properties as a consequence of the characterization of Reedy cofibrations in the previous section.

Theorem 3.7 *Let P be an operad and Q be a reduced and well-pointed operad. The Reedy model category $\Lambda\text{Bimod}_{P;Q}$ is right proper. If $P(0) \in \text{Top}$ is a cofibrant space, $P_{>0} \in \Sigma\text{Operad}$ is a cofibrant operad and Q is componentwise cofibrant, then $\Lambda\text{Bimod}_{P;Q}$ is left proper relative to the class \mathcal{S} of componentwise cofibrant bimodules for which the arity zero left action map γ_0 is a cofibration. In the latter case, cofibrations with domain in \mathcal{S} are componentwise cofibrations, implying that the class of objects \mathcal{S} is closed under cofibrations. If in addition Q is Σ -cofibrant, then cofibrations with domain in the subclass $\mathcal{S}^\Sigma \subset \mathcal{S}$ of Σ -cofibrant objects are Σ -cofibrations, and \mathcal{S}^Σ is also closed under cofibrations.*

Proof The first statement is proved in exactly the same way as Theorem 2.14 using the facts that a Reedy fibration is always a projective (componentwise) fibration and that pullbacks

(and more generally limits) in the category of bimodules are defined componentwise and the fact that the category of spaces is right proper. For the other statements, we first recall that the forgetful functor $\mathcal{U}: \Lambda\text{Bimod}_{P; Q} \rightarrow \Sigma\text{Bimod}_{P; Q_{>0}}$ preserves colimits.³ In particular, pushouts are sent to pushouts. By Theorem 3.6, it also creates cofibrations. The statements then follow from the analogous statements Theorems 2.16 and 2.18 for $\Sigma\text{Bimod}_{P; Q_{>0}}$. \square

Let $\phi_1: P \rightarrow P'$ be a weak equivalence of operads and $\phi_2: Q \rightarrow Q'$ be a weak equivalence of reduced operads. Similarly to Sect. 2.2.2, we show that, under some conditions, the Reedy model categories of $(P-Q)$ -bimodules and $(P'-Q')$ -bimodules are Quillen equivalent. By abuse of notation, we denote by ϕ^* and $\phi_!$ the restriction functor and the extension functor, respectively, between the Reedy model categories:

$$\phi_!: \Lambda\text{Bimod}_{P; Q} \rightleftarrows \Lambda\text{Bimod}_{P'; Q'}: \phi^*. \tag{72}$$

In the same way as in Sect. 2.2.2, for any $M \in \Lambda\text{Bimod}_{P; Q}$ and $M' \in \Lambda\text{Bimod}_{P'; Q'}$, one has

$$\begin{aligned} \phi_!(M) &= \{\phi_!(M)(n) = \mathcal{F}_{P', Q'}^\Lambda(\mathcal{U}^\Lambda(M))(n) / \sim, n \geq 0\}, \\ \phi^*(M') &= \{\phi^*(M')(n) = M'(n), n \geq 0\}. \end{aligned}$$

Theorem 3.8 *Let $\phi_1: P \rightarrow P'$ be a weak equivalences of Σ -cofibrant operads and $\phi_2: Q \rightarrow Q'$ be a weak equivalence between reduced and componentwise cofibrant operads. Then one has Quillen equivalences*

$$\phi_!: \Lambda\text{Bimod}_{P; Q} \rightleftarrows \Lambda\text{Bimod}_{P'; Q'}: \phi^*, \tag{73}$$

$$\phi_!: T_r\Lambda\text{Bimod}_{P; Q} \rightleftarrows T_r\Lambda\text{Bimod}_{P'; Q'}: \phi^*. \tag{74}$$

Proof Since the restriction functor creates weak equivalences, one has a Quillen equivalence if, for any Reedy cofibrant object M in $\Lambda\text{Bimod}_{P; Q}$, the adjunction unit

$$M \longrightarrow \phi^*(\phi_!(M)) \tag{75}$$

is a weak equivalence. To prove it, we consider the adjunction $\tilde{\phi}_!: \Sigma\text{Bimod}_{P; Q_{>0}} \rightleftarrows \Sigma\text{Bimod}_{P'; Q'_{>0}}: \tilde{\phi}^*$ induced by the weak equivalences of operads $\phi_1: P \rightarrow P'$ and $\phi_2^{>0}: Q_{>0} \rightarrow Q'_{>0}$. By construction, one has the identity

$$\phi^*(\phi_!(M)) = \tilde{\phi}^*(\tilde{\phi}_!(M)).$$

By Theorem 2.19, the extension/restriction adjunction $(\tilde{\phi}_!, \tilde{\phi}^*)$ is a Quillen equivalence. Moreover, thanks to the characterization of Reedy cofibrations, M is also cofibrant in the projective model category $\Sigma\text{Bimod}_{P, Q_{>0}}$ and the map $M \longrightarrow \tilde{\phi}^*(\tilde{\phi}_!(M)) = \phi^*(\phi_!(M))$ is a weak equivalence. \square

3.2 The Connection Between the Model Category Structures on Bimodules

Similarly to the operadic case in [21], we build a Quillen adjunction between the projective model category of $(P-Q)$ -bimodules and the Reedy model category of $(P-Q)$ -bimodules,

³ In fact the forgetful functor from the category of $(P-Q)$ -bimodules to left P -modules creates colimits. It preserves colimits as it has a right adjoint by Proposition 3.14b for $Q_1 = Q, Q_2 = \mathbb{1}, Y = Q$. The fact that a cocone in bimodules is a colimit if it is a colimit in the category of left modules is easily verified.

where P is an operad and Q is a reduced operad. Furthermore, if M and N are two bimodules, then we show that there is a weak equivalence between the derived mapping spaces:

$$\Sigma \text{Bimod}_{P; Q}^h(M; N) \simeq \Lambda \text{Bimod}_{P; Q}^h(M; N).$$

For completeness of exposition, the last subsection is devoted to adapt the Boardman–Vogt resolution (well known for operads, see [7]) to the context of bimodules. We refer the reader to [13] for a detailed account of this construction.

3.2.1 Quillen Adjunction Between the Model Category Structures

Let P be an operad and Q be a well-pointed reduced operad. The projective and the Reedy model categories of bimodules have the same class of weak equivalences and induce the same homotopy category. Consequently, one has the following statement about the adjunctions

$$\begin{aligned} id : \Sigma \text{Bimod}_{P; Q} &\rightleftarrows \Lambda \text{Bimod}_{P; Q} : id, \\ id : T_r \Sigma \text{Bimod}_{P; Q} &\rightleftarrows T_r \Lambda \text{Bimod}_{P; Q} : id. \end{aligned} \tag{76}$$

Theorem 3.9 *For any operad P and any well-pointed reduced operad Q , the pairs of functors (76) form Quillen equivalences. Furthermore, for any pair $M, N \in \Lambda \text{Bimod}_{P; Q}$, one has*

$$\Sigma \text{Bimod}_{P; Q}^h(M, N) \simeq \Lambda \text{Bimod}_{P; Q}^h(M, N). \tag{77}$$

Moreover, if $M, N \in T_r \Lambda \text{Bimod}_{P; Q}$, with $r \geq 0$, then one has

$$T_r \Sigma \text{Bimod}_{P; Q}^h(M, N) \simeq T_r \Lambda \text{Bimod}_{P; Q}^h(M, N). \tag{78}$$

Proof In order to prove that the pairs of functors (76) form Quillen adjunctions we check that the right adjoint functors preserve fibrations and acyclic fibrations. Let $f : M \rightarrow N$ be an (acyclic) fibration in the Reedy model category of bimodules. By definition, the map f is a fibration if the corresponding map $\mathcal{U}^\Delta(f)$ in the category of Λ -sequences is a fibration. In other words, it means that the maps

$$M(n) \longrightarrow \mathcal{M}(M)(n) \times_{\mathcal{M}(N)(n)} N(n), \quad \text{with } n \in \mathbb{N}, \tag{79}$$

are (acyclic) Serre fibrations. On the other hand, the map $id(f)$ is a fibration in the projective model category of bimodules if the maps $M(n) \rightarrow N(n)$, with $n \in \mathbb{N}$, are Serre fibrations. According to the notation introduced in Sect. 1.2, the pair of functors

$$\Lambda[-] : \Sigma \text{Seq} \rightleftarrows \Lambda \text{Seq} : \mathcal{U}(-)$$

forms a Quillen adjunction (see [19, Theorem 8.3.20]). In particular, the forgetful functor \mathcal{U} preserves (acyclic) fibrations. As a consequence, if the maps (79) are (acyclic) Serre fibrations, then the induced maps $M(n) \rightarrow N(n)$, with $n \in \mathbb{N}$, are (acyclic) Serre fibrations. Furthermore, the Quillen adjunction so obtained is obviously a Quillen equivalence since we consider identity functors. Finally, the identities (77) and (78) are induced by taking a projective cofibrant replacement of M and a Reedy fibrant replacement of N and using the fact that any projectively cofibrant object is Reedy cofibrant and any Reedy fibrant object is projectively fibrant. □


3.2.2 Boardman–Vogt Type Resolution in the Projective/Reedy Model Category

From a $(P-Q)$ -bimodule M , we build a $(P-Q)$ -bimodule $\mathcal{B}_{P;Q}(M)$. The points of $\mathcal{B}_{P;Q}(M)(n)$, $n \geq 0$, are equivalence classes $[T; \{t_v\}; \{p_v\}; \{m_v\}; \{q_v\}]$, where $T \in \mathcal{S}\mathbb{P}_n$ (see Sect. 2.1.2) is a tree with section while $\{p_v\}_{v \in V^d(T)}$, $\{m_v\}_{v \in V^p(T)}$ and $\{q_v\}_{v \in V^u(T)}$ are families of points labelling the vertices below the section, on the section and above the section, respectively, by points in P , M and Q , respectively. Furthermore, $\{t_v\}_{v \in V(T) \setminus V^p(T)}$ is a family of real numbers in the interval $[0, 1]$ indexing the vertices which are not pearls. If e is an inner edge above the section, then $t_{s(e)} \geq t_{t(e)}$. Similarly, if e is an inner edge below the section, then $t_{s(e)} \leq t_{t(e)}$. In other words, closer to a pearl is a vertex, smaller is the corresponding number. The space $\mathcal{B}_{P;Q}(M)(n)$ is a quotient of the subspace of


$$\coprod_{T \in \mathcal{S}\mathbb{P}_n} \prod_{v \in V^p(T)} M(|v|) \times \prod_{v \in V^d(T)} [P(|v|) \times [0, 1]] \times \prod_{v \in V^u(T)} [Q(|v|) \times [0, 1]] \quad (80)$$

determined by the restrictions on the families $\{t_v\}$. The equivalence relation is generated by the conditions:


- (i) If a vertex is labelled by a unit in $P(1)$ or $Q(1)$, then locally one has the identity




- (ii) If a vertex is indexed by $a \cdot \sigma$, with $\sigma \in \Sigma$, then



- (iii) If two consecutive vertices, connected by an edge e , are indexed by the same real number $t \in [0, 1]$, then e is contracted using the operadic structures of P and Q . The vertex so obtained is indexed by the real number t .



- (iv) If a vertex above the section is indexed by 0, then its output edge is contracted by using the right module structure. Similarly, if a vertex below the section is indexed by 0, then all its incoming edges are contracted by using the left module structure. In both cases the new vertex becomes a pearl.







Fig. 15 Illustration of the left module structure $\gamma_{\mathcal{L}} : P(2) \times \mathcal{B}_{P; Q}(M)(3) \times \mathcal{B}_{P; Q}(M)(0) \rightarrow \mathcal{B}_{P; Q}(M)(3)$

- (v) If a univalent pearl is indexed by a point of the form $\gamma_0(x)$, with $x \in P(0)$, then we contract its output edge by using the operadic structure of P . In particular, if all the pearls connected to a vertex v are univalent and of the form $\gamma_0(x_v)$, then the vertex is identified to the pearled corolla with no input.



Let us describe the $(P-Q)$ -bimodule structure. Let $q \in Q(n)$ and $[T; \{t_v\}; \{p_v\}; \{m_v\}; \{q_v\}]$ be a point in $\mathcal{B}_{P; Q}(M)(m)$. The right operation $[T; \{t_v\}; \{p_v\}; \{m_v\}; \{q_v\}] \circ^i q$ consists in grafting the n -corolla labelled by q to the i -th incoming edge of T and indexing the new vertex by 1. Similarly, let $[T_i; \{t_v^i\}; \{p_v^i\}; \{m_v^i\}; \{q_v^i\}]$ be a family of points in the spaces $\mathcal{B}_{P; Q}(M)(m_i)$. The left module structure over P is defined as follows: each tree of the family is grafted to a leaf of the n -corolla labelled by $p \in P(n)$ from left to right. The new vertex, coming from the n -corolla, is indexed by 1 (see Fig. 15).

One has an obvious inclusion of Σ -sequences $\iota : M \rightarrow \mathcal{B}_{P; Q}(M)$, where each element $m \in M(n)$ is sent to an n -corolla labelled by m , whose only vertex is a pearl. Furthermore, the map

$$\mu : \mathcal{B}_{P; Q}(M) \rightarrow M ; [T; \{t_v\}; \{p_v\}; \{m_v\}; \{q_v\}] \mapsto [T; \{0\}; \{p_v\}; \{m_v\}; \{q_v\}], \tag{81}$$

is defined by sending the real numbers indexing the vertices to 0. The obtained element is identified to the pearled corolla labelled by a point in M . It is easy to see that μ is a $(P-Q)$ -bimodule map.

In order to get resolutions for truncated bimodules, one considers a filtration in $\mathcal{B}_{P; Q}(M)$ according to the number of *geometrical inputs* which is the number of leaves plus the number of univalent vertices above the section. A point in $\mathcal{B}_{P; Q}(M)$ is said to be *prime* if the real numbers indexing the vertices are strictly smaller than 1. Otherwise, a point is said to be *composite* and can be decomposed into *prime components* as shown in Fig. 16. More precisely, the prime components are obtained by removing the vertices indexed by 1.

A prime point is in the r -th filtration layer $\mathcal{B}_{P; Q}(M)_r$ if the number of its geometrical inputs is at most r . Similarly, a composite point is in the r -th filtration layer if its all prime components are in $\mathcal{B}_{P; Q}(M)_r$. For instance, the composite point in Fig. 16 is in the filtration layer $\mathcal{B}_{P; Q}(M)_6$. For each r , $\mathcal{B}_{P; Q}(M)_r$ is a $(P-Q)$ -bimodule and one has the following filtration of $\mathcal{B}_{P; Q}(M)$:

$$\mathcal{B}_{P; Q}(M)_0 \longrightarrow \mathcal{B}_{P; Q}(M)_1 \longrightarrow \dots \longrightarrow \mathcal{B}_{P; Q}(M)_{r-1} \longrightarrow \mathcal{B}_{P; Q}(M)_r \longrightarrow \dots \longrightarrow \mathcal{B}_{P; Q}(M). \tag{82}$$




Fig. 16 A composite point and its prime components




Fig. 17 Illustration of the right action by $*_0$

Theorem 3.10 (Theorem 2.12 in [13]) *Assume that P and Q are Σ -cofibrant operads, and M is a Σ -cofibrant $(P-Q)$ -bimodule for which the arity zero left action map $\gamma_0 : P(0) \rightarrow M(0)$ is a cofibration. Then the objects $\mathcal{B}_P;_Q(M)$ and $T_r\mathcal{B}_P;_Q(M)_r$ are cofibrant replacements of M and T_rM in the categories $\Sigma\text{Bimod}_P;_Q$ and $T_r\Sigma\text{Bimod}_P;_Q$, respectively. In particular, the maps μ and $T_r\mu|_{T_r\mathcal{B}_P;_Q(M)_r}$ are weak equivalences.*

Now we change slightly the above construction in order to produce Reedy cofibrant replacements for $(P-Q)$ -bimodules when Q is a reduced operad. Let M be a $(P-Q)$ -bimodule. We consider the Σ -sequence

$$\mathcal{B}_P^\Lambda;_Q(M) := \mathcal{B}_P;_{Q_{>0}}(M).$$

The superscript Λ is to emphasize that we get a cofibrant replacement in the Reedy model category structure. The left action and the positive arity right action are defined in the same way as for $\mathcal{B}_P;_{Q_{>0}}(M)$. The right action by $*_0 \in Q(0)$ is defined in the obvious way as the right action by $*_0$ on a in the vertex (a, t) connected to the leaf labelled by i as illustrated in the Fig. 17.

Note that since the arity zero component of $Q_{>0}$ is empty, in the union (80) we can consider only trees whose all non-pearl vertices have arities ≥ 1 . We denote this set by $s\mathbb{P}_n^{\geq 1}$. In other words, the space $\mathcal{B}_P^\Lambda;_Q(M)$ can be obtained as the restriction of the coproduct (80) to this set.

Proposition 3.11 *Assume that P and Q are Σ -cofibrant topological operads with $Q(0) = *$, and M is a Σ -cofibrant $(P-Q)$ -bimodule for which the arity zero left action map $\gamma_0 : P(0) \rightarrow M(0)$ is a cofibration. Then the objects $\mathcal{B}_P^\Lambda;_Q(M)$ and $T_r\mathcal{B}_P^\Lambda;_Q(M)$ are cofibrant replacements of M and T_rM in the categories $\Lambda\text{Bimod}_P;_Q$ and $T_r\Lambda\text{Bimod}_P;_Q$, respectively. In particular, the maps μ and $T_r\mu$ are weak equivalences.*

Proof The map $\mu : \mathcal{B}_P^\Lambda;_Q(M) = \mathcal{B}_P;_{Q_{>0}}(M) \rightarrow M$, which sends the real numbers indexing the vertices to 0, is a homotopy equivalence as shown by Theorem 3.10. Furthermore, by the same theorem, $\mathcal{B}_P^\Lambda;_Q(M) = \mathcal{B}_P;_{Q_{>0}}(M)$ is cofibrant in the projective model category of $(P-Q_{>0})$ -bimodules. Due to Theorem 3.6, $\mathcal{B}_P^\Lambda;_Q(M)$ is also Reedy cofibrant and it gives rise to a cofibrant resolution of M in the Reedy model category $\Lambda\text{Bimod}_P;_Q$. The same

arguments work for the truncated case. Note that $T_r \mathcal{B}_{P; Q}^\Lambda(M)_r = T_r \mathcal{B}_{P; Q}^\Lambda(M)$ since arity zero vertices above the section are not permitted. \square

3.2.3 A Functorial Cofibrant Replacement in the Projective/Reedy Model Category

In the previous subsection we described a construction of projective and Reedy cofibrant replacements provided the bimodule and the acting operads are Σ -cofibrant. In this subsection we explain how that construction can be used to functorially define a cofibrant replacement without any assumption on the bimodule, while assuming that the right-acting operad Q has cofibrant components and the left-acting operad P is Σ -cofibrant.

Given a $(P-Q)$ -bimodule M , we first replace it by the Σ -sequence $M' := |S_\bullet M|$, whose n -th space is the realization of the simplicial set of singular simplices in $M(n)$. The obtained object is a $(P'-Q')$ -bimodule, where the operads P' and Q' are similarly defined as $P' := |S_\bullet P|$, $Q' := |S_\bullet Q|$. Let E_∞ as usual denote a reduced Σ -cofibrant model of the commutative operad. Define Σ -sequences $M'_\infty, P'_\infty, Q'_\infty$ as objectwise products

$$M'_\infty(n) := M'(n) \times E_\infty(n), \quad P'_\infty(n) := P'(n) \times E_\infty(n), \quad Q'_\infty(n) := Q'(n) \times E_\infty(n).$$

We get that M'_∞ is a Σ -cofibrant bimodule over a pair (P'_∞, Q'_∞) of Σ -cofibrant operads. We can therefore apply the construction from the previous subsection.

For the following theorem denote by

$$\phi_1: P'_\infty \xrightarrow{\cong} P' \xrightarrow{\cong} P, \quad \phi_2: Q'_\infty \xrightarrow{\cong} Q' \xrightarrow{\cong} Q$$

the natural induced equivalences of operads. Note that

$$\phi_0: M'_\infty \xrightarrow{\cong} M' \xrightarrow{\cong} M$$

is an equivalence of $(P'_\infty-Q'_\infty)$ -bimodules.

Proposition 3.12 (a) *Assume that P is a Σ -cofibrant operad and Q is a componentwise cofibrant operad. Let M be any $(P-Q)$ -bimodule for which the arity zero left action map $\gamma_0: P(0) \rightarrow M(0)$ is injective. Then the objects $\phi_!(\mathcal{B}_{P'_\infty; Q'_\infty}(M'_\infty))$ and $\phi_!(T_r \mathcal{B}_{P'_\infty; Q'_\infty}(M'_\infty)_r)$ are cofibrant replacements of M and $T_r M$ in the categories $\Sigma \text{Bimod}_{P; Q}$ and $T_r \Sigma \text{Bimod}_{P; Q}$, respectively.*

(b) *Assume in addition that Q is reduced. Then the objects $\phi_!(\mathcal{B}_{P'_\infty; Q'_\infty}^\Lambda(M'_\infty))$ and $\phi_!(T_r \mathcal{B}_{P'_\infty; Q'_\infty}^\Lambda(M'_\infty))$ are cofibrant replacements of M and $T_r M$ in the categories $\Lambda \text{Bimod}_{P; Q}$ and $T_r \Lambda \text{Bimod}_{P; Q}$, respectively.*

Proof The result is an immediate consequence of Theorems 2.19, 3.8, 3.10 and Proposition 3.11. \square

3.3 The Reedy Fibrant Replacement as an Internal Hom

In this subsection, we provide a more conceptual understanding of the Reedy fibrant replacement described in Sect. 3.1.1. More precisely we explain this construction in terms of internal hom in the category of Σ -sequences, see Proposition 3.15.

3.3.1 The Right Closed Monoidal Category of Symmetric Sequences

It is well known and appears in almost any textbook on the theory of operads that the category ΣSeq of Σ -sequences has a monoidal structure $(\Sigma Seq, \circ, \mathbb{1})$ with the unit

$$\mathbb{1}(k) = \begin{cases} *, & k = 1; \\ \emptyset, & k \neq 1; \end{cases}$$

and the composition product

$$(X \circ Y)(k) = \coprod_{n \geq 0} X(n) \times_{\Sigma_n} \coprod_{\beta: [k] \rightarrow [n]} \prod_{i=1}^n Y(|\beta^{-1}(i)|).$$

Monoids in $(\Sigma Seq, \circ, \mathbb{1})$ are usual topological operads.

It is less known that ΣSeq is *closed* with respect to this monoidal structure. This fact is true for the category of Σ -sequences in any bicomplete closed symmetric monoidal category and was noticed by G. M. Kelly back in the 1970s [30]. More recently this also appeared in [35, Section 2.2] and in [24, Section 3].⁴ This means that ΣSeq is endowed with an internal hom functor

$$[-, -]: \Sigma Seq^{op} \times \Sigma Seq \rightarrow \Sigma Seq,$$

such that for any $X \in \Sigma Seq$, the functor $(-) \circ X$ is left adjoint to $[X, -]$. Sometimes this structure on a category is called *right* closed monoidal instead of just closed monoidal as a monoidal product with an object on the *right* has an adjoint. Explicitly,

$$[X, Y](k) = \prod_{n \geq 0} \left[\prod_{\alpha: [n] \rightarrow [k]} \text{Map} \left(\prod_{i=1}^k X(|\alpha^{-1}(i)|), Y(n) \right) \right]^{\Sigma_n}. \tag{83}$$

3.3.2 The Tensor-Hom Adjunction

For concreteness all the statements in this subsection are made for the category ΣSeq . One should mention, however, that they hold true for any bicomplete right closed monoidal category. (One only needs to replace the word “operad” by “monoid”).

Lemma 3.13 *Let $P, Q \in \Sigma Operad$ and $X, Y \in \Sigma Seq$. Consider the internal hom object $[X, Y] \in \Sigma Seq$.*

- (a) *If X is a left Q -module, then $[X, Y]$ is a right Q -module.*
- (b) *If Y is a left P -module, then $[X, Y]$ is a left P -module.*
- (c) *If X is a left Q -module and Y is a left P -module, then $[X, Y]$ is a $(P-Q)$ -bimodule.*

Sketch of the proof For (a), the right action map $[X, Y] \circ Q \rightarrow [X, Y]$ is the adjoint to the composition

$$[X, Y] \circ Q \circ X \xrightarrow{id_{[X, Y]} \circ \mu_X} [X, Y] \circ X \xrightarrow{ev_{X, Y}} Y.$$

Here $\mu_X: Q \circ X \rightarrow X$ is the left Q -action on X , and $ev_{X, Y}$ is the adjoint to the identity map $id_{[X, Y]}$.

⁴ We also refer to [2, Definition 1.20], where this structure appears implicitly and from where our formula (83) is borrowed.

For (b), the left action map $P \circ [X, Y] \rightarrow [X, Y]$ is adjoint to the composition

$$P \circ [X, Y] \circ X \xrightarrow{id_P \circ ev_{X,Y}} P \circ Y \xrightarrow{\mu_Y} Y.$$

Here μ_Y is the left P -action on Y .

The facts that these formulae correctly define P and Q actions are easily checked as well as the fact that these actions commute in case of (c). □

If X is a right module over an operad Q and Y is a left Q -module, one defines $X \circ_Q Y \in \Sigma Seq$ as the coequalizer

$$X \circ_Q Y = \text{coeq}(X \circ Q \circ Y \rightrightarrows X \circ Y),$$

where both arrows are $\mu_X \circ id_Y$ and $id_X \circ \mu_Y$.

In case X and Y are both right modules over an operad Q , one defines $[X, Y]_Q \in \Sigma Seq$ as the equalizer

$$[X, Y]_Q = \text{eq}([X, Y] \rightrightarrows [X \circ Q, Y]),$$

where the upper map is induced by the right Q -action $\mu_X : X \circ Q \rightarrow X$, and the lower map is the adjoint to the composition

$$[X, Y] \circ X \circ Q \xrightarrow{ev_{X,Y} \circ id_Q} Y \circ Q \xrightarrow{\mu_Y} Y.$$

Proposition 3.14 ([24, Proposition 5.22]) *Let $Q_1, Q_2, P \in \Sigma Operad$ and $Y \in \Sigma Bimod_{Q_1, Q_2}$.*

(a) *One has an adjunction between the categories of right Q_1 and Q_2 modules*

$$(-) \circ_{Q_1} Y : \Sigma RMod_{Q_1} \rightleftarrows \Sigma RMod_{Q_2} : [Y, -]_{Q_2}. \tag{84}$$

(b) *One has an adjunction between the categories of $(P-Q_1)$ and $(P-Q_2)$ bimodules*

$$(-) \circ_{Q_1} Y : \Sigma Bimod_{P, Q_1} \rightleftarrows \Sigma Bimod_{P, Q_2} : [Y, -]_{Q_2}. \tag{85}$$

Sketch of the proof The statements are general and hold in any bicomplete (right) closed monoidal category. The proof is essentially a categorification of the tensor-hom adjunction between the categories of (bi)modules over rings. □

3.3.3 The Reedy Fibrant Replacement

Let N be a Σ -cofibrant right module over a Σ -cofibrant operad Q . Viewed as a $(\mathbb{1}-Q)$ -bimodule, we consider its resolution $\mathcal{B}_{\mathbb{1};Q}(N)$, see Sect. 3.2.2, which is its cofibrant replacement as a right Q -module. It is easy to see that in case N is a $(P-Q)$ -bimodule for some operad P , the sequence $\mathcal{B}_{\mathbb{1};Q}(N)$ has also a natural structure of a $(P-Q)$ -bimodule. Denote by $Q^c := \mathcal{B}_{\mathbb{1};Q}(Q)$. One can show that Q^c is a cofibrant replacement of Q as a Q -bimodule. Roughly speaking it is because even before taking its resolution, Q is cofibrant as a left module over itself.

We leave the following proposition as an exercise to the reader. For this proposition we do not assume that Q is Σ -cofibrant.

Proposition 3.15 *Let $M \in \Lambda Bimod_{P,Q}$, let M^f be its Reedy fibrant replacement as defined in Sect. 3.1.1, and let $Q^c := \mathcal{B}_{\mathbb{1};Q}(Q)$, see Sect. 3.2.2. One has an isomorphism of $(P-Q)$ -bimodules:*

$$M^f = [Q^c, M]_Q.$$

Moreover, the fibrant replacement map $M \rightarrow M^f$ is induced by the (cofibrant, in case Q is Σ -cofibrant) replacement map $Q^c \rightarrow Q$:

$$M = [Q, M]_Q \rightarrow [Q^c, M]_Q.$$

As an interesting consequence we have the following.

Corollary 3.16 *For any $M \in \Lambda\text{Bimod}_{P;Q}$, one has a homeomorphism $(M^f)^f \cong M^f$ of fibrant replacements.*

Proof It follows from Proposition 3.15 and adjunction (85) that

$$(M^f)^f = [Q^c, [Q^c, M]_Q]_Q \cong [Q^c \circ_Q Q^c, M]_Q.$$

On the other hand, it is easy to see that $Q^c \circ_Q Q^c \cong Q^c$. To define an explicit homeomorphism $Q^c \xrightarrow{\cong} Q^c \circ_Q Q^c$, for each tree in Q^c we draw a horizontal line $t = 1/2$ and then replace all the real parameters in the vertices below the horizontal section $t \in [0, 1/2]$ by $2t$ and all the real parameters of the vertices above the section $t \in (1/2, 1]$ by $2t - 1$.⁵ □

3.4 The Subcategory of Reduced Bimodules

3.4.1 Main Properties

Let P and Q be two reduced operads. We can then consider the category $\Lambda_*\text{Bimod}_{P;Q}$ of reduced $(P\text{-}Q)$ -bimodules. This category has been used by the first and third authors [15] in order to get delooping theorems for the Taylor tower approximations of mapping spaces avoiding multi-singularities, i.e. singularities depending on several points.

The present section is devoted to adapt the constructions and theorems introduced in the previous sections to the category $\Lambda_*\text{Bimod}_{P;Q}$. By considering only reduced bimodules, we can simplify some constructions. Since the proofs are almost the same, we list altogether the main statements. Note that the statements are slightly improved for the functorial cofibrant resolution and for the extension/restriction adjunction.

Theorem 3.17 *Let P and Q be reduced operads and, moreover, assume that Q is well-pointed.*

- (i) **Reedy model structure.** *The category $\Lambda_*\text{Bimod}_{P;Q}$ admits a cofibrantly generated model category structure, called the Reedy model category structure, transferred from the Reedy model category $\Lambda_{>0}\text{Seq}$ along the adjunction*

$$\mathcal{F}_{P;Q}^{\Lambda_*} : \Lambda_{>0}\text{Seq} \rightleftarrows \Lambda_*\text{Bimod}_{P;Q} : \mathcal{U}^{\Lambda},$$

where the free functor is given by

$$\mathcal{F}_{P;Q}^{\Lambda_*}(M)(n) := \begin{cases} \mathcal{F}_{P>0;Q>0}^{\Sigma}(M)(n), & n \geq 1, \\ *, & n = 0, \end{cases} \tag{86}$$

The model category so obtained makes the adjunction $(\mathcal{F}_{P;Q}^{\Lambda_*}, \mathcal{U}^{\Lambda})$ into a Quillen adjunction. Moreover, the fibrant coresolution functor introduced in Sect. 3.1.1 restricts

⁵ In fact for any right Q -module N , one has $\mathcal{B}_{\mathbb{1};Q}(N) = N \circ_Q Q^c$ and it is always true that $\mathcal{B}_{\mathbb{1};Q}(\mathcal{B}_{\mathbb{1};Q}(N)) \cong \mathcal{B}_{\mathbb{1};Q}(N)$.

to the category $\Lambda_*\text{Bimod}_{P; Q}$ giving rise to a functorial fibrant replacement. In case P and Q are componentwise cofibrant, the functor $M \mapsto \phi_! \left(\mathcal{B}_{P'_\infty; Q'_\infty}^\Lambda(M'_\infty) \right)$ of Proposition 3.12 gives rise to a functorial cofibrant resolution in $\Lambda_*\text{Bimod}_{P; Q}$.

- (ii) **Characterization of Reedy cofibrations.** A morphism $\phi : M \rightarrow N$ in the category $\Lambda_*\text{Bimod}_{P; Q}$ is a Reedy cofibration if and only if the corresponding map $\phi_{>0} : M_{>0} \rightarrow N_{>0}$ is a cofibration in the projective model category of $(P_{>0}-Q_{>0})$ -bimodules.
- (iii) **Left and Right properness.** The Reedy model category $\Lambda_*\text{Bimod}_{P; Q}$ is right proper. If P is Reedy cofibrant and Q is componentwise cofibrant, then $\Lambda_*\text{Bimod}_{P; Q}$ is left proper relative to the class of componentwise cofibrant bimodules. In the latter case, the class of componentwise cofibrant bimodules is closed under cofibrations. In particular, cofibrant bimodules are componentwise cofibrant. If in addition Q is Σ -cofibrant, the class of Σ -cofibrant bimodules is also closed under cofibrations and cofibrant bimodules are Σ -cofibrant.
- (iv) **Extension/restriction adjunctions.** Let $\phi_1 : P \rightarrow P'$ and $\phi_2 : Q \rightarrow Q'$ be weak equivalences between componentwise cofibrant reduced operads. One has a Quillen equivalence

$$\phi_! : \Lambda_*\text{Bimod}_{P; Q} \rightleftarrows \Lambda_*\text{Bimod}_{P'; Q'} : \phi^*$$

All the above statements admit truncated versions for the categories $T_r \Lambda_*\text{Bimod}_{P; Q}$ with $r \geq 0$.

Proof The proof of (i) is exactly the same as for the construction of the usual Reedy model category of $(P-Q)$ -bimodules and the construction of the fibrant coresolution. Note that for the cofibrant replacement functor we do not require P to be Σ -cofibrant, but only that it is componentwise cofibrant. The reason is that the argument of Proposition 3.12 uses the extension/restriction adjunction. The latter has also the same improved requirement that P and P' are componentwise cofibrant, see statement (iv) above.

Statement (ii) follows from the fact that the forgetful functor $\iota : \Lambda_*\text{Bimod}_{P; Q} \rightarrow \Lambda\text{Bimod}_{P; Q}$ creates cofibrations, see Proposition 3.18. Applying the characterization of Reedy cofibrations for bimodules of Theorem 3.6, a map $M \rightarrow N$ in $\Lambda_*\text{Bimod}_{P; Q}$ is a cofibration if and only if it is a cofibration in $\Sigma\text{Bimod}_{P; Q_{>0}}$. Since $P(0) = M(0) = N(0) = *$, the arity zero component can be naturally ignored, and the map in question is a cofibration if and only if $M_{>0} \rightarrow N_{>0}$ is one in $\Sigma\text{Bimod}_{P_{>0}; Q_{>0}}$ (or, equivalently, in $\Sigma_{>0}\text{Bimod}_{P_{>0}; Q_{>0}}$).

Statement (iii) follows from Theorem 3.7 and the fact that the inclusion functor ι creates weak equivalences, fibrations, pullbacks, and also cofibrations and pushouts, see Proposition 3.18.

We now prove (iv). Since the restriction functor creates weak equivalences, one has to check that, for any Reedy cofibrant object M in $\Lambda_*\text{Bimod}_{P; Q}$, the adjunction unit

$$M \longrightarrow \phi^*(\phi_!(M))$$

is a weak equivalence. Due to the characterization of Reedy cofibrations, $M_{>0}$ is also cofibrant in the projective model category $\Sigma_{>0}\text{Bimod}_{P_{>0}; Q_{>0}}$. Since $(\phi_1)_{>0} : P_{>0} \rightarrow P'_{>0}$ and $(\phi_2)_{>0} : Q_{>0} \rightarrow Q'_{>0}$ are still weak equivalences between componentwise cofibrant operads, by Theorem 2.22, the pair of functors $((\phi_{>0})_!; (\phi_{>0})^*)$ gives rise to a Quillen equivalence

$$(\phi_{>0})_! : \Sigma_{>0}\text{Bimod}_{P_{>0}; Q_{>0}} \rightleftarrows \Sigma_{>0}\text{Bimod}_{P'_{>0}; Q'_{>0}} : (\phi_{>0})^*$$

Therefore, the map $M_{>0} \rightarrow (\phi_{>0})^*((\phi_{>0})!(M_{>0}))(n)$ is a weak equivalence. The statement is a consequence of the identity

$$\phi^*(\phi!(M))_{>0} = (\phi_{>0})^*((\phi_{>0})!(M_{>0})).$$

□

3.4.2 Quillen Adjunction Between the Reedy Model Categories

Let P and Q be two reduced operads with Q well-pointed. The inclusion functor ι from the category of (possibly truncated) reduced bimodules into the category of (possibly truncated) bimodules has a left adjoint τ called the unitarization functor. This latter one consists in collapsing the arity zero component to a point and adjusting the other components according to the equivalence relation induced by this collapse:

$$\begin{aligned} \tau : \Lambda \text{Bimod}_{P; Q} &\rightleftarrows \Lambda_* \text{Bimod}_{P; Q} : \iota, \\ \tau : T_r \Lambda \text{Bimod}_{P; Q} &\rightleftarrows T_r \Lambda_* \text{Bimod}_{P; Q} : \iota. \end{aligned} \tag{87}$$

Proposition 3.18 *The pairs of functors (87) form Quillen adjunctions. Moreover, the inclusion functor ι creates weak equivalences, fibrations, cofibrations, limits, and colimits.*

Proof By construction, ι creates equivalences and fibrations. As a consequence, the adjunction (87) is a Quillen one. Since the limits are taken objectwise, and the limit of any point-constant diagram is a point, ι creates limits. Note that τ , as a left adjoint, preserves colimits and cofibrations, and $\tau \circ \iota = id$. Therefore, one only needs to check that ι preserves colimits and cofibrations.

For colimits, we again have to show that the colimit of any diagram in $\Lambda \text{Bimod}_{P; Q}$ with values in reduced bimodules is a reduced bimodule. The truncation functor $T_r : \Lambda \text{Bimod}_{P; Q} \rightarrow T_r \Lambda \text{Bimod}_{P; Q}$ preserves colimits as it admits a right adjoint, see Sect. 3.1.2. In the case $r = 0$, the category $T_0 \Lambda \text{Bimod}_{P; Q}$ is equivalent to the category of P -algebras in Top . The one-point space is a free P -algebra generated by the empty set $F_P(\emptyset)$. On the other hand, the free P -algebra functor $F_P : Top \rightarrow Alg_P$ preserves colimits and the colimit of any empty set constant diagram is the empty set. The statement follows.

Finally, for cofibrations, since ι creates colimits, it is enough to check that the generating cofibrations in $\Lambda_* \text{Bimod}_{P; Q}$ are cofibrations in $\Lambda \text{Bimod}_{P; Q}$. The generating cofibrations in the former are the maps

$$\mathcal{F}_P^{\Lambda_*; Q}(\partial X) \rightarrow \mathcal{F}_P^{\Lambda_*; Q}(X), \tag{88}$$

where $\partial X \rightarrow X$ is one of the generating cofibrations of $\Lambda_{>0}Seq$. For a $\Lambda_{>0}$ -sequence Y , denote by Y_+ the Λ -sequence that agrees with Y in positive arities and has one point in arity zero. From the formulas (86) and (49), it follows that $\mathcal{F}_P^{\Lambda_*; Q}(Y) = \mathcal{F}_P^{\Lambda; Q}(Y_+)$. The inclusion $(\partial X)_+ \rightarrow X_+$ is a Σ -cofibration and, as a consequence, is a $\Lambda SeqP$ -cofibration. Together with the fact that the functor $\mathcal{F}_P^{\Lambda; Q}$ preserves cofibrations, we conclude that the inclusion (88), or equivalently $\mathcal{F}_P^{\Lambda; Q}((\partial X)_+) \rightarrow \mathcal{F}_P^{\Lambda; Q}(X_+)$, is a cofibration in $\Lambda \text{Bimod}_{P; Q}$. □

Theorem 3.19 *Let P and Q be as in Theorem 3.17. The inclusion functors ι from (87) induce fully faithful inclusions of homotopy categories. Moreover, for any pair $M, N \in \Lambda_* \text{Bimod}_{P; Q}$, one has*

$$\Lambda_* \text{Bimod}_{P; Q}^h(M, N) \simeq \Lambda \text{Bimod}_{P; Q}^h(\iota M, \iota N) \simeq \Sigma \text{Bimod}_{P; Q}^h(\iota M, \iota N). \tag{89}$$

Furthermore, if $M, N \in T_r \Lambda_* \text{Bimod}_{P; Q}$, with $r \geq 0$, then one has

$$T_r \Lambda_* \text{Bimod}_{P; Q}^h(M, N) \simeq T_r \Lambda \text{Bimod}_{P; Q}^h(\iota M, \iota N) \simeq T_r \Sigma \text{Bimod}_{P; Q}^h(\iota M, \iota N). \tag{90}$$

Proof The first statement follows from the fact that the functors ι themselves are fully faithful inclusions of model categories preserving equivalences, fibrations, and cofibrations, see Proposition 3.18. The equivalences (89) and (90) follow from Theorem 3.9 and the fact that ι preserve cofibrant and fibrant replacements and are fully faithful. \square

4 The Projective Model Category of O Infinitesimal Bimodules

Let O be an operad. An O infinitesimal bimodule, or just an O -Ibimodule, is a Σ -sequence $M \in \Sigma \text{Seq}$ together with operations

$$\begin{aligned} \circ^i &: M(n) \times O(m) \longrightarrow M(n+m-1), \text{ right infinitesimal operations with } i \in \{1, \dots, n\}, \\ \circ_i &: O(n) \times M(m) \longrightarrow M(n+m-1), \text{ left infinitesimal operations with } i \in \{1, \dots, n\}, \end{aligned}$$

$$\tag{91}$$

satisfying compatibility relations with the symmetric group action as well as associativity and unit axioms. More precisely, for any integers $i \in \{1, \dots, n\}$, $j \in \{i+1, \dots, n\}$, $k \in \{1, \dots, m\}$ and any permutations $\sigma \in \Sigma_n$ and $\tau \in \Sigma_m$, one has the following commutative diagrams:

$$\begin{array}{ccc} M(n) \times O(m) \times O(\ell) & \xrightarrow{\circ_k} & M(n) \times O(m+\ell-1) \\ \circ^j \downarrow & & \downarrow \circ^j \\ M(n+m-1) \times O(\ell) & \xrightarrow{\circ^{k+i-1}} & M(n+m+\ell-2) \end{array}$$

Linear associativity for the right infinitesimal operations

$$\begin{array}{ccc} M(n) \times O(m) \times O(\ell) & \xrightarrow{\circ^j} & M(n+m-1) \times O(\ell) \\ \circ^j \downarrow & & \downarrow \circ^{j+m-1} \\ M(n+\ell-1) \times O(m) & \xrightarrow{\circ^j} & M(n+m+\ell-2) \end{array}$$

Ramified associativity for the right infinitesimal operations

$$\begin{array}{ccc} O(n) \times M(m) \times O(\ell) & \xrightarrow{\circ_i} & M(n+m-1) \times O(\ell) \\ \circ_j \downarrow & & \downarrow \circ^{j+m-1} \\ O(n+\ell-1) \times M(m) & \xrightarrow{\circ_i} & M(n+m+\ell-2) \end{array}$$

Ramified compatibility between the left and right operations 1

$$\begin{array}{ccc} O(n) \times O(m) \times M(\ell) & \xrightarrow{\circ_i} & O(n+m-1) \times M(\ell) \\ \circ_j \downarrow & & \downarrow \circ^{j+m-1} \\ M(n+\ell-1) \times O(m) & \xrightarrow{\circ^j} & M(n+m+\ell-2) \end{array}$$

Ramified compatibility between the left and right operations 2

$$\begin{array}{ccc} O(n) \times M(m) \times O(\ell) & \xrightarrow{\circ_i} & M(n+m-1) \times O(\ell) \\ \circ^k \downarrow & & \downarrow \circ^{k+i-1} \\ O(n) \times M(m+\ell-1) & \xrightarrow{\circ_i} & M(n+m+\ell-2) \end{array}$$

Linear compatibility between the left and right operations

$$\begin{array}{ccc} O(n) \times O(m) \times M(\ell) & \xrightarrow{\circ_k} & O(n) \times M(m+\ell-1) \\ \circ_i \downarrow & & \downarrow \circ_i \\ O(n+m-1) \times M(\ell) & \xrightarrow{\circ^{k+i-1}} & M(n+m+\ell-2) \end{array}$$

Linear associativity for the left infinitesimal operations

$$\begin{array}{ccc} O(n) \times M(m) & \xrightarrow{\circ_i} & M(n+m-1) \\ \sigma^* \times \tau^* \downarrow & & \downarrow (\sigma \circ_{\sigma(i)} \tau)^* \\ O(n) \times M(m) & \xrightarrow{\circ_{\sigma(i)}} & M(n+m-1) \end{array}$$

Compatibility with the symmetric group action 1

$$\begin{array}{ccc} M(n) \times O(m) & \xrightarrow{\circ^j} & M(n+m-1) \\ \sigma_* \times \tau_* \downarrow & & \downarrow (\sigma \circ_{\sigma(i)} \tau)_* \\ M(n) \times O(m) & \xrightarrow{\circ^{\sigma(i)}} & M(n+m-1) \end{array}$$

Compatibility with the symmetric group action 2

$$\begin{array}{ccc}
 M(n) \times O(1) & \longleftarrow M(n) \longrightarrow & O(1) \times M(n) \\
 \searrow \circ^i & \parallel & \swarrow \circ_1 \\
 & M(n) &
 \end{array}$$

Compatibility with the unit of the operad

Note that the *ramified compatibility between the left and right operations 2* follows from the *ramified compatibility between the left and right operations 1* and the *compatibility with the symmetric group action* and was given for clarity.

A map between O -bimodules should preserve these operations. We denote by ΣIbimod_O the category of O -bimodules. Given an integer $r \geq 0$, we also consider the category of r -truncated bimodules $T_r \Sigma\text{Ibimod}_O$. An object of this category is an r -truncated Σ -sequence endowed with left and right operations (91) which are defined under the conditions $n \leq r$ and $n + m - 1 \leq r$. One has an obvious truncation functor

$$T_r(-) : \Sigma\text{Ibimod}_O \longrightarrow T_r \Sigma\text{Ibimod}_O.$$

In the rest of the paper, we use the notation

$$\begin{aligned}
 x \circ^i \theta &= \circ^i(x; \theta), \text{ for } x \in M(n) \text{ and } \theta \in O(m), \\
 \theta \circ_i x &= \circ_i(\theta; x), \text{ for } \theta \in O(n) \text{ and } x \in M(m),
 \end{aligned}$$

Example 4.1 If $\eta : O \rightarrow M$ is a map of O -bimodules, then η is also a map of O -bimodules. Indeed, any operad is an infinitesimal bimodule over itself. Since the right operations and the right infinitesimal operations are the same, the O -bimodule structure on M is given by the following left infinitesimal operations:

$$\begin{aligned}
 \circ_i : O(n) \times M(m) &\longrightarrow M(n + m - 1); \\
 (\theta, x) &\longmapsto \gamma_\ell(\theta; \eta(*_1), \dots, \eta(*_1), x, \eta(*_1), \dots, \eta(*_1)).
 \end{aligned}$$

4.1 Properties of the Category of Infinitesimal Bimodules

In this subsection we introduce some basic properties related to the category of O -bimodules where O is a fixed operad. First, we show that the category of O -bimodules is equivalent to the category of algebras over an explicit colored operad denoted by O_+ . Thereafter, we build the free bimodule functor using the language of trees. Finally, we give a combinatorial description of the pushout for infinitesimal bimodules.

4.1.1 Infinitesimal Bimodules as Algebras over a Colored Operad

From an operad O , we build a colored operad O_+ such that the category of O -bimodules is equivalent to the category of O_+ -algebras. More precisely, the colored operad O_+ , with set of colors $S = \mathbb{N}$, is concentrated in arity 1 and it is given by the formula

$$O_+(n; m) := \coprod_{\alpha_+ : [m]_+ \rightarrow [n]_+} \prod_{i \in [n]_+} O(|\alpha_+^{-1}(i)|), \tag{92}$$

where $[n]_+$ is the set obtained from $[n]$ by adding a basepoint denoted by 0. The map α_+ is a map of sets preserving the basepoint. In order to define operadic compositions

$$\circ_1 : O_+(n; m) \times O_+(k; n) \longrightarrow O_+(k; m),$$

we fix two pointed maps $\alpha_+ : [m]_+ \rightarrow [n]_+$ and $\beta_+ : [n]_+ \rightarrow [k]_+$ and we build a map of the form

$$\prod_{i \in [n]_+} \mathcal{O}(|\alpha_+^{-1}(i)|) \times \prod_{j \in [k]_+} \mathcal{O}(|\beta_+^{-1}(j)|) \longrightarrow \prod_{j \in [k]_+} \mathcal{O}(|(\beta_+ \circ \alpha_+)^{-1}(j)|).$$

For this purpose, we rewrite the left hand side term as follows:

$$\underbrace{\mathcal{O}(|\alpha_+^{-1}(0)|) \times \mathcal{O}(|\beta_+^{-1}(0)|) \times \prod_{i \in \beta_+^{-1}(0) \setminus \{0\}} \mathcal{O}(|\alpha_+^{-1}(i)|)}_{\text{Part 1}} \times \underbrace{\prod_{j \in [k]} \mathcal{O}(|\beta_+^{-1}(j)|) \times \prod_{i \in \beta_+^{-1}(j)} \mathcal{O}(|\alpha_+^{-1}(i)|)}_{\text{Part 2}}.$$

In Part 2, as in formula (24), we use the operadic structure of \mathcal{O} in order to get an element in $\mathcal{O}(|(\beta_+ \circ \alpha_+)^{-1}(j)|)$ with $j \neq 0$. For Part 1, let $\beta_+^{-1}(0) \setminus \{0\} = \{i_1, \dots, i_p\}$, $a_0 \in \mathcal{O}(|\alpha_+^{-1}(0)|)$, $b_0 \in \mathcal{O}(|\beta_+^{-1}(0)|)$, $a_r \in \mathcal{O}(|\alpha_+^{-1}(i_r)|)$, $1 \leq r \leq p$. Then the element $(a_0, b_0, a_1, \dots, a_p)$ in the Part 1 product is sent to $a_0 \circ_1 b_0(id, a_1, \dots, a_p)$.

Proposition 4.2 (Proposition 4.9 in [3]) *The category of \mathcal{O} -bimodules is equivalent to the category of \mathcal{O}_+ -algebras.*

Note that since the operad \mathcal{O}_+ has only unary operations, it can be viewed as an enriched in *Top* category. One has that an \mathcal{O}_+ -algebra is the same thing as an \mathcal{O}_+ -shaped diagram in *Top*:

$$Alg_{\mathcal{O}_+} = Top^{\mathcal{O}_+}.$$

4.1.2 The Free Bimodule Functor

In what follows, we introduce the left adjoints of the forgetful functors

$$\mathcal{U}^\Sigma : \Sigma\text{Ibimod}_{\mathcal{O}} \longrightarrow \Sigma\text{Seq} \quad \text{and} \quad \mathcal{U}^{T_r\Sigma} : T_r\Sigma\text{Ibimod}_{\mathcal{O}} \longrightarrow T_r\Sigma\text{Seq},$$

denoted by $\mathcal{IF}_{\mathcal{O}}^\Sigma$ and $\mathcal{IF}_{\mathcal{O}}^{T_r\Sigma}$, respectively. As usual in the operadic theory, the free functor can be described as a coproduct indexed by a particular set of trees. In that case, we use the set of *pearled trees* which are pairs $T = (T; p)$ where T is a planar rooted tree, with leaves labelled by a permutation, and p is a particular vertex, called the *pearl*. A pearled tree is said to be *reduced* if each vertex is connected to the pearl by an inner edge. We denote by $p\mathbb{P}_n$ and $rp\mathbb{P}_n$ the sets of pearled trees and reduced pearled trees, respectively, having exactly n leaves (see Fig. 18).




Fig. 18 Examples of a pearled tree $T_1 \in p\mathbb{P}_5$ and a reduced pearled tree $T_2 \in rp\mathbb{P}_5$

Construction 4.3 Let $M = \{M(n)\}$ be a Σ -sequence. The space $\mathcal{IF}_O^\Sigma(M)(n)$ is obtained from the set of reduced pearled trees by indexing the pearl by a point in M whereas the other vertices are indexed by points in the operad O . More precisely, one has

$$\mathcal{IF}_O^\Sigma(M)(n) = \left(\prod_{T \in rP_n} M(|p|) \times \prod_{v \in V(T) \setminus \{p\}} O(|v|) \right) / \sim. \tag{93}$$

A point in $\mathcal{IF}_O^\Sigma(M)$ is denoted by $[T ; x_p ; \{\theta_v\}]$ where T is a reduced pearled tree, x_p is a point in M and $\{\theta_v\}_{v \in V(T) \setminus \{p\}}$ is a family of points in O . The equivalence relation is generated by the following relations:

- (i) *The unit relation:* if a vertex is indexed by the unit of the operad O , then we can remove it (see Fig. 19).
- (ii) *The compatibility with the symmetric group action:* if a vertex is labelled by $x \cdot \sigma$, with x a point in $O(n)$ or $M(n)$ and $\sigma \in \Sigma_n$, then we can remove σ by permuting the incoming edges (see Fig. 20).

The right infinitesimal operation \circ^i (respectively the left infinitesimal operation \circ_i) of a point $[T ; x_p ; \{\theta_v\}]$ with an element $\theta \in O(m)$ consists in grafting the m -corolla indexed by θ (respectively the reduced pearled tree T) into the i -th leaf of the reduced tree with section T (respectively the m -corolla indexed by θ). If the obtained element contains an inner edge joining two consecutive vertices other than a pearl, then we contract it using the operadic structure of O (see Fig. 21).

Similarly, the free r -truncated bimodule functor $\mathcal{IF}_O^{T_r, \Sigma}$ is obtained from the formula (28) by taking the restriction of the coproduct to the reduced pearled trees having at most r leaves




Fig. 19 Illustration of the unit relation




Fig. 20 Illustration of the compatibility with the symmetric group




Fig. 21 Illustration of the right infinitesimal operation $\circ^3 : \mathcal{IF}_O^\Sigma(M)(5) \times O(3) \rightarrow \mathcal{IF}_O^\Sigma(M)(7)$

and such that the pearl has at most r incoming edges. The equivalence relation, the left and right infinitesimal operations are defined in the same way. Finally, one has two functors:

$$\mathcal{IF}_O^\Sigma : \Sigma Seq \longrightarrow \Sigma \text{Ibimod}_O \quad \text{and} \quad \mathcal{IF}_O^{T_r \Sigma} : T_r \Sigma Seq \longrightarrow T_r \Sigma \text{Ibimod}_O.$$

Theorem 4.4 *One has the following adjunctions:*

$$\mathcal{IF}_O^\Sigma : \Sigma Seq \rightleftarrows \Sigma \text{Ibimod}_O : \mathcal{U}^\Sigma \quad \text{and} \quad \mathcal{IF}_O^{T_r \Sigma} : T_r \Sigma Seq \rightleftarrows T_r \Sigma \text{Ibimod}_O : \mathcal{U}^{T_r \Sigma}. \tag{94}$$

Proof The proof is similar to the proof of Theorem 2.8. For any O -Ibimodule M' and for any morphism of Σ -sequences $f : M \rightarrow M'$, we can build a unique map of O -Ibimodules $\tilde{f} : \mathcal{IF}_O^\Sigma(M) \rightarrow M'$ by induction on the number of vertices, such that $f = \tilde{f} \circ i$. We refer the reader to [12, Proposition 2.3] for more details. \square

4.1.3 Combinatorial Description of the Pushout

Let O be a topological operad. Contrary to the bimodule case, the left infinitesimal operations are unary. As a consequence, the pushout in the category of O -Ibimodules coincides with the pushout in the underlying category of sequences. More precisely, for any pushout diagrams

$$\begin{array}{ccc} A & \xrightarrow{f_1} & C \\ f_2 \downarrow & & \\ B & & \end{array} \quad \text{(respectively)} \quad \begin{array}{ccc} A_r & \xrightarrow{f_1} & C_r \\ f_2 \downarrow & & \\ B_r & & \end{array} \tag{95}$$

in the category of O -Ibimodules (respectively, r -truncated O -Ibimodules) we introduce the Σ -sequences

$$D(n) = \left(B(n) \bigsqcup_{A(n)} C(n) \right) \quad \text{and} \quad D_r(n) = \left(B_r(n) \bigsqcup_{A_r(n)} C_r(n) \right). \tag{96}$$

The above sequences inherit a (possibly truncated) infinitesimal bimodule structure over O and one has the following statement:

Proposition 4.5 *One has the following identities in the category of (possibly truncated) O -Ibimodules:*

$$D = \text{colim}_{\Sigma \text{Ibimod}_O} (B \leftarrow A \rightarrow C) \quad \text{and} \\ D_r = \text{colim}_{T_r \Sigma \text{Ibimod}_O} (B_r \leftarrow A_r \rightarrow C_r).$$

Proof Let $h_1 : C \rightarrow D'$ and $h_2 : B \rightarrow D'$ be two maps of O -Ibimodules such that $h_1 \circ f_1 = h_2 \circ f_2$. Then there exists a unique map of O -Ibimodule $\delta : D \rightarrow D'$ given by

$$\delta(x) = \begin{cases} h_1(x) & \text{if } x \in C, \\ h_2(x) & \text{if } x \in B. \end{cases}$$

This map is well defined and proves that D satisfies the universal property of the pushout. \square

4.2 Model Category Structure

By applying the transfer principle 1.1 to the adjunctions introduced in Sect. 4.1.2

$$\mathcal{IF}_O^\Sigma : \Sigma Seq \rightleftarrows \Sigma \text{Ibimod}_O : \mathcal{U}^\Sigma \quad \text{and} \quad \mathcal{IF}_O^{T_r \Sigma} : T_r \Sigma Seq \rightleftarrows T_r \Sigma \text{Ibimod}_O : \mathcal{U}^{T_r \Sigma}, \tag{97}$$

we get the following statement:

Theorem 4.6 *Let O be any topological operad. The category of (truncated) infinitesimal bimodules ΣIbimod_O (respectively, $T_r \Sigma \text{Ibimod}_O$, $r \geq 0$) inherits a cofibrantly generated model category structure, called the projective model category structure, in which all objects are fibrant. The model structure in question makes the adjunctions (97) into Quillen adjunctions. More precisely, a (possibly truncated) Ibimodule map f is a weak equivalence (respectively, a fibration) if and only if the induced map $\mathcal{U}^\Sigma(f)$ is a weak equivalence (respectively, a fibration) in the category of (possibly truncated) Σ -sequences.*

Proof Similar to the proof of Theorem 2.11. □

4.2.1 Left Properness of the Projective Model Category

Theorem 4.7 *For any topological operad O , the projective model category ΣIbimod_O is right proper. It is left proper provided O is componentwise cofibrant.*

Proof Since all the objects in ΣIbimod_O are fibrant, this category is right proper (we refer the reader to the proof of Theorem 2.2.1 for more details). Furthermore, as explained in Sect. 4.1.3, the pushout in the category of O -Ibimodules coincides with the pushout in the category of (non- Σ) sequences, which is obviously left proper as the category Top is such. By part (a) of the next theorem, cofibrations in ΣIbimod_O are componentwise cofibrations provided O is componentwise cofibrant. We conclude that in this case, ΣIbimod_O is also left proper. □

Theorem 4.8 (a) *If O is a componentwise cofibrant operad, then cofibrations in the category ΣIbimod_O are cofibrations componentwise. In particular, the class of componentwise cofibrant objects is closed under cofibrations and cofibrant objects have cofibrant components.*

(b) *If O is a Σ -cofibrant operad, then cofibrations in the category ΣIbimod_O are Σ -cofibrant. In particular, the class of Σ -cofibrant objects is closed under cofibrations and cofibrant objects are Σ -cofibrant.*

Proof It is enough to check that the generating cofibrations in ΣIbimod_O are cofibrations in Top componentwise, in the case (a), and are Σ -cofibrations, in the case (b). The first is a consequence of the pushout-product axiom, while the second is proved in exactly the same way as the similar statement for operads [6, Corollary 5.2]. □

4.2.2 Extension/Restriction Adjunction for the Projective Model Category of Infinitesimal Bimodules

Let $\phi : O \rightarrow O'$ be a map of operads. Similarly to the category of algebras (see Theorem 1.7) and bimodules (see Sect. 2.2.2), we show that the projective model categories of O -Ibimodules and O' -Ibimodules are Quillen equivalent under some conditions on the

operads. For this purpose, we recall the construction of the *restriction* functor ϕ^* and of the *extension* functor $\phi_!$ in the context of infinitesimal bimodules:

$$\phi_! : \Sigma \text{Ibimod}_O \rightleftarrows \Sigma \text{Ibimod}_{O'} : \phi^*.$$

• *The restriction functor:* The restriction functor ϕ^* sends an O' -Ibimodule M to the O -Ibimodule $\phi^*(M) = \{\phi^*(M)(n) = M(n), n \geq 0\}$ in which the O -bimodule structure is defined using the O' -bimodule structure of M as follows:


$$\begin{aligned} \circ^i : \phi^*(M)(n) \times O(m) &\longrightarrow \phi^*(M)(n + m - 1); \\ x ; \theta &\longmapsto x \circ^i \phi(\theta), \end{aligned}$$

$$\begin{aligned} \circ_i : O(n) \times \phi^*(M)(m) &\longrightarrow \phi^*(M)(n + m - 1); \\ \theta ; x &\longmapsto \phi(\theta) \circ_i x. \end{aligned}$$

• *The extension functor:* The extension functor $\phi_!$ is obtained as a quotient of the free O' -Ibimodule functor introduced in Sect. 4.1.2. More precisely, if M is an O -Ibimodule, then the extension functor is given by the formula

$$\phi_!(M)(n) = \mathcal{IF}^{\Sigma}_{O'}(\mathcal{U}^{\Sigma}(M))(n) / \sim$$

where the equivalence relation is generated by the axiom which consists in contracting inner edges having a vertex v indexed by a point of the form $\phi_1(\theta)$ using the O infinitesimal bimodule structure of M as illustrated in the following picture:



The O' -infinitesimal bimodule structure on the free object is compatible with the equivalence relation and provides an O' -Ibimodule structure on $\phi_!(M)$. Let us remark that, similarly to the bimodule case, the O -Ibimodule map $M \rightarrow \phi^*(\phi_!(M))$, sending a point $x \in M(n)$ to the n -corolla indexed by x , is not necessarily injective.

Theorem 4.9 *Let $\phi : O \rightarrow O'$ be a weak equivalence between operads with cofibrant components. The extension and restriction functors, as well as their truncated versions, give rise to Quillen equivalences:*

$$\phi_! : \Sigma \text{Ibimod}_O \rightleftarrows \Sigma \text{Ibimod}_{O'} : \phi^*, \tag{98}$$

$$\phi_! : T_r \Sigma \text{Ibimod}_O \rightleftarrows T_r \Sigma \text{Ibimod}_{O'} : \phi^*. \tag{99}$$

Proof As explained in Sect. 4.1.1, the projective model category of infinitesimal bimodules over an operad is equivalent to the projective model category of algebras over a specific colored operad. We denote by O_+ and O'_+ the corresponding colored operads associated to O and O' , respectively. One has that the adjunction (98) is induced by the extension/restriction adjunction between the categories of algebras:

$$\phi_! : \Sigma \text{Ibimod}_O = \text{Alg}_{O_+} \rightleftarrows \text{Alg}_{O'_+} = \Sigma \text{Ibimod}_{O'} : \phi^*.$$

From the fact that O and O' have cofibrant components and from the explicit formula (92) for the components of O_+ and O'_+ , we obtain that the colored operads O_+ and O'_+ have cofibrant components. Since they are concentrated in arity one, they are Σ -cofibrant. Applying [6,

Theorem 4.4], we conclude that the extension/restriction adjunction between the categories of algebras is a Quillen equivalence.

The truncated case is done similarly by restricting the operads O_+ and O'_+ to their subsets of colors. □

5 The Reedy Model Category of O Infinitesimal Bimodules

Let O be a reduced operad. From now on, we denote by Λbimod_O and $T_r \Lambda\text{bimod}_O$ the categories of O -bimodules and truncated O -bimodules, respectively, equipped with the Reedy model category structures. Note that as categories, $\Lambda\text{bimod}_O = \Sigma\text{bimod}_O$. Only the model structure is different. Similarly to the case of reduced operads and bimodules, this structure is transferred from the categories ΛSeq and $T_r \Lambda\text{Seq}$ along the adjunctions

$$\begin{aligned} \mathcal{IF}_O^\Lambda : \Lambda\text{Seq} &\rightleftarrows \Lambda\text{bimod}_O : \mathcal{U}^\Lambda, \\ \mathcal{IF}_O^{T_r \Lambda} : T_r \Lambda\text{Seq} &\rightleftarrows T_r \Lambda\text{bimod}_O : \mathcal{U}^\Lambda, \end{aligned} \tag{100}$$

where both free functors are obtained from the functors \mathcal{IF}_O^Σ and $\mathcal{IF}_O^{T_r \Sigma}$ by taking the restriction of the coproduct (93) to the reduced pearled trees without univalent vertices other than the pearl. In other words, one has

$$\mathcal{IF}_O^\Lambda(M) := \mathcal{IF}_{O_{>0}}^\Sigma(M), \quad \text{and} \quad \mathcal{IF}_O^{T_r \Lambda}(M) := \mathcal{IF}_{O_{>0}}^{T_r \Sigma}(M).$$

By construction, the above Σ -sequences are equipped with a (truncated) infinitesimal bimodule structure over $O_{>0}$. We can extend this structure in order to get a (truncated) O infinitesimal bimodule structure using the operadic structure of O and the Λ structure of M (see Fig. 22).

Theorem 5.1 *Let O be a reduced well-pointed operad. The categories Λbimod_O and $T_r \Lambda\text{bimod}_O$, with $r \geq 0$, admit cofibrantly generated model category structures, called Reedy model category structures, transferred from ΛSeq and $T_r \Lambda\text{Seq}$, respectively, along the adjunctions (100). In particular, these model category structures make the pairs of functors (100) into Quillen adjunctions.*

Proof The proof is similar to the proof of Theorem 3.1. The path object is given by the same formula (52) and the functorial fibrant coresolution (for which we need the assumption Q is well-pointed) is defined in Sect. 5.1.1. □

5.1 Properties of the Reedy Σ Model Category of Infinitesimal Bimodules

This subsection is divided into two parts. The first one is devoted to the construction of an explicit fibrant coresolution functor for infinitesimal bimodules. In the second part, we




Fig. 22 Illustration of the right action by $*_0$

characterize (acyclic) cofibrations in the Reedy model category of Ibimodules, we prove properness and we study extension/restriction adjunctions.

5.1.1 Fibrant Replacement Functor for Infinitesimal Bimodules

Let O be a reduced operad. In this subsection we produce a construction of a Reedy fibrant coresolution for infinitesimal bimodules if the operad O is well-pointed. More precisely, we show that the coresolution introduced in Sect. 3.1.1 gives rise to a coresolution for infinitesimal bimodules too. In other words, in both cases, bimodules and infinitesimal bimodules, we only need the right module part of the structures in order to get Reedy fibrant replacements. In the following, we overuse the notation introduced in Sect. 3.1.1. Given an infinitesimal bimodule M over O , we consider the space $M^f(n)$ from (53) taking $Q = O$. The family $M^f = \{M^f(n), n \geq 0\}$ admits a Σ -structure and a right module structure from operations (59).

In order to introduce the left infinitesimal operations, we need the following notation. For a tree $T \in \mathbb{P}[n]$, we denote by T_1, \dots, T_n the sub-trees grafting to the root of T according to the planar order. In particular, one has the identity $T = c_n(T_1, \dots, T_n)$ where c_n is the n -corolla. The number of leaves of each tree T_j , with $1 \leq j \leq n$, is denoted by $n_j[T]$. For any tree $T \in \mathbb{P}[n]$ and $1 \leq j \leq n$, we also consider the operations

$$\beta_j : D(T) \longrightarrow O(n_j[T]);$$

$$\{\theta_v\}_{v \in V(T) \setminus \{r\}} \longmapsto \eta(\{\theta_v\}_{v \in V(T_j)}),$$

given by composing the points of O indexing the vertices of the sub-tree T_j .

Finally, by using the operation Γ_k^m introduced in Sect. 3.1.1, one can define the left infinitesimal action

$$\circ_i : O(n) \times M^f(m) \longrightarrow M^f(n + m - 1);$$

$$\theta, \{f_T\}_{T \in \mathbb{P}[n]} \longmapsto \{(\theta \circ_i f)T\}_{T \in \mathbb{P}[n+m-1]},$$

where $(\theta \circ_i f)_T$ is the composite map:

$$\begin{array}{ccc}
 H(T) \times D(T) & \xrightarrow{(\theta \circ_i f)_T} & M([T]) \\
 \downarrow & & \uparrow \\
 \left(\prod_{\substack{1 \leq j \leq n+m-1 \\ j \notin \{i, \dots, i+m-1\}}} O(n_j[T]) \right) \times (H(\Gamma_{i-1}^m(T)) \times D(\Gamma_{i-1}^m(T))) & \longrightarrow & O \left(\sum_{\substack{1 \leq j \leq n+m-1 \\ j \notin \{i, \dots, i+m-1\}}} n_j[T] \right) \times M(\Gamma_{i-1}^m(T))
 \end{array}$$

If we denote by I the set $\{1, \dots, i - 1\} \cup \{i + m, \dots, n + m - 1\}$, then the left vertical map is given by the product $\prod_{j \in I} \beta_j$ to get the first factor and the operation induced by Γ_{i-1}^m , removing the incoming edges of the root of T corresponding to the set I , in order to get the second factor. The lower horizontal map is given by the map $f_{\Gamma_{i-1}^m(T)}$ on the second factor

and the following map

$$\begin{aligned} \theta(-, \dots, id, \dots, -) : & \left(\prod_{j \in \{1, \dots, i-1\}} O(n_j[T]) \right) \times \left(\prod_{j \in \{i+m, \dots, n+m-1\}} O(n_j[T]) \right) \\ \longrightarrow & O \left(\sum_{\substack{1 \leq j \leq n+m-1 \\ j \notin \{i, \dots, i+m-1\}}} n_j[T] \right); \\ (\theta_1, \dots, \theta_{i-1}); & (\theta'_1, \dots, \theta'_{n-i-1}) \longmapsto \theta(\theta_1, \dots, \theta_{i-1}, id, \theta'_1, \dots, \theta'_{n-i-1}), \end{aligned}$$

using the operadic structure of O , on the first factor. Finally, the right vertical map is obtained using the left infinitesimal operation \circ_ℓ , with $\ell = \sum_{1 \leq j \leq i-1} n_j[T] + 1$. The fact that the compatibility relations between the left and the right infinitesimal operations (defined at the beginning of Sect. 4) are satisfied is a consequence of the following observations:

- ▶ *Ramified compatibility between the left and right operations:* for $T \in \mathbb{P}[n + m + \ell - 2]$, $i \in \{1, \dots, n - 1\}$ and $j \in \{i + 1, \dots, n\}$, one has

$$\Gamma_{i-1}^m(T) = \Gamma_{i-1}^m(\delta_{j+m-1; \ell}(T)).$$

- ▶ *Linear compatibility between the left and right operations:* for $T \in \mathbb{P}[n + m + \ell - 2]$, $i \in \{1, \dots, n\}$ and $k \in \{1, \dots, m\}$, one has

$$\Gamma_{i-1}^m(\delta_{k+i-1; \ell}(T)) = \delta_{k; \ell}(\Gamma_{i-1}^{m+\ell-1}(T)).$$

Proposition 5.2 *The map $\eta : M \rightarrow M^f$ is a weak equivalence of O -Ibimodules. Furthermore, if the operad O is well-pointed, then the O -Ibimodule M^f is Reedy fibrant.*

Proof The reader can easily check that η is a map of O infinitesimal bimodules. The other statements are consequences Propositions 3.3 and 3.4. □

Remark 5.3 The same strategy can be used in order to get a fibrant replacement functor for r -truncated reduced O -Ibimodules. The fibrant replacement should be defined as a subspace of the product with an additional restriction $|T| \leq r$. The constraints are the same.

5.1.2 Characterization of Cofibrations/Left properness/Extension-restriction adjunction

In this section, we show that the properties related to the Reedy model category of reduced bimodules introduced in Sect. 3.1 admit counterparts in the context of infinitesimal bimodules. It means that we are able to give a characterization of Reedy cofibrations and we prove that ΛIbimod_O is left proper relative to componentwise cofibrant objects. We also prove that the extension/restriction adjunction gives rise to a Quillen equivalence between Reedy model categories of Ibimodules under some conditions on the operads.

Theorem 5.4 *Let O be a reduced well-pointed operad. A morphism $\phi : M \rightarrow N$ in the category of (possibly truncated) O -Ibimodules is a Reedy cofibration if and only if ϕ is a cofibration in the projective model category of (possibly truncated) $O_{>0}$ -Ibimodules.*

Idea of the proof The strategy used for the proof of Theorem 3.6 works with no change in the context of infinitesimal bimodules. Again, we introduce an adjunction $ar_s : \Lambda Ibimod_O \rightleftarrows \Lambda Ibimod_O : cosk_s$ where ar_s and $cosk_s$ are the arity filtration and the coskeleton functors, respectively. More precisely, if L_s and R_s denote the left adjoint and the right adjoint, respectively, to the truncation functor $T_s : \Lambda Ibimod_O \rightarrow T_s \Lambda Ibimod_O$, then one has the following identities:

$$ar_s = L_s \circ T_s \quad \text{and} \quad cosk_s = R_s \circ T_s.$$

In particular, the coskeleton functor is given by the formula (65) and inherits an infinitesimal bimodule structure over O . The Λ -structure and the right infinitesimal operations are given by (66) and (67), respectively. In order to define the left infinitesimal operations, we recall the following notation. Let $n, m > 0, \ell \in \{1, \dots, n\}$ and $h \in \Lambda_+([i]; [n + m - 1])$. If we denote by $\ell_1 \in \Lambda_+([m]; [n + m - 1])$ and $\ell_2 \in \Lambda_+([n]; [n + m - 1])$ the morphisms

$$\begin{aligned} \ell_1 : [m] &\longrightarrow [n + m - 1]; & \text{and } \ell_2 : [n] &\longrightarrow [n + m - 1]; \\ \alpha &\longmapsto \alpha + \ell, & \alpha &\longmapsto \begin{cases} \alpha & \text{if } \alpha \leq \ell, \\ \alpha + m & \text{if } \alpha > \ell, \end{cases} \end{aligned}$$

then there exist unique morphisms h_1 and h_2 making the following diagrams commute:

$$\begin{array}{ccc} [i] & \xrightarrow{h} & [n + m - 1] \\ \uparrow & & \uparrow \ell_1 \\ [[\text{Im}(\ell_1) \cap \text{Im}(h)]] & \xrightarrow{h_1} & [m] \end{array} \qquad \begin{array}{ccc} [i] & \xrightarrow{h} & [n + m - 1] \\ \uparrow & & \uparrow \ell_2 \\ [[\text{Im}(\ell_2 \setminus \{\ell\}) \cap \text{Im}(h)]] & \xrightarrow{h_2} & [n] \end{array}$$

Finally, if we denote by $\bar{\ell} = \ell - |\{\alpha \in [i] \mid h(\alpha) < \ell\}|$, then the left infinitesimal operations are given by

$$\begin{aligned} \circ_i : O(n) \times cosk_s(M)(m) &\longrightarrow cosk_s(M)(n + m - 1); \\ \theta ; \{x_u\}_{u \in \Lambda_+([i]; [m])}^{0 \leq i \leq s} &\longmapsto \{h_2^*(\theta) \circ_{\bar{\ell}} x_{h_1}\}_{h \in \Lambda_+([i]; [n + m - 1])}^{0 \leq i \leq s}. \end{aligned}$$

The rest of the proof is the same as the proof of Theorem 3.6. It consists in using the adjunction $(ar_s, cosk_s)$ in order to define by induction a solution to the lifting problem. \square

Theorem 5.5 *For any reduced well-pointed operad O , the Reedy model category $\Lambda Ibimod_O$ is right proper. It is left proper provided O is componentwise cofibrant. In the latter case, cofibrations are componentwise cofibrations, and, as a consequence, the class of componentwise cofibrant objects is closed under cofibrations and cofibrant Ibimodules are componentwise cofibrant. If O is Σ -cofibrant, then the cofibrations are Σ -cofibrations, the class of Σ -cofibrant objects is closed under cofibrations, and cofibrant objects are Σ -cofibrant.*

Proof Right properness immediately follows from Theorem 4.7. Indeed, $\Lambda Ibimod_O = \Sigma Ibimod_O$ as categories. Therefore, they have the same pullbacks. Moreover, a Reedy fibration is always a projective fibration. For left properness and the properties of cofibrations, we use the characterization of cofibrations of Theorem 5.4 together with the analogous properties of $\Sigma Ibimod_{O_{>0}}$ established in Theorems 4.7 and 4.8. \square

Let $\phi : O \rightarrow O'$ be a weak equivalence of reduced operads. Similarly to Sect. 4.2.2, we show that the Reedy model categories of O -Ibimodules and O' -Ibimodules are Quillen

equivalent. By abuse of notation, we denote by ϕ^* and $\phi_!$ the restriction functor and the extension functor, respectively, between the Reedy model categories:

$$\phi_! : \Delta\text{Ibimod}_O \rightleftarrows \Delta\text{Ibimod}_{O'} : \phi^*.$$

In the same way as in Sect. 4.2.2, for any $M \in \Delta\text{Ibimod}_O$ and $M' \in \Delta\text{Ibimod}_{O'}$, one has

$$\begin{aligned} \phi_!(M) &= \{\phi_!(M)(n) = \mathcal{F}_O^\wedge(\mathcal{U}^\wedge(M))(n) / \sim, \ n \geq 0\}, \\ \phi^*(M') &= \{\phi^*(M')(n) = M'(n), \ n \geq 0\}. \end{aligned}$$

Theorem 5.6 *Let $\phi : O \rightarrow O'$ be a weak equivalence between reduced componentwise cofibrant operads. One has Quillen equivalences*

$$\phi_! : \Delta\text{Ibimod}_O \rightleftarrows \Delta\text{Ibimod}_{O'} : \phi^*, \tag{101}$$

$$\phi_! : T_r \Delta\text{Ibimod}_O \rightleftarrows T_r \Delta\text{Ibimod}_{O'} : \phi^*. \tag{102}$$

Proof Since the restriction functor creates weak equivalences, one has to check that, for any Reedy cofibrant object M in ΔIbimod_O , the adjunction unit

$$M \longrightarrow \phi^*(\phi_!(M))$$

is a weak equivalence. Due to the characterization of Reedy cofibrations, M is also cofibrant in the projective model category of $O_{>0}$ -Ibimodules. Since $\phi_{>0} : O_{>0} \rightarrow O'_{>0}$ is still a weak equivalence between componentwise cofibrant operads, by Theorem 2.19, the pair of functors $((\phi_{>0})_!; (\phi_{>0})^*)$ gives rise to a Quillen equivalence and the map $M(n) \rightarrow (\phi_{>0})^*((\phi_{>0})_!(M))(n)$ is a weak equivalence. The statement is a consequence of the identity

$$\phi^*(\phi_!(M)) = (\phi_{>0})^*((\phi_{>0})_!(M)).$$

□

5.2 The Connection Between the Model Category Structures on Infinitesimal Bimodules

Similarly to the operadic case in [21], we build a Quillen adjunction between the projective and the Reedy model categories of infinitesimal bimodules over a reduced operad O . Furthermore, if M and N are two infinitesimal bimodules, then we show that there is a weak equivalence between the derived mapping spaces:

$$\Sigma\text{Ibimod}_O^h(M; N) \simeq \Delta\text{Ibimod}_O^h(M; N).$$

For completeness of exposition, at the end of the subsection, we explain how to adapt the Boardman–Vogt resolution (well known for operads, see [7]) to the context of infinitesimal bimodules. We refer the reader to [15] where this construction was defined by the first and third authors. Using this construction we define a functorial cofibrant replacement in the categories ΣIbimod_O and ΔIbimod_O provided O is componentwise cofibrant.

5.2.1 Quillen Adjunction Between the Model Category Structures

Let O be a reduced operad. The projective and the Reedy model categories of infinitesimal bimodules over O have the same set of weak equivalence and induce the same homotopy

category. Consequently, one has the following statement about the adjunctions

$$\begin{aligned}
 id : \Sigma \text{Ibimod}_O &\rightleftarrows \Delta \text{Ibimod}_O : id, \\
 id : T_r \Sigma \text{Ibimod}_O &\rightleftarrows T_r \Delta \text{Ibimod}_O : id.
 \end{aligned}
 \tag{103}$$

Theorem 5.7 *For any well-pointed reduced operad O , the pairs of functors (103) form Quillen equivalences. Furthermore, for any pair $M, N \in \Delta \text{Ibimod}_O$, one has*

$$\Sigma \text{Ibimod}_O^h(M, N) \simeq \Delta \text{Ibimod}_O^h(M, N).
 \tag{104}$$

Moreover, if $M, N \in T_r \Delta \text{Ibimod}_O$, with $r \geq 0$, then one has

$$T_r \Sigma \text{Ibimod}_O^h(M, N) \simeq T_r \Delta \text{Ibimod}_O^h(M, N).
 \tag{105}$$

Proof The proof is similar to that of Theorem 3.9. □


5.2.2 Cofibrant Resolution in the Projective/Reedy Model Category

Let O be an operad not necessarily reduced. From an O -Ibimodule M , we build an O -Ibimodule $\mathcal{I}b_O(M)$. The points of $\mathcal{I}b_O(M)(n)$, $n \geq 0$, are equivalence classes $[T; \{t_v\}; x_p; \{\theta_v\}]$, where $T \in p\mathbb{P}_n$ (see Sect. 4.1.2) is a pearled tree, x_p is a point in M labelling the pearl and $\{\theta_v\}_{v \in V(T) \setminus \{p\}}$ is a family of points in O labelling the vertices other than the pearl. Furthermore, $\{t_v\}_{v \in V(T) \setminus \{p\}}$ is a family of real numbers in the interval $[0, 1]$ indexing the vertices which are not pearls. According to the orientation toward the pearl, if e is an inner edge, then $t_{s(e)} \geq t_{t(e)}$. In other words, the closer to the pearl is a vertex, the smaller is the corresponding number. The space $\mathcal{I}b_O(M)(n)$ is a quotient of the subspace of


$$\coprod_{T \in p\mathbb{P}_n} M(|p|) \times \prod_{v \in V(T) \setminus \{p\}} [O(|v|) \times [0, 1]]
 \tag{106}$$

determined by the restrictions on the families $\{t_v\}$. The equivalence relation is generated by the unit condition (i) and the compatibility with the symmetric group relation (ii) of Construction 4.3 as well as the following conditions:

- (iii) If two consecutive vertices, connected by an edge e , are indexed by the same real number $t \in [0, 1]$, then e is contracted using the operadic structure of O . The vertex produced by this edge contraction is indexed by the real number t .



- (iv) If a vertex connected to the pearl is indexed by 0, then we contract the inner edge connecting them using the infinitesimal bimodule structure of M . In that case the new vertex, produced by the contraction, becomes the pearl.



Let us describe the O -Bimodule structure. Let $\theta \in O(n)$ and $[T; \{t_v\}; x_p; \{\theta_v\}]$ be a point in $\mathcal{I}b_O(M)(m)$. The right infinitesimal operation $[T; \{t_v\}; x_p; \{\theta_v\}] \circ^i \theta$ consists in grafting the n -corolla labelled by θ to the i -th incoming edge of T and indexing the new vertex by 1. Similarly, the left infinitesimal operation $\theta \circ_i [T; \{t_v\}; x_p; \{\theta_v\}]$ consists in grafting the pearled tree T to the i -th incoming edge of n -corolla labelled by θ and indexing the new vertex by 1 (see Fig. 23).

One has an obvious inclusion of Σ -sequences $\iota: M \rightarrow \mathcal{I}b_O(M)$, where each element $x \in M(n)$ is sent to an n -corolla, labelled by x , whose only vertex is a pearl. Furthermore, the following map:

$$\mu : \mathcal{I}b_O(M) \rightarrow M ; [T; \{t_v\}; x_p; \{\theta_v\}] \mapsto [T; \{0\}; x_p; \{\theta_v\}], \tag{107}$$

is defined by sending the real numbers indexing the vertices other than the pearl to 0. The so obtained element is identified to the pearled corolla labelled by a point in M . It is easy to see that μ is an O -Bimodule map.

In order to get resolutions for truncated infinitesimal bimodules, one considers a filtration in $\mathcal{I}b_O(M)$ according to the number of *geometrical inputs* which is the number of leaves plus the number of univalent vertices other than the pearl. A point in $\mathcal{I}b_O(M)$ is said to be *prime* if the real numbers indexing the vertices are strictly smaller than 1. Otherwise, a point is said to be *composite* and can be associated to a *prime component* as shown in Fig. 24. More precisely, the prime component is obtained by removing the vertices indexed by 1.

A prime point is in the r -th filtration layer $\mathcal{I}b_O(M)_r$ if the number of its geometrical inputs is at most r . Similarly, a composite point is in the r -th filtration layer if its prime component is in $\mathcal{I}b_O(M)_r$. For instance, the composite point in Fig. 24 is in the filtration




Fig. 23 Illustration of the right infinitesimal operation




Fig. 24 A composite point and its prime components

layer $\mathcal{I}b_O(M)_6$. For each r , $\mathcal{I}b_O(M)_r$ is an O -Ibimodule and one has the following filtration of $\mathcal{I}b_O(M)$:

$$\mathcal{I}b_O(M)_0 \longrightarrow \mathcal{I}b_O(M)_1 \longrightarrow \dots \longrightarrow \mathcal{I}b_O(M)_{r-1} \longrightarrow \mathcal{I}b_O(M)_r \longrightarrow \dots \longrightarrow \mathcal{I}b_O(M). \tag{108}$$

Theorem 5.8 (Theorem 3.10 in [15]) *Assume that O is a Σ -cofibrant operad, and M is a Σ -cofibrant O -Ibimodule. Then the objects $\mathcal{I}b_O(M)$ and $T_r\mathcal{I}b_O(M)_r$ are cofibrant replacements of M and T_rM in the categories ΣIbimod_O and $T_r\Sigma\text{Ibimod}_O$, respectively. In particular, the maps μ and $T_r\mu|_{T_r\mathcal{I}b_O(M)_r}$ are weak equivalences.*

Now we slightly change the above construction in order to produce Reedy cofibrant replacements for O -Ibimodules when O is a reduced operad. Let M be an infinitesimal bimodule over O . As a Σ -sequence, we set

$$\mathcal{I}b_O^\Delta(M) := \mathcal{I}b_{O_{>0}}(M)$$

The superscript Δ is to emphasize that we get a cofibrant replacement in the Reedy model category structure. The right and left action by the positive arity components is defined as it is on $\mathcal{I}b_{O_{>0}}(M)$. The right action by $*_0 \in O(0)$ is defined in the obvious way as the right action by $*_0$ on a in the vertex (a, t) connected to the leaf labelled by i as illustrated in Fig. 17.

Note that since the arity zero component of $O_{>0}$ is empty, in the union (106) we can consider only trees whose all non-pearl vertices have arities ≥ 1 . We denote this set by $p\mathbb{P}_n^{\geq 1}$. In other words, the space $\mathcal{I}b_O^\Delta(M)$ can be obtained as the restriction of the coproduct (106) to this set.

Proposition 5.9 [15, Proposition 3.12] *Assume that O is a reduced Σ -cofibrant operad, and M is a Σ -cofibrant O -Ibimodule. Then the objects $\mathcal{I}b_O^\Delta(M)$ and $T_r\mathcal{I}b_O^\Delta(M)$ are cofibrant replacements of M and T_rM in the categories ΔIbimod_O and $T_r\Delta\text{Ibimod}_O$, respectively. In particular, the maps μ and $T_r\mu$ are weak equivalences.*

Proof The map $\mu : \mathcal{I}b_O^\Delta(M) \rightarrow M$, which changes the assignment of the real numbers indexing the vertices to 0, is a homotopy equivalence. More precisely, it is a deformation retract in the category of Σ -sequences in which the homotopy consists in bringing the real numbers to 0. Furthermore, as a consequence of Theorem 5.8, $\mathcal{I}b_O^\Delta(M) = \mathcal{I}b_{O_{>0}}(M)$ is cofibrant in the projective model category of $O_{>0}$ -Ibimodules. Due to Theorem 5.4, $\mathcal{I}b_O^\Delta(M)$ is also Reedy cofibrant and it gives rise to a cofibrant resolution of M in the Reedy model category ΔIbimod_O . The same arguments work for the truncated case. Note that $T_r\mathcal{I}b_O^\Delta(M)_r = T_r\mathcal{I}b_O^\Delta(M)$, since arity zero non-pearl vertices are not permitted. \square

By means of Theorem 5.8 and Proposition 5.9, we construct a functorial cofibrant replacement in ΣIbimod_O and ΔIbimod_O assuming that O is componentwise cofibrant. We adapt notation from Sect. 3.2.3. Given an O -Ibimodule M , we define $M'_\infty := |S_\bullet M| \times E_\infty$, $O'_\infty := |S_\bullet| \times E_\infty$. One has that both M'_∞ and O'_∞ are Σ -cofibrant and M'_∞ is an O'_∞ -bimodule. Let

$$\phi : O'_\infty \xrightarrow{\cong} O, \quad \text{and} \quad \phi_0 : M'_\infty \xrightarrow{\cong} M$$

be the natural projections. Note that ϕ_0 can be viewed as a map of O'_∞ -Ibimodules.

Proposition 5.10 (a) Assume that O is a componentwise cofibrant operad. Let M be any O -bimodule. Then the objects $\phi_! (\mathcal{I}b_{O'_\infty} (M'_\infty))$ and $\phi_! (T_r \mathcal{I}b_{O'_\infty} (M'_\infty)_r)$ are cofibrant replacements of M and $T_r M$ in the categories ΣIbimod_O and $T_r \Sigma \text{Ibimod}_O$, respectively.

(b) Assume in addition that the operad O is reduced. Then the objects $\phi_! (\mathcal{I}b_{O'_\infty}^\Delta (M'_\infty))$ and $\phi_! (T_r \mathcal{I}b_{O'_\infty}^\Delta (M'_\infty))$ are cofibrant replacements of M and $T_r M$ in the categories ΔIbimod_O and $T_r \Delta \text{Ibimod}_O$, respectively.

Proof The result is an immediate consequence of Theorems 4.9, 5.6, 5.8 and Proposition 5.9. □

A Equivariant Homotopy Theory

A.1 Projective Cofibrations for Monoidal Action

In what follows, for any topological monoid G , we consider the projective model structure on the category of G -spaces denoted $G\text{-Top}$. We refer the reader to Sect. 1.1 for more details. We start by recalling the statement of Berger-Moerdijk concerning the pushout product axiom.

Lemma A.1 [7, Lemma 2.5.3] *Let $1 \rightarrow G_1 \rightarrow G \rightarrow G_2 \rightarrow 1$ be a short exact sequence of discrete groups. Let $A \rightarrow B$ be a G_2 -cofibration and $X \rightarrow Y$ be a G -equivariant G_1 -cofibration. Then the pushout product map $(A \times Y) \cup_{A \times X} (B \times X) \rightarrow B \times Y$ is a G -cofibration. Moreover, the latter is acyclic if $A \rightarrow B$ or $X \rightarrow Y$ is.*

We use an analogue of this result (namely, Lemmas A.5 and A.8 below) that can be applied to $\Sigma_n \wr O(1)$ and to $(\Sigma_k \times \Sigma_{n-k}) \wr O(1)$, which are not groups, but monoids, and also are not discrete. In fact non-discreteness is not a big problem. It is not hard to see that Berger-Moerdijk’s proof of [7, Lemma 2.5.3] works for topological groups as well. On the contrary, for their proof it is critical that the action is by groups. In fact the statement of Lemma A.1 in general does not hold when G, G_1 , and G_2 are monoids. Indeed, consider $G_1 = 1$ and $G_2 = G = \mathbb{N}$ the monoid of non-negative natural numbers with the natural addition operation. Then for $A \rightarrow B$ being $S^{k-1} \times \mathbb{N} \rightarrow D^k \times \mathbb{N}$, and $X \rightarrow Y$ being $\mathbb{N} \rightarrow \mathbb{N}$, $i \mapsto i + 1$, the pushout product map $(A \times Y) \cup_{A \times X} (B \times X) \rightarrow B \times Y$ is not an \mathbb{N} -cofibration.

Lemma A.2 *Let Γ be a topological monoid, $A \rightarrow B$ be a Γ -cofibration, $X \rightarrow Y$ be a cofibration (in Top), then the pushout-product map $(A \times Y) \cup_{A \times X} (B \times X) \rightarrow B \times Y$ is a Γ -cofibration, where X and Y are regarded as spaces endowed with a trivial action of Γ . Moreover, the latter is acyclic if $A \rightarrow B$ or $X \rightarrow Y$ is.*

The proof of this lemma is identical to that of [7, Lemma 2.5.2] in which Γ is not a topological monoid but a discrete group. For the convenience of the reader the argument is given below.

Proof Let $Z \rightarrow W$ be a trivial Γ -fibration. One has to show that any square

$$\begin{array}{ccc}
 (A \times Y) \cup_{A \times X} (B \times X) & \longrightarrow & Z \\
 \downarrow & & \downarrow \\
 B \times Y & \longrightarrow & W
 \end{array} \tag{109}$$

has a Γ -equivariant lift. Since Top is cartesian closed, the existence of such lift is equivalent to the existence of a Γ -equivariant lift of the square

$$\begin{array}{ccc}
 A & \longrightarrow & Map(Y, Z) \\
 \downarrow & & \downarrow \\
 B & \longrightarrow & Map(Y, W) \times_{Map(X, W)} Map(X, Z).
 \end{array}
 \tag{110}$$

Here the Γ -action on the mapping spaces $Map(Y, Z)$, $Map(Y, W)$, etc. is defined through the action on the target: $(f \cdot \gamma)(y) := f(y) \cdot \gamma$. Since the left vertical arrow is a Γ -cofibration, it is enough to show that the right arrow is a trivial Γ -fibration. On the other hand, since the forgetful functor $\Gamma-Top \rightarrow Top$ creates fibrations and weak equivalences, it suffices to verify that the right arrow is a trivial fibration in Top , in other words that any square

$$\begin{array}{ccc}
 S^{k-1} & \longrightarrow & Map(Y, Z) \\
 \downarrow & & \downarrow \\
 D^k & \longrightarrow & Map(Y, W) \times_{Map(X, W)} Map(X, Z)
 \end{array}$$

has a lift in Top . The latter is equivalent to the pushout-product property that the map $(S^{k-1} \times Y) \cup_{S^{k-1} \times X} (D^k \times X) \rightarrow D^k \times Y$ is a cofibration.

The acyclicity statement is proved similarly by starting with any not-necessarily trivial Γ -fibration $Z \rightarrow W$. □

Given a homomorphism of monoids $j: \Gamma \rightarrow \Gamma'$, one gets an extension-restriction adjunction

$$j_! : \Gamma-Top \rightleftarrows \Gamma'-Top : j^*
 \tag{111}$$

Lemma A.3 *For any morphism of topological monoids $j: \Gamma \rightarrow \Gamma'$, the extension-restriction adjunction (111) is a Quillen adjunction. In particular, the extension functor $j_!$ preserves cofibrations and acyclic cofibrations. The restriction functor j^* preserves fibrations and weak equivalences. Moreover, if Γ' is cofibrant as a right Γ -module, then j^* also preserves cofibrations.*

Proof All the statements follow from definition except the last one, which is implied by the fact that the restriction functor preserves colimits and sends the generating cofibrations $S^{k-1} \times Y \rightarrow D^k \times Y$ to cofibrations. The latter assertion is a consequence of Lemma A.2 applied to $A \rightarrow B$ being $\emptyset \rightarrow \Gamma'$, and $X \rightarrow Y$ being $S^{k-1} \rightarrow D^k$. □

Definition A.4 (a) A sequence of morphisms of topological monoids

$$1 \rightarrow \Gamma_1 \xrightarrow{i} \Gamma \xrightarrow{p} \Gamma_2 \rightarrow 1
 \tag{112}$$

is called *short exact sequence* if Γ_2 is a quotient of Γ as a topological space (p is a quotient map), and i is a homeomorphism onto $p^{-1}(1)$.

(b) A short exact sequence (112) of topological monoids is called *split-surjective*, if p admits a continuous section $s: \Gamma_2 \rightarrow \Gamma$ which is a morphism of monoids, and the map $\Gamma_2 \times \Gamma_1 \rightarrow \Gamma$, $(\gamma_2, \gamma_1) \mapsto s(\gamma_2) \cdot i(\gamma_1)$, is surjective.

For a split-surjective short exact sequence (112), thanks to the inclusions i and s , the monoids Γ_1 and Γ_2 can be viewed as subobjects of Γ . For this reason in the sequel, we will be omitting i and s considering $\Gamma_1, \Gamma_2 \subset \Gamma$. Note also that if Γ_2 is a group, then the map $\Gamma_2 \times \Gamma_1 \rightarrow \Gamma, (\gamma_2, \gamma_1) \mapsto \gamma_2 \cdot \gamma_1$, is always surjective (and, in fact, bijective).

Lemma A.5 *Let (112) be a split-surjective short exact sequence of monoids. Let also Γ_2 fit into a split-surjective short exact sequence*

$$1 \rightarrow \Gamma_2^0 \rightarrow \Gamma_2 \xrightarrow{p_2} G_2 \rightarrow 1 \tag{113}$$

with G_2 being a group. Also assume that all elements of Γ_2^0 commute with those from Γ_1 inside Γ . Let $A \rightarrow B$ be a Γ_2 -cofibration and $X \rightarrow Y$ be a Γ -equivariant Γ_1 -cofibration with both X and Y having trivial action of Γ_2^0 . Then the pushout product map $(A \times Y) \cup_{A \times X} (B \times X) \rightarrow B \times Y$ is a Γ -cofibration. Moreover, the latter is acyclic if $A \rightarrow B$ or $X \rightarrow Y$ is.

Example A.6 If $A \rightarrow B$ is a Γ_2 -cofibration and $X \rightarrow Y$ is a Γ_1 -cofibration, then the pushout product map $(A \times Y) \cup_{A \times X} (B \times X) \rightarrow B \times Y$ is a $\Gamma_1 \times \Gamma_2$ -cofibration. (Take $\Gamma = \Gamma_1 \times \Gamma_2$ and $G_2 = 1$.)

Proof of Lemma A.5 Let $Z \rightarrow W$ be a trivial Γ -fibration. One has to show that the square (109) has a Γ -equivariant lift. By adjunction such lift defines a lift in the square (110). If Γ were a group then the mapping spaces $Map(Y, Z), Map(Y, W)$, etc., would be endowed with a natural right Γ -action: $(f \cdot \gamma)(y) := f(y \cdot \gamma^{-1}) \cdot \gamma$. The lift in question would arise from a lift in (109) if and only if it were Γ -equivariant. However, in our more general situation the mapping spaces do not get a natural Γ -action and the condition on the induced lift of (110) is less obvious. Denote by $Map_{\Gamma_1}(Y, Z)$ ($Map_{\Gamma_1}(Y, W)$, etc.) the subspace of $Map(Y, Z)$ of Γ_1 -equivariant maps. Since the action of Γ_1 is trivial on A and on B , the induced lift of (110) must factor through a lift in the square

$$\begin{array}{ccc}
 A & \xrightarrow{\quad\quad\quad} & Map_{\Gamma_1}(Y, Z) \\
 \downarrow & & \downarrow \\
 B & \xrightarrow{\quad\quad\quad} & Map_{\Gamma_1}(Y, W) \times_{Map_{\Gamma_1}(X, W)} Map_{\Gamma_1}(X, Y).
 \end{array} \tag{114}$$

On the other hand, the spaces $Map_{\Gamma_1}(Y, Z), Map_{\Gamma_1}(Y, W)$, etc., have a natural Γ_2 -action defined as follows:

$$(f \cdot \gamma_2)(y) := f(y \cdot p_2(\gamma_2)^{-1}) \cdot \gamma_2,$$

where $\gamma_2 \in \Gamma_2$. One checks that if $f \in Map_{\Gamma_1}(Y, Z)$, then so is $f \cdot \gamma_2$:

$$\begin{aligned}
 (f \cdot \gamma_2)(y \cdot \gamma_1) &= f(y \cdot \gamma_1 \cdot p_2(\gamma_2)^{-1}) \cdot \gamma_2 \\
 &= f(y \cdot p_2(\gamma_2)^{-1} \cdot p_2(\gamma_2) \cdot \gamma_1 \cdot p_2(\gamma_2)^{-1}) \cdot \gamma_2 \\
 &= f(y \cdot p_2(\gamma_2)^{-1}) \cdot p_2(\gamma_2) \cdot \gamma_1 \cdot p_2(\gamma_2)^{-1} \cdot \gamma_2 \\
 &= f(y \cdot p_2(\gamma_2)^{-1}) \cdot p_2(\gamma_2) \cdot p_2(\gamma_2)^{-1} \cdot \gamma_2 \cdot \gamma_1 \\
 &= f(y \cdot p_2(\gamma_2)^{-1}) \cdot \gamma_2 \cdot \gamma_1 = (f \cdot \gamma_2)(y) \cdot \gamma_1.
 \end{aligned}$$

The third equation is obtained using the fact that $p_2(\gamma_2) \cdot \gamma_1 \cdot p_2(\gamma_2)^{-1} \in \Gamma_1$ and $f \in Map_{\Gamma_1}(Y, Z)$. The fourth equation uses that $p_2(\gamma_2)^{-1} \cdot \gamma_2 \in \Gamma_2^0$ and that Γ_1 commutes with Γ_2^0 .

We claim that a map $B \rightarrow \text{Map}(Y, Z)$ is adjoint to a Γ -equivariant map $F : B \times Y \rightarrow Z$ if and only if it factors through $\text{Map}_{\Gamma_1}(Y, Z)$ and the map $B \rightarrow \text{Map}_{\Gamma_1}(Y, Z)$ is Γ_2 -equivariant. (The same holds for the maps $A \rightarrow \text{Map}(Y, Z)$, $B \rightarrow \text{Map}(Y, W)$, etc.)

Indeed, let $F : B \times Y \rightarrow Z$ be a Γ -equivariant map. Since Γ_1 acts trivially on B ,

$$F(b, y) \cdot \gamma_1 = F(b \cdot \gamma_1, y \cdot \gamma_1) = F(b, y \cdot \gamma_1).$$

Thus, $F(b, -) \in \text{Map}_{\Gamma_1}(Y, Z)$. To check that the induced map $B \rightarrow \text{Map}_{\Gamma_1}(Y, Z)$ is Γ_2 -equivariant, we need to make sure that $F(b \cdot \gamma_2, y) = F(b, y \cdot p_2(\gamma_2)^{-1}) \cdot \gamma_2$. One has

$$F(b, y \cdot p_2(\gamma_2)^{-1}) \cdot \gamma_2 = F(b \cdot \gamma_2, y \cdot p_2(\gamma_2)^{-1} \cdot \gamma_2) = F(b \cdot \gamma_2, y).$$

The last equation uses the fact that $p_2(\gamma_2)^{-1} \cdot \gamma_2 \in \Gamma_2^0$ and that Γ_2^0 acts trivially on Y .

In the other direction, let $F : B \times Y \rightarrow Z$ be the adjoint of a Γ_2 -equivariant map $B \rightarrow \text{Map}_{\Gamma_1}(Y, Z) \rightarrow \text{Map}(Y, Z)$. One has to check that F is Γ -equivariant. Since the product map $\Gamma_2 \times \Gamma_1 \rightarrow \Gamma$ is surjective, each $\gamma \in \Gamma$ can be written as $\gamma = \gamma_2 \cdot \gamma_1$, $\gamma_1 \in \Gamma_1$, $\gamma_2 \in \Gamma_2$. We need to check that $F(b \cdot \gamma_2 \cdot \gamma_1, y \cdot \gamma_2 \cdot \gamma_1) = F(b, y) \cdot \gamma_2 \cdot \gamma_1$. One has

$$\begin{aligned} F(b \cdot \gamma_2 \cdot \gamma_1, y \cdot \gamma_2 \cdot \gamma_1) &= F(b \cdot \gamma_2, y \cdot \gamma_2 \cdot \gamma_1) && \text{since } \Gamma_1 \text{ acts trivially on } B, \\ &= F(b \cdot \gamma_2, y \cdot \gamma_2) \cdot \gamma_1 && \text{since } F(b \cdot \gamma_2, -) \in \text{Map}_{\Gamma_1}(Y, Z), \\ &= F(b, y \cdot \gamma_2 \cdot p_2(\gamma_2)^{-1}) \cdot \gamma_2 \cdot \gamma_1 && \text{since } F \text{ is } \Gamma_2\text{-equivariant,} \\ &= F(b, y) \cdot \gamma_2 \cdot \gamma_1 && \text{since } \Gamma_2^0 \text{ acts trivially on } Y. \end{aligned}$$

As a consequence of the above, the square (109) has a Γ -equivariant lift if and only if the square (114) has a Γ_2 -equivariant lift. On the other hand, the left vertical map in (114) is a Γ_2 -cofibration. Therefore a lift exists provided the right vertical map in (114) is a trivial Γ_2 -fibration. The latter holds provided that the map is a trivial fibration in Top , which follows from the fact that $(S^{k-1} \times Y) \cup_{S^{k-1} \times X} (D^k \times X) \rightarrow D^k \times Y$ is a Γ_1 -cofibration, see Lemma A.2. □

Lemma A.7 *Let Γ be a topological monoid and Z be a Γ -cofibrant space. Then the functor*

$$Z \times_{\Gamma} - : \Gamma^{op}\text{-Top} \rightarrow \text{Top},$$

from left Γ -modules to spaces, sends left Γ -spaces that are cofibrant in Top to cofibrant spaces, and preserves the weak equivalences between such objects.

Proof Since a retract of a cofibrant space is a cofibrant space and a retract of a weak equivalence is a weak equivalence, we can assume that Z is Γ -cellular: $Z = \text{colim}_{\alpha < \lambda} Z_{\alpha}$, where each map $Z_{<\alpha} := \text{colim}_{\beta < \alpha} Z_{\beta} \rightarrow Z_{\alpha}$ is a pushout of a generating Γ -cofibration. One has to check two statements.

- If the statement of the lemma holds for $Z_{<\alpha}$, then it does for Z_{α} .
- If the statement of the lemma holds for each Z_{β} , $\beta < \alpha$, then it does for $Z_{<\alpha}$.

For any generating Γ -cofibration $S^{k-1} \times \Gamma \rightarrow D^k \times \Gamma$ and any cofibrant in Top left Γ -space A , the induced map

$$(S^{k-1} \times \Gamma) \times_{\Gamma} A \rightarrow (D^k \times \Gamma) \times_{\Gamma} A \tag{115}$$

is $S^{k-1} \times A \rightarrow D^k \times A$. Since A is cofibrant, this map (115) is a cofibration. On the other hand, for any weak equivalence of Γ -spaces $A \rightarrow B$, the induced maps $S^{k-1} \times A \rightarrow S^{k-1} \times B$ and $D^k \times A \rightarrow D^k \times B$ are weak equivalences. The first statement above follows by applying Proposition 2.12 and recalling that Top is left proper.

To prove the second statement, we notice that the sequence $Z_\alpha \times_\Gamma A, \alpha < \lambda$, is a sequence of cofibrations, being pushouts of cofibrations of the form (115). Therefore,

$$\operatorname{colim}_{\alpha < \lambda} (Z_\alpha \times_\Gamma A) \simeq \operatorname{hocolim}_{\alpha < \lambda} (Z_\alpha \times_\Gamma A).$$

One has the same weak equivalence of spaces for B . Since $Z_\alpha \times_\Gamma A \rightarrow Z_\alpha \times_\Gamma B, \alpha < \lambda$, are all weak equivalences,

$$\operatorname{hocolim}_{\alpha < \lambda} (Z_\alpha \times_\Gamma A) \simeq \operatorname{hocolim}_{\alpha < \lambda} (Z_\alpha \times_\Gamma B).$$

On the other hand, the functors $(-)\times_\Gamma A$ and $(-)\times_\Gamma B$ preserve colimits. We conclude

$$Z \times_\Gamma A = \operatorname{colim}_{\alpha < \lambda} (Z_\alpha \times_\Gamma A) \simeq \operatorname{colim}_{\alpha < \lambda} (Z_\alpha \times_\Gamma B) = Z \times_\Gamma B.$$

□

Compatible action of a monoid and a group: We say that a topological monoid Γ is endowed with a right action of a group K if one is given a map $\Gamma \times K \rightarrow \Gamma, (\gamma, k) \mapsto \gamma^k$, which is a right K -action on the set Γ and for every $k \in K$, the map $(-)^k: \Gamma \rightarrow \Gamma$ is a monoid homomorphism. Given such an action, one defines the semi-direct product monoid $\Gamma \rtimes K$. Its underlying set is $\Gamma \times K$, while multiplication is as follows

$$(\gamma_1, k_1) \cdot (\gamma_2, k_2) = (\gamma_1 \cdot \gamma_2^{k_1^{-1}}, k_1 \cdot k_2).$$

A right $\Gamma \rtimes K$ -space A can equivalently be seen as a right Γ -module with a right K -action compatible in the sense

$$(a \cdot \gamma) \cdot k = (a \cdot k) \cdot \gamma^k, \quad \text{for any } a \in A, \gamma \in \Gamma, k \in K.$$

Note that the same map $\Gamma \times K \rightarrow \Gamma$, defines a right K -action on Γ^{op} – the monoid with the reversed multiplication. A right $\Gamma^{op} \rtimes K$ -space X can equivalently be seen as a left Γ -module with a right K -action compatible in the sense

$$(\gamma \cdot x) \cdot k = \gamma^k \cdot (x \cdot k), \quad \text{for any } x \in X, \gamma \in \Gamma, k \in K.$$

Lemma A.8 *Let $1 \rightarrow K_1 \rightarrow K \rightarrow K_2 \rightarrow 1$ be a short exact sequence of topological groups. Let Γ be a topological monoid endowed with a right K_2 -action. If $A \rightarrow B$ is a $\Gamma \rtimes K_2$ -cofibration and $X \rightarrow Y$ is a $\Gamma^{op} \rtimes K$ -equivariant K_1 -cofibration, then the pushout-product map $(A \times_\Gamma Y) \cup_{A \times_\Gamma X} (B \times_\Gamma X) \rightarrow B \times_\Gamma Y$ is a K -cofibration. Moreover, the latter is acyclic if $A \rightarrow B$ or $X \rightarrow Y$ is.*

Proof Let $Z \rightarrow W$ be a trivial K -fibration. One has to show that any square

$$\begin{CD} (A \times_\Gamma Y) \cup_{A \times_\Gamma X} (B \times_\Gamma X) @>>> Z \\ @VVV @VVV \\ B \times_\Gamma Y @>>> W \end{CD} \tag{116}$$

has a K -equivariant lift. One has a homeomorphism of mapping spaces

$$\operatorname{Map}_K(B \times_\Gamma Y, Z) = \operatorname{Map}_{\Gamma \rtimes K_2}(B, \operatorname{Map}_{K_1}(Y, Z)).$$

(One has similar homeomorphisms for $\operatorname{Map}_K(A \times_\Gamma Y, Z)$, etc.) The K -action on $B \times_\Gamma Y$ is the diagonal one: $(b, y) \cdot k = (b \cdot k, y \cdot k)$. The action of $\Gamma \rtimes K_2$ on $\operatorname{Map}_{K_1}(Y, W)$ is defined as follows:

$$(f \cdot (\gamma, k_2))(y) = f(\gamma \cdot (y \cdot k_2^{-1})) \cdot k,$$

where $k \in K$ is any point in the preimage of $k_2 \in K_2$.

Using this, the existence of a lift in (116) is equivalent to the existence of a $\Gamma \times K_2$ -equivariant lift of the square

$$\begin{array}{ccc}
 A & \longrightarrow & \text{Map}_{K_1}(Y, Z) \\
 \downarrow & & \downarrow \\
 B & \longrightarrow & \text{Map}_{K_1}(Y, W) \times_{\text{Map}_{K_1}(X, W)} \text{Map}_{K_1}(X, Z).
 \end{array} \tag{117}$$

Since $A \rightarrow B$ is a $\Gamma \times K_2$ -cofibration, one has to check that the right vertical arrow is a trivial $\Gamma \times K_2$ -fibration, or, equivalently a fibration in *Top*. In other words, one has to check that any square below

$$\begin{array}{ccc}
 S^{k-1} & \longrightarrow & \text{Map}_{K_1}(Y, Z) \\
 \downarrow & & \downarrow \\
 D^k & \longrightarrow & \text{Map}_{K_1}(Y, W) \times_{\text{Map}_{K_1}(X, W)} \text{Map}_{K_1}(X, Z)
 \end{array}$$

has a lift in *Top*. The latter is equivalent to the pushout-product property that the map $(S^{k-1} \times Y) \cup_{S^{k-1} \times X} (D^k \times X) \rightarrow D^k \times Y$ is a K_1 -cofibration, true by Lemmas A.1 or A.2. □

A.2 Cellularly Equivariant Cofibrations

Definition A.9 Let G be a discrete group.

- (a) A G -equivariant map $X_0 \rightarrow X_1$ is called a G -equivariant cell attachment if it fits into a pushout diagram

$$\begin{array}{ccc}
 S^{k-1} \times (H \backslash G) & \longrightarrow & D^k \times (H \backslash G) \\
 \downarrow & & \downarrow \\
 X_0 & \longrightarrow & X_1,
 \end{array} \tag{118}$$

where $H \subset G$ is a subgroup of G .

- (b) A G -equivariant map $X_0 \rightarrow X$ is called a *cellularly G -equivariant cofibration* if it is a G -equivariant retract of a possibly transfinite sequence of G -equivariant cell attachments.

In fact there exists a model structure on G -*Top* for which cofibrations are exactly the cellularly G -equivariant cofibrations [17]. This model structure produces a different (from projective) homotopy category as it has a smaller class of equivalences for which one has to take into account all orbit subspaces. Any G -space is still fibrant in this model structure. We do not use this more subtle model structure on G -*Top*. We just need a few technical lemmas below.

Lemma A.10 For any cellularly G -equivariant cofibration $X_0 \rightarrow X$ and any G -space Y , the induced map

$$\text{Map}_G(X, Y) \rightarrow \text{Map}_G(X_0, Y)$$

is a Serre fibration.

Proof A retract of a Serre fibration is a Serre fibration as well as is the limit of a tower of Serre fibrations. Therefore it is enough to consider the case of a G -equivariant cell attachment $X_0 \rightarrow X_1$ as in (118). One has a pullback square

$$\begin{array}{ccc} \text{Map}_G(X_1, Y) & \longrightarrow & \text{Map}_G(D^k \times (H \setminus G), Y) \\ \downarrow & & \downarrow \\ \text{Map}_G(X_0, Y) & \longrightarrow & \text{Map}_G(S^{k-1} \times (H \setminus G), Y). \end{array}$$

The left vertical map is a Serre fibration provided the right vertical map is one. One has

$$\begin{aligned} \text{Map}_G(D^k \times (H \setminus G), Y) &= \text{Map}_H(D^k, Y) = \text{Map}(D^k, Y^H); \\ \text{Map}_G(S^{k-1} \times (H \setminus G), Y) &= \text{Map}_H(S^{k-1}, Y) = \text{Map}(S^{k-1}, Y^H). \end{aligned}$$

Therefore, the right vertical map in the square above is the map $\text{Map}(D^k, Y^H) \rightarrow \text{Map}(S^{k-1}, Y^H)$, which is a Serre fibration. This follows from the fact that the invariant space Y^H is fibrant (like any topological space) and the map from S^{k-1} to D^k is a cofibration. □

Lemmas A.11–A.14 below help to recognize cellularly equivariant cofibrations.

Lemma A.11 *The realization of any G -equivariant inclusion of simplicial G -sets is a cellularly G -equivariant cofibration.*

Proof Obvious. □

Lemma A.12 *For any homomorphism $\phi : G_1 \rightarrow G_2$ of discrete groups, both the restriction and extension functors*

$$\phi_! : G_1\text{-Top} \rightleftarrows G_2\text{-Top} : \phi^*$$

preserve cellularly equivariant cofibrations.

Proof Obvious. □

Lemma A.13 *If*

$$X_0 \rightarrow X \quad \text{and} \quad Y_0 \rightarrow Y \tag{119}$$

are cellularly G -equivariant cofibrations, then so is the pushout-product map

$$(X \times Y_0) \cup_{X_0 \times Y_0} (X_0 \times Y) \rightarrow X \times Y. \tag{120}$$

Proof For generating G -equivariant cell attachments $S^{k_1-1} \times (H_1 \setminus G) \rightarrow D^{k_1} \times (H_1 \setminus G)$ and $S^{k_2-1} \times (H_2 \setminus G) \rightarrow D^{k_2} \times (H_2 \setminus G)$, the map (120) becomes

$$S^{k_1+k_2-1} \times (H_1 \setminus G) \times (H_2 \setminus G) \rightarrow D^{k_1+k_2} \times (H_1 \setminus G) \times (H_2 \setminus G).$$

The G -set $(H_1 \setminus G) \times (H_2 \setminus G)$ is isomorphic to a disjoint union of identical G -sets $(H_1 \cap H_2) \setminus G$. Thus, the statement of the lemma holds in this case. Similarly it is true for an arbitrary pair of G -equivariant cell attachments.

In case $X = \text{colim}_{\alpha < \lambda_1} X_\alpha$ and $Y = \text{colim}_{\alpha < \lambda_2} Y_\alpha$ are (possibly transfinite) sequences of G -equivariant cell attachments, then the inclusion (120) is also a (transfinite) sequence of G -equivariant cell attachments

$$\text{colim}_{(\alpha_1, \alpha_2) < (\lambda_1, \lambda_2)} Z_{\alpha_1, \alpha_2},$$

where the set $\lambda_1 \times \lambda_2$ is given the lexicographical order and therefore is also an ordinal. The spaces Z_{α_1, α_2} are defined recursively:

$$Z_{\alpha_1, \alpha_2} = \left(\operatorname{colim}_{(\beta_1, \beta_2) < (\alpha_1, \alpha_2)} Z_{\beta_1, \beta_2} \right) \cup (X_{\alpha_1} \times Y_{\alpha_2}),$$

with $Z_{0,0}$ being the left-hand side of (120).

Finally, if (119) are retracts of (transfinite) sequences of cell attachments then so is the pushout-product. \square

Lemma A.14 *Let $\partial X \rightarrow X$ be a cofibration in Top. Then $\partial(X^{\times n}) \rightarrow X^{\times n}$ is a cellularly Σ_n -equivariant cofibration.*

Proof One needs to check it first for the inclusion $S^{k-1} \rightarrow D^k$, which is done by stratifying $(D^k)^{\times n}$ into Σ_n -orbits and then decomposing the orbits into cells. Applying previous Lemmas A.12 and A.13 we can conclude that the inclusion

$$\partial \prod_{i=1}^{\ell} (D^{k_i})^{\times n_i} \rightarrow \prod_{i=1}^{\ell} (D^{k_i})^{\times n_i}$$

is a cellularly $(\prod_{i=1}^{\ell} \Sigma_{n_i})$ -equivariant cofibration. The rest of the argument is similar to the proof of the previous lemma. Assuming that $\partial X \rightarrow X$ is a possibly transfinite sequence of cell attachments $\operatorname{colim}_{\alpha < \lambda} X_{\alpha}$, we decompose $X_{\alpha}^{\times n}$ extending the cellular structure of $\operatorname{colim}_{\beta < \alpha} X_{\beta}^{\times n}$. \square

Acknowledgements The authors would like to thank C. Berger, P. Bressie, R. Campos, M. Ching, D. E. Fajou, N. Idrissi, M. Weber, and T. Willwacher for communication.

Funding Open Access funding enabled and organized by Projekt DEAL.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Angeltveit, V.: Enriched Reedy categories. Proc. Am. Math. Soc. **136**(7), 2323–2332 (2008)
2. Arone, G., Ching, M.: Cross-effects and the classification of Taylor towers. Geom. Topol. **20**(3), 1445–1537 (2016)
3. Arone, G., Turchin, V.: On the rational homology of high dimensional analogues of spaces of long knots. Geom. Topol. **18**(3), 1261–1322 (2014)
4. Barwick, C.: On Reedy Model Categories, Unpublished. [arXiv:0708.2832](https://arxiv.org/abs/0708.2832) (2007)
5. Bayeh, M., Hess, K., Karpova, V., Kedziorek, M., Riehl, E., Shipley, B.: Left-induced model structures and diagram categories. In: Women in Topology: Collaborations in Homotopy Theory, Contemporary Mathematics, vol. 641, pp. 49–81. American Mathematical Society, Providence (2015)
6. Berger, C., Moerdijk, I.: Axiomatic homotopy theory for operads. Comment. Math. Helv. **78**(4), 805–831 (2003)
7. Berger, C., Moerdijk, I.: The Boardman–Vogt resolution of operads in monoidal model categories. Topology **45**, 807–849 (2006)

8. Berger, C., Moerdijk, I.: Resolution of coloured operads and rectification of homotopy algebras. *Contemp. Math.* **431**, 31–58 (2007)
9. Berger, C., Moerdijk, I.: On an extension of the notion of Reedy category. *Math. Z.* **269**(3–4), 977–1004 (2008)
10. Brown, R.: *Topology and groupoids*. Third edition of Elements of modern topology [McGraw-Hill, New York, 1968]. BookSurge, LLC, Charleston, SC, 2006. xxvi+512 pp
11. Dundas, B., Röndigs, O., Østvær, P.A.: Enriched functors and stable homotopy theory. *Doc. Math.* **8**, 409–488 (2003)
12. Ducoulombier, J.: Swiss-Cheese action on the totalization of action-operads. *Algebr. Geom. Topol.* **16**(3), 1683–1726 (2016)
13. Ducoulombier, J.: From maps between coloured operads to Swiss-Cheese algebras. *Ann. Inst. Fourier (Grenoble)* **68**(2), 661–724 (2018)
14. Ducoulombier, J.: Delooping of high-dimensional spaces of string links. [arXiv:1809.00682](https://arxiv.org/abs/1809.00682) (2018)
15. Ducoulombier, J., Turchin, V.: Delooping the functor calculus tower. In: *The Proceedings of the London Mathematical Society*. [arXiv:1708.02203](https://arxiv.org/abs/1708.02203) (2022) (to appear)
16. Ducoulombier, J., Turchin, V., Willwacher, T.: On the delooping of (framed) embedding spaces. *Trans. Am. Math. Soc.* **374**, 7657–7677 (2021)
17. Farjoun, E.D.: Homotopy theories for diagrams of spaces. *Proc. Am. Math. Soc.* **101**(1), 181–189 (1987)
18. Fresse, B.: Homotopy of operads and Grothendieck–Teichmüller groups. Part 1. The algebraic theory and its topological background. In: *Mathematical Surveys and Monographs*, vol. 217. American Mathematical Society, Providence (2017)
19. Fresse, B.: Homotopy of operads and Grothendieck–Teichmüller groups. Part 2. The applications of (rational) homotopy theory methods. In: *Mathematical Surveys and Monographs*, vol. 217. American Mathematical Society, Providence (2017)
20. Fresse, B.: *Modules Over Operads and Functors*. *Lecture Notes in Mathematics*, vol. 1967. Springer, Berlin (2009)
21. Fresse, B., Turchin, V., Willwacher, T.: The homotopy theory of operad subcategories. *J. Homotopy Relat. Struct.* **13**(4), 689–702 (2018)
22. Garner, R., Kedziorek, M., Riehl, E.: Lifting accessible model structures. *J. Topol.* **13**(1), 59–76 (2020)
23. Goerss, P.G., Jardine, J.F.: *Simplicial Homotopy Theory*. In: *Progress in Mathematics*, vol. 174. Birkhäuser Verlag, Basel (1999). xvi+510 pp. ISBN: 3-7643-6064-X
24. Harper, J.E.: Homotopy theory of modules over operads and non- Σ operads in monoidal model categories. *J. Pure Appl. Algebra* **214**, 1407–1434 (2010)
25. Hess, K., Kedziorek, M., Riehl, E., Shipley, B.: A necessary and sufficient condition for induced model structures. *J. Topol.* **10**(2), 324–369 (2017)
26. Hirschhorn, P.S.: *Model Categories and Their Localizations*. *Mathematical Surveys and Monographs*, vol. 99, p.xvi+457 (2003)
27. Hackney, P., Robertson, M., Yau, D.: Relative left properness of colored operads. *Algebr. Geom. Topol.* **16**(5), 2691–2714 (2016)
28. Hovey, M.: *Model Categories*. *Mathematical Surveys and Monographs*, vol. 63. American Mathematical Society, Providence (1999)
29. Hovey, M.: *Monoidal model categories*. [arXiv:math/9803002](https://arxiv.org/abs/math/9803002) (1998)
30. Kelly, G.M.: On the operads of J. P. May. *Repr. Theory Appl. Categ.* **13**, 1–13 (2005)
31. Lewis, G.: *Compactly generated spaces*, appendix A of *The Stable Category and Generalized Thom Spectra*, PhD thesis Chicago (1978)
32. Lurie, J.: *Higher Topos Theory*. *Annals of Mathematics Studies*, vol. 170. Princeton University Press, Princeton (2009)
33. Moser, L.: Injective and projective model structures on enriched diagram categories. *Homol. Homotopy Appl.* **21**(2), 279–300 (2019)
34. Reedy, C.: *Homotopy theory of model categories*, Unpublished. <http://www-math.mit.edu/~psh/> (1974)
35. Rezk, C.: *Spaces of algebra structures and cohomology of operads*. PhD thesis, MIT (1996)
36. Rezk, C.: A model for the homotopy theory of homotopy theory. *Trans. Am. Math. Soc.* **353**(3), 973–1007 (2001)
37. Riehl, E., Verity, D.: The theory and practice of Reedy categories. *Theory Appl. Categ.* **29**, 256–301 (2014)