


RESEARCH

Open Access



A β oligomers trigger necroptosis-mediated neurodegeneration via microglia activation in Alzheimer's disease

Natalia Salvadores^{1,2,9}, Ines Moreno-Gonzalez^{3,4,5}, Nazaret Gamez^{3,5}, Gabriel Quiroz², Laura Vegas-Gomez³, Marcela Escandón¹, Sebastian Jimenez^{4,6,7}, Javier Vitorica^{4,6,7}, Antonia Gutierrez^{3,4}, Claudio Soto⁵ and Felipe A. Court^{1,2,8*} 

Abstract

Alzheimer's disease (AD) is a major adult-onset neurodegenerative condition with no available treatment. Compelling reports point amyloid- β (A β) as the main etiologic agent that triggers AD. Although there is extensive evidence of detrimental crosstalk between A β and microglia that contributes to neuroinflammation in AD, the exact mechanism leading to neuron death remains unknown. Using postmortem human AD brain tissue, we show that A β pathology is associated with the necroptosis effector pMLKL. Moreover, we found that the burden of A β oligomers (A β o) correlates with the expression of key markers of necroptosis activation. Additionally, inhibition of necroptosis by pharmacological or genetic means, reduce neurodegeneration and memory impairment triggered by A β o in mice. Since microglial activation is emerging as a central driver for AD pathogenesis, we then tested the contribution of microglia to the mechanism of A β o-mediated necroptosis activation in neurons. Using an in vitro model, we show that conditioned medium from A β o-stimulated microglia elicited necroptosis in neurons through activation of TNF- α signaling, triggering extensive neurodegeneration. Notably, necroptosis inhibition provided significant neuronal protection. Together, these findings suggest that A β o-mediated microglia stimulation in AD contributes to necroptosis activation in neurons and neurodegeneration. As necroptosis is a druggable degenerative mechanism, our findings might have important therapeutic implications to prevent the progression of AD.

Keywords: Alzheimer's disease, Amyloid- β oligomers, Necroptosis, Microglia, Neurodegeneration, Neuroprotection

Introduction

Alzheimer's disease (AD) represents the most frequent brain disorder in humans, affecting mainly the elderly population. AD is characterized by progressive neurodegeneration leading to impaired cognition, memory loss, and dementia [1]. The neuropathological hallmarks of AD include the accumulation of intraneuronal deposits of hyperphosphorylated tau (pTau) protein and intracellular

and extracellular aggregates of amyloid- β (A β), which are associated with dystrophic neurites and activated microglia [2]. Increasing studies have demonstrated that soluble oligomeric A β structures contribute importantly to neurodegeneration in AD and correlate with cognitive alterations [2–7].

Familial AD, which represents less than 5% of the cases, is caused by mutations in the amyloid precursor protein (APP) gene or mutations in the presenilin 1 or 2 genes, leading to increased deposition of A β [1, 2]. The most common form of the disease is sporadic AD with unknown origin, where aging is the most important risk factor. Accumulating evidence has attributed a causative

*Correspondence: felipe.court@umayor.cl

¹ Center for Integrative Biology, Faculty of Sciences, Universidad Mayor, Santiago, Chile

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

role to A β in the etiopathogenesis of AD [2, 8]. Neuroinflammation is a critical component of the pathogenesis of A β , involving abnormal activation of glial cells and the production of various proinflammatory cytokines [9]. Following the first description of activated microglial cells in close contact with amyloid plaques in AD brains, extensive research focusing on the role of microglia in AD pathogenesis has been conducted [10–14]. Upon interaction with A β , activated microglia initiate an innate immune response that leads to sustained production and secretion of pro-inflammatory mediators including interleukin (IL)-6, IL-1 β , and tumor necrosis factor- α (TNF- α), contributing to neuronal damage and cognitive decline [15–19]. Interestingly, mutations in the Triggering Receptor Expressed on Myeloid cells 2 (TREM2), which is expressed in microglia, increase the risk of AD by about 3- to 5-fold [20, 21]. TREM2 is involved in microglial signaling and plays an important role in A β phagocytosis as well as suppressing cytokine production and neuroinflammation [22–24]. These studies support an important contribution of inflammatory responses in the mechanism of A β -induced neurodegeneration.

Necroptosis is a mechanism of programmed cell death that participates in the pathogenesis of several diseases [25]. Necroptosis is activated in postmortem human AD brains and the expression of necroptotic markers correlates with the disease stage and progression at the cognitive level [26, 27]. Nevertheless, the mechanism that triggers necroptosis in AD remains to be established. Different stimuli induce necroptosis activation, including TNF ligand family members [28], interferons [29], and toll-like receptor 3 stimulation [30]. Stimulation of TNF type 1 receptor (TNFR1) leads to the recruitment of receptor-interacting protein kinase 1 (RIPK1) [31, 32], which triggers the activation of either caspase-8 or RIPK3, leading to apoptosis or necroptosis, respectively [33]. Following a stimulus such as TNF- α ligation, and in the absence of caspase-8 activity, the interaction of RIPK1 and RIPK3 induces the formation of the necrosome complex and RIPK3 phosphorylation, a key event in necroptosis activation [34–36]. RIPK3 activation leads to the recruitment and phosphorylation of the pseudokinase mixed lineage kinase domain-like protein (MLKL) within the necrosome complex. Upon activation, MLKL is oligomerized and executes necroptosis leading to cell death [37, 38]. Interestingly, recent studies suggest that necroptosis may operate as a relevant degenerative pathway in models of amyotrophic lateral sclerosis (ALS) [39], multiple sclerosis (MS) [40], and Parkinson's disease (PD) [41].

Here, we investigated the role of A β pathology in the mechanism underlying necroptosis activation in AD. We found that A β oligomers (A β o) are associated with

key necroptosis drivers in the brains of AD patients and that targeting this death mechanism reduces A β o-related neurodegeneration and memory impairment in mice. Furthermore, we report that in vitro, A β o elicit necroptosis-mediated neurodegeneration via microglia activation through the engagement of TNF- α signaling. This study sheds light on the role of necroptosis in the crosstalk between glial and neuronal cells in AD and provides evidence indicating that necroptosis constitutes a potential therapeutic target for the disease.

Methods

Human samples

Histological analyses: Hippocampal brain samples were obtained from the Neurological Tissue Bank of IDIBELL-Hospital of Bellvitge (Barcelona, Spain) fixed in formaldehyde and embedded in paraffin. Research on human samples was performed following the code of ethics of the World Medical Association (Declaration of Helsinki). Samples were manipulated following the universal precautions for working with human samples and as directed by the supplier biobank and local ethic committees following Spanish legislation.

Molecular and biochemical analyses: autopsy specimens from the medial temporal lobe (hippocampal/parahippocampal regions) were obtained from the tissue bank Fundación CIEN (BT-CIEN; Centro de Investigación de Enfermedades Neurológicas; Madrid, Spain) and from the Neurological Tissue Bank of IDIBELL-Hospital of Bellvitge (Barcelona, Spain). The utilization of postmortem human samples was approved by the corresponding biobank ethics committees and by the “Comite de Etica de la Investigacion (CEI), Hospital Virgen del Rocio”, Seville, Spain. We considered the age-matched Braak II individuals as controls in our experiments. Only Braak V–VI cases were clinically classified as demented (AD) patients.

Mice

MLKL knockout mice were kindly provided by Dr. Douglas Green (St. Jude Children's Research Hospital, Memphis, TN, USA) and have been described previously [42]; male (4–6 months old) mice were used, wild-type littermates were used as controls. For the GSK'872 experiment, C57BL/6 mice were used and were obtained from Jackson Laboratory (Bar Harbor, ME, USA), male (13–14 months old) mice were used. Animals were housed in groups of a maximum of five in individually ventilated cages under standard conditions (22 °C, 12 h light–dark cycle) receiving food and water ad libitum. All animal manipulations were carried out in accordance with standard regulations and approved by the Animal

Care and Use Scientific Ethics Committee of Mayor University.

Intracerebroventricular injection

Mice were anesthetized with isoflurane and placed in a stereotaxic frame (David Kopf Instruments, USA). A single cerebroventricular injection was performed in both lateral ventricles according to the following stereotaxic coordinates: -1.0 ± 0.06 mm posterior to bregma, 1.8 ± 0.1 mm lateral to the sagittal suture, and 2.4 mm in depth, as previously described [43]. Five μ l of a solution of A β o (100 μ M) or vehicle (PBS) were injected in each ventricle at a rate of 0.5 μ l/min. Behavioural and histopathological analysis were performed at 14 and 18 days after the injection, respectively.

GSK'872 administration

GSK'872 (Tocris Bioscience) was dissolved in DMSO to a final concentration of 100 mM (stock). The working solution was prepared by diluting the stock solution with DMSO and then sterile PBS at room temperature to 0.3 mM (20% DMSO). The vehicle consisted of a solution of 20% DMSO in PBS. GSK'872 or vehicle were administered via intraperitoneal injection (1 mg/Kg) at the day of the intracerebroventricular injection of A β o (day 0), and then at day 7 and 14 post-injection.

Behavioral assessment

Two weeks after surgery (day 14), cognitive performance was tested using the Morris Water Maze as previously described [44]. In this test, using visual cues for orientation, animals learn to swim to a hidden platform under the water. During the training, mice are placed into the pool and allowed to explore for 1 min. If the animals do not find the platform, they are guided to it. This training procedure was performed 4 times a day for four consecutive days per animal. On day five, the platform was removed from the pool, and the time that the animals spent in the target quadrant was measured. At the end of the experiments, animals were euthanized by an overdose of anesthesia, and all efforts were made to minimize suffering. Then, animals were transcardially perfused with 50 mL of PBS.

Preparation of synthetic oligomers

A β o were prepared as previously described [45]. Briefly, lyophilized recombinant A β 42 peptide (rPeptide, A-1163) was re-suspended to 5 mM in DMSO and then further diluted to 100 μ M in cold Neurobasal medium for the in vitro experiments or in sterile PBS for the in vivo

experiments. The peptide was incubated at 4 °C for 24 h to generate oligomers, which were characterized by electron microscopy.

Total RNA extraction, retrotranscription, and quantitative real-time PCR

Total RNA was extracted from human samples using TriPure Isolation Reagent (Roche). RNA integrity (RIN) was determined by RNA Nano 6000 (Agilent). No differences between the Braak groups were observed (RIN: 6.95 ± 1.4). RNA was quantified using NanoDrop 2000 spectrophotometer (Thermo Fischer). Retrotranscription (RT) (4 μ g of total RNA) was performed with the High-Capacity cDNA Archive Kit (Applied Biosystems). For quantitative real-time PCR (qPCR), 40 ng of cDNA were mixed with 2 \times Taqman Universal Master Mix (Applied Biosystems) and 20 \times Taqman Gene Expression assay probes (*Ripk1*, Hs01041869_m1; *Ripk3*, Hs00179132_m1; *Mkl1*, Hs04188505_m1; Applied Biosystems). qPCR reactions were done using an ABI Prism 7900HT (Applied Biosystems). The cDNA levels were determined using GAPDH and beta-actin. We observed a highly significant linear correlation between the cycle threshold (Ct) of both genes (beta-actin vs GAPDH, $r=0.912$, $F[1, 46]=157.73$, $p<0.0001$). Thus, normalization using either beta-actin (Hs99999903_1, Applied Biosystems) or GAPDH (Hs03929097_g1, Applied Biosystems) produced identical results. Routinely, we used GAPDH as housekeeper. Results were expressed using the comparative double-delta Ct method (2- $\Delta\Delta$ Ct). Δ Ct values represent GAPDH normalized expression levels. $\Delta\Delta$ Ct was calculated using Braak II.

Preparation of soluble S1 fractions

Soluble S1 fractions from human samples were prepared as described [46]. Briefly, human tissue was homogenized (Dounce homogenizer) in TBS (20 mM Tris-HCl, 140 mM NaCl, pH 7.5) containing protease and phosphatase inhibitors (Roche). Homogenates were ultracentrifuged (4 °C for 60 min) at 100,000 $\times g$ (Optima MAX Preparative Ultracentrifuge, Beckman Coulter). Supernatants (S1 fractions), were aliquoted and stored at -80 °C.

Tissue processing for histopathology

For the study of human samples, deparaffinized 10- μ m-thick sections were used. After auto-fluorescence blocking (Millipore), sections were subjected to antigen retrieval by heating at 80 °C for 30 min in 10 mM citrate buffer pH 6.0. Samples were blocked with 5% BSA diluted in PBS Tx-100 (0.2%). For double staining, samples were incubated with anti-PMLKL (1:200, Sabbiotech) for 48 h and stained with

Thioflavin-S (0.025% in PBS) for 8 min. Signal was detected incubating for 1.5 h with the corresponding Alexa secondary antibody (1:500, Invitrogen). For double immunostaining with anti-pMLKL (1:200, Sabbiotech) and anti-MAP2 (1:5000, Thermo Fisher) or anti-Iba-1 antibodies (1:1000, Wako), samples were incubated overnight at 4 °C. Then, sections were incubated with the corresponding Alexa secondary antibodies (1:1000). For triple staining, samples were incubated for 10 min in Amylo-Glo[®] RTD (Biosensis), washed, and consecutively incubated with anti-pMLKL and anti-Iba-1 antibodies. Signal was detected as described above. Finally, samples were treated with an antifade mounting medium (VECTASHIELD). Fluorescent microscopy images were taken in a Leica SP8 confocal laser microscope coupled to a computer with TCS NT software (Leica).

For the histological study on mice, following perfusion, the whole brain was quickly removed, the right hemisphere was fixed with 4% PFA and then incubated in 30% sucrose before for OCT-embedding. Brains were sagittally cut into 20- μ m-thick serial slices using a cryostat. Tissue sections were rinsed with TBS and then mounted onto superfrost plus slides. Antigen retrieval was performed by submerging slices in 10 mM citrate buffer pH 6.0 at 100 °C for 20 min. Samples were blocked with 5% BSA and then incubated with the antibody solution (anti-pMLKL 1:200, Abcam; anti-Iba1 1:700, Wako), overnight at 4 °C. Finally, antigens were visualized by incubating with the corresponding Alexa secondary antibodies (1:500, Thermo Fisher). Fluoro-Jade C staining: the FJC ready-to-dilute staining kit (Biosensis) was used. Briefly, sections were immersed in a basic alcohol solution consisting of 1% sodium hydroxide in 80% ethanol for 5 min, followed by 2 min in 70% ethanol and 2 min in distilled water. Then, sections were incubated for 10 min in a solution of 0.06% potassium permanganate. Samples were rinsed in distilled water and stained for 10 min with the FJ-C solution. After rinsing, the air-dried slides were cleared in xylene for 1 min and then coverslipped. Images were obtained using a Leica DMi8 microscope.

Image analysis

All analyzes were carried out using Image J software (v1.42q). Colocalization analysis: The Coloc2 plugin was used to calculate the Manders Correlation values for intensities above Costes threshold. These values were used to obtain the ratio of pixels from one channel that colocalizes with total pixels in the second channel. The percentage of stained area was determined in thresholded images, by quantifying the signal above threshold divided

by the total image area. Total stained area in a specific ROI was determined in thresholded images, by quantifying the signal above threshold in that ROI divided by the area of the ROI.

Cell culture

Hippocampal neuronal cultures: Primary neuronal cultures were prepared from embryonic Sprague–Dawley rats (E18). Briefly, meninges-free hippocampi were isolated, trypsinized, and plated onto poly-L-lysine (0.05 mg/ml)-coated glass coverslips on tissue culture wells (150 cells/mm²). Neurons were grown in Neurobasal media with L-glutamine and B27 supplements to consistently provide neuronal cultures >95% pure. Cultures were treated with 1 μ M AraC to inhibit glial cell proliferation. For all experiments, neurons were treated on in vitro day 7.

Microglial cell culture: Microglial cells were cultured from postnatal day 3 Sprague–Dawley rat brains. Briefly, meninges-free cortices were dissected, and cells were mechanically dissociated. Cells were plated onto tissue culture flasks in DMEM media containing 10% heat-inactivated fetal bovine serum. Culture medium was replaced on in vitro days 1 and 5. After 10 days, the cultures were shaken for 2 h at 180 rpm on a rotary shaker to remove microglia.

Co-cultures: Hippocampal and microglial cell cultures were performed as described above. Microglial cells were added to hippocampal neuronal cultures at a ratio of 1:1, at in vitro day 7, 4 h before A β treatments.

Preparation of conditioned medium

Conditioned medium was prepared as previously described [47]. Microglial cells (180 cells/mm²) were plated onto immobilized A β (7.3 pmol/mm²) to prevent their subsequent collection in the conditioned medium. To immobilize the oligomers, tissue culture dishes were coated with nitrocellulose that was obtained by dissolving a piece of membrane in methanol. Oligomers were distributed on the dish and allowed to dry for 10 min. Then, the dish was washed twice and blocked with 5% BCA in DMEM. After 48 h of stimulation, the microglial conditioned medium was collected and stored at – 80 °C. To prepare the control medium, the same process was performed except that oligomers were not added. Additionally, a negative control was prepared by collecting medium from dishes containing immobilized A β alone.

Immunofluorescence

Neurons were fixed in 4% PFA at room temperature for 20 min. Cells were then blocked using 5% fish gelatin,

0,1% triton-X-100 for 2 h. Then, cells were incubated with the primary antibody solution (anti-acetylated tubulin 1:1000, Sigma-Aldrich; anti-Iba1 1:000, Wako) over-night at 4 °C. After washing with PBS, cells were incubated with the secondary antibody solution (goat anti-mouse alexa 488 or 546 1:500, Thermo Fisher) for 1 h at room temperature. Cells were washed and cover-slipped with Mowiol mounting medium (Sigma).

Protein extraction, quantification, and co-immunoprecipitation

Briefly, cells were collected with a cell scraper and centrifuged at $5000 \times g$ for 5 min. The pellet was re-suspended in RIPA buffer (50 mM Tris pH 7.5, 150 mM NaCl, 1% NP-40, 0.5% sodium deoxycholate, 0.1% SDS, 100 µg/ml PMSF) with phosphatase and protease inhibitor cocktail and stored at -80 °C. Protein levels were quantified with Pierce BCA Protein Assay Kit (Thermo Scientific). Immunoprecipitation experiments were performed by incubating 100 µg of protein with 2.5 µg of anti-RIPK1 (BD Transduction Laboratories) with rotation at 4 °C for 48 h. Then, 50 µl of protein G magnetic beads (Biorad) were added to each sample and incubated with rotation at 4 °C for 3 h. Following magnetic separation, beads were mixed with loading buffer and boiled at 90 °C for 5 min. After the elution step, samples were analyzed by western blotting as described.

Western blot

Proteins were fractionated by electrophoresis using 10% sodium dodecyl sulphate (SDS) polyacrylamide gels, electroblotted into PVDF membranes (Hybond-P, GE Healthcare), and blocked with 5% BCA in TBS. Membranes were then incubated with the different antibodies (anti-RIPK3, abcam ab16090; anti-pMLKL, abcam ab196436, and anti-HSP90, Santa Cruz sc7947) overnight at 4 °C, followed by incubation with a horseradish peroxidase-conjugated antibody. The immunoreactive bands were visualized using ECL (Invitrogen). The images were obtained and analyzed with ChemiDoc™ Touch Imaging System (Bio-Rad). For the immunoprecipitation experiments, detection of the IgG was done following the detection of pMLKL, using membranes that were stripped with a solution containing glycine 0,2 M, SDS 0,1%, Tween 20 0,1%, pH 2,2 (2×5 min incubation). After confirming that no bands could be visualized with ECL, membranes were blocked and incubated with a horseradish peroxidase-conjugated anti-mouse antibody. To analyze Aβ in human samples, protein samples were loaded onto 16% SDS-Tris-Tricine-PAGE and transferred to PVDF (Inmobilon-P, Millipore). After blocking, using 5% non-fat milk, the membranes were incubated overnight, at 4°C, with a mixture of N-terminus (clone 82E1) mouse

monoclonal (1:1000, IBL) and anti-Aβ (clone 6E10) mouse monoclonal (1:1000, Signet).

Dot blot

Dot blots were done as described previously [48]. Briefly, 1 µg of protein from the different S1 fractions was first treated with PureProteome Protein G Magnetics Beads (Millipore) for contaminant IgG depletion and then directly applied to dry nitrocellulose, in a final volume of 2 µl. Then, blots were blocked for 1 h and incubated overnight at 4 °C with anti-Amyloid fibrils (OC, 1:5000, Merck-Millipore), anti-prefibrillar oligomers (A11, 1:5000 InvitroGen), anti-pMLKL (1:1000, abcam) or anti-GAPDH (1:10,000, Cell Signalling) antibodies. Blots were visualized using the Pierce ECL 2 Western Blotting Substrate detection method (0.5 pg, lower limit sensitivity; Thermo Scientific). The images were obtained and analyzed with ChemiDoc™ Touch Imaging System (Bio-Rad). Data were always normalized by the specific signal observed in the Braak II group.

Statistical analyses

All experiments were analyzed in a blinded manner. To analyze differences in the data obtained from molecular and biochemical analyses of human samples, data were compared by Kruskal–Wallis test followed by Dunn post-hoc test. For the association analyses, linear regression was done using the Spearman rho-test. Data from all other studies were statistically analyzed using Student's *t*-test (for two groups comparisons) or one-way ANOVA (for more than two groups) followed by Bonferroni's post hoc test. For the association analyses, linear regression was done using Pearson correlation. Data are shown as mean \pm SEM. All analyses were performed using Graph Pad Prism 5.0 software. Differences were considered significant with $p < 0.05$.

Results

Aβ pathology is associated with necroptosis activation in human AD brains

Although evidence for necroptosis activation has been reported in the brain of patients with AD [26, 27], the possible association between Aβ pathology and necroptosis activation has not been established. Therefore, we studied the association between Aβ and phosphorylated MLKL at Ser358 (pMLKL), which defines the activation of a canonical necroptotic process, in brain samples derived from the hippocampus of sporadic AD, mild cognitive impairment (MCI), and control (CTRL) cases (clinical data of the subjects used in the histopathological study are listed in Table 1). The histopathological assessment revealed low pMLKL immunoreactivity in the hippocampus of control samples (Fig. 1A). Of the four

Table 1 Clinical data of cases investigated by histological analysis of postmortem brain tissue

Condition	Gender	Age	Braak stage	MMSE
CTRL	M	80	I	30
CTRL	F	>89	II	30
CTRL	N/A	N/A	II	N/A
CTRL	N/A	N/A	II	N/A
MCI	N/A	N/A	N/A	N/A
MCI	N/A	N/A	N/A	N/A
MCI	N/A	N/A	N/A	N/A
AD	M	65	III	N/A
AD	M	84	III	N/A
AD	F	79	IV	N/A
AD	M	75	VI	0
AD	F	76	VI	1
AD	F	>89	VI	6
AD	M	84	VI	N/A
AD	M	81	VI	N/A
AD	F	56	VI	N/A
AD	F	66	VI	N/A
AD	F	76	VI	N/A
AD	F	86	VI	N/A


CTRL control, MCI mild cognitive impairment, AD Alzheimer's disease, M male, F female, N/A not available, MMSE mini-mental state examination

controls analyzed, three were A β plaque-free and did not present immunoreactivity for pMLKL (see CTRL1). The control sample that exhibited A β pathology presented a low extent of A β deposits, and some of them were associated with pMLKL (CTRL2, see arrows). Brain samples from AD cases showed considerable levels of A β plaques, and most of them depicted some degree (moderate to high) of positive pMLKL immunoreactivity. As shown in the merged images, pMLKL signal was observed within the A β plaques, and also surrounding the deposits. Remarkably, these findings were also observed in brain samples derived from patients with MCI, which is considered a precursor for AD [49]. The association between A β plaques and pMLKL was additionally studied using the specific probe for amyloid plaques, Amylo-Glo (Fig. 1B). The pMLKL immunoreactivity was validated using DAB staining (Additional file 1: Fig. S1). These data demonstrate a spatial correlation between A β pathology and the necroptosis effector in AD, which is consistent with the fact that A β plaques are often surrounded by dystrophic neurites and associated with synaptic loss [50].

Microglial cells are commonly observed around amyloid deposits [11] (Fig. 1C). Since previous studies in other degenerative conditions have shown necroptosis activation in microglial cells [51, 52], we then tested whether pMLKL immunoreactivity localizes to neurons

or microglia in AD. To this aim, we conducted colocalization analyses between pMLKL and MAP2 or Iba1, respectively, using AD brain tissue (Fig. 1D, E). The results revealed that most of the pMLKL immunoreactivity colocalized with MAP2. This result suggests that necroptosis activation occurs mainly in neurons of the AD brain.

Several lines of investigation indicate that A β oligomers (A β o) contribute importantly to neurodegeneration in AD [2, 3]. Indeed, previous studies have demonstrated that soluble oligomeric A β correlates better with cognitive deficits compared to plaques, which can actually sequester oligomers, acting as reservoirs from which oligomers can diffuse, reaching cellular targets in their vicinity [4–7]. Thus, we next moved forward to determine the relationship between the burden of A β o and necroptosis activation in AD brains. To this end, we evaluated the association between *Ripk1*, *Ripk3*, and *Mkl1* mRNA levels and the levels of soluble A β o in postmortem brain samples of AD patients at different stages of the disease (clinical data of the subjects used in this study are summarized in Table 2). Samples were divided into three groups: samples derived from Braak II, non-demented individuals, which did not accumulate A β in the hippocampus; samples from Braak III-IV patients, which presented A β accumulation; and samples from Braak V-VI patients, which presented significantly higher levels of A β compared to the latter group (Fig. 2A, B). In agreement with previously reported data [26], we observed that the expression levels of *Ripk1* and *Mkl1*, but not those of *Ripk3*, are elevated in the brains of individuals with higher Braak stages (Fig. 2C). The levels of oligomeric A β aggregates were assessed by dot blot using OC and A11 antibodies to detect soluble fibrillar and prefibrillar A β o, respectively (Fig. 2D). The dynamic range and the coefficient of variation of the dot-blot assays are shown in Additional file 2: Fig. S2. Quantitative analysis revealed that the extent of both types of structures was significantly higher in the brains of AD patients compared to healthy controls (Fig. 2E). We found a significant and positive correlation between A β o load and the levels of *Ripk1* but not *Ripk3* or *MLKL* mRNA (Fig. 2F, G). Since MLKL activation is post-translationally regulated by phosphorylation, we then measured the association of pMLKL levels with the load of A β o in AD brain samples. To achieve this, pMLKL levels were determined by dot blot (Fig. 2H). Quantitative measurement showed significantly increased pMLKL levels in AD subjects compared to controls (Fig. 2I). Notably, the levels of pMLKL correlate significantly and positively with the burden of A β o (Fig. 2J, K). These results suggest a link between oligomeric A β structures and activation of the necroptotic machinery.



Together, these data provide the first evidence that Aβ pathology is associated with necroptosis activation in AD.

Targeting necroptosis ameliorates neurodegeneration and memory impairment induced by Aβ in mice

Given the role of Aβ aggregates as major pathological mediators of both familial and sporadic AD and considering that Aβ are generally regarded as the most pathogenic form of Aβ, animal models based on intracerebral Aβ inoculation have been established as useful tools to

Table 2 Demographics of the individuals investigated by molecular and biochemical analyses of postmortem brain tissue

Braak stage	Gender (male, %)	Gender (female, %)	Age (Years, mean age \pm SD)	Postmortem delay (mean hours \pm SD)
II (n = 10)	61.54	38.46	78 \pm 8.5	7 \pm 4
III–IV (n = 9)	44.44	55.56	80 \pm 11	6 \pm 5
V–VI (n = 10)	38.89	61.11	79 \pm 10	8 \pm 4

study AD pathogenesis [3]. To assess whether A β can induce activation of the necroptosis pathway, we conducted an acute intracerebroventricular injection of synthetic A β to wild-type mice. As expected, A β injection triggered significant microgliosis compared to control mice (Fig. 3A, B). Moreover, A β treated mice underwent neurodegeneration in the hippocampus, as demonstrated by an increase in non-phosphorylated neurofilaments (Fig. 3C, D). Staining for pMLKL demonstrated significant necroptosis activation in A β injected mice compared to vehicle-injected ones (Fig. 3E, F). To validate the phospho-specific MLKL antibody, we compared the immunoreactivity of brain sections from wild-type and MLKL knock-out mice, treated with A β (Additional file 3: Fig. S3 A, B). In addition, to assess the cell specificity of the pMLKL immunostaining, we performed double labeling using antibodies to pMLKL and MAP2 or CD11b to observe neurons and microglia, respectively. As shown in Additional file 3: Fig. S3 C, D, most of the pMLKL signal observed in the hippocampus was associated with neurons. Notably, pMLKL expression positively correlated with neurodegeneration determined by Fluoro-Jade C staining (Additional file 4: Fig. S4) and negatively correlated with cognitive performance evaluated using the Morris water maze (MWM) behavioral test (Fig. 3G, H).

To evaluate the potential protective effect of MLKL inhibition against A β -induced neurodegeneration and memory alterations, we used genetically modified mice lacking *Mkl1*. To test the effects of targeting MLKL in the

progression of AD pathogenesis, we evaluated neurodegeneration in the brain of these mice by histopathological assessment using Fluoro-Jade C staining. As shown in Fig. 3I, J, A β administration elicited significant neurodegeneration compared to control mice, evidenced by increased Fluoro-Jade C stained area in the hippocampus. Notably, ablation of *Mkl1* expression decreased the levels of A β -related neurodegeneration. To determine whether targeting MLKL influences the microgliosis observed following A β treatments, immunostaining was done using Iba1. Although similar levels of microglial activation were observed following A β infusion in transgenic and wild-type mice, no statistically significant differences were found between vehicle and A β -treated mice in the knock-out group (Additional file 5: Fig. S5). In order to study the possible consequences of eliminating MLKL on the behavioral impairment observed in this AD model, we evaluated the memory capacity of mice using the MWM test. In agreement with previous reports [53, 54] A β -injected mice exhibited altered memory performance in the MWM test, as this group of animals spent less time in the target quadrant compared to control mice (Fig. 3K). In sharp contrast, *Mkl1* deletion significantly inhibited cognitive alterations observed in this AD model as the time spent in the target quadrant by MLKL^{-/-} mice treated with A β was significantly longer compared to MLKL^{+/+} mice injected with the oligomers (Fig. 3K). The protection provided by *Mkl1* deletion was also reflected when measuring the distance that mice traveled in the target quadrant during the test (Fig. 3L). The time that mice spent in the target quadrant during the test is graphically represented in the heat map (Fig. 3M). Importantly, swim speed and total traveled distance were not modified by A β treatment (Fig. 3N, O), suggesting this model does not cause locomotor or visual impairment.

To increase the translational potential of these findings, we next investigated the possible protective effect of targeting necroptosis by pharmacological inhibition in this AD model. To this end, we treated mice with the RIPK3 inhibitor GSK'872. Histopathological exploration of brain

(See figure on next page.)

Fig. 2 Activation of RIPK1 and MLKL correlate with soluble A β oligomer burden in AD brains. **A** Western blot analysis of total protein extracts from hippocampal samples using a mix of 6E10 and 82E1 antibodies was performed to determine the A β content. **B** The specific band was quantified by densitometric analysis, and represented in the graphs. **C** The graph represents the expression levels of *Ripk1*, *Ripk3*, and *Mkl1* in human brain samples from patients at different AD stages (Braak stage II (n = 10), III–IV (n = 9), V–VI (n = 10)). **D** Dot blots of soluble proteins extracted from postmortem brains of individuals with Braak II (n = 4), Braak III–IV (n = 5), and Braak V–VI (n = 8), probed with the indicated antibodies. **E** Blots were quantified by densitometric analysis and expressed relative to GAPDH protein, represented in the graph. **F, G** Association analyses of oligomeric A β load and mRNA levels were done by linear regression. **H** Dot blots of protein extracts from postmortem brains of individuals with Braak II (n = 4), Braak III–IV (n = 5), and Braak V–VI (n = 8), probed with the indicated antibodies. **I** Blots were quantified by densitometric analysis and expressed relative to GAPDH protein, represented in the graph. **J, K** Association analyses of oligomeric A β load and pMLKL levels were done by linear regression. Data are presented as mean \pm S.E.M. Data in **B, C, E**, and **I** were analyzed by Kruskal–Wallis test followed by Dunn post-hoc test. Data in **F, G, J** and **K** were analyzed by linear regression using Spearman rho-test. * p < 0.05; ** p < 0.01; *** p < 0.001; **** p < 0.0001




Fig. 2 (See legend on previous page.)

pMLKL levels revealed that intraperitoneal administration of GSK'872 significantly decreased necroptosis activation induced by A β (Fig. 4A, B). Moreover, evaluation of neurodegeneration using Fluoro-Jade C showed that GSK'872 significantly attenuated A β -related neurodegeneration to levels comparable to those detected in control mice (Fig. 4C,D). To determine whether targeting RIPK3 influences the microgliosis triggered by A β treatment, immunostaining was done using Iba1. As shown in Additional file 6: Fig. S6, *Ripk3* ablation did not alter the microglial activation induced by A β , suggesting neuronal necroptosis engagement occurs downstream microglial activation. To evaluate the functional consequences of inhibiting RIPK3, cognitive performance was tested using the MWM. As expected, and similarly to the result described above, mice subjected to A β infusion underwent significant memory impairment. Remarkably, the results of the test revealed that treatment with GSK'872 provided significant protection against A β -induced memory loss, as mice challenged with A β and treated with the drug, behaved similarly to control animals (Fig. 4E, F). The time that mice spent in the target quadrant during the test is graphically represented in the heat map (Fig. 4G). The results shown in Fig. 4H, I suggest that A β treatment did not cause locomotor or visual damage in these mice. Noteworthy, a negative correlation between neurodegeneration assessed by Fluoro-Jade C and cognitive performance was observed in these mice (Fig. 4).

Together, these findings suggest that neurodegeneration induced by A β is mediated by the necroptotic machinery and that targeting necroptosis provides significant neuroprotection and results in cognitive improvement in experimental AD.

Neurodegeneration triggered by A β -stimulated microglia occurs through necroptosis activation

As indicated above, AD brains are characterized by the presence of reactive microglial cells associated with A β aggregates [10–12, 14]. This can be observed in human

AD as well as in murine AD models (see Fig. 1C). Moreover, a role for microglia as mediators of A β -induced neurodegeneration has been extensively demonstrated [9]. Interestingly, it has been reported that rather than fibrillar amyloid, soluble oligomeric A β structures can more effectively potentiate a proinflammatory microglial response leading to neurodegeneration [55–59]. We then sought to explore whether microglial activation triggered by A β can elicit activation of the necroptotic pathway in neurons. We used synthetic A β , and their cytotoxic properties were first assessed in primary neurons. Oligomers triggered progressive neurodegeneration, evidenced by beading and retraction of neuronal processes, that was extensive after 72 h of incubation (Additional file 7: Fig. S7A–C). Notably, the addition of microglial cells to neuronal cultures significantly enhanced A β neurotoxicity (Additional file 7: Fig. S7D, E).

To test the possible involvement of microglia-secreted factors to the induction of necroptosis-mediated neuronal death, hippocampal neurons were treated with conditioned medium from primary rat microglial cells stimulated with A β (CM^{A β} , Fig. 5A). Treatment with CM^{A β} induced broad neuronal degeneration and the neurotoxic effects were dose-dependent (Fig. 5B, C). We then performed western blot analyses to determine the activation of necroptosis by analyzing the levels of RIPK3 and pMLKL in neurons upon treatment with CM^{A β} . Densitometric analysis showed increased levels of RIPK3 (Fig. 5D) and pMLKL (Fig. 5E) in neurons exposed to CM^{A β} compared with neurons treated with CM^{ctrl} (full blots are shown in Additional file 8: Fig. S8A, B). The physical interaction between RIPK1, RIPK3, and MLKL within the necrosome complex leads to MLKL phosphorylation, resulting in its oligomerization and execution of necroptosis [33]. In order to further determine if CM^{A β} triggers the formation of the necrosome in neurons, co-immunoprecipitation experiments were performed followed by western blot analysis. These results revealed a significantly higher RIPK1-pMLKL interaction in neurons treated with CM^{A β} compared with neurons treated

(See figure on next page.)

Fig. 3 MLKL ablation ameliorates neurodegeneration and memory loss induced by A β in mice. **A, B** Representative micrographs and quantitative analysis of brain sections from wild-type mice subjected to intracerebroventricular infusion of A β or vehicle at 18 days post-injection, immunostained with an Iba1 specific antibody (scale bar, 150 μ m). **C, D** Representative images and quantitative analysis of brain sections immunolabeled with an anti-non-phosphorylated neurofilament antibody at 18 days post A β injection (scale bar, 150 μ m). **E, F** Representative images and quantitative analysis of brain sections immunolabeled with an anti-pMLKL antibody at 18 days post A β injection (scale bar, 150 μ m). Association analyses of **(G)** pMLKL and Fluoro-Jade C stained area and **(H)** pMLKL stained area and cognitive performance, were done by linear regression. **I, J** Representative micrographs and quantitative measurement of brain sections from mice, treated as indicated in the images at 18 days post Vehicle or A β injection, stained with Fluoro-Jade C (scale bar, 150 μ m). n = 13 per group, 4 sections per brain, one image of the hippocampus per section was analyzed. **K, L, N, O** The graphs represent quantitative analyzes of the indicated parameters, obtained from the MWM test. **M** The heat map shows a graphic representation of the time that mice, treated as indicated in the image, spent in the target quadrant (TQ) during the MWM test. Data are presented as mean \pm S.E.M. The data in **B, D, and F** were analyzed by Student's *t*-test. Data in **G** and **H** were analyzed by linear regression using Pearson correlation. Data in **J–L, N, and O** were analyzed by one-way ANOVA followed by Bonferroni post-test. **p* < 0.05; ***p* < 0.01; ****p* < 0.001




Fig. 3 (See legend on previous page.)




Fig. 4 RIPK3 inhibition reduces neurodegeneration and memory loss induced by Aβ in mice. **A, B** Representative images and quantitative analysis of brain sections from wild-type mice at 18 days post-injection, treated as indicated in the images and immunolabeled with an anti-pMLKL antibody (scale bar, 150 µm). **C, D** Representative micrographs and quantitative measurement of brain sections from wild-type mice at 18 days post-injection, treated as indicated in the images, and stained with Fluoro-Jade C (scale bar, 150 µm). n = 10 per group, 4 sections per brain, one image of the hippocampus per section was analyzed. **E, F, H, I** Graphs represent quantitative analyses of the indicated parameters, obtained from the MWM test. **G** The heat map shows a graphic representation of the time that mice, treated as indicated in the image, spent in the target quadrant (TQ) during the MWM test. **J** Association analysis of Fluoro-Jade C stained area and cognitive performance was done by linear regression. Data are presented as mean ± S.E.M. Data in **B, D-F, H, and I** were analyzed by one-way ANOVA followed by Bonferroni post-test. Data in **J** were analyzed by linear regression using Pearson correlation. *p < 0.05; **p < 0.01; ***p < 0.001; ****p < 0.0001

with CM^{ctrl}, indicating that CM^{Aβ} was able to enhance the formation of the necrosome complex (Fig. 5E, full blots are shown in Additional file 8: Fig. S8C). Finally, to define the involvement of necroptosis in neuronal death in this setting, we treated cells with the RIPK1 antagonist nec-1 s or with the RIPK3 inhibitor GSK'872, prior to

the addition of CM^{Aβ} to the cultures. Importantly, neurons treated with CM^{Aβ} together with nec-1 s or GSK'872 showed significant protection from neurodegeneration compared with neurons treated with CM^{Aβ} alone (Fig. 5G–J).

Together, these data suggest that in response to stimulation with A β , neurotoxic factors released by activated microglia drive the formation of the necrosome and the initiation of the necroptosis signaling cascade in neurons.

TNF- α released by A β -stimulated microglia contributes to necroptosis activation

The secretion of TNF- α by A β -stimulated microglial cells has been previously described [47]. It has been demonstrated that activation of TNFR1 by TNF- α can trigger cell death via activation of the necroptotic pathway [31, 60]. Hence, we next investigated the possible contribution of TNF- α to A β -induced necroptosis. To this end, CM^{A β} was incubated with the anti-TNF- α antibody Infliximab to neutralize the protein and was then used to treat neuronal cultures. Remarkably, Infliximab provided significant protection to neurons compared to neurons treated with control CM^{A β} , which showed a high proportion of neuritic beading (Fig. 5K, L). To test whether the neuroprotection provided by Infliximab was due to the inhibition of necroptosis activation, we monitored the levels of pMLKL. Western blot analysis of neurons treated with CM^{A β} together with Infliximab showed decreased pMLKL levels compared to neurons treated with CM^{A β} alone (Fig. 5M, N; full blots are shown in Additional file 8: Fig. S8D). To assess the sufficiency of TNF- α to trigger necroptosis-mediated neurodegeneration, we treated neuronal cultures with recombinant TNF- α . A concentration of 40 ng/ml was used, as it has been previously shown that when combined with apoptosis inhibitors, it is able to trigger necroptosis in vitro [61, 62]. As shown in Additional file 9: Fig. S9A, B, neither treatment with TNF- α alone nor in combination with CM^{ctrl} induced neurodegeneration.

Together, these data suggest that, although TNF- α is necessary, it acts in conjunction with other molecules released by A β -activated microglia to induce necroptosis-mediated neurodegeneration.

Discussion

Several studies have demonstrated the involvement of necroptosis in the pathogenesis of neurodegenerative diseases, including PD [41], MS [40], ALS [39], Huntington disease [63], and ischemic neuronal injury [64]. It has been recently shown that necroptosis is activated in human AD brains [26, 27]. However, the mechanism by which this pathway is activated in AD is currently unknown. Here, we define a role for A β pathology in necroptosis activation in AD and demonstrate that microglial cells mediate A β -induced necroptosis activation in neurons.

The abnormal deposition of A β results from an imbalance between its production and clearance, constituting a crucial and probably causative event in the development of AD [2]. Nevertheless, A β appears to interact with multiple receptors, different types of cells and affect many biological processes and still, the exact mechanism by which neurons die in AD is not clear. Our results demonstrate a spatial correlation between A β pathology and the necroptosis effector pMLKL in AD brains and a correlation between A β burden and the expression of key necroptosis drivers, suggesting a possible link between A β aggregation and the activation of the necroptosis signaling pathway. Whereas A β plaques have long been a signature of AD histopathology, increasing studies over the past decade have demonstrated that soluble oligomeric A β constitutes the most neurotoxic type of A β aggregate [3, 5, 65, 66]. Accordingly, Brody et al., hypothesized that plaques may serve as a reservoir for toxic oligomers,

(See figure on next page.)

Fig. 5 Neurodegeneration induced by conditioned medium from A β -stimulated microglia occurs via necroptosis activation. **A** Schematic representation of the preparation of the conditioned medium. **B, C** Representative images and quantitative analysis from neurons treated for 72 h with culture medium collected from a culture dish where A β were previously immobilized using nitrocellulose (control), neurons treated with conditioned medium from untreated microglia (CM^{ctrl}), and neurons treated with 10% and 50% conditioned medium from A β -stimulated microglia (CM^{A β}); immunostaining was performed with anti-acetylated tubulin antibody (scale bar, 50 μ m). Western blot analyses of protein extracts from neurons treated for 6 h with CM were performed to determine the expression levels of RIPK3 (**D**) and pMLKL (**E**). Specific bands were quantified by densitometric analysis and expressed relative to HSP90 protein, represented in the graphs. **F** Protein extracts from neurons treated for 6 h with CM were immunoprecipitated with an antibody against RIPK1 and probed for pMLKL. The graph represents the densitometric analysis of the western blots. **G, H** Representative micrographs and quantitative measurements of neurons treated for 72 h with CM^{ctrl}, CM^{A β} and with Nec-1 s (30 μ M) prior to the addition of CM^{A β} , immunolabeled with anti-acetylated tubulin antibody (scale bar, 200 μ m). **I, J** Representative images and quantitative analysis of neurons treated for 72 h with CM^{ctrl}, CM^{A β} and with GSK'872 (1 μ M) prior to the addition of CM^{A β} , immunolabeled with anti-acetylated tubulin antibody (scale bar, 200 μ m). **K, L** Representative images and quantitative analysis of neurons treated with CM^{ctrl}, CM^{A β} , and with CM^{A β} that was previously mixed with Infliximab (0.01 μ g/ μ l, Infliximab/CM^{A β}), immunostained with anti-acetylated tubulin antibody (scale bar, 200 μ m). **M, N** Western blot analysis of protein extracts from neurons treated for 6 h under the indicated conditions was performed to determine the expression levels of pMLKL. Bands were quantified by densitometric analysis and expressed relative to HSP90 protein, represented in the graph. Each experiment was performed at least three independent times, with three replicates per condition each time. Data are presented as mean \pm S.E.M. Data in **C, H, J, L**, and **N** were analyzed by one-way ANOVA followed by Bonferroni post-test. Data in **D-F** were analyzed by Student's *t*-test. **p* < 0.05; ***p* < 0.01; ****p* < 0.001, *****p* < 0.0001




Fig. 5 (See legend on previous page.)

thereby capturing them in the extracellular space. It was suggested that over time, plaques get saturated and therefore oligomers can diffuse and exert their toxicity to the surrounding cells [4]. In agreement with this hypothesis, Koffie and colleagues demonstrated that oligomeric A β associated with plaques colocalizes with postsynaptic densities and is associated with spine collapse and synapse loss in postmortem AD samples [7]. Consistent with this evidence, our data point to oligomeric A β as a driver of necroptosis activation in AD, leading to neurodegeneration and memory loss. Although a previous report found no correlation between A β and necroptosis markers in AD brains, in this study the researchers considered A β plaque load and not oligomers for the analysis [26]. Many studies have demonstrated that the aggregation process of A β follows a seeding-nucleation mechanism that involves the formation of multiple types of intermediates including oligomers and fibrils [67, 68]. Additional studies are necessary to test the role of insoluble fibrillar A β in the activation of necroptosis. In addition, previous studies have shown colocalization between pTau and necroptosis markers in AD brains [26, 27], which is not surprising considering the mounting evidence proving the synergistic contribution of both misfolded A β and tau proteins to neurodegeneration [61–64]. Further studies would be required in order to test a potential interplay between both proteins for necroptosis activation.

In agreement with the study by Caccamo et al. [26], our results suggest that inhibition of the necroptotic pathway prevents neurodegeneration in AD. Previous data from our group provided evidence for a functional role of necroptosis in PD, as necroptosis inhibition improved motor performance in a PD model [41]. Importantly, the present study is the first to unveil the functional implications of targeting necroptosis in an experimental AD model. As necroptosis is a druggable pathogenic mechanism, our findings suggest that necroptosis inhibition has an important translational potential for AD.

Several pathological processes may trigger activation of the necroptotic cascade in AD, including inflammation and oxidative stress. Our data suggest that upon interaction with A β , microglial cells secrete cytotoxic molecules that induce neurodegeneration via necroptosis activation. Among the secretory mediators released by microglia, TNF- α elicits necroptosis in vitro and in vivo [31, 60]. Accordingly, previous studies in other neurodegenerative conditions, including MS and ALS, demonstrated that sustained inflammation is associated with TNF- α -mediated necroptosis [73, 74]. In line with this evidence, our work suggests that TNF- α released from microglia following its activation by A β contributes to necroptosis-mediated neurodegeneration. A β can stimulate signaling responses in microglial cells that

induce the release of a number of molecules that can contribute to neurodegeneration. Interestingly, there is evidence showing that glutamate, which has been previously demonstrated to be released by activated microglia [75, 76], can also induce necroptosis activation [77, 78]. Similarly, interferon- γ , which is produced by reactive microglia [79], has also been shown to participate in the activation of the necroptotic signaling pathway [29]. Additional studies are needed to unveil the microglial protein expression profile in response to A β stimulation and define the exact molecules involved in the activation of the necroptotic pathway.

Overall, in the present study, we provide new insights into the mechanism by which necroptosis is activated in AD, adding a new layer of complexity involving the crosstalk between microglia and neurons. Our results may contribute to understanding the mechanisms that lead to neurodegeneration in AD and support the view of necroptosis as a potential therapeutic target for disease intervention.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s40478-022-01332-9>.

Additional file 1: Fig. S1. Validation of the pMLKL immunoreactivity by DAB staining. Representative micrographs of hippocampal brain areas of AD patients ($n = 3$) immunostained with anti-pMLKL antibody. Orange arrows indicate pMLKL-positive neurons, and green arrows show pMLKL-positive microglia. Magnification, 20X.

Additional file 2: Fig. S2. Coefficient of variation of the dot-blot assays. The coefficient of variation (CV) was calculated using triplicates of Braak V-VI samples. A CV of 11.63% and 15.06% was determined for OC and A11, respectively.

Additional file 3: Fig. S3. Validation of the pMLKL antibody and assessment of cell specificity. Representative micrographs of (A) the dentate gyrus (DG) and (B) CA1 brain regions from wild-type and MLKL knockout mice subjected to intracerebral injection of A β , immunostained with anti-pMLKL antibody (scale bar, DG: 100 μ m, CA1: 150 μ m). (C, D) Representative images of the hilus of wild-type mice treated with A β , labeled with the indicated antibodies (scale bar, 50 μ m and 25 μ m for magnifications).

Additional file 4: Fig. S4. Intracerebral administration of A β elicits neurodegeneration in wild-type mice. Representative micrographs of brain sections from wild-type mice treated as indicated in the images, stained with Fluoro-Jade C (scale bar, 150 μ m).

Additional file 5: Fig. S5. MLKL inhibition attenuates A β -induced microgliosis in mice. (A, B) Representative micrographs and quantitative analysis of brain sections from wild-type and MLK1 knockout mice treated as indicated in the images, immunostained with anti-Iba1 antibody (scale bar, 200 μ m). The data are presented as mean \pm S.E.M. and were analyzed by one-way ANOVA followed by Bonferroni post-test. * $p < 0.05$.

Additional file 6: Fig. S6. RIPK3 inhibition does not alter A β -induced microgliosis in wild-type mice. (A, B) Representative micrographs and quantitative analysis of brain sections from wild-type mice treated as indicated in the images, immunostained with anti-Iba1 antibody (scale bar, 200 μ m). The data are presented as mean \pm S.E.M. and were analyzed by one-way ANOVA followed by Bonferroni post-test. * $p < 0.05$; ** $p < 0.01$.

Additional file 7: Fig. S7. A β neurotoxicity is worsened when microglial cells are present. **(A)** To characterize A β , a negative stain with 1% uranyl acetate was performed followed by electron microscopy analysis. Arrows indicate small oligomeric structures (scale bar, 200 nm). **(B, C)** Representative images and quantitative analysis of neurons treated as indicated, immunostained with anti-acetylated tubulin antibody (scale bar, 100 μ m). **(D, E)** Representative micrographs and quantitative measurement of neurons and neuron/microglia cocultures treated as indicated, immunolabeled with anti-acetylated tubulin and anti-Iba1 antibodies as specified in the images (scale bar, 100 μ m in the middle panel; 50 μ m in the right panel). Each experiment was performed at least three independent times, with three replicates per condition each time. The data are presented as mean \pm S.E.M. and were analyzed by one-way ANOVA followed by Bonferroni post-test. * p < 0.05; ** p < 0.01; **** p < 0.0001.

Additional file 8: Fig. S8. Full blots of pMLKL and RIPK3 western blots. **(A)** The membrane was first probed with an anti-hsp90 antibody and then incubated with anti-RIPK3. **(B)** The membrane was cut, and each piece was incubated with the indicated antibody. Left panel: colorimetric image showing protein ladder. **(C)** The membrane was cut and incubated with an anti-pMLKL antibody. Following stripping, and after confirming that no bands could be visualized with ECL, membranes were blocked, and detection of the IgG was done. Left panel: colorimetric image showing protein ladder. **(D)** The membrane was cut, and each piece was incubated with the indicated antibody. Left panel: colorimetric image showing protein ladder.

Additional file 9: Fig. S9. TNF- α alone is not sufficient to trigger neuronal necroptosis. **(G, H)** Representative micrographs and quantitative measurements of neurons treated for 72 h with CMctrl, CMA β , TNF- α , and with CMctrl/TNF- α , immunolabeled with anti-acetylated tubulin antibody (scale bar, 200 μ m). The data are presented as mean \pm S.E.M. and were analyzed by one-way ANOVA followed by Bonferroni post-test. * p < 0.05; ** p < 0.01.

Acknowledgements

We thank Dra. Claudia Duran for suggestions and Dr. Claudio Hetz for critical review of the manuscript.

Authors' contributions

N.S. designed research; N.S., I.M.-G., N.G., G.Q., L.V., S.J., and M.E. performed research; N.S., I.M.-G. and J.V. analyzed data; C.S., A.G., I.M.-G. and J.V. contributed new reagents or analytic tools; N.S. wrote the paper; F.C. supervised research and edited the paper. All authors read and approved the final manuscript.

Funding

This study was funded by grants from FONDECYT No. 3180341 (to NS); FONDECYT No. 1190518 and FONDAP program 15150012 (to FC); Instituto de Salud Carlos III (ISCIII) of Spain, co-financed by FEDER funds from European Union, through grants PI18/01556 (to JV), PI21/00914 (to JV), PI18/01557 (to AG) and PI21/00915 (to AG), CIBERNED (to JV and AG) and by Junta de Andalucía Consejería de Economía y Conocimiento co-financed by FEDER 2014-2020 through grants US-1262734 (JV), UMA18-FEDERJA-211 (AG), UMA20-FEDERJA-104 (I-MG) and P18-RT-2233 (to AG). Ministry of Science and Innovation (MICIN) State Research Agency grants PID2019-107090RA-I00 and Ramon y Cajal Program RYC-2017-21879 (to IMG) and grants from NIH R01AG059321 and R01AG061069 (to CS).

Declarations

Competing interests

The authors report no competing interests in relation to this manuscript.

Author details

¹Center for Integrative Biology, Faculty of Sciences, Universidad Mayor, Santiago, Chile. ²FONDAP Geroscience Center for Brain Health and Metabolism, Santiago, Chile. ³Department of Cell Biology, Facultad de Ciencias, Universidad de Malaga, IBIMA, Malaga, Spain. ⁴Networking Research Center on Neurodegenerative Diseases (CIBERNED), Madrid, Spain. ⁵Mitchell Center for Alzheimer's Disease, Department of Neurology, McGovern Medical School,

The University of Texas Health Science Center at Houston, Houston, TX, USA. ⁶Dpto. Bioquímica Y Biología Molecular, Facultad de Farmacia, Universidad de Sevilla, 41012 Sevilla, Spain. ⁷Instituto de Biomedicina de Sevilla (IBiS)-Hospital Universitario Virgen del Rocío/CSIC, Universidad de Sevilla, 41013 Sevilla, Spain. ⁸Buck Institute for Research on Ageing, Novato, CA, USA. ⁹Present Address: Laboratory of Neurodegenerative Diseases, Faculty of Sciences, Universidad Mayor, Temuco, Chile.

Received: 17 February 2022 Accepted: 18 February 2022

Published online: 09 March 2022

References

1. Alzheimer's Association, 2020 Alzheimer's disease facts and figures. *Alzheimer's Dementia* **16**, 391–460 (2020).
2. Selkoe DJ, Hardy J (2016) The amyloid hypothesis of Alzheimer's disease at 25 years. *EMBO Mol Med* **8**:595–608
3. Cline EN, Bicca MA, Viola KL, Klein WL (2018) The amyloid- β oligomer hypothesis: Beginning of the third decade. *J Alzheimer's Dis* **64**:S567–S610
4. Brody DL, Jiang H, Wildburger N, Esparza TJ (2017) Non-canonical soluble amyloid-beta aggregates and plaque buffering: controversies and future directions for target discovery in Alzheimer's disease. *Alzheimer's Res Therapy* **9**:1–13
5. Esparza TJ et al (2013) Amyloid-beta oligomerization in Alzheimer dementia versus high-pathology controls. *Ann Neurol* **73**:104–119
6. Hong S et al (2014) Soluble A β oligomers are rapidly sequestered from brain ISF in vivo and bind GM1 ganglioside on cellular membranes. *Neuron* **82**:308–319
7. Koffie RM et al (2009) Oligomeric amyloid β associates with postsynaptic densities and correlates with excitatory synapse loss near senile plaques. *Proc Natl Acad Sci USA* **106**:4012–4017
8. Wang J, Gu BJ, Masters CL, Wang YJ (2017) A systemic view of Alzheimer disease—insights from amyloid- β metabolism beyond the brain. *Nat Rev Neurol* **13**:612–623
9. Heneka MT et al (2015) Neuroinflammation in Alzheimer's disease. *Lancet Neurol* **14**:388–405
10. Keren-Shaul H et al (2017) A unique microglia type associated with restricting development of Alzheimer's disease. *Cell* **169**:1276–1290.e17
11. Simard AR, Soulet D, Gowing G, Julien JP, Rivest S (2006) Bone marrow-derived microglia play a critical role in restricting senile plaque formation in Alzheimer's disease. *Neuron* **49**:489–502
12. Mosher KI, Wyss-Coray T (2014) Microglial dysfunction in brain aging and Alzheimer's disease. *Biochem Pharmacol* **88**:594–604
13. Lee S et al (2010) CX3CR1 deficiency alters microglial activation and reduces beta-amyloid deposition in two Alzheimer's disease mouse models. *Am J Pathol* **177**:2549–2562
14. McGeer PL, Itagaki S, Tago H, McGeer EG (1988) Occurrence of HLA-DR reactive microglia in Alzheimer's disease. *Ann N Y Acad Sci* **540**:319–323
15. Combs CK, Karlo JC, Kao SC, Landreth GE (2001) beta-Amyloid stimulation of microglia and monocytes results in TNF α -dependent expression of inducible nitric oxide synthase and neuronal apoptosis. *J Neurosci Off J Soc Neurosci* **21**:1179–1188
16. El Khoury JB et al (2003) CD36 mediates the innate host response to β -amyloid. *J Exp Med* **197**:1657–1666
17. Miao J (2005) Reducing cerebral microvascular amyloid- protein deposition diminishes regional neuroinflammation in vasculotropic mutant amyloid precursor protein transgenic mice. *J Neurosci* **25**:6271–6277
18. Patel NS et al (2005) Inflammatory cytokine levels correlate with amyloid load in transgenic mouse models of Alzheimer's disease. *J Neuroinflammation* **2**:9
19. Fan R et al (2007) Minocycline reduces microglial activation and improves behavioral deficits in a transgenic model of cerebral microvascular amyloid. *J Neurosci Off J Soc Neurosci* **27**:3057–3063
20. R. Guerreiro, et al., TREM-2 variants in AD. **368**, 117–127 (2013).
21. Jonsson T et al (2013) Variant of TREM2 associated with the risk of Alzheimer's disease. *N Engl J Med* **368**:107–116
22. Takahashi K, Rochford CDP, Neumann H (2005) Clearance of apoptotic neurons without inflammation by microglial triggering receptor expressed on myeloid cells-2. *J Exp Med* **201**:647–657

23. Ma J, Jiang T, Tan L, Yu JT (2015) TYROBP in Alzheimer's Disease. *Mol Neurobiol* 51:820–826
24. Wang Y et al (2015) TREM2 lipid sensing sustains the microglial response in an Alzheimer's disease model. *Cell* 160:1061–1071
25. Salvadores N, Court FA (2020) The necroptosis pathway and its role in age-related neurodegenerative diseases: will it open up new therapeutic avenues in the next decade? *Expert Opin Ther Targets* 24:679–693
26. Caccamo A et al (2017) Necroptosis activation in Alzheimer's disease. *Nat Neurosci*. <https://doi.org/10.1038/nn.4608>
27. Koper MJ et al (2019) Necrosome complex detected in granulovacuolar degeneration is associated with neuronal loss in Alzheimer's disease. *Acta Neuropathologica*. <https://doi.org/10.1007/s00401-019-02103-y>
28. Wang H et al. (2017) PELI1 functions as a dual modulator of necroptosis and apoptosis by regulating ubiquitination of RIPK1 and mRNA levels of c-FLIP. In: Proceedings of the National Academy of Sciences of the United States of America, <https://doi.org/10.1073/pnas.1715742114>
29. Thapa RJ et al (2013) Interferon-induced RIP1/RIP3-mediated necrosis requires PKR and is licensed by FADD and caspases. *Proc Natl Acad Sci* 110:E3109–E3118
30. Kaiser WJ et al (2013) Toll-like receptor 3-mediated necrosis via TRIF, RIP3, and MLKL. *J Biol Chem* 288:31268–31279
31. Remijsen Q et al (2014) Depletion of RIPK3 or MLKL blocks TNF-driven necroptosis and switches towards a delayed RIPK1 kinase-dependent apoptosis. *Cell Death Dis* 5:e1004
32. Kearney CJ et al (2015) Necroptosis suppresses inflammation via termination of TNF- or LPS-induced cytokine and chemokine production. *Cell Death Differ* 22:1313–1327
33. Grootjans S, Berghe TV, Vandenabeele P (2017) Initiation and execution mechanisms of necroptosis: an overview. *Cell Death Differ* 24:1184–1195
34. Degterev A et al (2008) Identification of RIP1 kinase as a specific cellular target of necrostatins. *Nat Chem Biol* 4:313–321
35. He S et al (2009) Receptor interacting protein kinase-3 determines cellular necrotic response to TNF- α . *Cell* 137:1100–1111
36. Cho YS et al (2009) Phosphorylation-driven assembly of the RIP1-RIP3 complex regulates programmed necrosis and virus-induced inflammation. *Cell* 137:1112–1123
37. Cai Z et al (2013) Plasma membrane translocation of trimerized MLKL protein is required for TNF-induced necroptosis. *Nat Cell Biol* 16:55–65
38. Wallach D, Kang T-B, Dillon CP, Green DR (2016) Programmed necrosis in inflammation: toward identification of the effector molecules. *Science* 352:aaf2154–aaf2154
39. Ito Y et al (2016) RIPK1 mediates axonal degeneration by promoting inflammation and necroptosis in ALS. *Science* 353:603–608
40. Ofengeim D et al (2015) Activation of necroptosis in multiple sclerosis. *Cell Rep* 10:1836–1849
41. Oñate M et al (2019) The necroptosis machinery mediates axonal degeneration in a model of Parkinson disease. *Cell Death Differ*. <https://doi.org/10.1038/s41418-019-0408-4>
42. Wu J et al (2013) MIK1 knockout mice demonstrate the indispensable role of MIK1 in necroptosis. *Cell Res* 23:994–1006
43. Kim HY, Lee DK, Chung BR, Kim HV, Kim Y (2016) Intracerebroventricular injection of amyloid- β peptides in normal mice to acutely induce alzheimer-like cognitive deficits. *J Vis Exp* 2016:1–6
44. Vorhees CV, Williams MT (2006) Morris water maze: procedures for assessing spatial and related forms of learning and memory. *Nat Protoc* 1:848–858
45. Sadleir KR et al (2016) Presynaptic dystrophic neurites surrounding amyloid plaques are sites of microtubule disruption, BACE1 elevation, and increased A β generation in Alzheimer's disease. *Acta Neuropathologica* 2:1–22
46. Jimenez S et al (2014) Disruption of amyloid plaques integrity affects the soluble oligomers content from Alzheimer disease brains. *PLoS ONE* 9:1–17
47. Floden AM (2005) β -amyloid-stimulated microglia induce neuron death via synergistic stimulation of tumor necrosis factor and NMDA receptors. *J Neurosci* 25:2566–2575
48. Jimenez S et al (2008) Inflammatory response in the hippocampus of PS1M146L/APP 751SL mouse model of Alzheimer's disease: age-dependent switch in the microglial phenotype from alternative to classic. *J Neurosci* 28:11650–11661
49. Petersen RC et al (2009) Mild cognitive impairment: Ten years later. *Arch Neurol* 66:1447–1455
50. Serrano-Pozo A, Frosch MP, Masliah E, Hyman BT (2011) Neuropathological alterations in Alzheimer disease. *Cold Spring Harbor Perspect Med* 1:1002
51. Lu J et al (2019) Melatonin suppresses microglial necroptosis by regulating deubiquitinating enzyme A20 after intracerebral hemorrhage. *Front Immunol* 10:1–16
52. Huang Z et al (2018) Necroptosis in microglia contributes to neuroinflammation and retinal degeneration through TLR4 activation. *Cell Death Differ* 25:180–189
53. Faucher P, Mons N, Micheau J, Louis C, Beracochea DJ (2016) Hippocampal injections of induce selective working memory deficits and long-lasting alterations of ERK. *Signaling Pathway* 7:1–15
54. Wang G et al (2016) The effect of resveratrol on beta amyloid-induced memory impairment involves inhibition of phosphodiesterase-4 related signaling. *Oncotarget* 7:17380–17392
55. Paranjape GS, Gouwens LK, Osborn DC, Nichols MR (2012) Isolated amyloid- β (1–42) protofibrils, but not isolated fibrils, are robust stimulators of microglia. *ACS Chem Neurosci* 3:238–247
56. Xu H et al (2016) Environmental enrichment potently prevents microglia-mediated neuroinflammation by human amyloid β -protein oligomers. *J Neurosci* 36:9041–9056
57. Lucinaitė A et al (2020) Soluble A β oligomers and protofibrils induce NLRP3 inflammasome activation in microglia. *J Neurochem* 155:650–661
58. Sondag CM, Dhawan G, Combs CK (2009) Beta amyloid oligomers and fibrils stimulate differential activation of primary microglia. *J Neuroinflammation* 6:1–13
59. Dhawan G, Floden AM, Combs CK (2012) Amyloid- β oligomers stimulate microglia through a tyrosine kinase dependent mechanism. *Neurobiol Aging* 33:2247–2261
60. Liu S et al (2014) Necroptosis mediates TNF-induced toxicity of hippocampal neurons. *BioMed Res Int* 20:14
61. He S et al (2009) Receptor interacting protein kinase-3 determines cellular necrotic response to TNF- α . *Cell* 137:1100–1111
62. Hou J et al (2019) Discovery of potent necroptosis inhibitors targeting RIPK1 kinase activity for the treatment of inflammatory disorder and cancer metastasis. *Cell Death Dis* 10:493
63. Zhu S, Zhang Y, Bai G, Li H (2011) Necrostatin-1 ameliorates symptoms in R6/2 transgenic mouse model of Huntington's disease. *Cell Death Dis* 2:6–9
64. Degterev A et al (2005) Chemical inhibitor of nonapoptotic cell death with therapeutic potential for ischemic brain injury. *Nat Chem Biol* 1:112–119
65. Viola KL, Klein WL (2015) Amyloid β oligomers in Alzheimer's disease pathogenesis, treatment, and diagnosis. *Acta Neuropathol* 129:183–206
66. Lacor PN et al (2007) A β oligomer-induced aberrations in synapse composition, shape, and density provide a molecular basis for loss of connectivity in Alzheimer's disease. *J Neurosci* 27:796–807
67. Caughey B, Lansbury PT (2003) Protofibrils, pores, fibrils, and neurodegeneration: separating the responsible protein aggregates from the innocent bystanders. *Annu Rev Neurosci* 26:267–298
68. Soto C, Estrada L, Castilla J (2006) Amyloids, prions and the inherent infectious nature of misfolded protein aggregates. *Trends Biochem Sci* 31:150–155
69. Bloom GS (2014) Amyloid- β and tau: the trigger and bullet in Alzheimer disease pathogenesis. *JAMA Neurol* 71:505–508
70. Nishioka C, Liang HF, Barsamian B, Sun SW (2019) Amyloid-beta induced retrograde axonal degeneration in a mouse tauopathy model. *Neuroimage* 189:180–191
71. Roberson ED et al (2007) Reducing endogenous tau ameliorates amyloid β -induced deficits in an Alzheimer's disease mouse model. *Science* 316:750–754
72. Salvadores N, Gerónimo-Olvera C, Court FA (2020) Axonal degeneration in AD: the contribution of A β and Tau. *Front Aging Neurosci* 12:1–16
73. Yuan J, Amin P, Ofengeim D (2019) Necroptosis and RIPK1-mediated neuroinflammation in CNS diseases. *Nat Rev Neurosci* 20:19–33
74. Wegner KW, Saleh D, Degterev A (2017) Complex pathologic roles of RIPK1 and RIPK3: moving beyond necroptosis. *Trends Pharmacol Sci* 38:202–225

75. Noda M, Nakanishi H, Akaike N (1999) Glutamate release from microglia via glutamate transporter is enhanced by amyloid-beta peptide. *Neuroscience* 92:1465–1474
76. Klegeris A, Walker DG, McGeer PL (1997) Regulation of glutamate in cultures of human monocytic THP-1 and astrocytoma U-373 MG cells. *J Neuroimmunol* 78:152–161
77. Xu X et al (2010) The role of PARP activation in glutamate-induced necroptosis in HT-22 cells. *Brain Res* 1343:206–212
78. Hernández DE et al (2018) Axonal degeneration induced by glutamate excitotoxicity is mediated by necroptosis. *J Cell Sci* 131:jcs214684
79. Kawanokuchi J et al (2006) Production of interferon-gamma by microglia. *Multiple Sclerosis* (Houndmills, Basingstoke, England) 12:558–564

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

