



## Original Article

## Factors associated with poor outcomes among hospitalized patients with COVID-19: Experience from a MERS-CoV referral hospital



Mazin Barry<sup>a,\*</sup>, Muath Alotaibi<sup>a</sup>, Abdulellah Almohaya<sup>a,b</sup>, Abdulwahab Aldrees<sup>a</sup>, Ali AlHijji<sup>a</sup>, Nouf Althabit<sup>a</sup>, Sara Alhasani<sup>a</sup>, Layan Akkielah<sup>a</sup>, Abdulaziz AlRajhi<sup>a</sup>, Thamer Nohh<sup>c</sup>, Mohamad-Hani Temsah<sup>d</sup>, Jaffar A. Al-Tawfiq<sup>e,f,g</sup>

<sup>a</sup> Division of Infectious Diseases, Department of Internal Medicine, College of Medicine, King Saud University, Riyadh, Saudi Arabia

<sup>b</sup> Internal Medicine Department, Ad-Dariya Hospital, Ministry of Health, Riyadh, Saudi Arabia

<sup>c</sup> Trauma and Acute Care Surgery Unit, College of Medicine, King Saud University, Riyadh, Saudi Arabia

<sup>d</sup> Critical Care Unit, Pediatric Department, College of Medicine, King Saud University, Riyadh, Saudi Arabia

<sup>e</sup> Specialty Internal Medicine and Quality Department, Johns Hopkins Aramco Healthcare, Dhahran, Saudi Arabia

<sup>f</sup> Infectious Disease Division, Department of Medicine, Indiana University School of Medicine, Indianapolis, IN, USA

<sup>g</sup> Infectious Disease Division, Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, MD, USA

## ARTICLE INFO

## Article history:

Received 27 May 2021

Received in revised form

27 September 2021

Accepted 29 September 2021

## Keywords:

COVID-19

Risk factors

ICU

Invasive mechanical ventilation

## ABSTRACT

**Background:** Coronavirus disease 2019 (COVID-19) has resulted in millions of deaths, including more than 6000 deaths in the Kingdom of Saudi Arabia (KSA). Identifying key predictors of intensive care unit (ICU) admission and mortality among infected cases would help in identifying individuals at risk to optimize their care. We aimed to determine factors of poor outcomes in hospitalized patients with COVID-19 in a large academic hospital in Riyadh, KSA that serves as a Middle East Respiratory Syndrome coronavirus (MERS-CoV) referral center.

**Methods:** This is a single-center retrospective cohort study of hospitalized patients between March 15 and August 31, 2020. The study was conducted at King Saud University Medical City (KSUMC). COVID-19 infection was confirmed using real-time reverse transcriptase polymerase chain reaction (RT-PCR) for SARS-CoV-2. Demographic data, clinical characteristics, laboratory, radiological features, and length of hospital stay were obtained. Poor outcomes were admission to ICU, need for invasive mechanical ventilation (IMV), and in-hospital all-cause mortality.

**Results:** Out of 16,947 individuals tested in KSUMC, 3480 (20.5%) tested positive for SARS-CoV-2 and of those 743 patients (21%) were hospitalized. There were 62% males, 77% were younger than 65 years. Of all cases, 204 patients (28%) required ICU admission, 104 (14%) required IMV, and 117 (16%) died in hospital. In bivariate analysis, multiple factors were associated with mortality among COVID-19 patients. Further multivariate analysis revealed the following factors were associated with mortality: respiratory rate more than 24/min and systolic blood pressure <90 mmHg in the first 24 h of presentation, lymphocyte count <1 × 10<sup>9</sup>/L and aspartate transaminase level >37 units/L in the first 48 h of presentation, while a RT-PCR cycle threshold (Ct) value ≤24 was a predictor for IMV.

**Conclusion:** Variable factors were identified as predictors of different outcomes among COVID-19 patients. The only predictor of IMV was a low initial Ct values of SARS-CoV-2 PCR. The presence of tachypnea, hypotension, lymphopenia, and elevated AST in the first 48 h of presentation were independently associated with mortality. This study provides possible independent predictors of mortality and invasive mechanical ventilation. The data may be helpful in the early identification of high-risk COVID-19 patients in areas endemic with MERS-CoV.

© 2021 The Author(s). Published by Elsevier Ltd on behalf of King Saud Bin Abdulaziz University for Health Sciences. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

\* Corresponding author at: Division of Infectious Diseases, Department of Internal Medicine, College of Medicine, King Saud University, PO Box 2925, Riyadh 11461, Saudi Arabia.

E-mail address: [mbarry@ksu.edu.sa](mailto:mbarry@ksu.edu.sa) (M. Barry).

<https://doi.org/10.1016/j.jiph.2021.09.023>

1876-0341/© 2021 The Author(s). Published by Elsevier Ltd on behalf of King Saud Bin Abdulaziz University for Health Sciences. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

As of February 2021, just over one year has passed since the severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) was first discovered. This virus is the cause of the coronavirus disease 2019 (COVID-19) pandemic, which has infected almost 100 million people worldwide with almost two million deaths [1,2]. It has caused almost 370,000 cases with >6000 deaths in the Kingdom of Saudi Arabia (KSA) [2]. Thus, potential predictors of COVID-19-related mortality among different populations should be investigated to help optimize patient care and appropriately utilize health care resources [3].

Recent reports from several countries have suggested that older age, male gender, history of heart disease [3,4], and baseline chronic conditions [5] are significantly associated with poor outcomes. Hypertension (HTN) is associated with poor outcomes, and anticoagulant treatment may be associated with clinical benefits in terms of survival in patients with COVID-19 [6]. Thus, identifying predictors of poor outcomes across different populations is essential. In KSA, limited studies have investigated the predictors of morbidity and mortality in patients with COVID-19. A retrospective study of 352 critically ill patients with COVID-19 in KSA showed various predictors of 28-day mortality including old age, active smoking, increased lactate and D-dimers [7], similarly another study from Mecca showed that advanced age, end-stage renal disease, low oxygen saturations and high D-dimers were all associated with mortality [8]. A case series of 768 patients at a single center in Riyadh revealed that lung disease, obesity, smoking, diabetes mellitus were all predictors of intensive care admission [9], while another study from Madina showed higher mortality odds in older groups above 65 years with high C-reactive protein [10]. Within the same region, a study from Oman showed that in-hospital mortality was associated with older age, heart disease, liver disease, high ferritin, and in-patients requiring intensive care admission [11], while a study from Kuwait showed that age above 50 years, smoking, elevated C-reactive protein and elevated procalcitonin levels were all associated with admission to intensive care [12].

In addition, KSA is currently the only country in the world where Middle East Respiratory Syndrome Coronavirus (MERS-CoV) still causes sporadic community cases and nosocomial outbreaks [14]. Therefore, we aimed to study numerous cases and investigate their clinical and laboratory predictors for poor outcomes.

## Methodology

### *Design, inclusion, and exclusion criteria*

This work is a retrospective cohort single-center study carried out at King Saud University Medical City (KSUMC) in Riyadh, Saudi Arabia, which serves as a MERS-CoV referral hospital. During the study period from March 15 to August 31, 2020, we included patients aged 18 years or older who tested positive for SARS-CoV-2 infection and required hospitalization for at least 24 h. Patients who were still hospitalized by the last day of study inclusion were followed until discharge or in-hospital mortality, with the last outcome monitored until October 15, 2020. We aimed to study factors associated with a poor outcome, which is defined as: admission to the intensive care unit (ICU), the use of invasive mechanical ventilation (IMV), or in hospital all-cause mortality.

Based on the definition of severity by the Saudi Ministry of Health [15], disease severity was classified into asymptomatic, upper respiratory tract infection (URTI), mild to moderate disease (i.e., no oxygen requirement, no clinical evidence of pneumonia but with other symptoms, such as fever and diarrhea), severe disease (in which oxygen therapy was needed), and critical cases (in which

acute respiratory distress syndrome, overt sepsis, an altered level of consciousness, or multiorgan dysfunction were documented). All patients were admitted to single rooms under droplet and contact precautions as per hospital policy. They were placed in airborne infection isolation rooms with 6–12 air changes per hour, or, when not available, in a single room with high-efficiency particulate air filter. If aerosol-generating medical procedures were required, healthcare workers (HCWs) used fit-tested N95 masks [13].

### *Laboratory testing*

All the hospitalized patients underwent a nasopharyngeal or throat swab upon admission. The swabs were sent in viral transport media (Copan, Brescia, Italy). A confirmed COVID-19 was based on a positive result for both SARS-CoV-2 *E* and *S* genes using the RealStar® SARS-CoV-2 real-time reverse transcriptase PCR (RT-PCR) kit (Altona-Diagnostics, Hamburg, Germany) and Rotor-gene Q system (Qiagen®, Santa Clarita, CA, USA).

### *Data collection*

Patient data, which included demographic data, previous and current medical history, clinical findings, vital signs upon admission, laboratory results (i.e., molecular tests, microbiological tests), and radiological investigations (i.e., chest X-rays (CXR), computed tomographic scans), were collected directly from electronic health records. The hospital course, outcomes, and complications were extracted.

### *Statistical analysis*

Categorical variables (i.e., gender, occupation, nationality, smoking history, presenting complaints, CXR findings, treatment) were studied utilizing the chi-square test and logistic regression models. The SPSS software (version 25, IBM Corp.) was used for statistical analysis. Simple univariate logistic regression was applied to obtain associations, and a P value of <.05 was considered significant. A multivariate analysis was carried out using significant variables from univariate analysis to determine significant variables.

## Results

### *Demography*

Between March 15 and August 31, 2020, 16,974 adult patients were tested for SARS-CoV-2 by RT-PCR at KSUMC, Riyadh. Of those patients, 3481 (20.50%) were SARS-CoV-2 positive (Fig. 1). Of the positive cases, 744 (21.37%) were hospitalized, and one patient was transferred to another hospital and was excluded from further analysis. Thus, 743 patients were included in the study and reached an outcome of either discharge or in-hospital mortality. Among these patients, 459 (62%) were male, and 284 (38%) were female. The median age was 53 years, with an interquartile range of 24 (39–63), and 572 (77%) were below the age of 65 years. Among these patients, 56% were Saudi nationals, and 7% were HCWs. Underlying medical conditions included obesity in 43%, diabetes mellitus (DM) in 42%, and HTN in 40% (Table 1).

The source of SARS-CoV-2 infection was not identified in 58% of cases, whereas close contact with SARS-CoV-2-infected individuals was reported in 26%. Only .4% involved travel-related infections.

### *Clinical presentations*

Fever was the most common presentation (66%,  $n = 489$ ), followed by dyspnea (62%,  $n = 462$ ), cough (60%,  $n = 447$ ), diarrhea and/or vomiting (36%,  $n = 266$ ), and easy fatiguability (25%,  $n = 185$ ).

**Table 1**  
Demographic and clinical characteristics of 743 hospitalized patients with SARS-CoV-2 at King Saud University Medical City.

	(n)	(%)		
Gender	Female	284	38.2	
	Male	459	61.8	
Age	<65 years	572	77	
	≥65 year	171	23	
Median age (IQR)	53 (39–63)			
Occupation	HCWs	52	7	
	Non-HCWs	691	93	
Nationality	Saudi Nationals	423	57	
	Arab countries nationals	116	16	
	Indian Subcontinent nationals.	85	11	
	Other countries nationals.	119	16	
BMI, kgm <sup>2</sup>	Obese (≥30)	316	42.5	
	Overweight (25–29.9)	220	29.6	
	Normal (18.5–24.9)	143	19.2	
	Underweight (<18.5)	13	1.7	
Smoking	NA (not documented)	51	6.9	
	Active or former smoker	33	4.4	
	Life-time non-smoker	674	90.7	
Comorbid conditions	NA (not documented)	36	4.9	
	Diabetes mellitus	308	41.5	
	Hypertension	295	39.7	
	Dyslipidemia	104	14	
	Ischemic heart diseases	72	9.7	
	Lung diseases <sup>a</sup>	63	8.5	
	Heart failure	50	6.7	
	Chronic kidney disease	44	5.9	
	Hypothyroidism	41	5.5	
	Stroke	27	3.6	
	Autoimmune diseases	18	2.4	
	Active cancer	17	2.3	
	Pregnancy	Yes	42	14.8
No		242	85.2	
Blood group	A (Rh +/-)	100	32	
	B (Rh +/-)	71	22.8	
	AB (Rh +/-)	13	4.2	
	O (Rh +/-)	128	41	
	NA (not documented)	431		
Epidemiological link	Travel history to a country with SARS-CoV-2 cases	2	.3	
	Close contact with confirmed SARS-CoV-2 cases	193	26	
	Visit to a healthcare facility within 14 days	116	15.6	
	Not identified	432	58.1	
Presenting complaints	Fever	489	65.8	
	Dyspnea	462	62.2	
	Cough	447	60.2	
	GI symptoms	266	35.8	
	Diarrhea (n = 116, 44%)			
	Other GI symptoms (n = 150, 56%)			
	Fatigue	185	24.9	
	URTI symptoms	120	16.2	
	Headache	77	10.4	
	Anosmia	50	6.7	
	Asymptomatic	87	11.7	
	Duration of symptoms	7 days or less	480	64.6
		More than 7 days	176	23.7
NA (not applicable)		87	11.7	
Critical disease		141	19	
Severity <sup>b</sup>	Severe disease	123	16.5	
	Mild/moderate disease	312	42	
	URTI	80	10.8	
	Asymptomatic	87	11.7	
Initial admission site	Intensive care unit	104	14	
	Ward	639	86	
First poor outcome	Death	117	15.7	
	Discharge	626	84.3	
Median length of stay (IQR): 8 days (4–15 days)				
Second poor outcome	IMV need	103	13.9	
	No IMV need	640	86.1	
Median MV days (IQR): 8 days (4–19 days)				
Third poor outcome	ICU admission	204	27.5	
	No ICU admission	539	72.5	
Median ICU days (IQR): 8 days (5–16 days)				

HCWs; healthcare workers, GI; gastrointestinal, URTI; upper respiratory tract infection, ICU; intensive care unit, IMV; invasive mechanical ventilation.

<sup>a</sup> Including chronic obstructive pulmonary disease, bronchial asthma, interstitial lung disease, obstructive sleep apnea–obesity hypoventilation syndrome, and bronchiectasis.

<sup>b</sup> According to the Saudi Ministry of Health protocol for patients suspected of or confirmed to have COVID-19 [9].




Fig. 1. Diagram showing the study population and outcomes.

Most patients presented to a hospital within 7 days of symptom onset ( $n = 480$ , 65%). In relation to the severity of the disease, the disease was critical in 19%, severe in 17%, and mild/moderate in 42%, and 11% had URTI. The remaining 11.7% had asymptomatic infections and tested positive during active screening strategy. Among the patients, 104 (14%) were directly admitted to the ICU, and the remaining 639 (86%) were admitted to the general ward. From the total cohort, 281, 263, and 264 underwent testing for MERS-CoV, influenza A, and influenza B, respectively, via PCR. None were positive, indicating no co-infection with these viruses in the tested cohort.

#### Factors associated with poor outcomes (univariate analysis)

By the end of the study period, 117 (15.7%) had died. The association of different risk factors with mortality was tested using univariate analysis. Significant risk factors were old age ( $\geq 65$  years), non-HCW, and the presence of a comorbid-illness. Age was not a significant risk factor for IMV or ICU admission. The presence of lung infiltration as shown in chest X-ray upon hospital admission was significantly associated with mortality ( $P$  value  $< .001$ , OR 2.5, 95% CI [1.5–4.1]).

Clinical and laboratory risk factors in relation to ICU admission, IMV, or mortality are detailed in Table 2.

#### Factors associated with poor outcomes (multivariate analysis)

All significant factors in univariate analysis were reanalyzed using multivariate logistic regression with each poor outcome, namely, mortality, IMV, and ICU admission. The analysis showed that an initial respiratory rate  $>24/\text{min}$  ( $P = .031$ , OR 2.8, 95% CI [1.1–7.2]), a systolic blood pressure (SBP)  $<90$  mmHg ( $P = .041$ , OR 6.0, 95% CI [1.1–34.0]) in the first 24 h of presentation, a lymphocyte count  $<1 \times 10^9/\text{L}$  ( $P = .020$ , OR 2.9, 95% CI [1.2–7.2]), and an aspar-

tate transaminase (AST) level  $>37$  units/L ( $P = .026$ , OR 3.3, 95% CI [1.2–9.4]) in the first 48 h of presentation, were all independently associated with mortality.

A SBP  $<90$  mmHg and AST  $>37$  units/L were also independent risk factors for ICU admission. Ct values that are  $\leq 24$  of the first positive RT-PCR was the only risk factor for IMV in the multivariate analysis (Table 3).

## Discussion

In this study, we described the clinical presentations and outcomes of patients with COVID-19 who required ICU admission. In addition, we examined the risk factors associated with poor outcomes of death, ICU admission, and IMV. Most of our patients were male, accounting for 61.8% of the study population. A similar finding was observed in a systematic review and meta-analysis showing that among 13,654 patients, 8557 (62.6%) were male [16]. Approximately 77% of our patients were younger than 65 years, the finding that younger adults represent the majority of hospitalized patients with COVID-19 was also found by another local study from a center in the capital of Saudi Arabia, where 66% of patients were younger than 60 years [17]. The median age of our patients was 53 years, which is similar to a systemic review of six studies conducted in China, where the median age was 55.5 years [18]. Of the included patients, 42.5% were obese (i.e., with a BMI of  $\geq 30$ ), which is much less than the results in a case series of 463 patients in Detroit, US (58%) [19] and greater than the result observed by Argenziano MG et al. (36.3%) [20].

Importantly, the most prevalent chronic disease was DM at 42%, followed by HTN at 40%. Active cancer and autoimmune diseases were not prevalent in our study population, which is different from a retrospective case series in Spain that included 1549 patients; the study showed that 55% of patients hospitalized with COVID 19 had HTN, 45% had a history of cerebrovascular disease, and only

**Table 2**  
Predictors of poor outcomes among hospitalized patients with SARS-CoV-2 infection at King Saud University Medical City using univariate analysis.

	Univariate OR (95% CI) (For mortality)	P	Univariate OR (95% CI) (For invasive mechanical ventilation)	P	Univariate OR (95% CI) (For ICU admission)	P
Demographic data						
Gender	–	.441	–	.340	–	.091
Age ≥65 years (vs. <65 years)	2.5 (1.6–3.8)	<.001	–	.279	–	.077
Non-HCWs (vs. HCWs)	10.3 (1.4–75.2)	.005	8.8 (1.2–64.6)	.010	10.3 (2.5–42.9)	<.001
BMI	–	.170	–	.501	–	.220
Smoking history	–	.077	–	.064	–	.084
Blood group	–	.817	–	.773	–	.582
Past medical history (each against no history of that disease)						
Presence of ≥1 comorbidities (vs. none)	2.9 (1.8–4.8)	<.001	2.0 (1.3–3.2)	.003	1.9 (1.3–2.7)	<.001
Use of ACEIs or ARBs (vs. no use)	–	.266	–	.928	–	.364
Hypertension	2.7 (1.8–4.0)	<.001	1.8 (1.2–2.8)	.004	1.8 (1.3–2.5)	<.001
Diabetes mellitus	2.2 (1.5–3.3)	<.001	1.8 (1.2–2.8)	.004	1.8 (1.3–2.5)	<.001
Heart failure	2.5 (1.3–4.7)	.004	–	.381	2.4 (1.4–4.3)	.002
Stroke	–	.415	–	.249	–	.486
Chronic kidney disease	2.1 (1.1–4.3)	.030	–	.192	1.9 (1.0–3.6)	.039
Dyslipidemia	2.1 (1.3–3.5)	.002	–	.088	–	.078
Any lung disease <sup>a</sup>	–	.066	–	.213	–	.836
Autoimmune disease	–	.753	–	.731	–	.968
Hypothyroidism	–	.810	–	.541	–	.530
Active cancer	–	.166	–	.800	–	.582
History and vital signs in the first 24 h						
History of fever (vs. no fever)	–	.189	–	.148	–	.051
History of dyspnea (vs. no dyspnea)	1.9 (1.2–3.0)	.005	2.3 (1.4–3.7)	.001	2.5 (1.7–3.6)	<.001
History of Cough (vs. no cough)	–	.571	–	.357	–	.432
History of Anosmia (vs. no Anosmia)	–	.473	–	.726	–	.833
History of GI symptoms (vs. no GI symptoms)	–	.414	–	.391	–	.603
Symptoms duration less than 7 days (vs. ≥7 days)	2.6 (1.5–4.6)	<.001	1.8 (1.0–3.0)	.037	–	.321
Initial temperature of ≥38 °C (vs. below 38)	1.6 (1.0–2.2)	.031	1.6 (1.1–2.5)	.026	1.5 (1.0–2.0)	.030
Initial systolic blood pressure <90 mmHg (vs. ≥90 mmHg)	4.1 (2.0–8.4)	<.001	4.9 (2.4–10.0)	<.001	3.2 (1.6–6.3)	.001
Initial respiratory rate >24 breaths/min (vs. ≤24 breaths/min)	2.9 (1.9–4.3)	<.001	3.6 (2.3–5.5)	<.001	5.0 (3.5–7.0)	<.001
Initial oxygen saturation below 94% (vs. ≥94%)	1.9 (1.3–2.9)	.002	2.2 (1.4–3.4)	<.001	3.0 (2.1–4.2)	<.001
Laboratory and radiological findings in the first 48 h						
RT-PCR Ct value ≤24 (vs. >24)	1.8 (1.2–2.7)	.004	1.7 (1.1–2.6)	.018	–	.072
White blood cell counts of <4 × 10 <sup>9</sup> cells/L (vs. ≥4 × 10 <sup>9</sup> cells/L)	–	.471	–	.756	–	.460
Lymphocytes count less than 1 × 10 <sup>9</sup> cells/L (vs. ≥1 × 10 <sup>9</sup> cells/L)	3.7 (2.5–5.7)	<.001	4.8 (3.0–7.6)	<.001	4.2 (3.0–6.0)	<.001
Platelets count less than 150 × 10 <sup>9</sup> cells/L (vs. ≥150 × 10 <sup>9</sup> cells/L)	–	.293	–	.161	–	.292
D-dimer >.45 mg/dL (vs. ≤.45 mg/dL)	2.2 (1.0–4.6)	.044	2.7 (1.1–6.3)	.021	2.3 (1.3–4.0)	.005
AST >37 units/L (vs. ≤37 units/L)	3.4 (2.1–5.3)	<.001	3.7 (2.2–6.0)	<.001	3.0 (2.0–4.2)	<.001
ALT >61 units/L (vs. ≤61 units/L)	–	.627	1.6 (1.0–2.6)	.035	1.5 (1.0–2.2)	.028
Creatinine >115 μmol/L (vs. ≤115)	3.9 (2.5–6.0)	<.001	3.2 (2.0–5.0)	<.001	2.8 (1.9–4.1)	<.001
Ferritin >400 μg/L (vs. ≤400 μg/L)	1.6 (1.0–2.4)	.037	2.4 (1.5–4.0)	<.001	2.3 (1.6–3.4)	<.001
CRP >20 mg/L (vs. ≤20 mg/L)	4.3 (1.7–10.8)	.001	4.0 (1.6–10.0)	.002	2.5 (1.4–4.4)	.001
PCT >.5 ng/mL (vs. ≤.5 ng/mL)	2.3 (1.2–4.4)	.008	2.0 (1.1–3.8)	.029	2.3 (1.5–3.8)	<.001
Trop I >100 ng/L (vs. ≤100 ng/L)	2.6 (1.3–5.2)	.006	2.8 (1.4–5.6)	.003	2.9 (1.5–5.7)	.001
CK >308 units/L (vs. ≤308 units/L)	2.0 (1.2–3.4)	.006	2.3 (1.4–3.8)	.001	1.6 (1.1–2.5)	.030
Lactate >2 mmol/L (vs. ≤2)	–	.080	1.9 (1.1–3.1)	.014	–	.075
IL-6 >7 pg/mL (vs. ≤7)	–	.052	–	.262	–	.390
CXR infiltrates (vs. no infiltrates)	2.5 (1.5–4.1)	<.001	3.0 (1.7–5.2)	<.001	3.0 (2.0–4.4)	<.001

HCWs; healthcare workers, RT-PCR; reverse transcription-polymerase chain reaction, ct; cycle threshold, AST; aspartate transaminase, CRP; C-reactive protein, PCT; procalcitonin, Trop I; troponin I, CK; creatinine kinase, CXR; chest X-ray, BMI; body mass index, ACEIs; angiotensin converting enzyme inhibitors, ARBs; angiotensin receptor blockers, GI; gastrointestinal, ALT; alanine transaminase, IL-6; interleukin-6, ICU; intensive care unit.

<sup>a</sup> Including chronic obstructive pulmonary disease, bronchial asthma, interstitial lung disease, obstructive sleep apnea-obesity hypoventilation syndrome, and bronchiectasis.

**Table 3**  
Multivariate analysis of predictors of poor outcomes (mortality, invasive mechanical ventilation, ICU admission) among hospitalized patients with SARS-CoV-2 infection at King Saud University Medical City.

	Multivariate OR (95% CI)	P
Predictors of mortality		
Initial respiratory rate >24 breaths/min (vs. ≤24 breaths/min)	2.8 (1.1–7.2)	.031
Initial systolic blood pressure <90 mmHg (vs. ≥90 mmHg)	6.0 (1.1–34.0)	.041
Lymphocyte count < 1 × 10 <sup>9</sup> cells/L (vs. ≥1 × 10 <sup>9</sup> cells/L)	2.9 (1.2–7.2)	.020
AST >37 units/L (vs. ≤37 units/L)	3.3 (1.2–9.4)	.026
Predictors of invasive mechanical ventilation		
RT-PCR Ct value ≤ 24 (vs. >24)	2.5 (1.1–5.4)	.023
Predictors of ICU admission		
Initial systolic blood pressure < 90 mmHg (vs. ≥90 mmHg)	5.9 (1.2–30.0)	.032
AST > 37 units/L (vs. ≤37 units/L)	3.1 (1.2–8.5)	.024

RT-PCR; reverse transcription-polymerase chain reaction, Ct; cycle threshold, AST; aspartate transaminase.



24.8% had DM [21]. Another retrospective study in New York, USA showed that among 1000 patients with COVID-19, 57% had HTN, and 34.3% had DM [20]. These findings may suggest the increasing prevalence of DM in Saudi Arabia; a recent report in KSA showed that 48% of admitted patients had DM which is almost similar to our finding [22]. Although no definite risk was significantly associated with blood type, the most frequent blood group (41%) in our study population of 743 hospitalized patients was ABO group O (Rh +−). This finding was different from an observation in a case control study that included over 1000 Iraqi patients, where blood group A (Rh +−) was the most common [23].

Pregnant women with COVID-19 infection comprised 14.8% of our population, which is similar to a reported analysis of COVID-19 cases in the UK [24]. The majority of the patients included in this study did not report a specific epidemiological link of COVID-19 infection acquisition, and only 26% reported contact with a confirmed case. This finding is different from the reported results of a study in Wuhan, China, where 92% of patients reported contact with a confirmed case [25]. The lack of any known contact may be attributed to unrecognized asymptomatic transmission [26,27].

Fever was the most reported symptom, followed by dyspnea and cough. Only 6.7% of the patients presented with anosmia, which was the least reported symptom. This result is consistent with the results of a systemic review and meta-analysis by Mesas et al. [5] and with studies in the US, UK, Turkey, and China [20,24,28,29]. However, in one study that included data from 18 European hospitals, headache, anosmia, and nasal obstruction were the most reported symptoms [30]. The duration of symptoms was less than 7 days in most of our patients, which is similar to the reported duration of illness in a study in Wuhan, China [31]. This result can be explained by most of our patients having mild to moderate disease. Patients with URTI or with asymptomatic infection were mainly admitted for other reasons (e.g., delivery). A majority of the patients were admitted to COVID-19 wards, with only 14% requiring direct ICU admission with an overall median length of hospital stay of 8 days. One study conducted in New York showed a higher percentage of ICU admissions (23.6%) with an overall median length of stay of 6 days [20]. The case fatality ratio (CFR) in our study was 15.7%, whereas those in two studies conducted in the US were 21.1% and 21% [20,32]. A study in Wuhan, China, showed an overall CFR of 28.27% [25]. A study in the UK including 20,133 patients showed a CFR of 26% with a median age of 80 years [24]. Our cohort was consistent with previous studies citing age as an important predictor of mortality [33,34].

Serum lactate was identified as a risk factor for the need for IMV in our study. Chen et al. showed that high serum lactate was found in mechanically ventilated patients; however, it was not adjusted to other factors [35]. Inconsistent with our study Wang et al. showed an association between serum lactate and mortality, with an odds ratio of 2.6 [36]. Most of our cohort did not have a lactate level upon admission, as it was analyzed only when the patient deteriorated, which may explain its association with the need for IMV.

Our univariate analysis showed that D-dimer and creatinine were associated with mortality, ICU admission, and the need for IMV. Figliozzi et al. have demonstrated in a meta-analysis of >3000 patients that higher D-dimer is associated with death, ICU admission, and/or IMV [16]. A study in Brazil that identified predictors of poor outcome showed that a creatinine level of >105 µmol/L is significantly associated with all three outcomes. Consistent with our cohort, their multivariate analysis showed that creatinine level was not associated with mortality [33].

Our cohort did not show a poor outcome associated with gender, similar to other studies [33,37]. On the contrary, a meta-analysis of >600,000 patients showed significantly worse outcomes with male gender; however, this result was not adjusted to other patients' characteristics [16].

A striking feature of our study is that BMI was not significantly associated with poor outcome. Obesity prevalence in Saudi Arabia is approximately 35% [38], and that in our cohort was slightly higher. A meta-analysis of >10,000 patients showed poor outcomes associated with obesity, with a pooled OR of 1.88 [39]. Obesity prevalence in that meta-analysis was 33%, which was less than that in our cohort. In addition, in this meta-analysis, studies from the USA that has an age-adjusted prevalence of obesity of 42.4% [40], which is almost similar to Saudi Arabia at 38.3% [41], did not show statistically significant obesity risk.

Smoking prevalence in Saudi Arabia has a median of 16.5% among students, 22.6% among adults, and 25% among the those above 65 years of age [42]. However, our cohort showed a much lower rate of only 4.4%, these differences may be due to the cited study being conducted more than a decade ago, since then, in 2017, a 100% excise tax on tobacco products was enforced which led to a significant reduction in smoking in the country [43]. A meta-analysis of 19 studies showed a progression of COVID-19 in patients with a history of smoking (OR 1.91) [44].

Our study did not show any association between poor outcome and ABO blood group. A meta-analysis of five studies showed a slightly increased association between mortality in patients with COVID-19 and blood group A [45]. Our institute performs blood grouping in patients that receive blood or blood products; hence, half of our patients had no documented blood group.

Furthermore, consistent with other studies [46,47], the use of angiotensin converting enzyme inhibitors or angiotensin receptor blockers was not associated with any changes in COVID-19 progression.

Cerebrovascular diseases, active malignancy, and lung diseases are linked to increased morbidity and mortality in patients with COVID-19 [16]. However, our cohort did not show a similar association. This result may be due to a younger cohort, which has a lower prevalence of diseases compared with older populations.

A recent meta-analysis of >5000 patients showed that in contrast with dyspnea, cough and gastrointestinal (GI) symptoms were not associated with higher mortality [5]. In another meta-analysis, presence of diarrhea on presentation was associated with more severe disease with an odd ratio of 1.63 (95% CI: 1.11–3.38) compared to those who did not have diarrhea [39]. A median duration of symptoms of 5 days has been associated with higher mortality [40]. By contrast, a study in China showed that COVID-19 survivors and non-survivors showed a similar median duration of 11 days [25]. Our study did not show a significant association between cough, GI symptoms, or the duration of symptoms and poor outcome.

Using multivariate logistic regression, we identified several clinical and laboratory predictors of poor outcomes. Although a cut-off value of 30 breaths/min is considered one of the criteria for severe disease as per the Saudi Ministry of Health guidelines [15], we found that a lower value of >24 breaths/min is an independent risk factor associated with mortality. A study in South Florida showed an OR of 2.54 for ICU admission and death using multivariate analysis [41]. A similar study in Louisiana showed a hazard ratio of 2 for in-hospital death with a respiratory rate of >24 breaths/min [34].

We report low SBP as an additional risk factor for mortality. A study in New York including >6000 patients admitted showed that hypotension is associated with mortality (HR: 1.38 [1.06–1.8]) [42].

Lymphopenia has been studied as a potential predictor of poor outcome in multiple publications. In one prospective study, lymphopenia had no effect on outcomes [43]. A large meta-analysis that included >20,000 patients [16] showed that lymphocytopenia was associated with an increased risk of mortality in hospitalized patients with COVID-19. Our study also showed it to be an independent risk factor for mortality (OR of 2.9).

AST is a marker for liver function. A meta-analysis of 21 studies showed that elevated AST, especially in older population, is asso-

ciated with increased mortality (effect size: 1.61) [5]. A study in China showed that AST is more related to mortality compared with other liver enzymes (HR of 4.81 if AST is between 40 and 120 U/L) [44]. Another meta-analysis of four studies showed an OR of 2.53 for ICU admission with elevated AST [42].

As with other viral illnesses, such as HIV, Ebola, and influenza, SARS-CoV-2 viral load is associated with disease severity and progression [45]. Reinforcing findings in recent reports [46], our study showed a significant association between high viral load and the need for IMV, with a Ct value of  $\leq 24$  serving as an independent predictor for the need for IMV and indicating more severe disease.

Our study has several limitations. Given that this work is a retrospective study from a single center, not all patients were tested for all inflammatory markers, and the effect of therapeutic measures was not studied.

In conclusion, variable factors were identified as predictors of different outcomes among COVID-19 patients in a MERS-CoV referral hospital. A low initial Ct value of SARS-CoV-2 PCR predicted IMV. The presence of tachypnea, hypotension, lymphopenia, and elevated AST in the first 48 h of presentation were independently associated with mortality. This study provides independent predictors of mortality and IMV. The data may be helpful in the early identification of high-risk patients with COVID-19. Large prospective studies are warranted to validate these results.

#### Funding

No funding sources.

#### Competing interests

None declared.

#### Ethical approval

The study was approved by the Institutional Review Board of the KSUMC (IRB# E20-4979).

#### CRediT authorship contribution statement

**Mazin Barry:** Conceptualization, Validation, Investigation, Resources, Writing - original draft, Writing - review & editing, Visualization, Supervision, Project administration. **Muath Alotaibi:** Methodology, Software, Validation, Formal analysis, Investigation, Data curation, Writing - original draft, Writing - review & editing. **Abdullah Almhaya:** Methodology, Software, Validation, Formal analysis, Investigation, Data curation, Writing - original draft, Writing - review & editing. **Abdulwahab Aldrees:** Investigation, Data curation, Writing - original draft, Writing - review & editing. **Ali AlHijji:** Investigation, Data curation, Writing - original draft, Writing - review & editing. **Nouf Althabit:** Investigation, Data curation, Writing - original draft, Writing - review & editing. **Sara Alhasani:** Investigation, Data curation, Writing - original draft, Writing - review & editing. **Layan Akkielah:** Investigation, Data curation, Writing - original draft. **Abdulaziz AlRajhi:** Investigation, Data curation, Writing - original draft, Writing - review & editing. **Thamer Noubh:** Writing - review & editing, Visualization, Project administration. **Mohamad-Hani Temsah:** Writing - review & editing, Visualization, Project administration. **Jaffar A. Al-Tawfiq:** Validation, Formal analysis, Investigation, Writing - review & editing, Visualization, Project administration.

#### Acknowledgement

We would like to thank Scribendi English editing service.

#### References

- [1] World Health Organization. WHO Director-General's opening remarks at the media briefing on COVID-19—11 March 2020; 2020.
- [2] Johns Hopkins University. Johns Hopkins Coronavirus Resource Center. 2020. COVID-19 Map; 2020.
- [3] Tian W, Jiang W, Yao J, Nicholson CJ, Li RH, Sigurslid HH, et al. Predictors of mortality in hospitalized COVID-19 patients: a systematic review and meta-analysis. *J Med Virol* 2020;92:1875–83.
- [4] Bhargava A, Sharma M, Akagi E, Szpunar SM, Saravolatz L. Predictors for in-hospital mortality from COVID-19 infection among adults 18–65 years. *Infect Control Hosp Epidemiol* 2020;1–13.
- [5] Mesas AE, Caverio-Redondo I, Alvarez-Bueno C, Sarria Cabrera MA, Maffei de Andrade S, Sequi-Dominguez I, et al. Predictors of in-hospital COVID-19 mortality: a comprehensive systematic review and meta-analysis exploring differences by age, sex and health conditions. *PLoS One* 2020;15:e0241742.
- [6] Boari GEM, Chiarini G, Bonetti S, Malerba P, Bianco G, Faustini C, et al. Prognostic factors and predictors of outcome in patients with COVID-19 and related pneumonia: a retrospective cohort study. *Biosci Rep* 2020;40.
- [7] Alharthy A, Aletreby W, Faqih F, Balhamar A, Alaklobi F, Alanezi K, et al. Clinical characteristics and predictors of 28-day mortality in 352 critically ill patients with COVID-19: a retrospective study. *J Epidemiol Glob Health* 2021;11:98–104.
- [8] Alwafi H, Naser AY, Qanash S, Brinji AS, Ghazawi MA, Alotaibi B, et al. Predictors of length of hospital stay, mortality, and outcomes among hospitalised COVID-19 patients in Saudi Arabia: a cross-sectional study. *J Multidiscip Healthc* 2021;14:839–52.
- [9] Abohamr SI, Abazid RM, Aldossari MA, Amer HA, Badhawi OS, Aljunaidi OM, et al. Clinical characteristics and in-hospital mortality of COVID-19 adult patients in Saudi Arabia. *Saudi Med J* 2020;41:1217–26.
- [10] Albalawi O, Alharbi Y, Bakouri M, Alqahani A, Alanazi T, Almutairi AZ, et al. Clinical characteristics and predictors of mortality among COVID-19 patients in Saudi Arabia. *J Infect Public Health* 2021;14:994–1000.
- [11] Khamis F, Memish Z, Bahrani MA, Dowaiqi SA, Pandak N, Bolushi ZA, et al. Prevalence and predictors of in-hospital mortality of patients hospitalized with COVID-19 infection. *J Infect Public Health* 2021;14:759–65.
- [12] Almazeedi S, Al-Youha S, Jamal MH, Al-Haddad M, Al-Muhaini A, Al-Ghimlas F, et al. Characteristics, risk factors and outcomes among the first consecutive 1096 patients diagnosed with COVID-19 in Kuwait. *EclinicalMedicine* 2020;24:100448.
- [13] Health Mo. Command and Control Center. National Public Health Events, Epi-week 06. 07/02/2021–13/02/2021. Saudi Arabia: Ministry of Health; 2021 <https://www.moh.gov.sa/en/CCC/Pages/default.aspx>.
- [14] Barry M, Phan MV, Akkielah L, Al-Majed F, Alhethel A, Somily A, et al. Nosocomial outbreak of the Middle East Respiratory Syndrome coronavirus: a phylogenetic, epidemiological, clinical and infection control analysis. *Travel Med Infect Dis* 2020;37:101807.
- [15] MOH. Saudi MoH Protocol for Patients Suspected of/Confirmed with COVID-19 Supportive care and antiviral treatment of suspected or confirmed COVID-19 infection (Version 2.2) December 13th, 2020; 2020.
- [16] Figliozzi S, Masci PG, Ahmadi N, Tondi L, Koutli E, Aimo A, et al. Predictors of adverse prognosis in COVID-19: a systematic review and meta-analysis. *Eur J Clin Invest* 2020;50:e13362.
- [17] Sarfraz S, Mohammed A, Reem A, Fahad A, Faisal A, Rayan A, et al. Clinical characteristics of patients with COVID-19 in Saudi Arabia — a single center experience. *Res Rev Infect Dis* 2020;3.
- [18] Jiang F, Deng L, Zhang L, Cai Y, Cheung CW, Xia Z. Review of the clinical characteristics of coronavirus disease 2019 (COVID-19). *J Gen Intern Med* 2020;35:1545–9.
- [19] Suleyman G, Fadel RA, Malette KM, Hammond C, Abdulla H, Entz A, et al. Clinical characteristics and morbidity associated with coronavirus disease 2019 in a series of patients in metropolitan Detroit. *JAMA Netw Open* 2020;3:e2012270.
- [20] Argenziano MG, Bruce SL, Slater CL, Tiao JR, Baldwin MR, Barr RG, et al. Characterization and clinical course of 1000 patients with coronavirus disease 2019 in New York: retrospective case series. *BMJ* 2020;369:m1996.
- [21] Jiménez E, Fontán-Vela M, Valencia J, Fernández-Jiménez I, Álvaro-Alonso EA, Izquierdo-García E, et al. Characteristics, complications and outcomes among 1549 patients hospitalised with COVID-19 in a secondary hospital in Madrid, Spain: a retrospective case series study. *BMJ Open* 2020;10:e042398.
- [22] Al Hayek AA, Robert AA, Matar AB, Algarni A, Alkubedan H, Alharbi T, et al. Risk factors for hospital admission among COVID-19 patients with diabetes. A study from Saudi Arabia. *Saudi Med J* 2020;41:1090–7.
- [23] Ad'hiah AH, Abdullah MH, Alsudani MY, Shnawa RMS, Al-Sa'ady AJR, Allami RH, et al. Association between ABO blood groups and susceptibility to COVID-19: profile of age and gender in Iraqi patients. *Egypt J Med Hum Genet* 2020;21:76.
- [24] Docherty AB, Harrison EM, Green CA, Hardwick HE, Pius R, Norman L, et al. Features of 20 133 UK patients in hospital with covid-19 using the ISARIC WHO Clinical Characterisation Protocol: prospective observational cohort study. *BMJ* 2020;369:m1985.
- [25] Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020;395:1054–62.
- [26] Al-Tawfiq JA. Asymptomatic coronavirus infection: MERS-CoV and SARS-CoV-2 (COVID-19). *Travel Med Infect Dis* 2020;35:101608.

- [27] Aljishi JM, Alhazzaj AH, Alkhabbaz FL, AlAbduljabar TH, Alsaif A, Alsaif H, et al. Clinical characteristics of asymptomatic and symptomatic COVID-19 patients in the Eastern Province of Saudi Arabia. *J Infect Public Health* 2021;14:6–11.
- [28] Aggarwal S, Garcia-Telles N, Aggarwal G, Lavie C, Lippi G, Henry BM. Clinical features, laboratory characteristics, and outcomes of patients hospitalized with coronavirus disease 2019 (COVID-19): early report from the United States. *Diagnosis (Berl)* 2020;7:91–6.
- [29] Zhou Y, He Y, Yang H, Yu H, Wang T, Chen Z, et al. Exploiting an early warning Nomogram for predicting the risk of ICU admission in patients with COVID-19: a multi-center study in China. *Scand J Trauma Resusc Emerg Med* 2020;28:106.
- [30] Lechien JR, Chiesa-Estomba CM, Place S, Van Laethem Y, Cabaraux P, Mat Q, et al. Clinical and epidemiological characteristics of 1420 European patients with mild-to-moderate coronavirus disease 2019. *J Intern Med* 2020;288:335–44.
- [31] Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020;395:497–506.
- [32] Richardson S, Hirsch JS, Narasimhan M, Crawford JM, McGinn T, Davidson KW, et al. Presenting characteristics, comorbidities, and outcomes among 5700 patients hospitalized with COVID-19 in the New York City area. *JAMA* 2020.
- [33] Brandao Neto RA, Marchini JF, Marino LO, Alencar JCG, Lazar Neto F, Ribeiro S, et al. Mortality and other outcomes of patients with coronavirus disease pneumonia admitted to the emergency department: a prospective observational Brazilian study. *PLoS One* 2021;16:e0244532.
- [34] Price-Haywood EG, Burton J, Fort D, Seoane L. Hospitalization and mortality among black patients and white patients with Covid-19. *N Engl J Med* 2020;382:2534–43.
- [35] Chen J, Zhu Y-F, Du Z-Q, Li W-F, Zhang M-J, Zhao S-D, et al. Predictors of mechanical ventilation for COVID-19: combined data from three designated hospitals. *Eur Rev Med Pharmacol Sci* 2020;24(24):13065–71.
- [36] Wang Z, Wang Z. Identification of risk factors for in-hospital death of COVID-19 pneumonia — lessons from the early outbreak. *BMC Infect Dis* 2021;21:113.
- [37] Holler JG, Eriksson R, Jensen TO, van Wijhe M, Fischer TK, Sogaard OS, et al. First wave of COVID-19 hospital admissions in Denmark: a Nationwide population-based cohort study. *BMC Infect Dis* 2021;21:39.
- [38] World Health Organization. Global Health Observatory (GHO) data: overweight and obesity by country; 2020.
- [39] Malik P, Patel U, Patel K, Martin M, Shah C, Mehta D, et al. Obesity a predictor of outcomes of COVID-19 hospitalized patients—a systematic review and meta-analysis. *J Med Virol* 2021;93:1188–93.
- [40] Hales CM, Carroll MD, Fryar CD, Ogden CL. Prevalence of obesity and severe obesity among adults: United States, 2017–2018. *NCHS Data Brief* 2020:1–8.
- [41] Al-Qahtani AM. Prevalence and predictors of obesity and overweight among adults visiting primary care settings in the Southwestern Region, Saudi Arabia. *Biomed Res Int* 2019;2019:8073057.
- [42] Bassiony MM. Smoking in Saudi Arabia. *Saudi Med J* 2009;30:876–81.
- [43] Alghamdi A, Fallatah A, Okal F, Felemban T, Eldigire M, Almodaimegh H. Smoking behaviour after enforcement of a 100% tax on tobacco products in Saudi Arabia: a cross-sectional study. *East Mediterr Health J* 2020;26:39–46.
- [44] Patanavanich R, Glantz SA. Smoking is associated with COVID-19 progression: a meta-analysis. *Nicotine Tob Res* 2020;22:1653–6.
- [45] Liu N, Zhang T, Ma L, Zhang H, Wang H, Wei W, et al. The impact of ABO blood group on COVID-19 infection risk and mortality: a systematic review and meta-analysis. *Blood Rev* 2021;48:100785. <http://dx.doi.org/10.1016/j.blre.2020.100785>. Epub 2020 Dec 8. PMID: 33309392; PMCID: PMC7834371.
- [46] Alamer AA, Almulhim AS, Alrashed AA, Abraham I. Mortality, severity, and hospital admission among COVID-19 patients with ACEI/ARB use: a meta-analysis stratifying countries based on response to the first wave of the pandemic. *Healthcare (Basel)* 2021;9.
- [47] Zhang P, Zhu L, Cai J, Lei F, Qin JJ, Xie J, et al. Association of inpatient use of angiotensin-converting enzyme inhibitors and angiotensin II receptor blockers with mortality among patients with hypertension hospitalized with COVID-19. *Circ Res* 2020;126:1671–81.