



Review

# The Effects of High-Speed Resistance Training on Health Outcomes in Independent Older Adults: A Systematic Review and Meta-Analysis

Alexandre Duarte Martins <sup>1,2,3,\*</sup> , Orlando Fernandes <sup>1</sup>, Ana Pereira <sup>3,4</sup>, Rafael Oliveira <sup>2,3,5</sup> ,  
Franco David Alderete Goñi <sup>6</sup> , Nilton João Chantre Leite <sup>1</sup> and João Paulo Brito <sup>2,3,5</sup>

- <sup>1</sup> Comprehensive Health Research Centre (CHRC), Departamento de Desporto e Saúde, Escola de Saúde e Desenvolvimento Humano, Universidade de Évora, Largo dos Colegiais, 7000-727 Évora, Portugal; orlandoj@uevora.pt (O.F.); niltonchleite@gmail.com (N.J.C.L.)
- <sup>2</sup> Sports Science School of Rio Maior, Polytechnic Institute of Santarém, 2040-413 Rio Maior, Portugal; rafaeloliveira@esdrm.ipsantarem.pt (R.O.); jbrito@esdrm.ipsantarem.pt (J.P.B.)
- <sup>3</sup> Life Quality Research Centre, 2040-413 Rio Maior, Portugal; ana.fatima.pereira@ese.ips.pt
- <sup>4</sup> School of Education, Polytechnic Institute of Setúbal, 2910-761 Setúbal, Portugal
- <sup>5</sup> Research Centre in Sport Sciences, Health Sciences and Human Development, 5001-801 Vila Real, Portugal
- <sup>6</sup> Department of Physical Education and Sports, University of Granada, 18010 Granada, Spain; francod.agoni@gmail.com
- \* Correspondence: af\_martins17@hotmail.com



**Citation:** Martins, A.D.; Fernandes, O.; Pereira, A.; Oliveira, R.; Alderete Goñi, F.D.; Leite, N.J.C.; Brito, J.P. The Effects of High-Speed Resistance Training on Health Outcomes in Independent Older Adults: A Systematic Review and Meta-Analysis. *Int. J. Environ. Res. Public Health* **2022**, *19*, 5390. <https://doi.org/10.3390/ijerph19095390>

Academic Editor: Paul B. Tchounwou

Received: 29 March 2022

Accepted: 27 April 2022

Published: 28 April 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

**Abstract:** Human ageing involves several physiological impairments—in particular, a decrease in sensorimotor function and changes in the nervous system reduce muscle strength, power, balance, and functional capacity performance. Preventive strategies are essential to ensure the quality of life of the elderly. High-speed resistance training (HSRT) may be an effective approach to muscle power development in this population, with significant short-term effects on neural adaptations and muscle power production. Therefore, the present study intends to analyze and systematize the studies focused on HSRT interventions and their effects on health outcomes in independent older adults. Four electronic databases (PubMed, Web of Science, EBSCO, and Scielo) were used for the purposes of searching randomized controlled trials that measured at least one key outcome measure focusing on velocity-based training and health outcomes in older adults on 7 March 2022 and identified 1950 studies. At the end of the process, fourteen studies were included in this systematic review and ten studies were included in the quantitative analysis. The main results showed that HSRT interventions would improve health measures, mostly cognitive function (large effects,  $p = 0.001$ , SMD = 0.94), neuromuscular function (moderate effects,  $p = 0.003$ , SMD = 0.70), and physical function (moderate effects,  $p = 0.04$ , SMD = 0.55 and  $p = 0.009$ , SMD =  $-0.59$ ). Additionally, the results suggested that interventions with ten weeks or more, performed three times a week, provide significant improvements in neuromuscular function. In this sense, HSRT is effective for improving overall health outcomes in older adults. Future studies should include proper follow-ups (e.g., minimum six months) to assess the durability of HSRT intervention effects on all health-related variables.

**Keywords:** older people; high-speed resistance training; power training; neuromuscular function; health measures

## 1. Introduction

In accordance with the latest data report from Eurostat, approximately 20% of the total population in Europe is over 65 years old, and by 2050, this figure will rise to around 30% [1]. Human ageing involves several physiological impairments [2,3]. In particular, the decrease in the sensorimotor function and changes in the nervous system [4] reduce muscle strength, power, balance, and functional capacity performance [5,6], which can increase the risk of falling [7]. Falls are the second leading cause of unintentional injury-related

deaths in the world population, and are becoming a serious public health problem for the elderly [8]. The United Kingdom spends an estimated 2.3 billion dollars a year in the treatment of injuries caused by falls [9]. Preventive strategies are essential to ensure the quality of life of the elderly and reduce healthcare costs, which threaten the stability of the public health system.

Falling or the fear of falling may impair physical activity levels and set in motion a vicious cycle in which the risk of future falls increases due to the deterioration of functional ability caused by not actively participating in daily activities [10,11]. The reduction in physical activity levels can have negative physiological effects for an ageing population (e.g., increased chronic diseases and the risk of non-communicable diseases) [12].

Due to reduced levels of physical activity and ageing, loss of muscle strength is associated with a decrease in the performance of daily life activities and negatively affects health, functional autonomy, survival, and quality of life, further increasing the risk of falling [13,14]. In this regard, the loss of muscle strength may be more closely associated with the impairment of muscle power than muscle mass [15–18]. On the other hand, muscle power has been reported to decline faster with ageing when compared to muscle strength [17,19].

High-speed resistance training (HSRT) may be an effective approach to muscle power development in this population [20–25], with significant short-term effects on neural adaptations and muscle power production [15,25,26]. Several studies [24,27–29] suggest that the above-mentioned intervention has an impact on the power generation capacity of older adults. In this sense, the HSRT approach is a key intervention strategy to counteract the harmful effects caused by ageing [28,30–33]. Some studies [34,35] reported that this method also makes it possible to mitigate capillary deterioration resulting from ageing, especially in the lower limbs. A suitable capillarization can improve protein synthesis metabolism by ensuring the transport of amino acids and growth factors to muscle fibers—in particular, growth factor-1 (IGF-1), hepatocyte growth factor (HGF), interleukin-6 (IL-6), and myostatin—thereby regulating satellite cell function and supporting muscle repair and/or remodeling/adaptation, which are essential after 65 years of age [36,37]. In addition, this intervention can offer advantages concerning other traditional resistance training (RT) interventions, mainly when the participants perform the concentric muscle contraction of each exercise repetition as rapidly as possible, showing improvements in morphological and neural adaptations [25,38] and functional capacity performance [39].

To our knowledge, solely two systematic reviews studied the effects of interventions of HSRT on health outcomes in older people, such as functional capacity [40] and power outcomes [41]. However, said studies were conducted in participants aged over 60, compared the effects caused by fast-intended velocity vs. moderate and low-velocity resistance training, and did not compare the RT program's effects on the experimental group (EG) vs. the control group (CG). Therefore, it is relevant to carry out a systematic review that summarizes the effects of HSRT interventions on a population over 65 years of age and compare the EG and the CG in the most diverse parameters related to health status, such as body composition parameters, phase angle, heart rate variability, balance, and gait variability. Another significant criterion used in this review is that it only includes independent participants (i.e., older people that are able to perform activities of daily living independently, notably eating, bathing, and mobility) [42,43].

Therefore, the present review intends to analyze and systematize the studies focused on HSRT interventions and their effects on health outcomes in independent older adults. In this regard, health outcomes are defined as events that occur with an end result after an intervention. These outcomes can be measured clinically through physical examinations, laboratory and imaging tests, self-reported through questionnaires, or observed, such as fluctuations in gait pattern [44]. Thus, the present review study and meta-analysis sought to systematize the effects of HSRT interventions on health outcomes. Finally, we hypothesized that the HSRT interventions would improve health status parameters and that

longer-duration HSRT interventions provide more significant benefits than short-duration HSRT interventions.

## 2. Materials and Methods

This systematic review and meta-analysis was conducted in accordance with the items of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) [45]. This review was also protocolled at the International Prospective Register of Systematic Reviews (PROSPERO), registered under number CRD42021272242.

### 2.1. Search Strategy and Data Sources

The following electronic databases were used to search for relevant publications on 7 March 2022: PubMed, Web of Science (WoS), Psychology and Behavioral Sciences Collection (EBSCO), and SciELO Citation Index (SciELO CI). In order to include the most recently published studies in the review, we set citation notifications for all these databases, but no restriction to date of publication was applied.

The data search was performed by using the U.S. National Library of Medicine's Medical Subject Headings terms and English language terms related to HSRT and the elderly population. The following word combinations were used in all databases: ("elderly" OR "older people" OR "older adult" OR "older subject") AND ("high-speed resistance training" OR "high-speed strength training" OR "high-speed power training" OR "power-training" OR "velocity-based training" OR "rapid-strength" OR "explosive-strength"). It was possible to perform 28 different searches in each of the mentioned databases (112 in total) through these word combinations. Additionally, the reference lists of the studies retrieved were manually searched to identify potentially eligible studies not captured by the electronic searches (1 study was found).

### 2.2. Eligibility Criteria

The inclusion of studies for revision was decided by consensus between the first and last authors. In case of disagreement regarding the inclusion of an article, the second author was consulted. The selected studies were then filtered according to the following inclusion criteria: (1) studies included participants with age  $\geq 65$  years old, from both sexes, without any clinical condition and physically independent; (2) at least one of the protocols under study was an HSRT program with maximal concentric velocity (i.e., as fast as possible,  $\leq 1$  s) lasting 2 weeks or longer; (3) studies included pre- and post-intervention measurements; (4) the study compared the EG with the CG at post-intervention; and (5) only original and full-text studies written in English, Portuguese, or Spanish.

Trials were excluded if: (1) studies included participants with age  $< 65$  years old; (2) participants performed an intervention other than HSRT; (3) did not compare the results of the EG with the CG; (4) did not clearly describe the concentric and/or eccentric velocity; (5) examined of the effects of combined training methods (i.e., fast-intended and moderate-velocity or balance training); (6) included participants with any associated disease (i.e., cancer, dementia, diabetes) or non-physically independent; (7) written in a language other than English, Portuguese, or Spanish; and (8) other article types than original (e.g., reviews, letters to editors, trial registrations, proposals for protocols, editorials, book chapters, and conference abstracts).

### 2.3. Data Extraction

All articles identified by the search strategy underwent an evaluation of the titles and abstracts, in duplicate, by two assigned researchers. All studies that did not meet the inclusion criteria were excluded. Abstracts that did not provide sufficient information regarding the inclusion and exclusion criteria were selected for full-text evaluation. In a second phase, the same two assigned researchers independently evaluated all selected full-text articles and conducted a second selection following the inclusion and exclusion

criteria. In a meeting with a third researcher present, disagreements between reviewers were solved by consensus.

Data extraction from the selected full-text articles was conducted by both assigned researchers independently, using a list of intended data: (i) first author; (ii) year of publication; (iii) the sample (size and age); (iv) intervention details—duration, frequency, intensity, volume (sets and repetitions), rest interval, concentric and eccentric velocity, and type of exercises; (v) control group characteristics; and (vi) main results (health status).

#### 2.4. Methodological Quality Assessment

The methodological study quality was assessed using the Physiotherapy Evidence Database (PEDro) scale [46], a valid and reliable instrument to assess eligibility, group allocation, blinding of allocation, and comparison between groups at baseline and its outcomes. This scale includes 11 questions with yes or no answers (yes = 1; no = 0), providing a total score that ranges between 0 (poor methodological quality) and 10 (excellent methodological quality) (the first item is not included in the rating).

Scores were obtained from the PEDro database and were therefore scored independently, avoiding any potential bias of the authors. When a study was not available on the PEDro database, two authors alone (A.D.M. and R.O.) rated the risk of bias. Disagreements between authors were solved by consensus in a meeting with the last author present (J.P.B.).

#### 2.5. Level of Evidence

Based on the physiotherapy evidence database scale and to assess the interventions' evidence, the Van Tulder criteria [47] were applied. Therefore, the selected studies were grouped by levels of evidence, according to their methodological quality. A study with a physiotherapy evidence database score of 6 or more is considered level 1 (high methodological quality) (6–8: good, 9–10: excellent) and a score of 5 or less is considered level 2 (low methodological quality) (4–5: moderate; <4: poor).

Due to the clinical and statistical heterogeneity of the results, a qualitative review was performed, conducting a best-evidence synthesis [48,49]. This classification indicates that if the number of studies displaying the same level of evidence for the same outcome measure or equivalent is lower than 50% of the total number of studies found, no evidence can be concluded regarding any of the methods involved in the study.

#### 2.6. Statistical Analysis

When the same variable was found in more than two articles, it was included in the meta-analysis. In this sense, the studies included in this analysis ( $n = 10$ ) were analyzed using the random effects model [50]. The forest plot was generated for some health outcomes. Studies' heterogeneity was assessed by calculating the following statistics: (i)  $\tau^2$ , (ii)  $\chi^2$ , and (iii)  $I^2$ . The following classification was used to evaluate the  $I^2$  (i.e., described inconsistency between trials): lower than 50% represents low heterogeneity; 50–74% represents substantial heterogeneity; and 75% and higher represents considerable heterogeneity [51].

The following scale was used to evaluate the standardized mean differences (SMDs) (i.e.,  $SMDs = ([\text{mean post-value intervention group} - \text{mean post-value control group}] / \text{pooled variance})$  [52], according to the following thresholds: 0–0.19 = negligible effect; 0.20–0.49 = small effect; 0.50–0.79 = moderate effect; and 0.80 and higher = large effect.


All statistical analyses were performed within the Cochrane Review Manager (RevMan) [Computer program], version 5.4.1, The Cochrane Collaboration, 2020 [53].

### 3. Results

#### 3.1. Studies Included

A total of 1950 studies were retrieved from the selected databases and one study was found in other sources, using the chosen keywords. These studies were then exported to reference manager software (EndNote™ 20.0.1, Clarivate Analytics, Philadelphia, PA,

USA). Duplicates (1540 references) were subsequently removed either automatically or manually. The remaining 411 articles were screened for their relevance based on titles and abstracts, resulting in the removal of a further 209 studies. At the end of the screening procedure, we had 202 eligible articles, which were read and analyzed in depth. After reading full texts, a further 188 studies were excluded due to not meeting the eligibility criteria (Figure 1). A total of fourteen studies were identified as meeting the criteria for inclusion and were assessed for quality using the PEDro scale. Finally, ten of these articles were included in the quantitative analysis (meta-analysis).



**Figure 1.** Flow chart of study design by PRISMA 2020.

### 3.2. Methodological Quality

The methodological assessment of the fourteen studies included can be found in Table 1. These studies obtained a score between 3 [54–57] and 6 [58,59] on the PEDro scale. From all studies

reviewed, two studies presented high methodological quality (level 1) [58,59], and twelve studies showed low methodological quality (level 2) [54–57,60–67]. Furthermore, the results of the PEDro scale showed that five studies did not specify the inclusion and exclusion criteria [55–57,65,66], three studies did not randomly allocate the participants into groups [54–56], two studies did not perform a baseline comparability [57,60], and all studies allocated the participants into groups in a concealed way.

**Table 1.** Analysis of the risk of bias of the studies included in this systematic review.

Authors	PEDro Scale											Total Score	Methodological Quality
	1	2	3	4	5	6	7	8	9	10	11		
Fang et al. [60] §	Y	1	0	0	0	0	0	1	0	1	1	4	Moderate
Lee et al. [63]	Y	1	0	1	0	0	0	1	0	1	1	5	Moderate
Yoon et al. [67]	N	1	0	1	0	0	0	0	0	1	1	4	Moderate
Yoon et al. [66]	Y	1	0	1	0	0	0	0	0	1	1	4	Moderate
Beijersbergen et al. [54] §	Y	0	0	1	0	0	0	1	0	0	1	3	Poor
Beijersbergen et al. [55] §	N	0	0	1	0	0	0	1	0	0	1	3	Poor
Beijersbergen et al. [56] §	N	0	0	1	0	0	0	1	0	0	1	3	Poor
Hvid et al. [61]	Y	1	0	1	0	0	1	0	0	1	1	5	Moderate
Sayers & Gibson [65]	N	1	0	1	0	0	0	0	0	1	1	4	Moderate
Marsh et al. [64]	Y	1	0	1	0	0	0	0	0	1	1	4	Moderate
Reid et al. [59]	Y	1	0	1	0	0	0	1	1	1	1	6	Good
Katula et al. [62]	Y	1	0	1	0	0	0	0	0	1	1	4	Moderate
Bean et al. [58]	Y	1	0	1	0	0	1	1	0	1	1	6	Good
Miszko et al. [57]	N	1	0	0	0	0	0	0	0	1	1	3	Poor

Abbreviations: 1, Eligibility; 2, Random allocation; 3, Concealed allocation; 4, Baseline comparability; 5, Blind subjects; 6, Blind therapists; 7, Blind assessors; 8, Adequate follow-up; 9, Intention-to-treat analysis; 10, Between-group comparisons; 11, Point estimates and variability; Y, yes; N, No; §, Scored by reviewers. Note: Eligibility criteria item does not contribute to total score.

Additionally, none of the studies included blinded the participants and the technicians responsible for the program's sessions, two studies blinded the technicians who measured at least one key outcome [58,61], seven studies could not perform assessments on at least one key outcome from more than 85% of subjects [57,61,62,64–67], only one study performed an intention-to-treat analysis [59], three studies did not report a between-group statistical comparison for at least one key outcome [54–56], and finally, all studies reported both point measures and measures of variability for at least one key outcome.

### 3.3. Studies' Characteristics

Table 2 shows the characteristics of the studies included in this systematic review in terms of country, sample, age, body fat, body mass, and BMI. Most of the included studies did not register the intervention, and only one study reported the protocol [61]. Two studies [58,66] only included females in their analysis and no study was conducted with only men.

Regarding the origin of the studies, six studies were conducted in the United States (North of America) [58–60,62,64,65], three were performed in Germany (Europe) [54–56], three were conducted in South Korea (Asia) [63,66,67], and the last two articles were carried out in two different countries on the same mainland (Europe)—Denmark [61] and Greece [57]. Fourteen studies included in this review randomized 408 participants into two groups; thus, CG included 199 participants and EG (were performed HSRT) included 209 participants. These participants ranged in age from 69 years old [54] to 82 years old [61].

**Table 2.** Characteristics of included studies.

Authors	Country	Sample by Gender (N)	Age (years) (M $\pm$ SD)	Groups (N)	Body Fat (%)	Body Mass (kg)	BMI (kg/m <sup>2</sup> )	Registered Protocol?
Fang et al. [60]	USA	9 F 11 M	CG = 71.8 $\pm$ 4.8 EG = 70.8 $\pm$ 5.8	CG = 8 EG = 12	NR	NR	NR	No
Lee et al. [63]	South Korea	24 F 16 M	CG = 74.2 $\pm$ 4.4 EG = 73.7 $\pm$ 4.6	CG = 22 EG = 18	NR	CG = 60.6 EG = 61.0	NR	No
Yoon et al. [67]	South Korea	30 F 13 M	CG = 74.0 $\pm$ 4.3 EG = 73.9 $\pm$ 4.4	CG = 23 EG = 22	NR	NR	CG = 24.4 EG = 24.9	No
Yoon et al. [66]	South Korea	30 F	CG = 78.0 $\pm$ 1.0 EG = 75.0 $\pm$ 0.9	CG = 7 EG = 14	CG = 33.4 EG = 38.8	CG = 51.2 EG = 58.4	CG = 22.9 EG = 25.5	No
Beijersbergen et al. [54]	Germany	25 F 16 M	CG = 69.1 $\pm$ 4.4 EG = 72.9 $\pm$ 5.4	CG = 14 EG = 15	NR	CG = 73.9 EG = 73.6	CG = 25.5 EG = 25.8	No
Beijersbergen et al. [55]	Germany	29 *	CG = 69.1 $\pm$ 4.4 EG = 72.9 $\pm$ 5.4	CG = 14 EG = 15	NR	CG = 73.9 EG = 73.6	CG = 25.5 EG = 25.8	No
Beijersbergen et al. [56]	Germany	25 *	CG = 69.7 $\pm$ 5.0 EG = 72.1 $\pm$ 5.4	CG = 13 EG = 12	NR	NR	CG = 25.1 EG = 26.2	No
Hvid et al. [61]	Denmark	23 F 14 M	CG = 81.6 $\pm$ 1.1 EG = 82.3 $\pm$ 1.3	CG = 21 EG = 16	NR	CG = 73.4 EG = 76.5	NR	Yes
Sayers and Gibson [65]	USA	24 F 14 M	CG = 72.8 $\pm$ 4.1 EG = 74.1 $\pm$ 6.4	CG = 12 EG = 13	NR	CG = 78.6 EG = 76.6	NR	No
Marsh et al. [64]	USA	25 F 11 M	CG = 74.4 $\pm$ 5.2 EG = 76.8 $\pm$ 6.4	CG = 15 EG = 15	NR	CG = 81.0 EG = 81.2	CG = 30.4 EG = 30.7	No
Reid et al. [59]	USA	31 F 26 M	CG = 79.7 $\pm$ 9.0 EG = 72.3 $\pm$ 6.0	CG = 12 EG = 23	NR	CG = 70.2 EG = 80.9	CG = 26.5 EG = 29.8	No
Katula et al. [62]	USA	36 *	CG = 74.3 $\pm$ 5.4 EG = 76.8 $\pm$ 6.5	CG = 13 EG = 12	NR	CG = 81.7 EG = 81.2	CG = 30.7 EG = 30.8	No
Bean et al. [58]	USA	21 F	CG = 78.9 $\pm$ 7.8 EG = 77.1 $\pm$ 5.7	CG = 10 EG = 11	NR	CG = 65.6 EG = 60.1	CG = 26.4 EG = 26.2	No
Miszko et al. [57]	Greece	22 F 17 M	CG = 72.4 $\pm$ 7.2 EG = 72.3 $\pm$ 6.7	CG = 15 EG = 11	CG = 26.8 EG = 29.1	CG = 68.2 EG = 79.7	NR	No

Abbreviations: N, number; M, mean; SD, standard deviation; CG, control group; EG, experimental group; F, female; M, male; USA, united states of America; BMI, body mass index; kg, kilograms; NR, not reported. \*, date unreported gender data.

### 3.4. Interventions' Characteristics

Table 3 presents the description of the interventions performed in the included studies, namely exercise modality, names of exercises, weekly frequency, intensity, sets, reps, duration of the intervention and sessions, eccentric velocity, and whether the sessions were supervised or not.

The duration of the RT interventions ranged from eight [60,63] to sixteen weeks [57,67], with two studies reporting for eight weeks [60,63], three reporting for ten weeks [54–56], seven reporting for twelve weeks [58,59,61,62,64–66], and two reporting for sixteen weeks [57,67]. Concerning the duration of sessions, these varied from 30 [58] to 60 min [64,66,67]; the intensity of the sessions mainly was quantified through the percentage of repetition maximum (RM) [57,59–62,64,65]. For the majority of the studies, the sessions were performed three times a week [54–60,62–65,67].





**Table 3.** Description of the interventions performed in the included studies.

Authors	Exercise Modality	Exercises (Names)	Frequency (Days/Week)	Intensity	Sets/Exercise (N)	Reps per Set (N)	Rest	Intervention Duration (weeks)	Session Duration (min)	Eccentric Velocity (s)	Supervised
Fang et al. [60]	Computerized Pneumatic Machines	Leg press; seated row; leg curl; chest press; hip adduction; lat pulldown; hip abduction; tricep extension; seated calf raise; bicep curl	3	40% 1 RM—lat pulldown 50% 1 RM—leg press, chest press, seated row 55% 1 RM—bicep curl 60% 1 RM—leg curl, tricep extension 65% 1 RM—calf raise, hip abduction, hip adduction	1 (wk 1) 2 (wk 2) 3 (wk 3–8)	10–12	1–2 min after each circuit	8	40–45	2	Yes
Lee et al. [63]	Elastic Bands	Seated row; one-leg press; applied pec deck flus or lateral raises; seated leg raises; squats; full squats; wide squats; bridging	3	12 to 13 from 6–20 RPE scale	2–3	10–12	30 s between exercises 1 min between sets	8	50	3	Yes
Yoon et al. [67]	Elastic Bands	Seated row; one leg press; applied pec deck flus; seated leg raises; lateral raise; semi-squats; wide squats; and bridging	3	12 to 13 from 6–20 RPE scale	2–3	12–15	NR	16	60	2	Yes
Yoon et al. [66]	Elastic Bands	NR	2	12 to 13 from 6–20 RPE scale	2–3	12–15	2 min between exercises 1 min between sets	12	60	2	Yes
Beijersbergen et al. [54]	Weightlifting Machine	Leg press; ankle press; knee extension; knee flexion	3	40–60% 3 RM	3	6–10	NR	10	NR	Normal pace	NR
Beijersbergen et al. [55]	Weightlifting Machine	Leg press; ankle press; knee extension; knee flexion.	3	40–60% 3 RM	3	6–10	NR	10	NR	Normal pace	NR
Beijersbergen et al. [56]	Weightlifting Machine	Leg press; ankle press; knee extension; knee flexion	3	40–60% 3 RM	3	6–10	NR	10	NR	Normal pace	NR
Hvid et al. [61]	Weightlifting Machine	NR	2	70–80% 1 RM	3	10 (wk 1–6) 8 (wk 7–12)	NR	12	NR	2–3	Yes
Sayers and Gibson [65]	Computerized Pneumatic Machines	Leg press; seated knee extension	3	40% 1 RM	3	12–14	NR	12	NR	2	Yes
Marsh et al. [64]	Computerized Pneumatic Machines	Leg press; knee extension	3	70% 1 RM	3	8–10	NR	12	60	2–3	Yes
Reid et al. [59]	Computerized Pneumatic Machines	Leg press; knee extension	3	70% 1 RM	3	8	NR	12	NR	>2	Yes
Katula et al. [62]	Computerized Pneumatic Machines	Leg press; knee extension	3	70% 1 RM	3	8–10	NR	12	NR	2–3	Yes
Bean et al. [58]	Weighted Vest	Chair stands; toe raises; pelvic raises; step ups; seated tricep dips; chest press	3	At 16 from 6–20 RPE scale	3	10	1–2 min between sets	12	30	>2	Yes
Miszko et al. [57]	Computerized Pneumatic Machines	Seated row; chest press; tricep extension; leg press; leg extension; seated leg curl; bicep curls; plantar flexion; jump squats	3	40–70% 1 RM	3	8–10	NR	16	NR	2	NR

Abbreviations: N, number; min, minutes; s, seconds; wk, week; RM, repetition maximum; RPE, rated perceived exertion; NR, not reported.

The research seems to be relatively diverse regarding the number of sets and repetitions. The number of sets performed in the interventions was as follows: ten interventions used three sets [54–59,61,62,64,65], three interventions used two to three sets [63,66,67], and only one intervention used a non-fixed number of sets [60]. Regarding the number of repetitions used per set of exercises, three studies used six to ten [54–56], three studies used eight to ten [57,62,64], two studies used ten to twelve [60,63], one study used twelve to fourteen [65], two studies used twelve to fourteen [66,67], one study only used eight [59], one study only used ten [58], and one study used a non-fixed number of repetitions [61].

The exercise interventions of the fourteen studies included in this systematic review were performed using elastic bands [63,66,67], in a weightlifting machine [54–56,61], using computerized pneumatic machines [57,59,60,62,64,65], and using a weighted vest [58].

Ten studies reported the dropout rates of their respective participants [54,56–58,61–66], with two studies stating that no participants dropped out of the EG [58,64]. Furthermore, seven studies presented the dropout rate of its EG with further explanations [54,56,57,61,63,65,66] and one study provided no explanation for said dropout [62]. Two studies did not report adverse events in both groups [58,63]. However, three studies reported several negative events in both groups, notably falls that did not result in fractures (three cases in EG and CG), and temporary localized joint pain (three cases in EG and one case in CG) [64]. One of the three studies mentioned above reported sacral-iliac pain following completion of baseline testing in CG, as well as persistent chest pain in EG [59], and another study reported the occurrence of serious medical events [66]. Additionally, some participants from the groups mentioned above withdrew from the studies due to the diagnosis of several types of tumors and aortic aneurism [59,64]. Finally, supervised exercise interventions were used in most studies (Table 3).

### 3.5. Results of the Studies

Table 4 presents the aim, outcome variable, and main results of the fourteen studies in this review. Only outcomes addressed in more than three studies will be described here.

Physical function was analyzed in seven studies [54,57,58,61,63,66,67]. Four studies showed improvements in physical function in only the EG between pre- and post-training [58,61,63,67]. Outcomes of muscle strength, such as grip strength, were assessed in six studies [56,57,61,64,66,67]; three out of seven studies [61,64,67] reported differences between CG and EG at post-training in these outcomes ( $p < 0.05$ ). Lastly, outcomes of muscle power were described in four studies [54,55,59,65]; of these four studies, two [59,65] demonstrated significant differences between CG and EG at post-training in peak power outcomes at different %RM (40 to 90% 1 RM) ( $p < 0.05$ ).

**Table 4.** Main results of the included studies.

Authors	Aim	Outcome Variable	Group	Main Results (M ± SD)		p Value	p Value (CG vs. EG) &	Effect Size <sup>a</sup>
				Pre	Post			
Fang et al. [60]	To characterize changes in the retinal microvascular density and their relationship with cognitive function in cognitively normal older people	Outcomes of optical function						
		Vessel density in the total retinal vascular network	CG	1.79	1.79	>0.05	>0.05	NR
			EG	1.79	1.79	>0.05		NR
		Vessel density in the superficial vessel plexus	CG	1.79	1.78	>0.05	>0.05	NR
			EG	1.78	1.78	>0.05		NR
		Vessel density in the deep vessel plexus	CG	1.80	1.80	>0.05	>0.05	NR
			EG	1.80	1.80	>0.05		NR
		Outcome of cognitive function						
		MMSE (score)	CG	30 ± 0	29 ± 1	<b>0.08</b>	NR	NR
			EG	30 ± 1	30 ± 1	0.59		NR
Lee et al. [63]	To investigate the effects of HSRT on neuromuscular, executive, and gait performance	Outcomes of neuromuscular function						
		Movement time (ms)	CG	906.36 ± 36.27	1010.04 ± 65.89	>0.05	<0.05	NR
			EG	921.69 ± 40.10	799.51 ± 72.84	<0.05		NR
		Pre-motor time (ms)	CG	700.64 ± 36.62	757.34 ± 56.37	>0.05	>0.05	NR
			EG	663.82 ± 40.48	626.96 ± 62.32	>0.05		NR
		Motor time (ms)	CG	205.67 ± 17.45	252.70 ± 34.44	>0.05	<0.05	NR
			EG	271.40 ± 19.29	181.15 ± 38.08	<0.05		NR
		Antagonist co-activation (% p. EMG)	CG	25.28 ± 3.75	25.04 ± 3.72	>0.05	>0.05	NR
			EG	27.89 ± 4.14	26.32 ± 3.48	>0.05		NR
		Rate of EMG rise (% p. EMG/s <sup>-1</sup> )	CG	169.30 ± 12.04	161.82 ± 10.41	>0.05	<0.05	NR
			EG	166.48 ± 13.31	197.94 ± 11.51	<0.05		NR
		Normalized peak torque (N·m·kg <sup>-1</sup> )	CG	NR	NR	>0.05	>0.05	NR
			EG	NR	NR	<0.05		NR
		Rate of torque development	CG	NR	NR	>0.05	<0.05	NR
			EG	NR	NR	<0.01		NR
		Outcome of cognitive function						
FAB (score)	CG	11.72 ± 2.11	11.77 ± 2.22	>0.05	>0.05	NR		
	EG	11.38 ± 2.56	12.16 ± 1.79	>0.05		NR		
Outcomes of physical function								
4.44 m gait speed (s)	CG	NR	NR	>0.05	>0.05	NR		
	EG	NR	NR	<0.01		NR		
TUG (s)	CG	NR	NR	>0.05	>0.05	NR		
	EG	NR	NR	<0.01		NR		

Table 4. Cont.

Authors	Aim	Outcome Variable	Group	Main Results (M ± SD)		p Value	p Value (CG vs. EG) &	Effect Size <sup>a</sup>		
				Pre	Post					
Outcomes of cognitive function										
Yoon et al. [67]	To elucidate the effects of high-speed resistance exercise on cognitive function and physical performance	Memory (score)	CG	10.26 ± 2.85	10.52 ± 2.79	>0.05	0.445	0.09		
			EG	8.55 ± 2.39	10.00 ± 3.71	>0.05		0.46 #		
		Processing speed (s)	CG	43.04 ± 11.95	42.59 ± 15.92	>0.05	0.036	0.03		
			EG	54.15 ± 28.43	48.26 ± 27.33	<0.05		0.21 #		
		Cognitive flexibility (s)	CG	188.92 ± 81.38	187.20 ± 70.14	>0.05	0.532	0.45 #		
			EG	163.37 ± 62.45	140.82 ± 34.65	>0.05		0.02		
		Working memory (score)	CG	10.09 ± 2.04	10.39 ± 1.83	>0.05	0.448	0.14		
			EG	10.20 ± 1.54	10.70 ± 1.34	>0.05		0.35 #		
		FAB (score)	CG	11.87 ± 2.12	12.09 ± 2.00	>0.05	0.022	0.11		
			EG	12.00 ± 2.45	13.70 ± 2.11	<0.05		0.74 *		
		Outcomes of physical function								
		SPBB (score)	CG	10.04 ± 1.46	10.91 ± 1.20	>0.05	0.001	0.65 *		
			EG	9.25 ± 2.31	10.85 ± 1.60	<0.05		0.81 *		
		TUG (s)	CG	9.95 ± 1.51	9.89 ± 1.59	>0.05	<0.01	0.04		
			EG	10.66 ± 2.41	9.26 ± 2.03	<0.01		0.65 *		
		4.44 m gait speed (s)	CG	6.04 ± 0.82	5.58 ± 0.81	>0.05	0.027	0.56 #		
			EG	6.21 ± 1.04	5.34 ± 0.81	<0.01		0.93 *		
		Outcomes of neuromuscular function								
Grip strength (kg)	CG	21.81 ± 6.31	23.78 ± 7.14	>0.05	0.020	0.29 #				
	EG	21.41 ± 6.58	23.60 ± 7.76	<0.05		0.30 #				
Isokinetic 60°/s peak torque/BW	CG	70.77 ± 24.32	64.23 ± 20.72	>0.05	0.004	0.01				
	EG	65.05 ± 25.82	71.20 ± 36.68	<0.05		0.19				
Isokinetic 180°/s average power per rep (W)	CG	72.77 ± 23.82	66.59 ± 23.67	>0.05	0.001	0.26 #				
	EG	68.32 ± 40.60	82.09 ± 44.63	<0.05		0.32 #				

Table 4. Cont.

Authors	Aim	Outcome Variable	Group	Main Results (M ± SD)		p Value	p Value (CG vs. EG) &	Effect Size <sup>a</sup>		
				Pre	Post					
Outcomes of body composition										
Yoon et al. [66]	To compare the effects of two different exercise regimens on cognitive function, body composition, muscular strength, and functional ability	BW (kg)	CG	51.19 ± 4.13	50.14 ± 3.75	<0.01	0.001	NR		
			EG	58.39 ± 6.82	57.50 ± 7.01	<0.01		NR		
		BMI(kg/m <sup>2</sup> )	CG	22.90 ± 1.81	22.54 ± 1.71	>0.05	0.005	NR		
			EG	25.46 ± 2.47	25.02 ± 2.35	<0.01		NR		
		Skeletal muscle mass (kg)	CG	17.11 ± 1.45	17.49 ± 1.35	>0.05	0.008	NR		
			EG	18.84 ± 2.84	19.63 ± 2.63	<0.01		NR		
		Percent body fat (%)	CG	33.39 ± 8.26	33.27 ± 3.35	>0.05	0.004	NR		
			EG	38.83 ± 5.37	35.50 ± 4.72	<0.01		NR		
		WHR	CG	0.88 ± 0.04	0.86 ± 0.03	>0.05	0.014	NR		
			EG	0.91 ± 0.07	0.89 ± 0.06	<0.05		NR		
		Arm circumference (cm)	CG	26.71 ± 2.75	26.29 ± 2.69	>0.05	0.494	NR		
			EG	29.21 ± 2.91	28.04 ± 2.32	>0.05		NR		
		Thigh circumference (cm)	CG	49.14 ± 4.60	47.57 ± 4.71	>0.05	0.115	NR		
			EG	51.14 ± 4.00	50.57 ± 4.69	>0.05		NR		
		Outcomes of cognitive function								
		MMSE (score)	CG	22.29 ± 1.11	21.14 ± 1.57	<0.05	<0.05	-0.85		
EG	21.00 ± 1.04		25.36 ± 1.78	<0.01	2.99 §					
MoCA (score)	CG	18.71 ± 2.63	18.14 ± 2.97	>0.05	<0.05	0.19				
	EG	18.29 ± 2.81	24.29 ± 2.58	<0.01		2.22 §				
Outcomes of physical function										
SPPB (score)	CG	7.14 ± 1.77	7.57 ± 0.98	>0.05	<0.05	0.30 #				
	EG	8.14 ± 2.48	10.79 ± 1.58	<0.01		1.27 §				
TUG (s)	CG	11.48 ± 1.02	10.59 ± 1.03	>0.05	>0.05	-0.87				
	EG	0.51 ± 1.86	9.14 ± 1.42	>0.05		-0.83				

Table 4. Cont.

Authors	Aim	Outcome Variable	Group	Main Results (M ± SD)		p Value	p Value (CG vs. EG) &	Effect Size <sup>a</sup>		
				Pre	Post					
Outcomes of neuromuscular function										
Beijersbergen et al. [54]	To determine the effects of lower-extremity power training and detraining on lower-limb muscle power and gait kinematics	Grip strength (kg)	CG	17.69 ± 0.91	18.99 ± 1.81	>0.05	<0.05	1.86 §		
			EG	19.26 ± 3.57	24.01 ± 4.14	<0.01		4.58 £		
		Isokinetic 60°/s peak torque/BW—Right extensor (Nm)	CG	105.71 ± 20.92	116.86 ± 31.74	>0.05	>0.05	0.41 #		
			EG	77.57 ± 29.03	115.43 ± 31.92	<0.01		1.24 §		
		Isokinetic 60°/s peak torque/BW—Right flexor (Nm)	CG	74.86 ± 17.44	58.14 ± 18.00	<0.05	>0.05	−0.94		
			EG	66.21 ± 20.77	67.43 ± 17.82	>0.05		0.06		
		Isokinetic 60°/s peak torque/BW—left extensor (Nm)	CG	96.86 ± 25.42	108.86 ± 27.67	>0.05	>0.05	0.45 #		
			EG	90.36 ± 31.70	90.57 ± 21.99	>0.05		0.01		
		Isokinetic 60°/s peak torque/BW—left flexor (Nm)	CG	69.57 ± 19.84	53.00 ± 16.34	<0.05	>0.05	−0.91		
			EG	64.14 ± 17.22	58.10 ± 17.60	>0.05		−0.35		
		Isokinetic 180°/s peak torque/BW—Right extensor (Nm)	CG	57.29 ± 13.88	67.43 ± 18.60	<0.05	<0.05	0.62 *		
			EG	49.07 ± 20.00	68.29 ± 17.55	<0.01		1.02 *		
		Isokinetic 180°/s peak torque/BW—Right flexor (Nm)	CG	54.43 ± 9.24	37.71 ± 9.93	>0.05	>0.05	−1.74		
			EG	44.14 ± 14.69	36.00 ± 15.11	>0.05		−0.55		
		Isokinetic 180°/s peak torque/BW—Left extensor (Nm)	CG	54.71 ± 14.85	66.57 ± 21.45	<0.05	>0.05	0.64 *		
			EG	50.29 ± 17.82	63.21 ± 19.88	<0.05		0.68 *		
		Isokinetic 180°/s peak torque/BW—Left flexor (Nm)	CG	51.43 ± 13.13	32.57 ± 13.01	<0.01	>0.05	−1.44		
			EG	45.71 ± 14.31	37.22 ± 13.79	<0.05		−0.60		
		Outcomes of neuromuscular function								
				Maximal muscle power of knee extension 60°/s (W)	CG	98.4 ± 39.4	93.6 ± 39.1	0.680	NR	−0.12
EG	97.5 ± 37.7				119.9 ± 43.2	<0.01	0.59 #			
Maximal muscle power of knee extension 120°/s (W)	CG			169.4 ± 61.7	164.7 ± 49.2	0.738	NR	−0.08		
	EG			161.7 ± 64.5	199.1 ± 72.8	<0.01		0.58 #		
Maximal muscle power of knee extension 180°/s (W)	CG			229.2 ± 77.6	234.0 ± 64.9	0.685	NR	0.06		
	EG			216.9 ± 93.5	256.9 ± 96.3	<0.01		0.43 #		
Maximal muscle power of knee flexion 60°/s (W)	CG			54.6 ± 24.6	51.1 ± 20.4	0.557	NR	−0.14		
	EG			54.6 ± 27.7	71.5 ± 37.5	<0.01		0.61 *		
Maximal muscle power of knee flexion 120°/s (W)	CG			103.4 ± 38.9	101.0 ± 35.5	0.794	NR	−0.06		
	EG			104.6 ± 59.0	126.8 ± 71.1	<0.01		0.38 #		
Maximal muscle power of knee flexion 180°/s (W)	CG			156.6 ± 59.0	161.7 ± 50.3	0.541	NR	0.09		
	EG			167.7 ± 92.2	186.3 ± 96.9	0.002		0.20 #		

Table 4. Cont.

Authors	Aim	Outcome Variable	Group	Main Results (M ± SD)		p Value	p Value (CG vs. EG) &	Effect Size <sup>a</sup>		
				Pre	Post					
Beijersbergen et al. [55]	To examine the effects of lower-extremity power training and detraining on gait kinetics	Maximal muscle power of knee plantar flexion 20°/s (W)	CG	13.6 ± 6.7	15.3 ± 6.9	0.095	NR	0.25		
			EG	12.6 ± 8.1	17.3 ± 9.8	<b>0.001</b>		0.57 #		
		Maximal muscle power of knee plantar flexion 40°/s (W)	CG	25.3 ± 11.2	29.1 ± 14.7	0.225	NR	0.34 #		
			EG	23.3 ± 13.6	30.1 ± 16.0	<b>&lt;0.01</b>		0.50 #		
		Maximal muscle power of knee plantar flexion 60°/s (W)	CG	35.8 ± 15.8	36.6 ± 19.6	0.734	NR	0.05		
			EG	32.4 ± 21.1	40.8 ± 23.2	<b>&lt;0.01</b>		0.40 #		
		Outcomes of physical function								
		Stair ascent power (W·kg <sup>-1</sup> )	CG	4.36 ± 0.65	4.41 ± 0.85	0.819	NR	0.07		
			EG	4.05 ± 0.84	4.36 ± 0.92	0.075		0.38 #		
		Stair descent power (W·kg <sup>-1</sup> )	CG	5.01 ± 0.91	5.21 ± 1.18	0.225	NR	0.22 #		
			EG	4.48 ± 0.87	4.88 ± 1.21	0.061		0.46 #		
		Six-min walk test (m/s)	CG	1.26 ± 0.14	1.27 ± 0.14	0.293	NR	0.03		
			EG	1.29 ± 0.14	1.31 ± 0.15	0.252		0.18		
		Gait velocity (habitual speed) (m/s)	CG	1.35 ± 0.14	1.34 ± 0.16	0.652	NR	-0.07		
			EG	1.32 ± 0.16	1.36 ± 0.15	0.220		0.25 #		
		Gait velocity (fast speed) (m/s)	CG	1.97 ± 0.35	1.93 ± 0.31	0.526	NR	-0.18		
			EG	1.85 ± 0.28	1.96 ± 0.38	<b>0.026</b>		0.39 #		
		Outcomes of neuromuscular function								
		Knee extensor power 60 deg/s (W/kg·m)	CG	NR	NR	>0.05	NR	NR		
			EG	NR	NR	<b>&lt;0.05</b>		NR		
		Knee extensor power 120 deg/s (W/kg·m)	CG	NR	NR	>0.05	NR	NR		
			EG	NR	NR	<b>&lt;0.05</b>		NR		
		Knee extensor power 180 deg/s (W/kg·m)	CG	NR	NR	>0.05	NR	NR		
			EG	NR	NR	<b>&lt;0.05</b>		NR		
Plantar flexor power 20 deg/s (W/kg·m)	CG	NR	NR	>0.05	NR	NR				
	EG	NR	NR	<b>&lt;0.05</b>		NR				
Plantar flexor power 40 deg/s (W/kg·m)	CG	NR	NR	>0.05	NR	NR				
	EG	NR	NR	<b>&lt;0.05</b>		NR				
Plantar flexor power 60 deg/s (W/kg·m)	CG	NR	NR	>0.05	NR	NR				
	EG	NR	NR	<b>&lt;0.05</b>		NR				
Outcomes of physical function										
Gait velocity (habitual speed) (m/s)	CG	1.35 ± 0.14	1.34 ± 0.16	0.652	NR	NR				
	EG	1.32 ± 0.16	1.36 ± 0.15	0.220		NR				
Gait velocity (fast speed) (m/s)	CG	1.97 ± 0.35	1.93 ± 0.31	0.526	NR	NR				
	EG	1.85 ± 0.28	1.96 ± 0.38	<b>0.026</b>		NR				

Table 4. Cont.

Authors	Aim	Outcome Variable	Group	Main Results (M ± SD)		p Value	p Value (CG vs. EG) &	Effect Size <sup>a</sup>		
				Pre	Post					
Outcomes of neuromuscular function										
Beijersbergen et al. [56]	To examine the effects of 10 weeks of lower-extremity power training on gait velocity and neuromuscular activation of lower-extremity muscles during level walking	Isometric muscle strength of the knee flexors	CG	NR	NR	>0.05	NR	≤0.31		
			EG	NR	NR	<b>0.002</b>		0.34 #		
		Isometric muscle strength of the knee extensors	CG	NR	NR	<b>0.021</b>	NR	0.50 #		
			EG	NR	NR	<b>&lt;0.01</b>		0.74 *		
		Isometric muscle strength of the plantar flexors	CG	NR	NR	>0.05	NR	NR		
			EG	NR	NR	<b>0.002</b>		0.80 *		
		EMG amplitudes of the knee flexors	CG	NR	NR	>0.05	NR	NR		
			EG	NR	NR	<b>0.004</b>		1.47 §		
		EMG amplitudes of the knee extensors	CG	NR	NR	<b>0.021</b>	NR	NR		
			EG	NR	NR	<b>0.013</b>		0.50 #		
		EMG amplitudes of the plantar flexors	CG	NR	NR	>0.05	NR	NR		
			EG	NR	NR	0.076		0.47 #		
		Outcomes of physical function								
				Gait velocity (habitual speed) (m/s)	CG	1.41 ± 0.19	1.39 ± 0.29	0.240	NR	-0.11
EG	1.28 ± 0.14				1.34 ± 0.17	0.079	0.42 #			
Gait velocity (fast speed) (m/s)	CG			1.91 ± 0.26	1.89 ± 0.26	0.396	NR	-0.07		
	EG			1.82 ± 0.18	1.88 ± 0.23	0.059		0.31 #		
Outcomes of neuromuscular function										
Hvid et al. [61]	To examine the effects of 12 weeks of progressive high-intensity power training on the outcomes of knee extensor voluntary muscle activation and maximal gait speed	Thickness (cm)	CG	2.52 ± 0.11	2.50 ± 0.11	>0.05	>0.05	NR		
			EG	2.69 ± 0.12	2.74 ± 0.15	>0.05		NR		
		Strength (N. m)	CG	101.9 ± 0.1	102.7 ± 8.3	>0.05	<b>&lt;0.05</b>	NR		
			EG	98.9 ± 7.7	113.1 ± 7.5	<b>&lt;0.05</b>		NR		
		Voluntary activation (%)	CG	79.1 ± 2.4	79.0 ± 2.6	>0.05	<b>&lt;0.05</b>	NR		
			EG	78.9 ± 3.5	84.9 ± 2.1	<b>&lt;0.05</b>		NR		
		Outcomes of physical function								
				2-MWT (m/s)	CG	1.06 ± 0.04	1.03 ± 0.04	<b>&lt;0.05</b>	<b>&lt;0.05</b>	NR
EG	1.00 ± 0.06				1.09 ± 0.07	<b>&lt;0.05</b>	NR			



Table 4. Cont.

Authors	Aim	Outcome Variable	Group	Main Results (M ± SD)		p Value	p Value (CG vs. EG) &	Effect Size <sup>a</sup>		
				Pre	Post					
Outcomes of neuromuscular function										
Sayers and Gibson [65]	To determine whether low-load HSPT and traditional high-load, slow-speed resistance training may have differing effects on power output obtained across a range of external resistances and to explore the impact of these RT regimens on the determinants of muscle power	Peak power 40% 1 RM (W)	CG	292.6 ± 165.9	283.5 ± 153.9	NR	<0.05	NR		
			EG	313.2 ± 118.4	383.3 ± 113.3	NR		NR		
		Peak power 50% 1 RM (W)	CG	315.5 ± 180.0	313.9 ± 169.1	NR	≤0.02	NR		
			EG	339.0 ± 138.1	426.6 ± 133.6	NR		NR		
		Peak power 60% 1 RM (W)	CG	315.4 ± 178.8	327.3 ± 180.0	NR	≤0.03	NR		
			EG	350.6 ± 144.1	456.6 ± 136.6	NR		NR		
		Peak power 70% 1 RM (W)	CG	304.4 ± 174.3	320.3 ± 180.5	NR	<0.01	NR		
			EG	342.8 ± 149.6	456.3 ± 150.2	NR		NR		
		Peak power 80% 1 RM (W)	CG	281.4 ± 155.3	302.9 ± 169.7	NR	≤0.02	NR		
			EG	317.2 ± 145.6	457.6 ± 149.3	NR		NR		
		Peak power 90% 1 RM (W)	CG	252.1 ± 150.7	242.4 ± 111.2	NR	≤0.04	NR		
			EG	272.9 ± 154.2	431.3 ± 162.4	NR		NR		
		Velocity at peak power (N·m·s <sup>-1</sup> )	CG	NR	NR	NR	<0.01	NR		
			EG	NR	NR	NR		NR		
Outcomes of neuromuscular function										
Marsh et al. [64]	To compare the effects of lower-extremity power training on muscle strength, physical function, and body composition	Knee extension strength (kg)	CG	25.03 ± 10.03	23.99 ± NR	NR	0.024	NR		
			EG	25.87 ± 11.77	29.33 ± NR	NR		NR		
		Knee extension power (W)	CG	161.46 ± 62.44	143.60 ± NR	NR	0.003	NR		
			EG	148.33 ± 84.50	221.30 ± NR	NR		NR		
		Leg press strength (kg)	CG	93.97 ± 31.17	100.12 ± NR	NR	0.026	NR		
			EG	94.07 ± 34.09	117.82 ± NR	NR		NR		
		Leg press power (W)	CG	185.47 ± 75.64	211.4 ± NR	NR	<0.01	NR		
			EG	171.06 ± 89.04	308.7 ± NR	NR		NR		
		Outcomes of neuromuscular function								
		Reid et al. [59]	To explore the effects of power training on muscle power and strength	Knee extension 1 RM—Δ (W)	CG	NR	NR	NR	<0.01	NR
EG	NR				NR	<0.01	NR			
Knee extension—absolute peak power at 40% 1 RM (W)	CG			NR	NR	NR	≤0.003	NR		
	EG			48 ± 15	62 ± 16	0.002		NR		
Knee extension—absolute peak power at 70% 1 RM (W)	CG			NR	NR	NR	≤0.003	NR		
	EG			77 ± 39	118 ± 54	<0.05		NR		
Leg press 1 RM—Δ (W)	CG			NR	NR	NR	>0.05	NR		
	EG			NR	NR	0.140		NR		
Leg press—absolute peak power at 40% 1 RM (W)	CG			NR	NR	NR	>0.05	NR		
	EG			NR	NR	0.190		NR		
Leg press peak power at 70% 1 RM—Δ (W)	CG			NR	NR	NR	>0.05	NR		
	EG			NR	NR	0.220		NR		


Table 4. Cont.

Authors	Aim	Outcome Variable	Group	Main Results (M ± SD)		p Value	p Value (CG vs. EG) &	Effect Size <sup>a</sup>		
				Pre	Post					
Outcomes of quality of life										
Katula et al. [62]	To compare the effects of strength training and power training to one another and to a wait list control group with respect to changes in quality of life	Self-efficacy for strength (score)	CG	25.07 ± 17.67	34.46 ± 19.88	>0.05	<b>&lt;0.05</b>	0.50 #		
			EG	25.07 ± 17.53	72.16 ± 22.58	<b>&lt;0.05</b>		2.34 £		
		Satisfaction with physical function (score)	CG	−0.73 ± 1.69	−0.35 ± 1.96	>0.05	<b>&lt;0.05</b>	0.21 #		
			EG	−0.85 ± 1.86	1.10 ± 1.20	<b>&lt;0.05</b>		1.21 §		
		Life satisfaction (score)	CG	22.93 ± 6.46	21.46 ± 6.06	>0.05	<b>&lt;0.05</b>	−0.23		
			EG	25.78 ± 7.47	29.25 ± 6.38	<b>&lt;0.05</b>		0.50 #		
Outcomes of physical function										
Bean et al. [58]	To evaluate the efficacy of another form of weighted vest exercise on muscle power and mobility function	SPBB (score)	CG	7.30 ± 1.50	NR	<b>0.009</b>	0.377	NR		
			EG	7.70 ± 1.30	NR	<b>&lt;0.01</b>		NR		
		Chair-5 time (s)	CG	19.60 ± 4.10	NR	<b>&lt;0.01</b>	<b>0.019</b>	NR		
			EG	18.50 ± 3.60	NR	<b>&lt;0.01</b>		NR		
		Gait speed (m/s)	CG	0.70 ± 0.16	NR	0.339	0.356	NR		
			EG	0.80 ± 0.15	NR	<b>0.006</b>		NR		
		Unilateral stance time (s)	CG	6.05 ± 5.90	NR	0.900	0.342	NR		
			EG	4.52 ± 5.40	NR	<b>0.028</b>		NR		
Outcomes of neuromuscular function										
Miszko et al. [57]	To determine whether power training was more efficacious than strength training for improving whole-body physical function and to examine the relationship between changes in anaerobic power and muscle strength and changes in physical function	Chest press—1 RM (kg)	CG	29.36 ± 12.20	29.18 ± 13.60	NR	>0.05	NR		
			EG	31.01 ± 12.90	34.81 ± 14.60	NR		NR		
		Leg press—1 RM (kg)	CG	75.61 ± 38.90	79.71 ± 37.50	NR	>0.05	NR		
			EG	95.45 ± 33.20	107.65 ± 32.20	NR		NR		
		Outcomes of physical function								
		CS-PFP test—total (score)	CG	55.5 ± 14	57.0 ± 18	NR	<b>&lt;0.05</b>	NR		
EG	58.2 ± 13		67.1 ± 13	NR	NR					

Abbreviations: Significant differences are highlighted in bold. CG, control group; EG, experimental group; M, mean; SD; standard deviation; NR, not reported; MMSE, mini-mental state test; FAB, frontal assessment battery; SPPB, short physical performance battery; TUG, time up and go test; ms, milliseconds; s, seconds; kg, kilograms; BW, body weight; Nm, newton meter; m/s, meter per second; deg/s, degree per second; mm, millimeter; cm, centimeters; BMI, body mass index; WHR, waist-to-hip ratio; MoCA, cognitive assessment; W, watts; EMG, surface electromyography for maximal muscle activation; RFD, rate of force development; CS-PFP, continuous-scale physical functional performance; N, newton; RM, repetition maximum; HSRT, high-speed resistance training; μV, microvolt; rep, repetitions; &, p value of comparison between CG and EG at post; <sup>a</sup>, effect size by Hopkins et al. [68]; #, small effect size (between 0.20 and 0.59); \*, moderate effect size (between 0.60 and 1.19); §, large effect size (between 1.20 and 1.99); £, very large effect size (≥2.00).


### 3.6. Meta-Analysis

The meta-analysis with ten studies showed moderate to large effects of the HSRT interventions on some health outcomes (cognitive function, neuromuscular function, and physical function) in older adults. Four studies revealed large effects on cognitive function (analyzed through frontal assessment battery (FAB) and mini-mental state examination (MMSE)) in favor of intervention groups compared to control groups ( $p = 0.001$ ,  $SMD = 0.94$  [0.20, 1.68],  $I^2 = 71%$ ); see Figure 2.




**Figure 2.** Forest plot presenting standardized mean difference and 95% confidence intervals from studies reporting high-speed resistance-training-induced changes in cognitive function between the intervention groups and the control groups. IV, independent variable; CI, confidence interval; SMD, standardized mean difference; FAB, frontal assessment battery; MMSE, mini-mental state examination [60,63,66,67].

In the same sense, six articles indicated moderate effects on neuromuscular function (analyzed through grip strength and muscle and power outcomes) in favor of intervention groups compared to control groups ( $p = 0.003$ ,  $SMD = 0.70$  [0.24, 1.15],  $I^2 = 53%$ ); see Figure 3. Finally, the analysis of the effects on physical function was divided into approaches in order to facilitate the interpretation of the results (Figure 4A,B). Consequently, this figure shows moderate effects on physical function in favor of intervention groups compared to control groups (Figure 4A,  $p = 0.004$ ,  $SMD = 0.55$  [0.02, 1.07],  $I^2 = 72%$ ; and Figure 4B,  $p = 0.009$ ,  $SMD = -0.59$  [-1.27, 0.08],  $I^2 = 35%$ ).



**Figure 3.** Forest plot presenting standardized mean difference and 95% confidence intervals from studies reporting high-speed resistance-training-induced changes in neuromuscular function between the intervention groups and the control groups. IV, independent variable; CI, confidence interval; SMD, standardized mean difference [54,57,61,65–67].



**Figure 4.** Forest plot presenting standardized mean difference and 95% confidence intervals from studies reporting high-speed resistance-training-induced changes in physical function ((A), through SPBB and gait velocity; (B), through TUG) between the intervention groups and the control groups. IV, independent variable; CI, confidence interval; SMD, standardized mean difference; SPPB, short physical performance battery; TUG, timed up and go test [54–56,61,66,67].

## 4. Discussion

The present systematic review and meta-analysis aimed to analyze and systematize the studies focused on HSRT interventions and their effects on health outcomes in independent older adults. The main findings suggest a lack of good-quality and pre-registered studies comparing HSRT interventions and CG. These analyses revealed moderate to substantial levels of heterogeneity and a wide range of predictive interval. There were moderate to substantial levels of heterogeneity between the studies. High heterogeneity may indicate that there are no benefits of HSRT over control groups [51,69]. Furthermore, all the variables show a prediction interval crossing the zero line.

The main finding of this review suggests that HSRT interventions would improve health measures—mostly cognitive function (large effects), neuromuscular function (moderate effects), and physical function (moderate effects)—thereby confirming the study hypotheses.

It is also important to note that there is a relationship between the duration of the intervention and the level of benefit. In this regard, interventions with ten weeks or more revealed significant improvements in muscle strength and power [54–56,59,61,64–67] and physical function [58,61,63,66,67]. These findings are of extreme importance for clinical practice, since muscle power is essential for physical performance during daily tasks [70,71].

For the sake of clarity, the Discussion section is organized into subsections according to the variables studied.

### 4.1. Cognitive Function

Regarding cognitive function, only one study investigated the retinal microvasculature density through changes occurring in the brain in cognitively normal older people [60]. The authors reported a negative correlation between retinal microvascular density and cognitive function ( $r = -0.54$ ;  $p = 0.007$ ), indicating that the improvement in cognitive function after HSRT could be mediated by the vascular effects. Nonetheless, regarding cognitive function, the studies by Yoon et al. [66] and Yoon et al. [67] addressed the effects of HSRT on cognitive function through the Montreal cognitive assessment (MoCA), FAB, and MMSE tests. The authors showed positive effects on processing speed, cognitive assessment, FAB, and MMSE test parameters. However, Lee et al. [63], who reported positive effects of 8-week HSRT on the neuromuscular function and gait performance in the elderly with mild cognitive impairment, failed to find any increase in the executive function of the frontal lobe. The sample characteristics and time of intervention may have compromised the expected results for these outcomes. Other studies have reported positive effects of HSRT on processing speed and executive function [67,72]. However, current evidence is limited.

The small number of included studies reduced the statistical power of this meta-analysis and did not allow an appropriate analysis of heterogeneity. Hence, more research on the role of exercise parameters (e.g., volume, types, and intensity) in specific cognitive functions is necessary. Addressing cognitive function is particularly important when considering the decline in executive and functional ability associated with ageing [73–76]. In this sense, cognitive function and muscular power [6,77,78] should be considered the primary factors influencing the decline in functional status in elderly populations. Consequently, it is fundamental to find ways to attenuate age-related executive function and muscular power decline due to the significant impact on life quality and the maintenance of an independent lifestyle.

Since neuromuscular deterioration accompanying the process of ageing is one of the first signs of decreased power output [78,79] and atrophy in areas associated with neuromuscular control [75], a potential relationship may exist between age-related declines in muscular power and cognitive function. Remarkably, research shows that exercise programs enhance a variety of functional and cognitive tasks, and that brain regions exclusively dedicated to executive function appear to be especially sensitive to exercise training [80–83]. The HSRT interventions showed promising results in cognitive executive function and memory, cognitive flexibility, and working memory [63,84]. The intensity

of these interventions can be controlled through changes in load and movement velocity. However, research that examined resistance training and cognition used loading to increase the intensity, which thus justifies further studies controlling the speed or using both load indicators [63].

#### 4.2. Neuromuscular Function

Concerning neuromuscular function with a particular focus on neural functions, some studies investigated the effects of HSRT on neural factors [54–56,61,63]. The effects of the HSRT intervention on neural factors showed positive results in two aspects: first, between the CG and EG in movement time, motor time, rate of electromyography (EMG) rise, rate of torque development, and percentage of voluntary activation; and second, between pre- and post-intervention in normalized parameters peak torque, and in the EMG amplitudes of the knee flexors, extensors, vastus lateralis (VL), vastus medialis (VM), and rectus femoris [56].

In the latter study [56], muscle activation determined by recording the surface electromyogram during a maximal isometric leg exercise demonstrated that the rates of EMG rose, similar to other studies. In this sense, Reid et al. [85] reported a significant increase in the rate of EMG rise among the elderly averaged 77-year old who performed HSRT at 40% of 1 RM for 16 weeks compared to the elderly who underwent HSRT at 70% of 1 RM. It can be inferred that the low-intensity HSRT may influence the apparent increase in the rate of EMG rise. This increase may occur because the abilities to recruit motor units and maximal motor unit-firing rate were developed in parallel to the increased rate of torque development associated with recruiting motor units [86].

In the last two decades, studies with pneumatic and isokinetic devices have shown that the peak muscle power of the leg extensors [54–56,59,63,65–67,87,88] and ankle plantar flexors [18] are stronger predictors of functional performance than muscle strength. The peak muscle power declines sooner and more rapidly than muscle strength [19,89] and is a more important predictor of functional performance than strength [18,87]. Therefore, peak muscle power could be a more critical variable on which to focus RT programs among older men and women.

Based on data from selected studies [56,61,63], this review reveals that HSRT was more effective than traditional resistance training with rising muscle activation, as reported by Reid et al. [85]. There was an increase in muscle activation of VM and VL muscles, suggesting increased recruitment and firing frequency of motor units [55,86,90,91]. In addition to gains in maximal EMG amplitudes and isometric strength, older adults showed elevated knee extensor activation and coactivation during early stance and elevated plantar flexor activation during push-off [55]. The data suggest that power training-induced increases in voluntary muscle activation and agonist muscle activation underlie increases in isometric muscle strength and gait velocity. Therefore, training based on HSRT can modify neuromuscular activation and increase older adults' leg muscle strength and walking performance.

The small number of included studies for grip strength reduced the statistical power of this meta-analysis and there were substantial levels of heterogeneity between studies. However, all the variables showed a prediction interval crossing the zero line, suggesting the possibility of the real beneficial effects of HSRT. For the four studies on the muscle and power outcomes, null heterogeneity may indicate that there are real benefits of HSRT over the CG.

The mechanism underlying the increased neuromuscular activation may involve, but is not limited to, increased neural drive via corticospinal pathways, increased motor neuron and/or muscle fiber excitability, an increased number of active motor units, and/or increased conduction velocity [92,93].

The neuromuscular function was evaluated through various strength parameters, such as grip strength and isokinetic peak torque/body weight [66,67], rate of torque development [63], isokinetic power in average at different execution speeds [67], isometric

strength [56], peak power at 40 to 80% 1 RM [65], and knee extension and leg press strength [64].

Moreover, the maximal isokinetic power at different execution speeds was evaluated in relation to body weight and joint range [54], and one study evaluated maximal isometric strength relative to leg length [61]. Two studies evaluated peak power at different percentages of 1 RM [64,65], as well as the velocity at peak power in air resistance equipment [65]. The maximal strength was evaluated through the 1 RM in leg extension, leg press [59,64], and chest press exercises [57]. Our findings evidenced that HSRT interventions are effective in improving strength and power. Despite the consistency in the results of the assessments on neuromuscular function in the comparison of pre–post or/and between CG and EG, some studies do not report positive effects in some parameters [57,59,66]. In the study by Reid et al. [59], the 1 RM of knee extension increased significantly from baseline in either the strength training group (41%) or the HSRT group (49%) when compared to CG ( $p < 0.01$ ). However, there was no significant time–group interaction for the execution of the 1 RM leg press exercise. Hence, the authors concluded that the principal finding of their study was that, in older people with mild–moderate mobility impairments, a 12-week HSRT intervention induced similar improvements in lower-extremity muscle power compared to traditional slow-velocity resistance training. The non-significant increases in absolute power on the leg press in response to HSRT are considerably lower than those reported by other studies in older adults [57,64,65,94]. These results suggest that gains in muscle quality, which occurred without measurable hypertrophy, may have been due to neural adaptation to this form of explosive resistance training [63,95]. The muscle power decreases more precipitously than muscle strength in older adults, causing a dramatic loss in the ability to produce force rapidly.

The current evidence emphasizes the challenges regarding the development of optimal exercise strategies for clinically relevant outcomes in older adults [96,97]. In this sense, strategies that promote rapid increases in strength development should be a top priority in resistance training interventions, followed by methods that stimulate muscle hypertrophy [98,99].

#### 4.3. Physical Function

Regarding physical function, namely gait function, over the past two decades, there has been a proliferation of research surrounding resistance training to increase functional activities. Specifically, studies have demonstrated that HSRT can promote significant improvements in functional outcomes [57,100–102] compared to standard resistance training programs.

Nine of the studies in this systematic review used some form of a timed walking test, namely the 2-min maximal walking test; 10-m walking test; gait velocity at usual speed and fast speed; 4.44 m gait speed test; timed up and go test (TUG test); and 8-foot up and go test. Most of the studies reported improvements in the pre–post within-group and between-group (EG vs. CG). Nevertheless, Lee et al. [63] only found improvements in the pre–post in EG (4.44 m gait speed test and TUG test). Some studies [54–56] did not improve gait velocity at the usual speed. However, two of the three studies [54,55] presented improvements when comparing pre–post to gait velocity at fast speed. However, the significant between-group change in gait speed suggested that the clinical relevance is 0.05–0.10 m/s [103].

According to Ramírez-Campillo et al. [24], the HSRT intervention in older people significantly improved walking ability and TUG test results compared to the CG. Ringsberg et al. [104] and Karttunen et al. [105] reported that the older men and women (average 75 years old) showed a statistically significant relationship between isometric knee extensor strength and gait speed. Thereby, in this population, the HSRT with heavy and light loads seems to mediate adaptations in the neural system and translates into improvements in physical function, namely in gait function. In the same sense, Harridge et al. [106] reported submaximal voluntary activation of the knee extensor muscles in frail older people.

The current knowledge indicates incomplete voluntary muscle activation as an essential mechanism for the loss of mechanical muscle function and physical function, contributing to the pathway of mobility disability [61,107,108].

In this meta-analysis, only two studies addressed the SPPB variable, which reduced the statistical power of these meta-analyses and did not allow an appropriate analysis of heterogeneity. They present high heterogeneity ( $I^2 = 91\%$ , 1.01 [−1.16, 3.18]); however, the variables show a prediction interval crossing the zero line. The same is observed in the gait velocity variable; although it was included in five studies, three of the studies had a major limitation, which was its origin in the same clinical trial [54–56], which could result in analysis bias. Finally, in the meta-analysis for the TUG test, only two studies reported this variable and presented values favorable to the intervention with the HSRT.

#### 4.4. Body Composition

Finally, regarding body composition, the HSRT seems to induce skeletal muscle hypertrophy and neuromuscular adaptations [66]. This intervention can also facilitate increases in muscle quality in response to power training [59] and promote significant reductions in several parameters, namely body weight, body mass index, percent body fat, and waist-to-hip ratio [66]. Some studies showed evident morphological adaptations [58,109,110] due to an increase in muscle thickness of VL in all strength training groups, which indicates that hypertrophy likely contributed to the increased maximal and dynamic strength.

According to Reid et al. [59], the studies that reported improvements in muscle strength and power without associated alterations in body composition may reflect enhanced neural adaptations in the early stages of training, as reported by other studies [91,111]. It is likely that the earlier motor unit activation and enhanced maximal firing rates associated with HSRT would be principle stimuli for neural adaptations to this explosive form of training [111]. The results of improvements in muscle strength and power without associated alterations in body composition may also be a consequence of the short duration of interventions and low weekly frequency, and the differences in neuromuscular adaptation should be investigated over longer periods (e.g., six months).

#### 4.5. Limitations

This study presents two limitations that should be pointed out. The first is the poor methodological quality of some of the included studies. Twelve of the fourteen included studies showed low methodological quality (level 2). This should be taken as a serious warning to the scientific community and may hinder the credibility and practical effect of said studies. Second, although our meta-analysis suggests very satisfactory results, it was conducted with a small number of studies in each function, so it is necessary that future studies be conducted in this area.

### 5. Conclusions

The present systematic review confirms that HSRT interventions that progressively increase in training intensity can improve several health outcomes in older people, mostly cognitive, neuromuscular, and physical function (moderate to large effects).

Furthermore, the results suggest that interventions with ten weeks or more and performed three times a week result in significant improvements in neuromuscular function.

### 6. Future Lines of Research

It should be noted that even though gait variability [97] and phase angle from bioelectrical impedance [112] have high clinical significance, none of the included studies verified the effect of the HSRT intervention on said outcomes. Therefore, we suggest that future studies conducted with older people and based on this approach (HSRT intervention) include the following variables. The first is gait variability—which refers to the magnitude of the stride-to-stride fluctuations and their changes over time during a walk—which may be useful in understanding the physiology of gait, in quantifying age-related and pathologic



alterations in the locomotor control system, and in augmenting the objective measurement of mobility and functional status [113,114]. A gait variability study through non-linear methods [97,115] can help us to understand the physiological changes associated with ageing induced by an increase in noise in the neuromotor system, promoting significant changes in the control of the locomotor system [113,116]. Multiple age-related physiological changes increase neuromotor noise, increasing gait variability. If older adults alter how they regulate their stride variables, this could further exacerbate this variability. The second is the phase angle from electrical bioimpedance—the assessment of this indicator represents an interesting measure for future research on healthy populations, because it can reflect cellular health and mass and cell membrane integrity [112,117], and in the elderly population, it is a predictor of muscle function [96,118] and sarcopenia [119] and an indicator of general health status [120,121].

Finally, we also recommend further research to include an adequate follow-up (e.g., minimum six months) to assess the durability of HSRT intervention effects on all health-related variables.

**Author Contributions:** Conceptualization: A.D.M., O.F., A.P., R.O., F.D.A.G., N.J.C.L. and J.P.B.; Methodology: A.D.M., R.O. and J.P.B.; Software: A.D.M., F.D.A.G. and N.J.C.L.; Validation: A.D.M., A.P. and J.P.B.; Formal Analysis: A.D.M., R.O. and J.P.B.; Investigation: A.D.M., O.F., A.P., R.O., F.D.A.G., N.J.C.L. and J.P.B.; Resources: A.D.M., R.O., N.J.C.L. and J.P.B.; Data Curation: A.D.M., O.F., A.P., R.O. and J.P.B.; Writing—Original Draft: A.D.M., O.F., A.P. and J.P.B.; Writing—Review and Editing: A.D.M., O.F., A.P., R.O. and J.P.B.; Visualization: A.D.M., O.F., A.P., R.O., F.D.A.G., N.J.C.L. and J.P.B.; Supervision: O.F. and J.P.B.; Project administration: A.D.M., O.F. and J.P.B.; Funding acquisition: A.D.M., A.P., R.O. and J.P.B. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by the Portuguese Foundation for Science and Technology, I.P., Grant/Award Number 2021.04598.BD and UIDP/04748/2020.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Tornero-Quiñones, I.; Sáez-Padilla, J.; Espina Díaz, A.; Abad Robles, M.T.; Sierra Robles, Á. Functional Ability, Frailty and Risk of Falls in the Elderly: Relations with Autonomy in Daily Living. *Int. J. Environ. Res. Public Health* **2020**, *17*, 1006. [CrossRef] [PubMed]
2. Orssatto, L.B.R.; Moura, B.M.; Bezerra, E.S.; Andersen, L.L.; Oliveira, S.N.; Diefenthaler, F. Influence of Strength Training Intensity on Subsequent Recovery in Elderly. *Exp. Gerontol.* **2018**, *106*, 232–239. [CrossRef] [PubMed]
3. Thomas, E.; Battaglia, G.; Patti, A.; Brusa, J.; Leonardi, V.; Palma, A.; Bellafiore, M. Physical Activity Programs for Balance and Fall Prevention in Elderly: A Systematic Review. *Medicine* **2019**, *98*, e16218. [CrossRef] [PubMed]
4. Paraskevoudi, N.; Balci, F.; Vatakis, A. “Walking” through the Sensory, Cognitive, and Temporal Degradations of Healthy Aging. *Ann. N. Y. Acad. Sci.* **2018**, *1426*, 72–92. [CrossRef]
5. Clark, D.; Pojednic, R.M.; Reid, K.F.; Patten, C.; Pasha, E.P.; Phillips, E.M.; Fielding, R.A. Longitudinal Decline of Neuromuscular Activation and Power in Healthy Older Adults. *J. Gerontol. Ser. A Biol. Sci. Med. Sci.* **2013**, *68*, 1419–1425. [CrossRef]
6. Izquierdo, M.; Cadore, E.L. Muscle Power Training in the Institutionalized Frail: A New Approach to Counteracting Functional Declines and Very Late-Life Disability. *Curr. Med. Res. Opin.* **2014**, *30*, 1385–1390. [CrossRef]
7. Vaishya, R.; Vaish, A. Falls in Older Adults Are Serious. *Indian J. Orthop.* **2020**, *54*, 69–74. [CrossRef]
8. Khanuja, K.; Joki, J.; Bachmann, G.; Cuccurullo, S. Gait and Balance in the Aging Population: Fall Prevention Using Innovation and Technology. *Maturitas* **2018**, *110*, 51–56. [CrossRef]
9. NICE. Falls in Older People: Assessing Risk and Prevention. 2016. Available online: <https://www.nice.org.uk/guidance/cg161/resources/falls-in-older-people-assessing-risk-and-prevention-pdf-35109686728645> (accessed on 25 January 2022).
10. Keating, C.J.; Cabrera-Linares, J.C.; Párraga-Montilla, J.A.; Latorre-Román, P.A.; Del Castillo, R.M.; García-Pinillos, F. Influence of Resistance Training on Gait & Balance Parameters in Older Adults: A Systematic Review. *Int. J. Environ. Res. Public Health* **2021**, *18*, 1759. [CrossRef]

11. Sherrington, C.; Fairhall, N.; Wallbank, G.; Tiedemann, A.; Michaleff, Z.A.; Howard, K.; Clemson, L.; Hopewell, S.; Lamb, S. Exercise for Preventing Falls in Older People Living in the Community: An Abridged Cochrane Systematic Review. *Br. J. Sports Med.* **2020**, *54*, 885–891. [[CrossRef](#)]
12. Woolford, S.J.; Sohan, O.; Dennison, E.M.; Cooper, C.; Patel, H.P. Approaches to the Diagnosis and Prevention of Frailty. *Aging Clin. Exp. Res.* **2020**, *32*, 1629–1637. [[CrossRef](#)] [[PubMed](#)]
13. Brady, A.O.; Straight, C.R.; Evans, E.M. Body Composition, Muscle Capacity, and Physical Function in Older Adults: An Integrated Conceptual Model. *J. Aging Phys. Act.* **2014**, *22*, 441–452. [[CrossRef](#)] [[PubMed](#)]
14. Manini, T.M.; Clark, B. Dynapenia and Aging: An Update. *J. Gerontol. Ser. A Biol. Sci. Med. Sci.* **2012**, *67*, 28–40. [[CrossRef](#)] [[PubMed](#)]
15. Cadore, E.L.; Casas-Herrero, A.; Zambom-Ferraresi, F.; Idoate, F.; Millor, N.; Gómez, M.; Rodriguez-Mañas, L.; Izquierdo, M. Multicomponent Exercises Including Muscle Power Training Enhance Muscle Mass, Power Output, and Functional Outcomes in Institutionalized Frail Nonagenarians. *Age* **2014**, *36*, 773–785. [[CrossRef](#)] [[PubMed](#)]
16. Casas-Herrero, A.; Cadore, E.L.; Zambom-Ferraresi, F.; Idoate, F.; Millor, N.; Martínez-Ramírez, A.; Gómez, M.; Rodriguez-Mañas, L.; Marcellán, T.; de Gordo, A.R.; et al. Functional Capacity, Muscle Fat Infiltration, Power Output, and Cognitive Impairment in Institutionalized Frail Oldest Old. *Rejuvenation Res.* **2013**, *16*, 396–403. [[CrossRef](#)]
17. Han, L.; Yang, F. Strength or Power, Which Is More Important to Prevent Slip-Related Falls? *Hum. Mov. Sci.* **2015**, *44*, 192–200. [[CrossRef](#)]
18. Suzuki, T.; Bean, J.F.; Fielding, R.A. Muscle Power of the Ankle Flexors Predicts Functional Performance in Community-Dwelling Older Women. *J. Am. Geriatr. Soc.* **2001**, *49*, 1161–1167. [[CrossRef](#)]
19. Izquierdo, M.; Ibañez, J.; Gorostiaga, E.; Garrues, M.; Zúñiga, A.; Antón, A.; Larrión, J.L.; Häkkinen, K. Maximal Strength and Power Characteristics in Isometric and Dynamic Actions of the Upper and Lower Extremities in Middle-Aged and Older Men. *Acta Physiol. Scand.* **1999**, *167*, 57–68. [[CrossRef](#)]
20. Cadore, E.L.; Izquierdo, M. How to Simultaneously Optimize Muscle Strength, Power, Functional Capacity, and Cardiovascular Gains in the Elderly: An Update. *Age Dordr. Neth.* **2013**, *35*, 2329–2344. [[CrossRef](#)]
21. De Vos, N.J.; Singh, N.A.; Ross, D.A.; Stavrinou, T.M.; Orr, R.; Fiatarone Singh, M.A. Optimal Load for Increasing Muscle Power during Explosive Resistance Training in Older Adults. *J. Gerontol. Ser. A Biol. Sci. Med. Sci.* **2005**, *60*, 638–647. [[CrossRef](#)]
22. Fielding, R.A.; LeBrasseur, N.K.; Cuoco, A.; Bean, J.; Mizer, K.; Fiatarone Singh, M.A. High-Velocity Resistance Training Increases Skeletal Muscle Peak Power in Older Women. *J. Am. Geriatr. Soc.* **2002**, *50*, 655–662. [[CrossRef](#)] [[PubMed](#)]
23. Henwood, T.R.; Riek, S.; Taaffe, D.R. Strength versus Muscle Power-Specific Resistance Training in Community-Dwelling Older Adults. *J. Gerontol. Ser. A Biol. Sci. Med. Sci.* **2008**, *63*, 83–91. [[CrossRef](#)] [[PubMed](#)]
24. Ramírez-Campillo, R.; Castillo, A.; de la Fuente, C.I.; Campos-Jara, C.; Andrade, D.C.; Álvarez, C.; Martínez, C.; Castro-Sepúlveda, M.; Pereira, A.; Marques, M.C.; et al. High-Speed Resistance Training Is More Effective than Low-Speed Resistance Training to Increase Functional Capacity and Muscle Performance in Older Women. *Exp. Gerontol.* **2014**, *58*, 51–57. [[CrossRef](#)] [[PubMed](#)]
25. Wallerstein, L.F.; Tricoli, V.; Barroso, R.; Rodacki, A.L.F.; Russo, L.; Aihara, A.Y.; da Rocha Correa Fernandes, A.; de Mello, M.T.; Ugrinowitsch, C. Effects of Strength and Power Training on Neuromuscular Variables in Older Adults. *J. Aging Phys. Act.* **2012**, *20*, 171–185. [[CrossRef](#)]
26. Unhjem, R.; Tøien, T.; Kvellestad, A.C.G.; Øren, T.S.; Wang, E. External Resistance Is Imperative for Training-Induced Efferent Neural Drive Enhancement in Older Adults. *J. Gerontol. Ser. A Biol. Sci. Med. Sci.* **2021**, *76*, 224–232. [[CrossRef](#)]
27. De Vos, N.J.; Singh, N.A.; Ross, D.A.; Stavrinou, T.M.; Orr, R.; Singh, M.A.F. Effect of Power-Training Intensity on the Contribution of Force and Velocity to Peak Power in Older Adults. *J. Aging Phys. Act.* **2008**, *16*, 393–407. [[CrossRef](#)]
28. Pereira, A.; Izquierdo, M.; Silva, A.; Costa, A.; Bastos, E.; González-Badillo, J.; Marques, M. Effects of High-Speed Power Training on Functional Capacity and Muscle Performance in Older Women. *Exp. Gerontol.* **2012**, *47*, 250–255. [[CrossRef](#)]
29. Sayers, S.P.; Gibson, K. High-Speed Power Training in Older Adults: A Shift of the External Resistance at Which Peak Power Is Produced. *J. Strength Cond. Res. Natl. Strength Cond. Assoc.* **2014**, *28*, 616–621. [[CrossRef](#)]
30. Holviala, J.H.S.; Sallinen, J.M.; Kraemer, W.J.; Alen, M.J.; Häkkinen, K.K.T. Effects of Strength Training on Muscle Strength Characteristics, Functional Capabilities, and Balance in Middle-Aged and Older Women. *J. Strength Cond. Res.* **2006**, *20*, 336–344. [[CrossRef](#)]
31. Marques, D.; Neiva, H.; Marinho, D.; Marques, M. Novel Resistance Training Approach to Monitoring the Volume in Older Adults: The Role of Movement Velocity. *Int. J. Environ. Res. Public Health* **2020**, *17*, 7557. [[CrossRef](#)]
32. Marques, D.; Neiva, H.; Faíl, L.; Gil, M.; Marques, M. Acute Effects of Low and High-Volume Resistance Training on Hemodynamic, Metabolic and Neuromuscular Parameters in Older Adults. *Exp. Gerontol.* **2019**, *125*, 7. [[CrossRef](#)] [[PubMed](#)]
33. Pereira, A.; Izquierdo, M.; Silva, A.J.; Costa, A.M.; González-Badillo, J.J.; Marques, M.C. Muscle Performance and Functional Capacity Retention in Older Women after High-Speed Power Training Cessation. *Exp. Gerontol.* **2012**, *47*, 620–624. [[CrossRef](#)] [[PubMed](#)]
34. Moro, T.; Brightwell, C.R.; Phalen, D.E.; McKenna, C.F.; Lane, S.J.; Porter, C.; Volpi, E.; Rasmussen, B.B.; Fry, C.S. Low Skeletal Muscle Capillarization Limits Muscle Adaptation to Resistance Exercise Training in Older Adults. *Exp. Gerontol.* **2019**, *127*, 110723. [[CrossRef](#)] [[PubMed](#)]

35. Snijders, T.; Nederveen, J.P.; Joannisse, S.; Leenders, M.; Verdijk, L.B.; van Loon, L.J.C.; Parise, G. Muscle Fibre Capillarization Is a Critical Factor in Muscle Fibre Hypertrophy during Resistance Exercise Training in Older Men. *J. Cachexia Sarcopenia Muscle* **2017**, *8*, 267–276. [[CrossRef](#)]
36. Nederveen, J.P.; Joannisse, S.; Snijders, T.; Ivankovic, V.; Baker, S.K.; Phillips, S.M.; Parise, G. Skeletal Muscle Satellite Cells Are Located at a Closer Proximity to Capillaries in Healthy Young Compared with Older Men. *J. Cachexia Sarcopenia Muscle* **2016**, *7*, 547–554. [[CrossRef](#)]
37. Snijders, T.; Nederveen, J.P.; McKay, B.R.; Joannisse, S.; Verdijk, L.B.; van Loon, L.J.C.; Parise, G. Satellite Cells in Human Skeletal Muscle Plasticity. *Front. Physiol.* **2015**, *6*, 283. [[CrossRef](#)]
38. Straight, C.R.; Lindheimer, J.B.; Brady, A.O.; Dishman, R.K.; Evans, E.M. Effects of Resistance Training on Lower-Extremity Muscle Power in Middle-Aged and Older Adults: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Sports Med. Auckl. NZ* **2016**, *46*, 353–364. [[CrossRef](#)]
39. Byrne, C.; Faure, C.; Keene, D.J.; Lamb, S.E. Ageing, Muscle Power and Physical Function: A Systematic Review and Implications for Pragmatic Training Interventions. *Sports Med. Auckl. NZ* **2016**, *46*, 1311–1332. [[CrossRef](#)]
40. Da Rosa Orssatto, L.B.; de la Rocha Freitas, C.; Shield, A.J.; Silveira Pinto, R.; Trajano, G.S. Effects of Resistance Training Concentric Velocity on Older Adults' Functional Capacity: A Systematic Review and Meta-Analysis of Randomised Trials. *Exp. Gerontol.* **2019**, *127*, 110731. [[CrossRef](#)]
41. Katsoulis, K.; Stathokostas, L.; Amara, C.E. The Effects of High- Versus Low-Intensity Power Training on Muscle Power Outcomes in Healthy, Older Adults: A Systematic Review. *J. Aging Phys. Act.* **2019**, *27*, 422–439. [[CrossRef](#)]
42. Bieńkiewicz, M.M.N.; Brandi, M.-L.; Goldenberg, G.; Hughes, C.M.L.; Hermsdörfer, J. The Tool in the Brain: Apraxia in ADL. Behavioral and Neurological Correlates of Apraxia in Daily Living. *Front. Psychol.* **2014**, *5*, 353. [[CrossRef](#)] [[PubMed](#)]
43. Edemekong, P.F.; Bomgaars, D.L.; Sukumaran, S.; Levy, S.B. Activities of Daily Living. In *StatPearls*; StatPearls Publishing: Treasure Island, FL, USA, 2021.
44. Oleske, D.M.; Islam, S.S. Chapter 5—Role of Epidemiology in the Biopharmaceutical Industry. In *Pharmacovigilance: A Practical Approach*; Doan, T., Renz, C., Bhattacharya, M., Lievano, F., Scarazzini, L., Eds.; Elsevier: Amsterdam, The Netherlands, 2019; pp. 69–87. ISBN 978-0-323-58116-5.
45. Page, M.J.; McKenzie, J.E.; Bossuyt, P.M.; Boutron, I.; Hoffmann, T.C.; Mulrow, C.D.; Shamseer, L.; Tetzlaff, J.M.; Akl, E.A.; Brennan, S.E.; et al. The PRISMA 2020 Statement: An Updated Guideline for Reporting Systematic Reviews. *BMJ* **2021**, *372*, 71. [[CrossRef](#)] [[PubMed](#)]
46. Maher, C.G.; Sherrington, C.; Herbert, R.D.; Moseley, A.M.; Elkins, M. Reliability of the PEDro Scale for Rating Quality of Randomized Controlled Trials. *Phys. Ther.* **2003**, *83*, 713–721. [[CrossRef](#)] [[PubMed](#)]
47. Van Tulder, M.; Furlan, A.; Bombardier, C.; Bouter, L.; Editorial Board of the Cochrane Collaboration Back Review Group. Updated Method Guidelines for Systematic Reviews in the Cochrane Collaboration Back Review Group. *Spine* **2003**, *28*, 1290–1299. [[CrossRef](#)]
48. Kollen, B.J.; Lennon, S.; Lyons, B.; Wheatley-Smith, L.; Scheper, M.; Buurke, J.H.; Halfens, J.; Geurts, A.C.H.; Kwakkel, G. The Effectiveness of the Bobath Concept in Stroke Rehabilitation: What Is the Evidence? *Stroke* **2009**, *40*, e89–e97. [[CrossRef](#)]
49. Vaughan-Graham, J.; Cott, C.; Wright, F.V. The Bobath (NDT) Concept in Adult Neurological Rehabilitation: What Is the State of the Knowledge? A Scoping Review. Part II: Intervention Studies Perspectives. *Disabil. Rehabil.* **2015**, *37*, 1909–1928. [[CrossRef](#)]
50. Borenstein, M.; Hedges, L.V.; Higgins, J.P.T.; Rothstein, H.R. A Basic Introduction to Fixed-Effect and Random-Effects Models for Meta-Analysis. *Res. Synth. Methods* **2010**, *1*, 97–111. [[CrossRef](#)]
51. Deeks, J.J.; Higgins, J.P.; Altman, D.G.; Group on behalf of the Cochrane Statistical Methods. Analysing Data and Undertaking Meta-Analyses. In *Cochrane Handbook for Systematic Reviews of Interventions*; John Wiley & Sons, Ltd.: Hoboken, NJ, USA, 2019; pp. 241–284. ISBN 978-1-119-53660-4.
52. Cohen, J.W. *Statistical Power Analysis for the Behavioral Sciences*, 2nd ed.; Lawrence Erlbaum Associates: Hillsdale, MI, USA, 1988.
53. Deeks, J.J.; Higgins, J.P.T. *Statistical Algorithms in Review Manager 5*; The Cochrane Collaboration: London, UK, 2016.
54. Beijersbergen, C.M.I.; Granacher, U.; Gäbler, M.; Devita, P.; Hortobágyi, T. Kinematic Mechanisms of How Power Training Improves Healthy Old Adults' Gait Velocity. *Med. Sci. Sports Exerc.* **2017**, *49*, 150–157. [[CrossRef](#)]
55. Beijersbergen, C.M.I.; Granacher, U.; Gäbler, M.; Devita, P.; Hortobágyi, T. Hip Mechanics Underlie Lower Extremity Power Training-Induced Increase in Old Adults' Fast Gait Velocity: The Potsdam Gait Study (POGS). *Gait Posture* **2017**, *52*, 338–344. [[CrossRef](#)]
56. Beijersbergen, C.M.I.; Granacher, U.; Gäbler, M.; Devita, P.; Hortobágyi, T. Power Training-Induced Increases in Muscle Activation during Gait in Old Adults. *Med. Sci. Sports Exerc.* **2017**, *49*, 2198–2025. [[CrossRef](#)]
57. Miszko, T.A.; Cress, M.E.; Slade, J.M.; Covey, C.J.; Agrawal, S.K.; Doerr, C.E. Effect of Strength and Power Training on Physical Function in Community-Dwelling Older Adults. *J. Gerontol. Ser. A Biol. Sci. Med. Sci.* **2003**, *58*, 171–175. [[CrossRef](#)] [[PubMed](#)]
58. Bean, J.F.; Herman, S.; Kiely, D.K.; Frey, I.C.; Leveille, S.G.; Fielding, R.A.; Frontera, W.R. Increased Velocity Exercise Specific to Task (InVEST) Training: A Pilot Study Exploring Effects on Leg Power, Balance, and Mobility in Community-Dwelling Older Women. *J. Am. Geriatr. Soc.* **2004**, *52*, 799–804. [[CrossRef](#)] [[PubMed](#)]
59. Reid, K.F.; Callahan, D.M.; Carabello, R.J.; Phillips, E.M.; Frontera, W.R.; Fielding, R.A. Lower Extremity Power Training in Elderly Subjects with Mobility Limitations: A Randomized Controlled Trial. *Aging Clin. Exp. Res.* **2008**, *20*, 337–343. [[CrossRef](#)] [[PubMed](#)]

60. Fang, M.; Strand, K.; Zhang, J.; Totillo, M.; Chen, Q.; Signorile, J.F.; Jiang, H.; Wang, J. Characterization of Retinal Microvasculature and Its Relations to Cognitive Function in Older People after Circuit Resistance Training. *Exp. Gerontol.* **2020**, *142*, 111114. [[CrossRef](#)]
61. Hvid, L.G.; Strotmeyer, E.S.; Skjødt, M.; Magnussen, L.V.; Andersen, M.; Caserotti, P. Voluntary Muscle Activation Improves with Power Training and Is Associated with Changes in Gait Speed in Mobility-Limited Older Adults—A Randomized Controlled Trial. *Exp. Gerontol.* **2016**, *80*, 51–56. [[CrossRef](#)]
62. Katula, J.A.; Rejeski, W.J.; Marsh, A.P. Enhancing Quality of Life in Older Adults: A Comparison of Muscular Strength and Power Training. *Health Qual. Life Outcomes* **2008**, *6*, 45. [[CrossRef](#)]
63. Lee, D.W.; Yoon, D.H.; Lee, J.-Y.; Panday, S.B.; Park, J.; Song, W. Effects of High-Speed Power Training on Neuromuscular and Gait Functions in Frail Elderly with Mild Cognitive Impairment Despite Blunted Executive Functions: A Randomized Controlled Trial. *J. Frailty Aging* **2020**, *9*, 179–184. [[CrossRef](#)]
64. Marsh, A.P.; Miller, M.E.; Rejeski, W.J.; Hutton, S.L.; Kritchevsky, S.B. Lower Extremity Muscle Function after Strength or Power Training in Older Adults. *J. Aging Phys. Act.* **2009**, *17*, 416–443. [[CrossRef](#)]
65. Sayers, S.P.; Gibson, K. A Comparison of High-Speed Power Training and Traditional Slow-Speed Resistance Training in Older Men and Women. *J. Strength Cond. Res.* **2010**, *24*, 3369–3380. [[CrossRef](#)]
66. Yoon, D.H.; Kang, D.; Kim, H.-J.; Kim, J.-S.; Song, H.S.; Song, W. Effect of Elastic Band-Based High-Speed Power Training on Cognitive Function, Physical Performance and Muscle Strength in Older Women with Mild Cognitive Impairment. *Geriatr. Gerontol. Int.* **2017**, *17*, 765–772. [[CrossRef](#)]
67. Yoon, D.H.; Lee, J.-Y.; Song, W. Effects of Resistance Exercise Training on Cognitive Function and Physical Performance in Cognitive Frailty: A Randomized Controlled Trial. *J. Nutr. Health Aging* **2018**, *22*, 944–951. [[CrossRef](#)] [[PubMed](#)]
68. Hopkins, W.G.; Marshall, S.W.; Batterham, A.M.; Hanin, J. Progressive Statistics for Studies in Sports Medicine and Exercise Science. *Med. Sci. Sports Exerc.* **2009**, *41*, 3–12. [[CrossRef](#)] [[PubMed](#)]
69. Rücker, G.; Schwarzer, G.; Carpenter, J.R.; Schumacher, M. Undue Reliance on I(2) in Assessing Heterogeneity May Mislead. *BMC Med. Res. Methodol.* **2008**, *8*, 79. [[CrossRef](#)] [[PubMed](#)]
70. Coelho-Júnior, H.J.; Uchida, M.C.; Picca, A.; Bernabei, R.; Landi, F.; Calvani, R.; Cesari, M.; Marzetti, E. Evidence-Based Recommendations for Resistance and Power Training to Prevent Frailty in Community-Dwellers. *Aging Clin. Exp. Res.* **2021**, *33*, 2069–2086. [[CrossRef](#)] [[PubMed](#)]
71. Fragala, M.S.; Cadore, E.L.; Dorgo, S.; Izquierdo, M.; Kraemer, W.J.; Peterson, M.D.; Ryan, E.D. Resistance Training for Older Adults: Position Statement From the National Strength and Conditioning Association. *J. Strength Cond. Res.* **2019**, *33*, 2019–2052. [[CrossRef](#)]
72. Fried, L.P.; Tangen, C.M.; Walston, J.; Newman, A.B.; Hirsch, C.; Gottdiener, J.; Seeman, T.; Tracy, R.; Kop, W.J.; Burke, G.; et al. Frailty in Older Adults: Evidence for a Phenotype. *J. Gerontol. Med. Sci.* **2001**, *56A*, 146–156. [[CrossRef](#)]
73. Bherer, L.; Erickson, K.I.; Liu-Ambrose, T. A Review of the Effects of Physical Activity and Exercise on Cognitive and Brain Functions in Older Adults. *J. Aging Res.* **2013**, *2013*, 657508. [[CrossRef](#)] [[PubMed](#)]
74. Johnson, J.K.; Lui, L.-Y.; Yaffe, K. Executive Function, More than Global Cognition, Predicts Functional Decline and Mortality in Elderly Women. *J. Gerontol. Ser. A Biol. Sci. Med. Sci.* **2007**, *62*, 1134–1141. [[CrossRef](#)]
75. Raz, N.; Rodrigue, K.M. Differential Aging of the Brain: Patterns, Cognitive Correlates and Modifiers. *Neurosci. Biobehav. Rev.* **2006**, *30*, 730–748. [[CrossRef](#)]
76. Royall, D.R.; Palmer, R.; Chiodo, L.K.; Polk, M.J. Declining Executive Control in Normal Aging Predicts Change in Functional Status: The Freedom House Study. *J. Am. Geriatr. Soc.* **2004**, *52*, 346–352. [[CrossRef](#)]
77. Foldvari, M.; Clark, M.; Laviolette, L.C.; Bernstein, M.A.; Kaliton, D.; Castaneda, C.; Pu, C.T.; Hausdorff, J.M.; Fielding, R.A.; Singh, M.A. Association of Muscle Power with Functional Status in Community-Dwelling Elderly Women. *J. Gerontol. Ser. A Biol. Sci. Med. Sci.* **2000**, *55*, M192–M199. [[CrossRef](#)] [[PubMed](#)]
78. Reid, K.F.; Pasha, E.; Doros, G.; Clark, D.J.; Patten, C.; Phillips, E.M.; Frontera, W.R.; Fielding, R.A. Longitudinal Decline of Lower Extremity Muscle Power in Healthy and Mobility-Limited Older Adults: Influence of Muscle Mass, Strength, Composition, Neuromuscular Activation and Single Fiber Contractile Properties. *Eur. J. Appl. Physiol.* **2014**, *114*, 29–39. [[CrossRef](#)] [[PubMed](#)]
79. McKinnon, N.B.; Montero-Odasso, M.; Doherty, T.J. Motor Unit Loss Is Accompanied by Decreased Peak Muscle Power in the Lower Limb of Older Adults. *Exp. Gerontol.* **2015**, *70*, 111–118. [[CrossRef](#)] [[PubMed](#)]
80. Hillman, C.H.; Erickson, K.I.; Kramer, A.F. Be Smart, Exercise Your Heart: Exercise Effects on Brain and Cognition. *Nat. Rev. Neurosci.* **2008**, *9*, 58–65. [[CrossRef](#)] [[PubMed](#)]
81. Kramer, A.F.; Colcombe, S.J.; McAuley, E.; Scalf, P.E.; Erickson, K.I. Fitness, Aging and Neurocognitive Function. *Neurobiol. Aging* **2005**, *26* (Suppl. S1), 124–127. [[CrossRef](#)]
82. Sáez de Astear, M.L.; Martínez-Velilla, N.; Zambom-Ferraresi, F.; Casas-Herrero, Á.; Izquierdo, M. Role of Physical Exercise on Cognitive Function in Healthy Older Adults: A Systematic Review of Randomized Clinical Trials. *Ageing Res. Rev.* **2017**, *37*, 117–134. [[CrossRef](#)] [[PubMed](#)]
83. Strassnig, M.T.; Signorile, J.F.; Potiaumpai, M.; Romero, M.A.; Gonzalez, C.; Czaja, S.; Harvey, P.D. High Velocity Circuit Resistance Training Improves Cognition, Psychiatric Symptoms and Neuromuscular Performance in Overweight Outpatients with Severe Mental Illness. *Psychiatry Res.* **2015**, *229*, 295–301. [[CrossRef](#)] [[PubMed](#)]

84. Cherup, N.; Roberson, K.; Potiaumpai, M.; Widdowson, K.; Jaghab, A.-M.; Chowdhari, S.; Armitage, C.; Seeley, A.; Signorile, J. Improvements in Cognition and Associations with Measures of Aerobic Fitness and Muscular Power Following Structured Exercise. *Exp. Gerontol.* **2018**, *112*, 76–87. [[CrossRef](#)]
85. Reid, K.F.; Martin, K.I.; Doros, G.; Clark, D.J.; Hau, C.; Patten, C.; Phillips, E.M.; Frontera, W.R.; Fielding, R.A. Comparative Effects of Light or Heavy Resistance Power Training for Improving Lower Extremity Power and Physical Performance in Mobility-Limited Older Adults. *J. Gerontol. Ser. A Biol. Sci. Med. Sci.* **2015**, *70*, 374–380. [[CrossRef](#)]
86. Laroche, D.P. Initial Neuromuscular Performance in Older Women Influences Response to Explosive Resistance Training. *Isokinet. Exerc. Sci.* **2009**, *17*, 197. [[CrossRef](#)]
87. Bean, J.F.; Kiely, D.K.; Herman, S.; Leveille, S.G.; Mizer, K.; Frontera, W.R.; Fielding, R.A. The Relationship between Leg Power and Physical Performance in Mobility-Limited Older People. *J. Am. Geriatr. Soc.* **2002**, *50*, 461–467. [[CrossRef](#)] [[PubMed](#)]
88. Sayers, S.P. High-Speed Power Training: A Novel Approach to Resistance Training in Older Men and Women. A Brief Review and Pilot Study. *J. Strength Cond. Res.* **2007**, *21*, 518–526. [[CrossRef](#)] [[PubMed](#)]
89. Metter, E.J.; Conwit, R.; Tobin, J.; Fozard, J.L. Age-Associated Loss of Power and Strength in the Upper Extremities in Women and Men. *J. Gerontol. Ser. A Biol. Sci. Med. Sci.* **1997**, *52*, 267–276. [[CrossRef](#)] [[PubMed](#)]
90. Häkkinen, K.; Kallinen, M.; Izquierdo, M.; Jokelainen, K.; Lassila, H.; Mälkiä, E.; Kraemer, W.J.; Newton, R.U.; Alen, M. Changes in Agonist-Antagonist EMG, Muscle CSA, and Force during Strength Training in Middle-Aged and Older People. *J. Appl. Physiol.* **1998**, *84*, 1341–1349. [[CrossRef](#)] [[PubMed](#)]
91. Häkkinen, K.; Kraemer, W.J.; Newton, R.U.; Alen, M. Changes in Electromyographic Activity, Muscle Fibre and Force Production Characteristics during Heavy Resistance/Power Strength Training in Middle-Aged and Older Men and Women. *Acta Physiol. Scand.* **2001**, *171*, 51–62. [[CrossRef](#)] [[PubMed](#)]
92. Aagaard, P.; Suetta, C.; Caserotti, P.; Magnusson, S.P.; Kjaer, M. Role of the Nervous System in Sarcopenia and Muscle Atrophy with Aging: Strength Training as a Countermeasure. *Scand. J. Med. Sci. Sports* **2010**, *20*, 49–64. [[CrossRef](#)]
93. McKinnon, N.B.; Connelly, D.M.; Rice, C.L.; Hunter, S.W.; Doherty, T.J. Neuromuscular Contributions to the Age-Related Reduction in Muscle Power: Mechanisms and Potential Role of High Velocity Power Training. *Ageing Res. Rev.* **2017**, *35*, 147–154. [[CrossRef](#)]
94. Earles, D.R.; Judge, J.O.; Gunnarsson, O.T. Velocity Training Induces Power-Specific Adaptations in Highly Functioning Older Adults. *Arch. Phys. Med. Rehabil.* **2001**, *82*, 872–878. [[CrossRef](#)]
95. Kamen, G. Aging, Resistance Training, and Motor Unit Discharge Behavior. *Can. J. Appl. Physiol. Rev. Can. Physiol. Appl.* **2005**, *30*, 341–351. [[CrossRef](#)]
96. Matias, C.N.; Nunes, C.L.; Francisco, S.; Tomeleri, C.M.; Cyrino, E.S.; Sardinha, L.B.; Silva, A.M. Phase Angle Predicts Physical Function in Older Adults. *Arch. Gerontol. Geriatr.* **2020**, *90*, 104151. [[CrossRef](#)]
97. Stergiou, N. *Biomechanics and Gait Analysis*; Elsevier: New York, NY, USA, 2020; ISBN 978-0-12-813440-5.
98. Ribeiro, A.S.; Aguiar, A.F.; Schoenfeld, B.J.; Nunes, J.P.; Cavalcante, E.F.; Cadore, E.L.; Cyrino, E.S. Effects of Different Resistance Training Systems on Muscular Strength and Hypertrophy in Resistance-Trained Older Women. *J. Strength Cond. Res.* **2018**, *32*, 545–553. [[CrossRef](#)] [[PubMed](#)]
99. Ribeiro, A.S.; Nunes, J.P.; Schoenfeld, B.J. Selection of Resistance Exercises for Older Individuals: The Forgotten Variable. *Sports Med. Auckl. NZ* **2020**, *50*, 1051–1057. [[CrossRef](#)] [[PubMed](#)]
100. Bottaro, M.; Machado, S.N.; Nogueira, W.; Scales, R.; Veloso, J. Effect of High versus Low-Velocity Resistance Training on Muscular Fitness and Functional Performance in Older Men. *Eur. J. Appl. Physiol.* **2007**, *99*, 257–264. [[CrossRef](#)] [[PubMed](#)]
101. Orr, R.; de Vos, N.J.; Singh, N.A.; Ross, D.A.; Stavrinou, T.M.; Fiatarone-Singh, M.A. Power Training Improves Balance in Healthy Older Adults. *J. Gerontol. Ser. A Biol. Sci. Med. Sci.* **2006**, *61*, 78–85. [[CrossRef](#)] [[PubMed](#)]
102. Reid, K.F.; Fielding, R.A. Skeletal Muscle Power: A Critical Determinant of Physical Functioning in Older Adults. *Exerc. Sport Sci. Rev.* **2012**, *40*, 4–12. [[CrossRef](#)]
103. Perera, S.; Mody, S.H.; Woodman, R.C.; Studenski, S.A. Meaningful Change and Responsiveness in Common Physical Performance Measures in Older Adults. *J. Am. Geriatr. Soc.* **2006**, *54*, 743–749. [[CrossRef](#)]
104. Ringsberg, K.; Gerdhem, P.; Johansson, J.; Obrant, K.J. Is There a Relationship between Balance, Gait Performance and Muscular Strength in 75-Year-Old Women? *Age Ageing* **1999**, *28*, 289–293. [[CrossRef](#)]
105. Karttunen, N.; Lihavainen, K.; Sipilä, S.; Rantanen, T.; Sulkava, R.; Hartikainen, S. Musculoskeletal Pain and Use of Analgesics in Relation to Mobility Limitation among Community-Dwelling Persons Aged 75 Years and Older. *Eur. J. Pain Lond. Engl.* **2012**, *16*, 140–149. [[CrossRef](#)]
106. Harridge, S.D.; Kryger, A.; Stensgaard, A. Knee Extensor Strength, Activation, and Size in Very Elderly People Following Strength Training. *Muscle Nerve* **1999**, *22*, 831–839. [[CrossRef](#)]
107. Clark, B.C.; Taylor, J.L.; Hong, S.L.; Law, T.D.; Russ, D.W. Weaker Seniors Exhibit Motor Cortex Hypoexcitability and Impairments in Voluntary Activation. *J. Gerontol. Ser. A Biol. Sci. Med. Sci.* **2015**, *70*, 1112–1119. [[CrossRef](#)]
108. Suetta, C.; Hvid, L.G.; Justesen, L.; Christensen, U.; Neergaard, K.; Simonsen, L.; Ortenblad, N.; Magnusson, S.P.; Kjaer, M.; Aagaard, P. Effects of Aging on Human Skeletal Muscle after Immobilization and Retraining. *J. Appl. Physiol.* **2009**, *107*, 1172–1180. [[CrossRef](#)] [[PubMed](#)]

109. Cadore, E.L.; Pinto, R.S.; Lhullier, F.L.R.; Correa, C.S.; Alberton, C.L.; Pinto, S.S.; Almeida, A.P.V.; Tartaruga, M.P.; Silva, E.M.; Kruegel, L.F.M. Physiological Effects of Concurrent Training in Elderly Men. *Int. J. Sports Med.* **2010**, *31*, 689–697. [[CrossRef](#)] [[PubMed](#)]
110. Hrada, K.V.; Hicks, A.L.; McCartney, N. Training for Muscle Power in Older Adults: Effects on Functional Abilities. *Can. J. Appl. Physiol.* **2003**, *28*, 178–189. [[CrossRef](#)] [[PubMed](#)]
111. Van Cutsem, M.; Duchateau, J.; Hainaut, K. Changes in Single Motor Unit Behaviour Contribute to the Increase in Contraction Speed after Dynamic Training in Humans. *J. Physiol.* **1998**, *513 Pt 1*, 295–305. [[CrossRef](#)]
112. Sardinha, L.B. Physiology of Exercise and Phase Angle: Another Look at BIA. *Eur. J. Clin. Nutr.* **2018**, *72*, 1323–1327. [[CrossRef](#)]
113. Hausdorff, J.M. Gait Dynamics, Fractals and Falls: Finding Meaning in the Stride-to-Stride Fluctuations of Human Walking. *Hum. Mov. Sci.* **2007**, *26*, 555–589. [[CrossRef](#)]
114. Stergiou, N.; Harbourne, R.; Cavanaugh, J. Optimal Movement Variability: A New Theoretical Perspective for Neurologic Physical Therapy. *J. Neurol. Phys. Ther. JNPT* **2006**, *30*, 120–129. [[CrossRef](#)]
115. Rahman, H.; Pipinos, I.I.; Johanning, J.M.; Myers, S.A. Gait Variability Is Affected More by Peripheral Artery Disease than by Vascular Occlusion. *PLoS ONE* **2021**, *16*, e0241727. [[CrossRef](#)]
116. Dingwell, J.B.; Salinas, M.M.; Cusumano, J.P. Increased Gait Variability May Not Imply Impaired Stride-to-Stride Control of Walking in Healthy Older Adults: Winner: 2013 Gait and Clinical Movement Analysis Society Best Paper Award. *Gait Posture* **2017**, *55*, 131–137. [[CrossRef](#)]
117. Norman, K.; Stobäus, N.; Pirlich, M.; Bösly-Westphal, A. Bioelectrical Phase Angle and Impedance Vector Analysis—Clinical Relevance and Applicability of Impedance Parameters. *Clin. Nutr.* **2012**, *31*, 854–861. [[CrossRef](#)]
118. Nunes, J.P.; Ribeiro, A.S.; Silva, A.M.; Schoenfeld, B.J.; dos Santos, L.; Cunha, P.M.; Nascimento, M.A.; Tomeleri, C.M.; Nabuco, H.C.G.; Antunes, M.; et al. Improvements in Phase Angle Are Related with Muscle Quality Index after Resistance Training in Older Women. *J. Aging Phys. Act.* **2019**, *27*, 515–520. [[CrossRef](#)] [[PubMed](#)]
119. Basile, C.; Della-Morte, D.; Cacciatore, F.; Gargiulo, G.; Galizia, G.; Roselli, M.; Curcio, F.; Bonaduce, D.; Abete, P. Phase Angle as Bioelectrical Marker to Identify Elderly Patients at Risk of Sarcopenia. *Exp. Gerontol.* **2014**, *58*, 43–46. [[CrossRef](#)] [[PubMed](#)]
120. Mattiello, R.; Amaral, M.A.; Mundstock, E.; Ziegelmann, P.K. Reference Values for the Phase Angle of the Electrical Bioimpedance: Systematic Review and Meta-Analysis Involving More than 250,000 Subjects. *Clin. Nutr.* **2020**, *39*, 1411–1417. [[CrossRef](#)] [[PubMed](#)]
121. Reljic, D.; Zarafat, D.; Jensen, B.; Herrmann, H.; Neurath, M.; Konturek, P.; Zopf, Y. Phase Angle and Vector Analysis from Multifrequency Segmental Bioelectrical Impedance Analysis: New Reference Data for Older Adults. *J. Physiol. Pharmacol.* **2020**, *71*. [[CrossRef](#)]