



AALBORG UNIVERSITY
DENMARK

Aalborg Universitet

A 20-year comparison of sex-specific fracture incidence between Type 1 and Type 2 diabetics and non-diabetics in Denmark

Nasser, Mohamad; Kvist, Annika; Vestergaard, Peter; Frost, Morten; Burden, Andrea M.

Published in:
Bone Reports

DOI (link to publication from Publisher):
[10.1016/j.bonr.2021.100771](https://doi.org/10.1016/j.bonr.2021.100771)

Creative Commons License
CC BY-NC-ND 4.0

Publication date:
2021

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Nasser, M., Kvist, A., Vestergaard, P., Frost, M., & Burden, A. M. (2021). A 20-year comparison of sex-specific fracture incidence between Type 1 and Type 2 diabetics and non-diabetics in Denmark. *Bone Reports*, 14(Suppl.), 4. [100771]. <https://doi.org/10.1016/j.bonr.2021.100771>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Methods: Mice stayed on a vibrating plate (2g/45Hz), 15 min/day for 6 weeks (plate kept still for CT). Bones were collected for microtomography, histomorphometry, RT-qPCR.

Results: Vibration reduced trabecular volume in OPN^{-/-} mice (vertebra BV/TV: -18%, $p < 0.05$) with increased osteoclast surfaces (Oc.S/BS: +72%, $p < 0.01$). OPN^{-/-} lost femoral cortical bone (Ct.Th: -9%, $p < 0.01$) under increased resorption (Ec.Oc.S/BS: +95%, $p < 0.05$). Strikingly, VB BSP^{-/-} mice gained cortical bone (Ct.Th: +7%, $p < 0.01$), with significant increase in formation, less osteoclasts (Ps.Oc.S/BS: -68%, $p < 0.05$) and lower RANKL and CathK expression. Sequential fluorochrome labeling (Figure 1) documented increased formation in VB BSP^{-/-} cortex (Figure 2). No effect was observed in other genotypes.




Figure 1: Sequential fluorochrome labeling on cortical bone using alizarin complexone (orange) and calcein blue (blue). VB BSP^{-/-} mouse.




Figure 2: Endocortical bone accretion in CT (empty circles) and VB (black circles) BSP^{-/-} mice. Values are mean \pm SEM of 6 mice.

Conclusion(s): BSP and OPN thus play crucial and opposite roles in bone response to mechanical stimulation. Because BSP^{-/-} mice overexpress OPN, and DKO mice showed no effect, interaction between these two factors seems required for a proper skeletal response to mechanical challenges.

doi:10.1016/j.bonr.2021.100770

Plenary Oral Presentations 2: Novel Aspects of Osteoporosis and Treatments

PLO07

A 20-year comparison of sex-specific fracture incidence between Type 1 and Type 2 diabetics and non-diabetics in Denmark

Mohamad Nasser^a, Annika Kvist^b, Morten Frost^a, Peter Vestergaard^{c,d}, Andrea M. Burden^b

^aUniversity of Southern Denmark, Department of Endocrinology and Metabolism- Molecular Endocrinology & Stem Cell Research Unit KMEB Odense University Hospital, Odense, Denmark

^bEth Zurich, Department of Chemistry and Applied Biosciences, Zurich, Switzerland

^cSteno Diabetes Centre, North Jutland, Aalborg, Denmark

^dAalborg University, Department of Endocrinology, Aalborg, Denmark

^eAalborg University, Department of Medicine, Aalborg, Denmark

Background/Introduction: Major osteoporotic fractures have declined in Nordic and western countries over the last two decades, despite an aging society of the general population. However, little is known if the decreasing trend of fractures is similar in patients with diabetes.

Purpose: To investigate the incidence rate (IR) of fractures in adult (age 18+) patients in Denmark between 1997-2017 with Type 1 Diabetes (T1D) and Type 2 Diabetes (T2D) compared to non-diabetics, stratified by sex.

Methods: All patients aged 18+ with a bone fracture (excluding skull and facial) between 1997 and 2017 were identified from the Danish National Health Service Register and linked to the Danish Medicines Agency Register of Medicinal Products Statistics (RMPS). Gender-specific IRs of fractures per 10,000 person years (PYs) were estimated. All analyses were stratified by diabetes diagnosis, defined as T1D, T2D, and non-diabetic.

Results: Overall, we identified a 25% decline in the IRs of fractures among T1D (from 571.6 to 427.0), a 58% decline among T2D (from 840.1 to 250.6), and a 10% decline among non-diabetics (from 179.2 to 161.5), between 1997 and 2017, respectively. Among males, the IRs declined 35% among T1D (from 499.3 to 322.9), 67% among T2D (from 786.9 to 263.8), and 10% among non-diabetics (from 191.8 to 186.9). Similarly, among women, we observed a 12% decline among T1D (from 666.3 to 586.3), a 49% decline among T2D (from 896.0 to 455.5), and 3% in non-diabetics (from 191.8 to 186.9).

Conclusion(s): In this 20-year population-based observation period, we identified a declining trend of fractures among diabetic patients, particularly among those with T2D. However, despite decreasing incidence, the IR of fractures in 2017 among patients with T1D or T2D was substantially higher than non-diabetics in Denmark, particularly among women, thereby highlighting the need to improve fracture management in these patients.

doi:10.1016/j.bonr.2021.100771

PLO08

Risk of fracture in patients with glucocorticoid requiring diseases

Giovanni Adami, Davide Gatti, Angelo Fassio, Alessandro Giollo, Ombretta Viapiana, Maurizio Rossini

University of Verona, Rheumatology Unit, Verona, Italy

Background/Introduction: Glucocorticoid-Induced Osteoporosis (GIOP) is the most common form of secondary osteoporosis. Nevertheless, the independent role of GCs and GC requiring diseases on fracture risk is still unclear.