



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

The Use of Cryotherapy in Conjunction with Surgical Removal of Mandibular Third Molars: a Single-Blinded Randomized Controlled Trial

Kjærgaard Larsen, Marie; Kofod, Thomas; Starch-Jensen, Thomas

Published in:
eJournal of Oral And Maxillofacial Research

DOI (link to publication from Publisher):
[10.5037/jomr.2021.12402](https://doi.org/10.5037/jomr.2021.12402)

Creative Commons License
CC BY-NC-ND 3.0

Publication date:
2021

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):
Kjærgaard Larsen, M., Kofod, T., & Starch-Jensen, T. (2021). The Use of Cryotherapy in Conjunction with Surgical Removal of Mandibular Third Molars: a Single-Blinded Randomized Controlled Trial. *eJournal of Oral And Maxillofacial Research*, 12(4), [e2]. <https://doi.org/10.5037/jomr.2021.12402>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

The Use of Cryotherapy in Conjunction with Surgical Removal of Mandibular Third Molars: a Single-Blinded Randomized Controlled Trial

Marie Kjærgaard Larsen¹, Thomas Kofod², Thomas Starch-Jensen³

¹Department of Oral and Maxillofacial Surgery, Aalborg University Hospital, Denmark.

²Department of Oral and Maxillofacial Surgery, Copenhagen University Hospital, Rigshospitalet, Denmark.

³Department of Oral and Maxillofacial Surgery, Aalborg University Hospital, Denmark.

Corresponding Author:

Marie Kjærgaard Larsen

Department of Oral and Maxillofacial Surgery

Aalborg University Hospital

18-22 Hobrovej, DK-9000 Aalborg

Denmark

Phone: +45 97 66 27 95

Fax: +45 97 66 28 25

E-mail: mkjaergaard@me.com

ABSTRACT

Objectives: Cryotherapy is frequently used to diminish postoperative sequelae following mandibular third molar surgery. The objective of this single-blinded randomized controlled trial was to assess the therapeutic efficiency of 30 minutes continuous cryotherapy on postoperative sequelae following surgical removal of mandibular third molars compared with no cryotherapy.

Material and Methods: Thirty patients (14 male and 16 female) including 60 mandibular third molars were randomly allocated to 30 minutes of immediately cryotherapy or no cryotherapy. Outcome measures included pain (visual analogue scale score), maximum mouth opening (trismus) and quality of life (oral health impact profile-14). Outcome measures were assessed preoperatively and one day, three days, seven days and one month following surgical removal of mandibular third molars. Descriptive and generalized estimating equation analyses were made. Level of significance was 0.05.

Results: No cryotherapy following surgical removal of mandibular third molars revealed a statistically significant lower visual analogue scale score of pain compared to thirty minutes of continuous cryotherapy after one day ($P < 0.05$). However, no statistically significant difference in trismus or oral health-related quality of life were revealed at any time point compared with no cryotherapy.

Conclusions: The therapeutic effect of 30 minutes continuous cryotherapy following surgical removal of mandibular third molars seem to be negligible. Thus, further randomized controlled trials assessing a prolonged application period of cryotherapy, alternative devices or use of intermittent cryotherapy are needed before definite conclusions and evidence-based clinical recommendations can be provided.

Keywords: cryotherapy; dentistry; mandible; pain; third molar; trismus.

Accepted for publication: 22 December 2021

To cite this article:

Kjærgaard Larsen M, Kofod T, Starch-Jensen T.

The Use of Cryotherapy in Conjunction with Surgical Removal of Mandibular Third Molars: a Single-Blinded Randomized Controlled Trial

J Oral Maxillofac Res 2021;12(4):e2

URL: <http://www.ejomr.org/JOMR/archives/2021/4/e2/v12n4e2.pdf>

doi: [10.5037/jomr.2021.12402](https://doi.org/10.5037/jomr.2021.12402)

INTRODUCTION

The most common surgical intervention undertaken in oral and maxillofacial surgery is surgical removal of mandibular third molars (SRM3). Postoperative pain, facial swelling, trismus and deterioration in postoperatively oral-health related quality of life are common and expected sequelae. Various therapies have been investigated to diminish postoperative sequelae including analgesics, antibiotics, corticosteroids, cryotherapy, low-laser therapy and compression [1-3]. Cryotherapy is a non-pharmacologic intervention that is frequently used following SRM3. Though, the therapeutic effect of cryotherapy seems to be controversy, which is emphasized in several systematic reviews [4]. Different techniques and devices are available for application of cryotherapy, including intermittent (applied for a shorter time and is reapplied several times over the duration of treatment) or continuous (applied for the duration of treatment and then removed) cryotherapy [4-6]. Frozen gel packs, ice packs wrapped in a washcloth or Hilotherm device are frequently used to reduce the skin and subcutaneous tissue temperature following SRM3 [5,7-11]. However, there are no evidence-based clinical recommendations regarding the most optimal technique, device or duration of treatment [4]. Cryotherapy is an easy, simple and non-pharmacologic intervention, which is generally accepted by most patients following SRM3. Use of continuous cryotherapy for a shorter period of time will have the least influence on the patient’s daily life, whereas long-lasting intermittent cryotherapy with the use of Hilotherm is time consuming, expensive and difficult to implement in dental practice. From a clinical and patient perspective, it would therefore be an advantage if continues cryotherapy for a limited period of time was capable of diminishing postoperative sequelae following SRM3. The objective of the present single-blinded randomized controlled trial was therefore to test the null-hypothesis of no difference in pain, trismus

and oral health-related quality of life following SRM3 with 30 minutes of immediate cryotherapy compared with no cryotherapy.

MATERIAL AND METHODS

Study design

The study was designed as a split-mouth single-blinded randomized controlled trial and conducted at the Department of Oral and Maxillofacial Surgery, Rigshospitalet, Copenhagen University Hospital, Denmark between October 2018 and January 2019. The study protocol was approved by Research Ethics Committee and the Danish Data Protection Agency (approval no.: N-20170016) and performed in accordance with the Declaration of Helsinki II and the Consolidated Standards of Reporting Trials (CONSORT) statement [12].

Patients scheduled for SRM3 prior to orthognathic surgery were screened. Patients with bilateral and comparable impacted mandibular third molars according to Pell and Gregory classification were invited to participate in the study [13]. Oral and written information regarding the study were explained to the patients, and written informed consent was obtained from every patient prior to enrolment. Participation in the study was voluntary and included patients could at any given time withdraw from the study.

Sample size calculation and study population

For obtaining a statistically significant treatment difference, a sample size was determined, expecting a difference of 20 mm in VAS score of pain at the first postoperative day, with an alpha value of 0.05 and a statistical power of 0.8. The sample size calculation was conducted using ClinCalc.com (<http://clincalc.com/stats/samplesize.aspx>, accessed 9th March 2017). Position of the mandibular third molars on panoramic radiographs were classified using Pell and Gregory system and Winter’s classification [13]. The inclusion and exclusion criteria are outlined in Table 1.

Table 1. Inclusion and exclusion criteria

Inclusion criteria	Exclusion criteria
<ul style="list-style-type: none"> - Bilateral symmetrical impacted mandibular third molars. - Indication for removal of mandibular third molars. - Age between 18 and 40 years. 	<ul style="list-style-type: none"> - Infections and inflammatory symptoms in the oral cavity at the time of surgery. - Previous maxillofacial trauma. - Craniofacial clefts or syndromes. - Systemic bone disease (i.e. arthritis) or diabetes mellitus. - Active acne vulgaris, viral, and fungal infections. - Psychological disease. - Pregnancy and breastfeeding. - Failure to attend follow-up.

Randomization and blinding

The mandibular third molars were randomly allocated to 30 minutes of immediate continuous cryotherapy or no cryotherapy using a computer-aided block randomization. A randomization sheet with a serial number from 1 to 30 were made (<http://www.randomization.com>, Randomization.com, date: 22nd August 2018). The numbers were stored in sealed envelopes, and every patient opened an envelope with a number. The assistant nurse combined the number with the randomization sheet to allocate the mandibular third molar to cryotherapy or no cryotherapy. The randomization sheet was kept by the assistant nurse until the study was unblinded.

The assistant nurse placed the cold gel pack on the patient's cheek following SRM3 and removed the gel pack after 30 minutes. Surgeon and assessor were therefore blinded in relation to the applied treatment.

Surgical procedure

The included patients underwent SRM3 in local anaesthesia by the same surgeon (MKL) using a standard technique. One mandibular third molar was removed at each time. Surgical removal of the second mandibular third molar was performed approximately one to six weeks after the first surgery.

Prior to the surgery, the patients received prophylactic analgesic including 400 mg ibuprofen (Ipre[®] - Takeda Pharma A/S; Hobro, Denmark) and 1,000 mg paracetamol (Pinex[®], Actavis Nordic A/S; Søborg, Denmark).

The inferior alveolar nerve and the lingual nerve were anaesthetized with 20 mg/mL mepivacaine hydrochloride and 5 µg/mL adrenaline (Carbocain-Adrenalin[®] - AstraZeneca; Copenhagen, Denmark). An incision from the anterior border of the ascending ramus of the mandible to the distal part of the lower first molar was performed. The mucosal flap was elevated and the bone around the mandibular third molar was removed with a round burr under irrigation with 0.9% saline solution. If necessary, the mandibular third molar was sectioned with a fissure bur. The tooth was elevated out, and the extraction socket and surrounding bone was irrigated with 0.9% saline solution. The surgical site was sutured with resorbable suture (Ethicon Vicryl Rapide[™] suture 4-0 - Johnson and Johnson Medical GmbH; Norderstedt, Germany). Time of surgery was measured from the incision until the last suture was made.

If the mandibular third molar was allocated for cryotherapy, a freezable cold gel pack (Cool Jaw[®] Soft Stretch Jaw Wrap with Cold Packs - Medico

International Inc.; Palmer, Pennsylvania, USA) and a jaw bra was applied on the cheek immediately following SRM3 for 30 minutes.

All patients received standard postoperative instructions and pain medication including mouth rinse with 0.12% chlorhexidine three times a day (Klorhexidin Mundskyl 0.12% - Faaborg Pharma; Faaborg, Denmark), 400 mg of ibuprofen three times a day (Ipre[®]) and 1,000 mg paracetamol four times a day (Pinex[®]).

Data collection

Clinical assessment was obtained preoperatively (T0), one day (T1), three days (T2), seven days (T3) and one month (T4) following SRM3. All data were collected by the same assessor (MKL).

Pain was evaluated by a 100-mm VAS-score obtained preoperatively (T0), one day (T1), three days (T2), seven days (T3), and one month (T4) postoperatively. All patients were carefully instructed in the use of VAS and had to mark on the line the point that represented their pain.

Mouth opening was measured as the maximum distance between the upper and lower incisal edges with a ruler preoperatively (T0), three days (T2), seven days (T3) and one month (T4) following SRM3. Oral health-related quality of life was evaluated by oral health impact profile 14 (OHIP-14) questionnaire obtained preoperatively (T0), seven days (T3), and one month (T4) following SRM3. OHIP-14 is composed of 14 questions and organised into seven dimensions including functional limitation, physical discomfort, psychological discomfort, physical disability, psychological disability, social disability and handicap (Table 2). The OHIP-14 scale ranged from 0 to 56 and dimension score ranged from 0 to 8. The values of the 14 items and each dimension were summed to calculate the OHIP-14 severity score with higher scores indicating poorer oral health-related quality of life. Patients were carefully instructed in the OHIP-14 questionnaire and completed the questionnaires by themselves.

Complications including infections, alveolitis sicca, lost sutures and dehiscence were registered three days (T2), seven days (T3) and one month (T4) following SRM3.

Statistical analysis

Anatomical position of the mandibular third molars was presented as counts and percentage on each treatment group. The time of surgery was presented with mean, standard deviation, minimum and maximum.

Table 2. OHIP-14 score

	Question ^a
Functional limitation	Have you had trouble pronouncing any words because of problems with your teeth, mouth or dentures?
	Have you felt that your sense of taste has worsened because of problems with your teeth, mouth or dentures?
Physical pain	Have you had painful aching in your mouth?
	Have you found it uncomfortable to eat any foods because of problems with your teeth, mouth or dentures?
Psychological discomfort	Have you been self-conscious because of your teeth, mouth or dentures?
	Have you felt tense because of problems with your teeth, mouth or dentures?
Physical disability	Has your diet been unsatisfactory because of problems with your teeth, mouth or dentures?
	Have you had to interrupt meals because of problems with your teeth, mouth or dentures?
Psychological disability	Have you found it difficult to relax because of problems with your teeth, mouth or dentures?
	Have you been a bit embarrassed because of problems with your teeth, mouth or dentures?
Social disability	Have you been a bit irritable with other people because of problems with your teeth, mouth or dentures?
	Have you had difficulty doing your usual jobs because of problems with your teeth, mouth or dentures?
Handicap	Have you felt that life in general was less satisfying because of problems with your teeth, mouth or dentures?
	Have you been totally unable to function because of problems with your teeth, mouth or dentures?

^aAnswers: 0 = never; 1 = hardly ever or nearly never; 2 = occasionally; 3 = fairly often or many times; 4 = very often.

Mean difference in pain and trismus were analysed with a generalized estimating equation (GEE) analysis for repeated observations. Oral health-related quality of life was presented with mean and standard deviation for OHIP-14 score. Missing observations in outcome variables were assumed to be missing randomly. The estimated mean value for pain, trismus and oral health-related quality of life were expressed with a 95% confidence interval (CI). Statistical significance of the P-value was set at 0.05. The analyses were descriptive and adjusted for age, sex, smoking and time of surgery. Parametric data were expressed as mean and standard deviation (M [SD]).

Data management and statistical analysis was performed with Excel version 2013 (Microsoft; Redmond, Washington, USA) and R software version 3.6.1 (R Foundation for Statistical Computing; Vienna, Austria; <http://www.r-project.org>).

RESULTS

Study population

Thirty patients (14 male and 16 female) with a mean age of 22.6 (SD 4.7) years were included resulting in 60 mandibular third molars (Figure 1). Length of surgery was 10.13 (SD 3.94) minutes with no statistically significant difference between test and control group (P = 0.438) (Table 3). The contralateral third molar was removed after 21 days (range 7 to 35 days).




Figure 1. Consort flow diagram.

Table 3. Anatomical position of mandibular third molars and time of surgery in the two groups and total

Variable	Level	No cryotherapy (n = 30)	Cryotherapy (n = 30)	Total (n = 60)
Anatomical position (Winter), N (%)	1	2 (6.7)	6 (20)	8 (13.3)
	2	5 (16.7)	7 (23.3)	12 (20)
	3	18 (60)	14 (46.7)	32 (53.3)
	4	5 (16.7)	3 (10)	8 (13.3)
Anatomical position (Pell and Gregory transversal), N (%)	1	0 (0)	0 (0)	0 (0)
	2	23 (76.7)	25 (83.3)	48 (80)
	3	7 (23.3)	5 (16.7)	12 (20)
Anatomical position (Pell and Gregory vertical), N (%)	1	3 (10)	5 (16.7)	8 (13.3)
	2	19 (63.3)	20 (66.7)	39 (65)
	3	8 (26.7)	5 (16.7)	13 (21.7)
Time of surgery (minutes)	Mean (SD)	9.83 (4.06) ^a	10.43 (3.87) ^a	10.13 (3.94) ^b
	Q1; Q3	7.25; 11	8.25; 12	7.75; 12
	Min; max	0; 20	0; 19	0; 20

n = number of wisdom teeth; Q1 = first quartile; Q3 = third quartile; SD = standard deviation.

^aMissing one mandibular third molar.

^bMissing two mandibular third molars.

Postoperative instructions were followed by all patients. Infection involving either fever, chills, sore lymph nodes and pus occurred following removal of five third molars, which were treated sufficiently with antibiotics involving phenoxymethylpenicillin 800 mg (Primcillin® - Meda A/S; Ballerup, Denmark) four times a day and metronidazole 500 mg (Metronidazol “DAK” - Takeda Pharma A/S; Hobro, Denmark) two times a day for seven days.

Pain

VAS score of pain with cryotherapy was 2.79 (SD 7.34) (T0), 60.97 (SD 21.66) (T1), 36.86 (SD 16.11) (T2), 11.30 (SD 11.96) (T3) and 5 (SD 100) (T4). VAS score of pain without cryotherapy was 2.71 (SD 6.95) (T0), 49.61 (SD 27.53) (T1), 29.64 (SD 22.1) (T2), 18.57 (SD 20.27) (T3) and 4 (SD 100) (T4). Patients without cryotherapy had a statistically significant lower VAS score of pain compared with 30 minutes of cryotherapy after one day (P < 0.05). There was no statistically significant difference between the two treatment modalities at any other time point (Table 4). In addition, there was no statistically significant difference in VAS score of pain between the two treatment modalities at any time points, when groups were adjusted for sex, smoking and time of surgery. However, VAS score of pain was 1.28 mm higher, when the age increased with 1 year after 1 month (T4), which was statistically significant (P < 0.05).

Mouth opening

Maximum mouth opening with cryotherapy was

48.72 (SD 9.25) mm (T0), 30.79 (SD 5.92) mm (T2), 41.26 (SD 8.31) mm (T3) and 48 (SD 11.22) (T4). Maximum mouth opening without cryotherapy was 50.03 (SD 8.82) mm (T0), 33.31 (SD 9.1) mm (T2), 40 (SD 8.18) mm (T3) and 42.6 (SD 10.97) mm (T4). There was no statistically significant difference between the two treatment modalities at any time point (Table 4).

There were statistically significant differences in maximum mouth opening between the two treatment modalities, when the groups were adjusted for age, sex, smoking and time of surgery. Mouth opening was restricted with 0.1 mm, when the age increased with one year after one month (T4), which was statistically significant (P < 0.05). Females had a statistically significant decreased mouth opening compared with males after one month (P < 0.05). Maximum mouth opening was statistically significant restricted after seven days (T3) and one month (T4), when the patients were smokers (P < 0.05). Maximum mouth opening was statistically significant decreased with increasing length of surgery (P < 0.05). The mouth opening decreased by 2.95 mm after seven days (T3), when length of surgery increased by one minute.

Oral health-related quality of life

Mean OHIP-14 score with 30 minutes of cryotherapy was 9.5 (SD 23.1) (T0) and 18.9 (SD 26.4) (T3), and mean OHIP-14 score without cryotherapy was 9.5 (SD 19.8) (T0) and 14.4 (SD 25.7) (T3) (Table 5). The mean OHIP-14 score was highest with 30 minutes of cryotherapy compared to no cryotherapy.

Table 4. Results before removal of M3 (T0) compared with one day (T1), three days (T2), seven days (T3) and one month (T4)

	Cryotherapy	T1-T0				T2-T0				T3-T0				T4-T0			
		Estimate	95% CI	SE	P	Estimate	95% CI	SE	P	Estimate	95% CI	SE	P	Estimate	95% CI	SE	P
Pain	without cryotherapy	Reference	-	-	-	Reference	-	-	-	Reference	-	-	-	Reference	-	-	-
	with cryotherapy	11.15	0.72; 21.57	5.32	< 0.05	6.87	-0.91; 14.66	3.971	0.083	-7.07	-15.34; 1.19	4.216	0.093	7.5	-5.23; 20.23	6.495	0.248
Trismus	without cryotherapy	Reference	-	-	-	Reference	-	-	-	Reference	-	-	-	Reference	-	-	-
	with cryotherapy	-	-	-	-	-1.4	-6.41; 3.61	2.554	0.584	3.68	-0.87; 8.23	2.321	0.113	1.45	-12.38; 15.28	7.054	0.837

CI = confidence interval; M3 = mandibular third molar; SE = standard error; VAS = visual analog scale.

Third molars allocated to no cryotherapy were used as reference for the group with cryotherapy.

Pain: assessed by VAS. Estimated value reveals differences in VAS score of pain in millimetres with cryotherapy compared with no cryotherapy.

Trismus: assessed by a ruler. Estimated value shows how many mm the incisal distance has increased or decreased compared to no cryotherapy.

Mean difference in pain and trismus were analyzed with a generalized estimating equation (GEE) analysis for repeated observations.

Table 5. OHIP-14 score

Question		T0											T3												
		Cryotherapy (n = 29)						No cryotherapy (n = 29)					Cryotherapy (n = 25)						No cryotherapy (n = 25)						
		0	1	2	3	4	SDS	0	1	2	3	4	SDS	0	1	2	3	4	SDS	0	1	2	3	4	SDS
Functional limitation	1	19	5	5	-	-	-	18	5	3	-	-	-	12	3	10	1	-	-	9	13	1	1	-	-
	2	23	2	1	-	-	19	24	3	-	-	-	14	16	5	2	3	-	44	14	6	1	2	-	32
Physical pain	3	9	7	12	1	-	-	7	8	12	-	-	-	1	3	12	6	4	-	1	5	8	4	5	-
	4	10	9	8	-	1	63	10	7	7	3	-	62	2	4	9	6	5	121	2	4	8	5	4	104
Psychological discomfort	5	12	6	5	4	2	-	12	7	4	4	1	-	14	3	5	3	1	-	18	-	3	3	-	-
	6	9	10	7	2	1	70	11	5	9	1	1	61	8	5	7	2	4	67	7	4	7	4	1	49
Physical disability	7	15	9	3	2	-	-	14	8	5	-	1	-	6	7	6	5	2	-	8	3	9	1	3	-
	8	17	9	2	-	1	38	15	8	3	-	2	44	7	8	7	4	-	76	9	8	2	3	1	61
Psychological disability	9	15	5	5	3	-	-	9	10	5	1	1	-	4	8	10	2	2	-	8	7	2	5	1	-
	10	16	2	7	3	1	53	15	7	4	1	1	49	14	4	7	0	1	64	14	4	5	1	-	47
Social disability	11	17	10	2	-	-	-	19	6	2	1	-	-	9	8	8	-	-	-	16	4	3	1	-	-
	12	26	3	-	-	-	17	21	5	2	-	-	22	11	6	4	2	3	56	12	4	5	2	1	37
Handicap	13	21	3	5	-	-	-	18	5	4	1	-	-	13	4	4	4	-	-	11	5	4	1	1	-
	14	27	2	-	-	-	15	23	4	1	-	-	22	15	5	3	3	-	44	18	5	-	-	1	29
Total OHIP-14 score		275						274					472						359						
Mean (SD)		9.5 (23.1)						9.5 (19.8)					18.9 (26.4)						14.4 (25.7)						

0 = never; 1 = hardly ever or nearly never; 2 = occasionally; 3 = fairly often or many times; 4 = very often.

n = number of wisdom teeth; OHIP-14 = oral health impact profile 14; SD = standard deviation; SDS = subscale dimension score.

Physical pain, psychological disability and discomfort presented highest OHIP-14-dimension score, while handicap and functional limitation exhibited the lowest score.

OHIP-14 questionnaires were missed in one case with 30 minutes of cryotherapy and in five cases with no cryotherapy.

DISCUSSION

The objective of the present single-blinded randomized controlled trial was to test the null-hypothesis of no difference in pain, trismus and oral health-related quality of life following SRM3 with 30 minutes of immediate cryotherapy compared with no cryotherapy.

A significant lower VAS score of pain was seen with no cryotherapy after one day compared with 30 minutes of cryotherapy ($P < 0.05$), whereas no significant differences were seen after three days ($P = 0.083$), seven days ($P = 0.093$) and one month ($P = 0.248$), respectively. Furthermore, no significant difference in trismus or oral-health related quality of life was observed at any time point. Consequently, 30 minutes of immediate continuous cryotherapy seems not to diminish postoperative sequelae following SRM3.

Pain is an expected sequelae following SRM3, which may interfere with patients' immediate oral health-related quality of life [14,15]. Within the first 24 hours postoperatively, the pain reaches highest intensity and gradually resolves after seven days [16,17]. Previous published systematic reviews have reported negligible effect of short-term continuous cryotherapy, which is in accordance with the results of the present study [18-20]. Consequently, continuous short-term cryotherapy seems not to diminish postoperative sequelae following SRM3. However, previous studies assessing long-lasting intermittent cryotherapy have demonstrated a significant beneficial effect on pain following SRM3 compared with the use of no cryotherapy [8,21]. Intermittent cryotherapy therefore appears to be more effective on postoperative pain following SRM3 compared with continuous cryotherapy [8,21]. During continuous cryotherapy with a gel pack, the temperature will not be held constant during the treatment period. The initial temperature will be low, and the end temperature will be warmer. In addition, the thickness of the subcutaneous adipose tissue may also influence the therapeutic effect of cryotherapy [22]. The therapeutic effect of continuous and intermittent cryotherapy on pain following SRM3 is inconclusive. Further studies are needed before final

conclusions can be made.

The most frequently used analgesics to control postoperative pain following SRM3 are paracetamol and non-steroidal anti-inflammatory drugs either alone or in combination [23]. A systematic review have shown, that non-steroidal anti-inflammatory drugs in conjunction with SRM3 diminish the physiological inflammatory response, reduce the average pain scores and the overall consumption of analgesics [24]. In the present study, patients' total consumption of analgesics was not systematically registered and completing VAS score of pain according to the time of ingestion of analgesics was not standardized, which potentially have influenced patient's perception of pain. Consequently, further randomized control trials assessing cryotherapy in conjunction with SRM3 should therefore monitor total consumption of analgesic including standardized regimes for ingestion of analgesic according to completion of VAS, questionnaires or alternative assessment methods.

Gender, age, smoking, time of surgery, surgical difficulty, flap design and intraoperative visibility of the inferior alveolar nerve are considered as predictive variables for increased pain following SRM3 [15,25]. In the present study, increasing age revealed a higher VAS score of pain compared with younger, whereas no significant difference in VAS score of pain was revealed according to smoking habits, length of surgery or gender. Uniformity of patient sample are therefore mandatory to diminish confounding variables. The included patients of the present study disclosed no significant difference in patient demographic, position of the mandibular third molars and time of surgery between the two treatment modalities.

Temporary trismus occurs frequently following SRM3 [26]. Linear measurement of the interincisal distance using various measurement tools is a simple and reliable method, which is commonly used for assessment of trismus following SRM3 [7,21]. In the present study, linear interincisal measurements revealed no significant difference in trismus following SRM3 with short term continuous cryotherapy compared with no cryotherapy. However, long-lasting intermittent cryotherapy have demonstrated a significant reduction in trismus compared with no cryotherapy [8]. Consequently, future studies assessing different application methods and treatment periods are needed before the beneficial effect of cryotherapy on trismus can be concluded.

Prolonged time of surgery results in increased trismus following SRM3, which is in accordance with the results of this randomized controlled trial [27-29].

Furthermore, smokers demonstrated more trismus compared to non-smokers, which previously has been reported [29]. Temporary trismus following SRM3 are influenced by the length of surgery, smoking habits and the surgical trauma, while the effect of cryotherapy seems negligible.

Deterioration in oral health-related quality of life is frequently seen following SRM3 [16,30]. In the present trial, 30 minutes of cryotherapy revealed a higher mean OHIP-14 score compared to no cryotherapy. Previous studies have demonstrated that continuous and intermittent cryotherapy improve oral health-related quality of life following SRM3 [5,8,31]. Oral health-related quality of life is influenced by age, gender, occlusion, present dental disease, previous dental experience, socioeconomic status, education, physical pain, psychological discomfort and psychological disability [32-34]. OHIP-14 evaluates patient's overall oral impairment and does not focus on a specific surgical intervention as i.e., SRM3. Consequently, further studies assessing oral health-related quality of life following SRM3 should include additional self-administrated questionnaires focusing on patient's perception of SRM3. Furthermore, self-administrated questionnaires are also recommended to include an association between oral health-related quality of life and demographic factors and socioeconomic status.

No statistically significant differences in pain and trismus between 30 minutes of continuous cryotherapy compared to no cryotherapy were seen. Though, the present randomized controlled trial is characterized by certain limitations including small sample size, no systematically registration of consumption of analgesics and no registration of the applied temperature of the jaw bra, which may have affected the outcome. In addition, socioeconomic status, educational background

and level of daily physical activity were not registered, which significantly influence patient's perception of recovery, pain and oral health-related quality of life following SRM3 [32,34]. Therefore, the conclusions drawn from the results of this study should be interpreted with caution.

CONCLUSIONS

The therapeutic effect of 30 minutes continuous cryotherapy following surgical removal of mandibular third molars seems to be negligible.

Further randomized controlled trials assessing longer therapy of cryotherapy, intermittent application and other devices for cryotherapy are therefore needed before definite conclusions and evidence-based clinical recommendations can be provided in diminishing postoperative sequelae following surgical removal of mandibular third molars.

ACKNOWLEDGMENTS AND DISCLOSURE STATEMENTS

A special thanks to Secretary Tanja Møller Andersen, Department of Oral and Maxillofacial Surgery, Rigshospitalet, Copenhagen University Hospital, Denmark for planning and booking of patients. Furthermore, a special thanks to Rikke Nørmark, Unit of Epidemiology and Biostatistics, Aalborg University Hospital, Aalborg, Denmark for statistic help.

The study protocol was approved by the Research Ethics Committee and the Danish Data Protection Agency.

The authors declare none competing interests. There was no external funding, Patient consent was not required.

REFERENCES

1. Atkinson HC, Currie J, Moodie J, Carson S, Evans S, Worthington JP, Steenberg LJ, Bisley E, Frampton C. Combination paracetamol and ibuprofen for pain relief after oral surgery: a dose ranging study. *Eur J Clin Pharmacol*. 2015 May;71(5):579-87. [Medline: [25778933](#)] [doi: [10.1007/s00228-015-1827-x](#)]
2. Juhl GI, Norholt SE, Tonnesen E, Hiesse-Provost O, Jensen TS. Analgesic efficacy and safety of intravenous paracetamol (acetaminophen) administered as a 2 g starting dose following third molar surgery. *Eur J Pain*. 2006 May;10(4):371-7. [Medline: [16085437](#)] [doi: [10.1016/j.ejpain.2005.06.004](#)]
3. Koçer G, Yuce E, Tuzuner Oncul A, Dereci O, Koskan O. Effect of the route of administration of methylprednisolone on oedema and trismus in impacted lower third molar surgery. *Int J Oral Maxillofac Surg*. 2014 May;43(5):639-43. [Medline: [24332587](#)] [doi: [10.1016/j.ijom.2013.11.005](#)]
4. Larsen MK, Kofod T, Starch-Jensen T. Therapeutic efficacy of cryotherapy on facial swelling, pain, trismus and quality of life after surgical removal of mandibular third molars: A systematic review. *J Oral Rehabil*. 2019 Jun;46(6):563-573. [Medline: [30869171](#)] [doi: [10.1111/joor.12789](#)]

5. Forouzanfar T, Sabelis A, Ausems S, Baart JA, van der Waal I. Effect of ice compression on pain after mandibular third molar surgery: a single-blind, randomized controlled trial. *Int J Oral Maxillofac Surg.* 2008 Sep;37(9):824-30. [Medline: [18603410](#)] [doi: [10.1016/j.ijom.2008.05.011](#)]
6. Greenstein G. Therapeutic efficacy of cold therapy after intraoral surgical procedures: a literature review. *J Periodontol.* 2007 May;78(5):790-800. [Medline: [17470011](#)] [doi: [10.1902/jop.2007.060319](#)]
7. Zandi M, Amini P, Keshavarz A. Effectiveness of cold therapy in reducing pain, trismus, and oedema after impacted mandibular third molar surgery: a randomized, self-controlled, observer-blind, split-mouth clinical trial. *Int J Oral Maxillofac Surg.* 2016 Jan;45(1):118-23. [Medline: [26597577](#)] [doi: [10.1016/j.ijom.2015.10.021](#)]
8. Ibikunle AA, Adeyemo WL. Oral health-related quality of life following third molar surgery with or without application of ice pack therapy. *Oral Maxillofac Surg.* 2016 Sep;20(3):239-47. [Medline: [27139019](#)] [doi: [10.1007/s10006-016-0558-1](#)]
9. Forsgren H, Heimdahl A, Johansson B, Krekmanov L. Effect of application of cold dressings on the postoperative course in oral surgery. *Int J Oral Surg.* 1985 Jun;14(3):223-8. [Medline: [3926665](#)] [doi: [10.1016/S0300-9785\(85\)80032-6](#)]
10. Glass GE, Waterhouse N, Shakib K. Hilotherapy for the management of perioperative pain and swelling in facial surgery: a systematic review and meta-analysis. *Br J Oral Maxillofac Surg.* 2016 Oct;54(8):851-856. [Medline: [27516162](#)] [doi: [10.1016/j.bjoms.2016.07.003](#)]
11. Bates AS, Knevil GJ. Systematic review and meta-analysis of the efficacy of hilotherapy following oral and maxillofacial surgery. *Int J Oral Maxillofac Surg.* 2016 Jan;45(1):110-7. [Medline: [26362489](#)] [doi: [10.1016/j.ijom.2015.08.983](#)]
12. Moher D, Hopewell S, Schulz KF, Montori V, Gøtzsche PC, Devereaux PJ, Elbourne D, Egger M, Altman DG; CONSORT. CONSORT 2010 explanation and elaboration: updated guidelines for reporting parallel group randomised trials. *Int J Surg.* 2012;10(1):28-55. [Medline: [22036893](#)] [doi: [10.1016/j.ijso.2011.10.001](#)]
13. MacGregor AJ. The radiological assessment of ectopic lower third molars. *Ann R Coll Surg Engl.* 1979 Mar;61(2):107-13. [Medline: [434748](#)] [PMC free article: [2492802](#)]
14. Colorado-Bonnin M, Valmaseda-Castellón E, Berini-Aytés L, Gay-Escoda C. Quality of life following lower third molar removal. *Int J Oral Maxillofac Surg.* 2006 Apr;35(4):343-7. [Medline: [16280233](#)] [doi: [10.1016/j.ijom.2005.08.008](#)]
15. Bortoluzzi MC, Guollo A, Capella DL. Pain levels after third molar surgical removal: an evaluation of predictive variables. *J Contemp Dent Pract.* 2011 Jul 1;12(4):239-44. [Medline: [22186857](#)] [doi: [10.5005/jp-journals-10024-1041](#)]
16. Duarte-Rodrigues L, Miranda EFP, Souza TO, de Paiva HN, Falci SGM, Galvão EL. Third molar removal and its impact on quality of life: systematic review and meta-analysis. *Qual Life Res.* 2018 Oct;27(10):2477-2489. [Medline: [29797177](#)] [doi: [10.1007/s11136-018-1889-1](#)]
17. Chopra D, Rehan HS, Mehra P, Kakkar AK. A randomized, double-blind, placebo-controlled study comparing the efficacy and safety of paracetamol, serratiopeptidase, ibuprofen and betamethasone using the dental impaction pain model. *Int J Oral Maxillofac Surg.* 2009 Apr;38(4):350-5. [Medline: [19168326](#)] [doi: [10.1016/j.ijom.2008.12.013](#)]
18. Larsen MK, Kofod T, Starch-Jensen T. Therapeutic efficacy of cryotherapy on facial swelling, pain, trismus and quality of life after surgical removal of mandibular third molars: A systematic review. *J Oral Rehabil.* 2019 Jun;46(6):563-573. [Medline: [30869171](#)] [doi: [10.1111/joor.12789](#)]
19. Osunde OD, Adebola RA, Omeje UK. Management of inflammatory complications in third molar surgery: a review of the literature. *Afr Health Sci.* 2011 Sep;11(3):530-7. [Medline: [22275950](#)] [PMC free article: [3261018](#)]
20. Hubbard TJ, Aronson SL, Denegar CR. Does Cryotherapy Hasten Return to Participation? A Systematic Review. *J Athl Train.* 2004 Mar;39(1):88-94. [Medline: [15085216](#)] [PMC free article: [385267](#)]
21. Laureano Filho JR, de Oliveira e Silva ED, Batista CI, Gouveia FM. The influence of cryotherapy on reduction of swelling, pain and trismus after third-molar extraction: a preliminary study. *J Am Dent Assoc.* 2005 Jun;136(6):774-8; quiz 807. [Medline: [16022042](#)] [doi: [10.14219/jada.archive.2005.0261](#)]
22. Otte JW, Merrick MA, Ingersoll CD, Cordova ML. Subcutaneous adipose tissue thickness alters cooling time during cryotherapy. *Arch Phys Med Rehabil.* 2002 Nov;83(11):1501-5. [Medline: [12422316](#)] [doi: [10.1053/apmr.2002.34833](#)]
23. Bailey E, Worthington HV, van Wijk A, Yates JM, Coulthard P, Afzal Z. Ibuprofen and/or paracetamol (acetaminophen) for pain relief after surgical removal of lower wisdom teeth. *Cochrane Database Syst Rev.* 2013 Dec 12;(12):CD004624. [Medline: [24338830](#)] [doi: [10.1002/14651858.CD004624.pub2](#)]
24. Cetira Filho EL, Carvalho FSR, de Barros Silva PG, Barbosa DAF, Alves Pereira KM, Ribeiro TR, Costa FWG. Preemptive use of oral nonsteroidal anti-inflammatory drugs for the relief of inflammatory events after surgical removal of lower third molars: A systematic review with meta-analysis of placebo-controlled randomized clinical trials. *J Craniomaxillofac Surg.* 2020 Mar;48(3):293-307. [Medline: [32070661](#)] [doi: [10.1016/j.jcms.2020.01.016](#)]
25. Benediktsdóttir IS, Wenzel A, Petersen JK, Hintze H. Mandibular third molar removal: risk indicators for extended operation time, postoperative pain, and complications. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2004 Apr;97(4):438-46. [Medline: [15088029](#)] [doi: [10.1016/j.tripleo.2003.10.018](#)]
26. Baqain ZH, Karaky AA, Sawair F, Khraisat A, Duaibis R, Rajab LD. Frequency estimates and risk factors for postoperative morbidity after third molar removal: a prospective cohort study. *J Oral Maxillofac Surg.* 2008 Nov;66(11):2276-83. [Medline: [18940492](#)] [doi: [10.1016/j.joms.2008.06.047](#)]

27. Grossi GB, Maiorana C, Garramone RA, Borgonovo A, Creminelli L, Santoro F. Assessing postoperative discomfort after third molar surgery: a prospective study. *J Oral Maxillofac Surg.* 2007 May;65(5):901-17. [Medline: [17448840](#)] [doi: [10.1016/j.joms.2005.12.046](#)]
28. Bello SA, Adeyemo WL, Bamgbose BO, Obi EV, Adeyinka AA. Effect of age, impaction types and operative time on inflammatory tissue reactions following lower third molar surgery. *Head Face Med.* 2011 Apr 28;7:8. [Medline: [21527036](#)] [PMC free article: [3114767](#)] [doi: [10.1186/1746-160X-7-8](#)]
29. Larsen MK, Kofod T, Duch K, Starch-Jensen T. Efficacy of methylprednisolone on pain, trismus and quality of life following surgical removal of mandibular third molars: a double-blind, split-mouth, randomised controlled trial. *Med Oral Patol Oral Cir Bucal.* 2021 Mar 1;26(2):e156-e163. [Medline: [32701926](#)] [PMC free article: [7980302](#)] [doi: [10.4317/medoral.24094](#)]
30. Phillips C, White RP Jr, Shugars DA, Zhou X. Risk factors associated with prolonged recovery and delayed healing after third molar surgery. *J Oral Maxillofac Surg.* 2003 Dec;61(12):1436-48. [Medline: [14663809](#)] [doi: [10.1016/j.joms.2003.08.003](#)]
31. van der Westhuijzen AJ, Becker PJ, Morkel J, Roelse JA. A randomized observer blind comparison of bilateral facial ice pack therapy with no ice therapy following third molar surgery. *Int J Oral Maxillofac Surg.* 2005 May;34(3):281-6. [Medline: [15741037](#)] [doi: [10.1016/j.ijom.2004.05.006](#)]
32. Conrad SM, Blakey GH, Shugars DA, Marciani RD, Phillips C, White RP Jr. Patients' perception of recovery after third molar surgery. *J Oral Maxillofac Surg.* 1999 Nov;57(11):1288-94; discussion 1295-6. [Medline: [10555792](#)] [doi: [10.1016/S0278-2391\(99\)90861-3](#)]
33. Phillips C, Gelesko S, Proffit WR, White RP Jr. Recovery after third-molar surgery: the effects of age and sex. *Am J Orthod Dentofacial Orthop.* 2010 Dec;138(6):700.e1-8; discussion 700-1. [Medline: [21130316](#)] [doi: [10.1016/j.ajodo.2010.06.013](#)]
34. Dimberg L, Arnrup K, Bondemark L. The impact of malocclusion on the quality of life among children and adolescents: a systematic review of quantitative studies. *Eur J Orthod.* 2015 Jun;37(3):238-47. [Medline: [25214504](#)] [doi: [10.1093/ejo/cju046](#)]

To cite this article:

Kjærgaard Larsen M, Kofod T, Starch-Jensen T.

The Use of Cryotherapy in Conjunction with Surgical Removal of Mandibular Third Molars: a Single-Blinded Randomized Controlled Trial

J Oral Maxillofac Res 2021;12(4):e2

URL: <http://www.ejomr.org/JOMR/archives/2021/4/e2/v12n4e2.pdf>

doi: [10.5037/jomr.2021.12402](#)

Copyright © Kjærgaard Larsen M, Kofod T, Starch-Jensen T. Published in the JOURNAL OF ORAL & MAXILLOFACIAL RESEARCH (<http://www.ejomr.org>), 31 December 2021.

This is an open-access article, first published in the JOURNAL OF ORAL & MAXILLOFACIAL RESEARCH, distributed under the terms of the [Creative Commons Attribution-Noncommercial-No Derivative Works 3.0 Unported License](#), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work and is properly cited. The copyright, license information and link to the original publication on (<http://www.ejomr.org>) must be included.