



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

The incidence of eosinophilic oesophagitis in 2007-2017 among children in North Denmark Region is lower than expected

Hollænder, Martin; Terkelsen, Jacob Holmen; Kramme, Frederik; Bredal, Kasper; Kragholm, Kristian; Dalby, Kasper; Hagstrøm, Søren; Melgaard, Dorte; Krarup, Anne Lund

Published in:
BMC Pediatrics

DOI (link to publication from Publisher):
[10.1186/s12887-022-03258-6](https://doi.org/10.1186/s12887-022-03258-6)

Creative Commons License
CC BY 4.0

Publication date:
2022

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):
Hollænder, M., Terkelsen, J. H., Kramme, F., Bredal, K., Kragholm, K., Dalby, K., Hagstrøm, S., Melgaard, D., & Krarup, A. L. (2022). The incidence of eosinophilic oesophagitis in 2007-2017 among children in North Denmark Region is lower than expected. *BMC Pediatrics*, 22(1), [183]. <https://doi.org/10.1186/s12887-022-03258-6>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

RESEARCH

Open Access



The incidence of eosinophilic oesophagitis in 2007–2017 among children in North Denmark Region is lower than expected

Martin Holl nder¹, Jacob Holmen Terkelsen¹, Frederik Kramme¹, Kasper Bredal¹, Kristian Kragholm², Kasper Dalby³, S ren Hagstr m⁴, Dorte Melgaard^{4,5} and Anne Lund Krarup^{6,7,8*}

Abstract

Background: In North Denmark Region (NDR), the incidence of Eosinophilic Oesophagitis (EoE) among adults has increased following a new biopsy protocol in 2011, whereas data on the incidence of EoE among children is lacking.

Aims: To describe the incidence of EoE in children aged 0–17 in NDR as well as diagnostic delay, clinical manifestations, treatment and complications.

Methods: This retrospective, register-based DanEoE cohort study included 18 children diagnosed with EoE between 2007–2017 in NDR. Medical files were reviewed with attention to symptoms, reason for referral, disease progress, treatment, symptomatic and histological remission as well as diagnostic delay.

Results: The median incidence per year (2007–2017) was 0.86/100,000 children in NDR aged 0–17 years. The median diagnostic delay among children was four years and six months. Sixty percent presented with food impaction at first hospital visit. After initial treatment, only one of 18 children achieved symptomatic and histologic remission and had a long-term treatment plan.

Conclusions: The calculated incidence among children was lower compared to similar studies. Combined with poor remission rates and lack of follow-up, it is likely that EoE is an underdiagnosed and insufficiently treated disease among children in NDR. Our findings suggest that more knowledge concerning EoE in children could lead to a higher incidence, shorter diagnostic delay and more effective treatment.

Keywords: Eosinophilic oesophagitis, Incidence, Child, Eosinophilia, Oesophagitis, Treatment

Introduction

Internationally, eosinophilic oesophagitis (EoE) is defined as a distinct clinicopathologic disease [1]. It is a chronic, immune/antigen-mediated oesophageal disease characterized by oesophageal dysfunction and eosinophil infiltration of the oesophageal epithelium (> 15 eosinophils per high-power field (EOS/HPF) [2, 3]. Adults with EoE

most often present with dysphagia and food impaction, whereas clinical manifestations among children are less specific and vary with age. Infants and toddlers may suffer from food refusal and failure to thrive, school-aged children from vomiting or pain located to the abdomen, chest, or throat and adolescents may display food avoidance and clinical manifestations may resemble that of adults [3, 4]. EoE symptoms in children thus differ from that of adults, which challenges the diagnostic process in children. Moreover, literature shows that EoE is associated with considerable diagnostic delay [5]. If untreated, EoE can lead to irreversible structural

*Correspondence: apslk@rn.dk

⁶ Department of Clinical Medicine, Aalborg University, Frederik Bajers Vej 7D, 9220 Aalborg, Denmark

Full list of author information is available at the end of the article



  The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

changes such as fibrosis, strictures and dysfunction of the oesophagus [6]. These changes may lead to reduced oesophageal distensibility, which is associated with a higher risk of food impaction [7].

In recent years, the incidence of EoE has been increasing in the western world [8]. In a geographical region in Denmark, North Denmark Region (NDR), the incidence of EoE among adults has increased 50-fold to 6–8/100,000 following implementation of a new biopsy protocol in 2011. This increase suggests a previous lack of attention to, or knowledge of, the disease [9]. For children, one study from 2009 found the incidence of EoE in Region of Southern Denmark to be 1.6/100,000 [10], which is low compared to other countries comparable to Denmark [8].

The aim of this study was to describe EoE incidence, diagnostic delay, clinical manifestations, treatment and complications in children aged 0–17 in NDR.

Methods

Study population

This retrospective study used the DanEoE cohort. DanEoE is a database including oesophageal biopsy results of all patients in NDR between 2007–2017 with SNOMED codes for both oesophagus mucosa (T62010) and > 15 EOS/HPF (M47150). All medical files, endoscopy, radiology and histology reports, biochemistry results and referral documents until 31st December 2017 were reviewed in detail by an experienced gastroenterologist and a gastroenterologist in training; subsequently, data were entered into the DanEoE database [11]. Details of DanEoE have been presented previously [9]. All management of data were performed in accordance with relevant guidelines and regulations.

In Denmark, all citizens have a unique civil registration number, which ensures complete linkage of data across national population-based medical registers as well as, including medical records, diagnostic codes, microbiologic, biochemistry, and pathologic findings [12, 13].

DanEoE children

Data on children under 18 years of age with > 15 EOS/HPF in the oesophagus were retrieved from the DanEoE database. One child suffered from a disease possibly causing EoE-like symptoms. However, since EoE was suspected and symptoms as well as > 15 EOS/HPF in the oesophagus were present, the child was included in the study cohort. The medical files of the included children were reviewed by the authors of this study with attention to symptoms, reason for referral, disease progress, treatment, symptomatic and histological remission as well as diagnostic delay. Diagnostic delay is defined as the time between first sign or symptom of EoE and diagnosis of

EoE. Second biopsy delay is the time from initial biopsy to second biopsy. Symptomatic remission delay is the time from initial treatment to symptomatic remission and histologic remission delay is the time from initial treatment to first oesophageal biopsy showing histologic remission (< 15 EOS/HPF). Findings were discussed with an experienced gastroenterologist before inclusion in the analysis. Children were divided into three age groups: 0–2, 3–12 and 13–17 years of age as symptoms present differently at different ages [3].

Statistics

For descriptive statistics median and 1st to 3rd quartiles (Q1–Q3) were used for continuous variables and counts and percentages for categorical variables. Incidence of our study population was calculated on the basis of data from Statistics Denmark. The DanEoE database contains data from 2007 to 2017. Due to structural changes in 2008, the number of inhabitants in the North Region of Denmark before 2008 cannot be obtained. Instead, the number of inhabitants in 2008 was used to calculate the incidence in 2007. This was deemed justified as fluctuations in population size in NDR are relatively low. Regarding diagnostic delay, four children were excluded due to incomplete or non-accessible data in the medical files. The SAS enterprise guide 71 (SAS institute inc., Cary, North Carolina, USA) was used for data management and statistical analysis.

Results

Incidence

A total of 18 children were diagnosed with EoE in the period 2007–2017 (1 in 2007, 2 in 2009, 2 in 2012, 5 in 2013, 4 in 2014, 1 in 2015, 3 in 2016). In the years of 2008, 2009, 2011, and 2017 no children were diagnosed with EoE in NDR. The median incidence per year (2007–2017) was 0.86/100,000 (Fig. 1). No children aged 0–2 years with EoE were found; no children below the age of four years were diagnosed with EoE in our study population.

Frequency of upper endoscopy and biopsy on any indication.

The total number of upper endoscopies (black line) in children, and the subgroup of those endoscopies where biopsies were obtained (gray line) in NDR in the period 2007–2017 is shown in Fig. 2. No indications were omitted.

Common clinical manifestations at first EoE-related hospital visit

The first EoE-related hospital visit in children aged 3–17 years were dominated by complaints of food impaction (Fig. 3). No children presented with signs of food avoidance; no children presented with failure to



Fig. 1 Incidence of Eosinophilic Oesophagitis in children aged 0–17 in North Denmark Region in the period 2007–2017




Fig. 2 Upper endoscopies in children in NDR (2007–2017) on any indication. The total number of endoscopies was obtained from the regional registry using the procedures codes for “upper endoscopies without biopsies”, “upper endoscopies with biopsies”, and age groups < 18 years

thrive or food refusal. The youngest child in our study population was three years and six months old at first EoE-related hospital visit.

At index endoscopy, two thirds of children had normal macroscopic findings (Table 1). A minority had macroscopic EoE signs. Edema and stenosis were not reported in our study population. Two children had other macroscopic findings: one had both ulcer and Mallory Weiss lesions, and one had food impaction. Oesophagitis was defined according to the LA classification [14]. In the endoscopy report of the patient with an ulcer, no LA-grade was reported.

Diagnostic delay

The median diagnostic delay among children was four years and six months (Fig. 4A). The time to second biopsy was one year and one month (Fig. 4A). The median symptomatic remission delay and histologic remission delay was five years and six months, respectively (Fig. 4B).

Treatment and follow-up

Fifteen of the 18 children were offered EoE-specific treatment. Treatment consisted of proton-pump inhibitor (PPI) or dietary treatment and was initiated within




Table 1 Study population demographics and macroscopic findings at index endoscopy

Number, (male/female)	18 (17/1)
Incidence (2007–2017)	0.86/100,000
Age at first EoE-related hospital visit, median years (Q1-Q3)	12 (11–14)
Age at first EoE-related hospital visit, n (%)	
0–2	0 (0%)
3–12	11 (61%)
13–17	7 (39%)
Macroscopic findings at index endoscopy, n (%)	
Macroscopic normal findings	12 (67%)
Macroscopic EoE findings, any type	4 (22%)
o Rings	2
o Furrows	2
o Edema	0
o White dots	1
o Stenosis	0
Scope laceration	0
Barretts oesophagus	0
Oesophagitis	0
Other macroscopic findings	2 (11%)

Abbreviations: EoE Eosinophilic Oesophagitis, Q1 1st Quartile, Q3 3rd quartile, n Number

1–2 months following diagnosis. After initial treatment, four children were both symptomatically and histologically evaluated, and two had both symptomatic and histologic remission. Only one of these two children had a long-term treatment plan, defined as a plan that maintains remission. In the children where initial treatment failed to reach documented symptomatic and histologic remission, only four received immediate second EoE treatment. The immediate second treatment failed to reach documented symptomatic and histologic remission in all of the four children. Three children received effective treatment by gastroenterologists after transition to adult care at the age of 18 years. In total, only one of 18 children received effective treatment defined as documented complete remission and a long-term treatment plan (Fig. 5).

Discussion

This retrospective register-based study of EoE in children in NDR in 2007–2017 based on the DanEoE cohort showed an incidence of 0.86/100,000 per year. The median diagnostic delay was four years and six months, and of the 15 children starting EoE treatment, one was effectively treated.




Fig. 4 Diagnostic and second biopsy delay, and symptomatic and histologic remission delay in children with Eosinophilic Oesophagitis in North Denmark Region. Diagnostic delay is the time from first symptom to diagnosis. Second biopsy delay is the time from initial biopsy to second biopsy. Symptomatic remission delay is the time from initial treatment to symptomatic remission and histologic remission delay is the time from initial treatment to first oesophageal biopsy showing histologic remission (< 15 EOS/HPF)




Fig. 5 Initial treatment and follow-up of Eosinophilic Oesophagitis in children in North Denmark Region. Effective treatment is defined as achieving both symptomatic and histologic remission and having a long-term treatment plan for maintenance. Long-term treatment is defined as treatment that maintains remission. Abbreviations: EoE, Eosinophilic Oesophagitis; n, Number

Incidence

The incidence of EoE among children in the NDR in 2007–2017 was much lower than expected. The incidence in similar studies in Europe is more than six times higher [8]. The incidence among children in the only other study in Denmark was calculated in 2008 [10] and was two times higher than the incidence calculated in

this study. As the incidence of EoE has been documented to increase in the entire western world, the low incidence found in the present study is most likely due to lack of detection. Our group found a 50-fold increase in EoE among adults in NDR in 2007–2017, following the introduction of a new biopsy protocol in 2011 [9]. According to this protocol, all patients with dysphagia should have

sampled at least eight biopsies regardless of macroscopic findings. This single intervention resulting in such a dramatic increase in incidence suggests that the awareness of EoE in NDR had been low. Moreover, the incidence of EoE reported in other European countries is very similar among adults and children [8]. The incidence found in this study was eight times lower in children compared to adults suggesting that not all children with EoE are diagnosed in NDR. Additionally, the male:female ratio of 17:1 and the lack of children aged 0–2 diagnosed with EoE in our study population differed from the literature. Spergel et al. 2009 included 620 children and found a male:female ratio of 3:1 [15], which is consistent with what has been reported in adults in NDR [16]. Furthermore, the results by Spergel et al. showed that 35% of children with EoE are diagnosed before the age of three and 68% before the age of 6 [15]. This suggests that many children with EoE aged 0–2 years are not found in NDR despite suffering from the disease [3, 15]. Regarding complications, food impaction was overrepresented in our study population compared to the literature [4]. This suggests that the children were diagnosed late, since food impaction is a late-stage manifestation of EoE [17]. We speculate that the diagnosis of EoE in these children was not considered until complications emerged, which may be explained by diffuse symptoms such as vomiting and abdominal pain being harder to attribute to the oesophagus and thus less likely to be associated with EoE.

Diagnostic delay

The median diagnostic delay in our study was four years and six months among all children in our cohort. This was in concordance with other studies in similar populations in Europe [18, 19]. Diagnostic delay may have serious consequences, as untreated EoE over time can cause irreversible changes in the oesophagus [20]. The diagnostic delay may also indicate that knowledge of EoE among general practitioners and pediatricians is lacking. An explanation might be that national guidelines on EoE in children have yet to be implemented for pediatricians or general practitioners. In 2014, the first Danish review on EoE was published in the journal read by most general practitioners [21]. The only official guidelines for EoE in Denmark is from 2015 concerning adults. In addition to this, the current biopsy protocol for children in NDR recommends two biopsies from the distal part of the oesophagus and two biopsies from the middle or proximal part of the oesophagus, which fails to meet 100% sensitivity [3].

Treatment and follow-up

Our results not only demonstrated a considerably prolonged diagnostic delay, but also suggests a problematic

course of treatment. Of the 15 children that received treatment, twelve were symptomatically evaluated, four were histologically evaluated, and four were both symptomatically and histologically evaluated. It is not surprising that most of the treated children had undergone symptomatic evaluation. It is, however, surprising that only four children were histologically evaluated, as this is necessary to assess the effectiveness of the EoE-specific treatment. Additionally, twice as many children achieved symptomatic than histologic remission, which may indicate lack of attention to the underlying cause of the symptoms. This is problematic as oesophageal inflammation does not necessarily subside when symptoms do. The lack of treatment may be explained by insufficient knowledge of the treatment guidelines for EoE. Lack of adherence to guidelines may explain the remission delay of almost 5 years. In our study, in the two children who achieved both histologic and symptomatic remission, only one of them had a long-term treatment plan. This is important as EoE will relapse without life-long treatment of the inflammation [8]. In summary, only one of the 18 children diagnosed with EoE in NDR from 2007–2017 achieved both symptomatic and histologic remission and had a long-term treatment plan.

Strengths and limitations

The generalizability of the results of this study is strong due to the complete and valid data in the DanEoE database covering the entire population of NDR. The NDR comprising one tenth of the Danish population, is well defined geographically and the composition of people reflects the Danish population. However, the small study population limits the external validity. In small data sets, outliers pose difficulties because they affect the median values more than would be the case in larger samples. On the other hand, the fact that the study population is small further adds to arguments that the EoE diagnosis in children is gravely underdiagnosed in Denmark.

Only children from 2007–2017 with biopsies confirming EoE were included in this study. The calculated incidence would have been more accurate if the study included more recent data, which unfortunately was not possible.

Because not all data with relevance to determining the diagnostic delay was available, four children were excluded in the calculation of diagnostic delay, which may have altered our results.

Conclusions

The calculated incidence among children in NDR with EoE was 0.86/100,000 children per year (2007–2017), which was lower compared with similar studies. The low incidence, poor remission rates and a lack of follow-up

suggest that EoE is underdiagnosed and insufficiently treated among children in NDR. Our findings suggest that more knowledge of and attention to EoE in children could increase incidence, reduce diagnostic delay and result in more effective treatment.

Abbreviations

EoE: Eosinophilic oesophagitis; PPI: Proton pump inhibitor; NDR: North Denmark Region; EOS/HPF: Eosinophils per high-power field.

Acknowledgements

Not applicable.

Authors' contributions

MH, JHT, FK, KB, DM and ALK gathered, analysed and interpreted relevant patient data from the DanEoE database, and drafted the manuscript. KK assisted in the statistical analysis of the data. SH and KD contributed to critically revise the drafted manuscript. All authors read and approved the final manuscript.

Funding

The study was funded by Marie Pedersen and Jensine Heiberg's Foundation, Center for Clinical Research, North Regional Hospital Hjørring, Denmark. The funding was used for language revision and publication fee.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request. Data is not shared up front, as they are part of a database in continuous research activity.

Declarations

Ethics approval and consent to participate

DanEoE was approved by the Danish Data Protection Agency, ID number 2018–59. The ethical committee evaluated the project as not being in need of ethical approval within Danish law.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹School of Medicine and Health, Aalborg University, Frederik Bajers Vej 7D, 9220 Aalborg, Denmark. ²Department of Cardiology, Unit of Clinical Biostatistics and Epidemiology, Aalborg University Hospital, Hobrovej 18-22, 9100 Aalborg, Denmark. ³Pediatric Center, Slotsgade 20C, 5000 Odense, Denmark. ⁴Department of Pediatrics, Aalborg University Hospital, Hobrovej 18-22, 9100 Aalborg, Denmark. ⁵Center for Clinical Research, North Denmark Regional Hospital, Bispensgade 37, 9800 Hjørring, Denmark. ⁶Department of Clinical Medicine, Aalborg University, Frederik Bajers Vej 7D, 9220 Aalborg, Denmark. ⁷Department of Acute Medicine and Trauma Care, Aalborg University Hospital, Hobrovej 18-20, 9000 Aalborg, Denmark. ⁸Department of Gastroenterology, Aalborg University Hospital, Hobrovej 18-22, 9100 Aalborg, Denmark.

Received: 12 September 2021 Accepted: 28 March 2022

Published online: 06 April 2022

References

- Attwood SE, Smyrk TC, Demeester TR, Jones JB. Esophageal eosinophilia with dysphagia. A distinct clinicopathologic syndrome. *Dig Dis Sci*. 1993;38(1):109–16.
- Capucilli P, Hill DA. Allergic Comorbidity in Eosinophilic Esophagitis: Mechanistic Relevance and Clinical Implications. *Clin Rev Allergy Immunol*. 2019 Aug [cited 2 Feb 2021];57(1):111–27. Available from: <https://sfx.aub.aau.dk/sfxaub?id=pmid%3A30903437&req.language=eng&sid=Entrez%3APubMed>
- Liacouras CA, Furuta GT, Hirano I, Atkins D, Attwood SE, Bonis PA, et al. Eosinophilic esophagitis: updated consensus recommendations for children and adults. *J Allergy Clin Immunol*. 2011;128(1):3–20.e6. <https://doi.org/10.1016/j.jaci.2011.02.040>.
- De Matteis A, Pagliaro G, Corleto VD, Pacchiarotti C, Di Giulio E, Villa MP, et al. Eosinophilic esophagitis in children: clinical findings and diagnostic approach. *Curr Pediatr Rev*. 2020;16(3):206–14.
- Hruz P. Epidemiology of eosinophilic esophagitis. *Dig Dis*. 2014;32(1–2):40–7. Available from: <https://www.embase.com/search/results?subaction=viewrecord&id=L603958957&from=export>
- Kumar S, Choi S, Gupta SK. Eosinophilic Esophagitis - a primer for otolaryngologists. *JAMA Otolaryngol Head Neck Surg*. 2019;145(4):373–80. Available from: <https://www.embase.com/search/results?subaction=viewrecord&id=L626517069&from=export>
- Lucendo AJ, Molina-Infante J, Arias Á, von Arnim U, Bredenoord AJ, Bussmann C, et al. Guidelines on eosinophilic esophagitis: evidence-based statements and recommendations for diagnosis and management in children and adults. *United Eur Gastroenterol J*. 2017;5(3):335–58. Available from: <https://www.embase.com/search/results?subaction=viewrecord&id=L616076628&from=export>
- Navarro P, Arias Á, Arias-González L, Laserna-Mendieta EJ, Ruiz-Ponce M, Lucendo AJ. Systematic review with meta-analysis: the growing incidence and prevalence of eosinophilic oesophagitis in children and adults in population-based studies. *Aliment Pharmacol Ther*. 2019;49(9):1116–25. Available from: <https://www.embase.com/search/results?subaction=viewrecord&id=L627203273&from=export>
- Krstrup AL, Drewes AM, Ejstrup P, Laurberg PT, Vyberg M. Implementation of a biopsy protocol to improve detection of esophageal eosinophilia: A Danish registry-based study. *Endoscopy*. 2020;53:15–24.
- Dalby K, Nielsen RG, Kruse-Andersen S, Fenger C, Bindslev-Jensen C, Ljungberg S, et al. Eosinophilic oesophagitis in infants and children in the region of southern Denmark: a prospective study of prevalence and clinical presentation. *J Pediatr Gastroenterol Nutr*. 2010;51(3):280–2.
- Erichsen R, Lash TL, Hamilton-Dutoit SJ, Bjerregaard B, Vyberg M, Pedersen L. Existing data sources for clinical epidemiology: the Danish national pathology registry and data bank. *Clin Epidemiol*. 2010;2:51–6.
- Frank L. Epidemiology. When an entire country is a cohort. Vol. 287, *Science*. New York: United States; 2000. 2398–9.
- Frank L. Epidemiology. The epidemiologist's dream: Denmark. Vol. 301, *Science*. New York: United States; 2003. 163.
- Shaheen NJ, Weinberg DS, Denberg TD, Chou R, Qaseem A, Shekelle P. Upper endoscopy for gastroesophageal reflux disease: best practice advice from the clinical guidelines committee of the American college of physicians. *Ann Intern Med*. 2012;157(11):808–16.
- Spergel JM, Brown-Whitehorn TF, Beausoleil JL, Franciosi J, Shuker M, Verma R, et al. 14 years of eosinophilic esophagitis: clinical features and prognosis. *J Pediatr Gastroenterol Nutr*. 2009 Jan [cited 4 June 2021];48(1):30–6. Available from: <https://sfx.aub.aau.dk/sfxaub?id=pmid%3A19172120&req.language=eng&sid=Entrez%3APubMed>
- Melgaard D, Westmark S, Laurberg PT, Krstrup AL. A diagnostic delay of ten years in the DanEoE Cohort calls for focus on education - a population-based cross-sectional study of incidence, diagnostic process and complications of eosinophilic oesophagitis in the North Denmark Region. *Eur J Gastroenterol*. 2021. Available from: <https://pubmed.ncbi.nlm.nih.gov/34117730/>
- Straumann A, Bussmann C, Zuber M, Vannini S, Simon H-U, Schoepfer A. Eosinophilic esophagitis: analysis of food impaction and perforation in 251 adolescent and adult patients. *Clin Gastroenterol Hepatol Off Clin Pract J Am Gastroenterol Assoc*. 2008;6(5):598–600.
- Oude Nijhuis R, Warners M, De Wijkerslooth L, Smout AJPM, Bredenoord AJ. The natural course and long-term consequences of untreated eosinophilic esophagitis in a large cohort. *Neurogastroenterol Motil*. 2018;30. Available from: <https://www.embase.com/search/results?subaction=viewrecord&id=L623598800&from=export>
- Schoepfer AM, Safroneeva E, Bussmann C, Kuchen T, Portmann S, Simon H-U, et al. Delay in diagnosis of eosinophilic esophagitis increases risk

for stricture formation in a time-dependent manner. *Gastroenterology*. 2013;145(6):1230–2.

20. Miehke S. Clinical features of eosinophilic esophagitis. *Dig Dis*. 2014;32(1–2):61–7. Available from: <https://www.embase.com/search/results?subaction=viewrecord&id=L603958920&from=export>
21. Krarup AL, Vyberg M, Ejstrup P. Eosinofil øsofagitis hos voksne. *Ugeskr læger*. 2014;176/19(september):1755–8.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

