

University of Windsor

Scholarship at UWindsor

Chemistry and Biochemistry Publications

Department of Chemistry and Biochemistry

1-3-2018

Bamboo-like Chained Cavities and Other Halogen-Bonded Complexes from Tetrahaloethynyl Cavitands with Simple Ditopic Halogen Bond Acceptors

Lotta Turunen
University of Jyväskylä

Fangfang Pan
Huazhong Normal University

Ngong Kodiah Beyeh
Aalto University

John F. Trant
University of Windsor

Robin H.A. Ras
Aalto University

See next page for additional authors

Follow this and additional works at: <https://scholar.uwindsor.ca/chemistrybiochemistrypub>



Part of the [Biochemistry, Biophysics, and Structural Biology Commons](#), and the [Chemistry Commons](#)

Recommended Citation

Turunen, Lotta; Pan, Fangfang; Beyeh, Ngong Kodiah; Trant, John F.; Ras, Robin H.A.; and Rissanen, Kari. (2018). Bamboo-like Chained Cavities and Other Halogen-Bonded Complexes from Tetrahaloethynyl Cavitands with Simple Ditopic Halogen Bond Acceptors. *Crystal Growth and Design*, 18 (1), 513-520. <https://scholar.uwindsor.ca/chemistrybiochemistrypub/200>

This Article is brought to you for free and open access by the Department of Chemistry and Biochemistry at Scholarship at UWindsor. It has been accepted for inclusion in Chemistry and Biochemistry Publications by an authorized administrator of Scholarship at UWindsor. For more information, please contact scholarship@uwindsor.ca.

Authors

Lotta Turunen, Fangfang Pan, Ngong Kodiah Beyeh, John F. Trant, Robin H.A. Ras, and Kari Rissanen

This document is the Accepted Manuscript version of a Published Work that appeared in final form in the Crystal Growth & Design, copyright © 2017 American Chemical Society after peer review and technical editing by the publisher. To access the final edited and published work see <https://doi.org/10.1021/acs.cgd.7b01517>.

Article

Bamboo-like Chained Cavities and Other Halogen-Bonded Complexes from Tetrahaloethynyl Cavitands with Simple Ditopic Halogen Bond Acceptors

Lotta Turunen, FangFang Pan, Ngong Kodiah Beyeh, John F. Trant, Robin H. A. Ras, and Kari Rissanen

Cryst. Growth Des., **Just Accepted Manuscript** • DOI: 10.1021/acs.cgd.7b01517 • Publication Date (Web): 06 Dec 2017Downloaded from <http://pubs.acs.org> on December 8, 2017**Just Accepted**

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23

Bamboo-like Chained Cavities and Other Halogen- Bonded Complexes from Tetrahaloethynyl Cavitands with Simple Ditopic Halogen Bond Acceptors

24
25
26
27
28
29

Lotta Turunen,^a Fangfang Pan,^{b} Ngong Kodiah Beyeh,^{c,d*} John F. Trant,^d Robin H. A. Ras^c and
Kari Rissanen^{a*}*

30
31
32
33
34

^aUniversity of Jyväskylä, Department of Chemistry, Nanoscience Center, P.O. Box 35, 40014
University of Jyväskylä, Finland.

35
36
37
38
39
40

^bKey Laboratory of Pesticides and Chemical Biology, Ministry of Education, College of
Chemistry, Central China Normal University, Wuhan 430079, China.

41
42
43
44
45

^cAalto University, School of Science, Department of Applied Physics, Puumiehenkuja 2, 02150
Espoo, Finland.

46
47
48
49
50

^dUniversity of Windsor, Department of Chemistry and Biochemistry, N9B 3P4 Windsor, ON,
Canada.

51
52
53
54
55
56
57
58
59
60

ABSTRACT: Halogen-bonding provides a useful complement to hydrogen-bonding and metal-
coordination as a tool for organizing supramolecular systems. Resorcinarenes, tetrameric bowl-

1
2
3 shaped cavitands, have been previously shown to function as efficient scaffolds for generating
4 dimeric capsules in both solution and solid-phase, and complicated one-, two-, and three-
5 dimensional frameworks in the solid phase. Tetrahaloethynyl resorcinarenes (bromide and
6 iodide) position the halogen atoms in a very promising “crown-like” orientation for acting as
7 organizing halogen-bond donors to help build capsules and higher-order networks. Symmetric
8 divalent halogen bond acceptors including bipyridines, 1,4 dioxane, and 1,4-
9 diazabicyclo[2.2.2]octane are very promising halogen-bond accepting partners for creating these
10 systems. This report describes the complex structures arising from combining these various
11 systems including self-included dimers, herringbone-packed architectures enclosing medium
12 (186 Å³) cavities, and a very intriguing bamboo-like 1-D rod with large (683 Å³) cavities
13 between adjacent dimeric units. These various structures, all organized through host-host, host-
14 acceptor, and host-solvent interactions highlight the emergent complexity of these types of
15 complexes. As halogen bonds are weaker than hydrogen-bonds, the resulting architectures are
16 harder to predict, and these results provide additional insight into the parameters requiring
17 consideration when designing crystalline supramolecular systems using halogen-bonds as the
18 core organizing principle.

41 Introduction

42
43
44
45 The design and construction of capsular assemblies is a growing area of focus within the
46 supramolecular chemistry community.¹ From a simple *post hoc* theorizing viewpoint, defined
47 capsular assemblies should be easily designable through a complex, but deterministic evaluation
48 of the supramolecular geometries emerging from the structure of the constituent units.¹⁻⁴
49
50 However, at the nanoscopic level, the processes leading to capsule formation are considerably
51
52
53
54
55
56
57
58
59
60

1
2
3 more complicated. In addition to obtaining the correct geometry by using the appropriate
4 connecting units, ligand design and subsequent crystal engineering requires a fine control and
5
6 balance between a multitude of interdependent structural, electronic and other parameters
7
8 including: the strength of the bonding interactions, the size of any capsular cavity, the spatial
9
10 orientation and robustness of the supramolecular synthons and interaction motifs, and the
11
12 potential complicating interference from competing solvent molecules or other possible species,
13
14 *viz.* cations and/or anions.¹⁻⁴ Predicting and controlling these supramolecular interactions within
15
16 a multi-component system is particularly challenging when specific capsular structures are
17
18 desired.¹⁻⁴ However, significant progress has been made, and the emergent behavior of these
19
20 interactions is quite well understood in hydrogen bond⁵⁻⁹ (HB) and metal-ligand¹⁰⁻¹⁴ mediated
21
22 capsular structures. The formation and control of purely halogen-bonded (XB)¹⁵⁻²¹
23
24 supramolecular capsules, on the other hand, are still poorly understood and only a few examples
25
26 are reported in the literature.²²⁻²⁸ As halogen bonds offer a very versatile and useful
27
28 complementary organizing principle to the H- and metal-ligand interactions, an improved
29
30 understanding of these systems is required to provide the necessary tools to access ever more
31
32 elaborate supramolecular architectures.
33
34
35
36
37
38
39

40
41 Resorcinarenes,²⁹⁻³⁷ their salts,³⁸⁻⁴⁴ and resorcinarene cavitands^{22-26,28,45-49} are well-studied
42
43 families of macrocyclic receptors. Resorcinarenes are particularly useful due to their ease of
44
45 preparation, simplicity of functionalization, and versatility as hosts for a wide variety of
46
47 guests.²⁹⁻⁴⁴ The known resorcinarene capsules formed through HBs, metal-ligand coordination
48
49 bonds and dynamic covalent bonds have recently been reviewed by Kobayashi and Yamanaka.⁵⁰
50
51 Concurrent with that review, we reported a novel dimeric solid-state capsule held together solely
52
53 by halogen bonds (XBs) based on a pre-organized tetratopic resorcinarene XB acceptor.²⁵ More
54
55
56
57
58
59
60

recently, we have extended these studies and have demonstrated that tetrahaloethynyl cavitands act as efficient multidentate XB donors.^{51,52} These methylene or ethylene bridged resorcinarene cavitands, with upper rims functionalized by either four bromo- or four iodoethynyl groups, are capable of forming XBs with themselves, additional guests, and/or with solvent molecules.^{51,52} We envisaged that under specific conditions and using structurally suitable XB acceptors, the tetrahaloethynyl resorcinarene cavitands could form halogen-bonded capsular assemblies. In this contribution, we report halogen-bonded solid-state complexes formed between tetrahaloethynyl cavitands (Figure 1) and heterocyclic ditopic XB acceptors, 2,2'-bipyridine (**2,2'-bipy**), 4,4'-bipyridine (**4,4'-bipy**), 1,4-diazabicyclo[2.2.2]octane (**DABCO**) and 1,4-dioxane (**dioxane**). These ligands are classical Lewis bases and have been used extensively in the construction of other XB architectures.¹⁶⁻²¹




Figure 1. The tetratopic tetrahaloethynyl cavitand XB donors **1** – **3** and the ditopic XB acceptors **2,2'-bipy**, **4,4'-bipy**, **DABCO** and **dioxane**.

1
2
3 Aakeröy and coworkers⁵³ have proposed simple guidelines to predict the halogen bonding
4 behavior of ditopic acceptors based on the complementarity of the electrostatic potential surfaces
5 of specific donor and guest binding sites. These guidelines work very well for systems
6 incorporating a few relatively strong XBs. Yet it may fail in complicated systems if
7 supramolecular organization relies on a large number of moderate XB interactions; this may be
8 particularly true when other competing non-covalent interactions are possible. In this work, 16
9 examined combinations of crystallization partners, seven co-crystals suitable for single crystal X-
10 ray crystallographic diffraction were obtained, *viz.* **1a&2,2'-bipy**, **1a&4,4'-bipy-A**, **1a&4,4'-**
11 **bipy-B**, **1b&4,4'-bipy**, **2&4,4'-bipy**, **3&DABCO** and **3&dioxane**. Contrary to our expectations,
12 many of the ditopic XB acceptors in these structures only act as monodentate XB acceptors. The
13 obtained structures demonstrate that XB interactions are sufficient to organize XB-acceptors,
14 solvent molecules, and tetratopic XB cavitand donors into complex halogen-bonded
15 supramolecular architectures in the solid state.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33

34 The co-crystals were obtained by mixing either 1:1 or 1:2 molar ratios of the
35 tetrahaloethynyl cavitand XB donor and the ditopic XB acceptor (**2,2'-bipy**, **4,4'-bipy** and
36 **DABCO**, respectively) in acetone. Slow evaporation of the solvent provided single crystals
37 suitable for single crystal X-ray crystallography. **Dioxane** was introduced as a potential guest by
38 adding only a few drops into the acetone solution containing cavitand **3**; 1,4-dioxane-
39 incorporating crystals emerged upon evaporation. Our previous crystallographic analysis^{51,52}
40 showed that cavitands **1 – 3**, with four haloethynyl XB donor groups have two distinct XB
41 acceptor sites: the eight ether oxygens at the cavitand bridges, and the four ethynyl (–C≡C–)
42 groups on the upper rim. In addition to these XB donor and acceptor sites, the cavitand has a
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

cavity appropriate for small guest encapsulation.^{45-49,51,52} All these features combine to allow for a multitude of non-covalent




Figure 2. The weaker intermolecular binding motifs **A** = $\text{CH}\cdots\pi$, **B** = $\text{X}\cdots\text{O}$, **C** = $(\text{X}\cdots\text{O})_2$, **D** = $\text{X}\cdots\pi$, **E** = $\pi\cdots\pi$ previously observed in tetrahaloethynyl cavitands-ligand crystals (the $\text{X}\cdots\text{N}$ XBs not shown).^{51,52}

interactions, especially the strong $\text{X}\cdots\text{N}$ XBs, but also the weaker $\text{X}\cdots\text{O}$ and $\text{X}\cdots\pi$ XBs and the $\text{CH}\cdots\pi$ and $\pi\cdots\pi$ interactions (Figure 2).^{51,52}

Based on our previous studies^{51,52} the major structural feature in all methylene-bridged tetrabromo/iodoethynyl cavitands (i.e., **1a**, **1b** and **2**) solid-state structures is that they form self-included dimers through $\text{C-H}\cdots\pi$ interactions and self-complementary association between the haloethynyl groups and the cavity (motif **A**, Figure 2). This inserts one of the bromo- or iodoethynyl groups of each cavitand into the cavity of its partner and *vice versa*. These self-included dimers combine for six polarized halogen atoms, uninvolved in the dimerization, available to interact with other halogen bond acceptors. Consequently, co-crystallization of **1a** with two equivalents of **2,2'-bipy** gave rise to a self-included complex **1a&2,2'-bipy** (Figures 3, S1 and S2). The asymmetric unit contains one tetrabromoethynyl cavitand and one and a half

2,2'-bipyridine molecules; this self-included dimer is formally a 2:3 complex (donor:acceptor, Figure S2A) and involves no solvent (acetone) molecules. Each dimer forms Br \cdots N XBs with four 2,2'-bipyridine molecules. One of them acts as a monodentate XB donor (Figure 3a, orange color) with one short XB ($R_{\text{BrN}} = 0.89, 3.04 \text{ \AA}$), while the second (Figure 3a, green) has two longer ($R_{\text{BrN}} = 0.92, 3.14 \text{ \AA}$) Br \cdots N XBs that incorporates the dimer into a 1-D cavitand-dimer chain along the diagonal of the a - and b -axis (Figure 3b). In the above discussion, R refers to the interaction ratio, also known as normalized interaction distance, $R_{\text{XB}} = d_{\text{XB}}/(X_{\text{vdw}} + B_{\text{vdw}})$,⁵⁴⁻⁵⁶ where X represents the XB donor (here Br or I) and B the XB acceptor (here N, O or π).





Figure 3. (a) The self-included dimer in **1a&2,2'-bipy** as a CPK presentation showing the mono-XB-bipy in orange and the di-XB-bipy in green. (b) The Br \cdots N XBs connecting the adjacent dimers shown as CPK model, diagonal of the a - and b -axis, bromine with brown and nitrogen with green color. (c) The Br \cdots O XBs, the Br \cdots O XB connects two other dimers along the b -axis, the XB interacting atoms shown as CPK, bromine with brown and oxygen with red color.

The two remaining Br atoms of the dimer participate in a Br \cdots O XB with the neighboring dimer ($R_{\text{BrO}} = 0.88, 2.98 \text{ \AA}$, Figure 3c) linking the dimers into a 1-D chain through interaction motif **B** (Figure 2), enhanced by additional inter-dimer $\pi\cdots\pi$ interactions (motif **E**, Figure 2), along the crystallographic *b*-axis, similar to those observed in the previously reported solvate structures.⁵² Together these Br \cdots N and Br \cdots O XB interactions stabilize the crystal lattice (Figure S1 and S2)

Two different structures were obtained from the co-crystallization of **1a** and **4,4'-bipy**, an isomer of **2,2'-bipy**. Depending on the XB donor to XB acceptor ratio, either **1a&4,4'-bipy-A** (1:2) or **1a&4,4'-bipy-B** (1:1) were obtained. The **1a&4,4'-bipy-A** has one cavitand and two **4,4'-bipy** molecules in the asymmetric unit. Yet, as in **1a&2,2'-bipy**, it is a self-included dimer but in a 2:4 ratio (Figure 4a). As in **1a&2,2'-bipy**, six Br atoms are available for XBs. Quite surprisingly, only one of the two **4,4'-bipy** acceptors is halogen-bonded ($R_{\text{BrN}} = 0.86, 2.93 \text{ \AA}$), and then only as a monodentate ligand (Figure 4a, highlighted in green). The non-XB coordinating nitrogen atom forms a weak N \cdots H-C non-classical hydrogen bond with the hexyl chain of an adjacent dimer (Figure S4).



1
2
3 **Figure 4.** (a) The self-included dimer in **1a&4,4'-bipy-A** as a CPK representation showing the
4 mono-XB-bipy in green and the non-XB-bipy in yellow. (b) The Br \cdots O XB connects two other
5 dimers along the *a*-axis, the XB Br (brown) and O (red) atoms are shown as CPK models for
6 clarity.
7

8
9 The second **4,4'-bipy** ligand (Figure 4a, highlighted in yellow) is not halogen-bonded,
10 but rather is trapped between the XB-bipy and the outer wall of the cavitand (Figure 4a). The
11 non-XB-bipy does have a weak hydrogen-bond with the C-H of the cavitand's methylene-bridge.
12 The other end forms a weak self-complementary C-H \cdots N HB dimer with the adjacent non-XB-
13 bipy, as the green bipy has the two aromatic rings twisted out of coplanarity (Figure S4C). These
14 interactions consequently account for two of the six available bromine atoms of the dimer. An
15 additional two of the remaining Br atoms both participate in two separate Br \cdots O XB to adjacent
16 dimers ($R_{\text{BrO}} = 0.88, 3.02 \text{ \AA}$, Figure 4b) through interaction motif **B** (Figure 2) creating a 1-D
17 chain of dimers along the crystallographic *a*-axis; this is similar to the systems we have
18 previously observed in the solvate structures.⁵² The remaining two bromine atoms of the
19 **1a&4,4'-bipy-A** dimer are surrounded by the adjacent **4,4'-bipy** molecules and dimers and do
20 not participate in any significant interactions.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35

36
37 We had expected that all of the halogens in the 1:2 (**1a: 4,4'-bipy**) would have been
38 involved in interactions, and that these would have been effectively bridged by the adaptable
39 bipy ligands. To further explore these systems, we conducted an additional experiment, using
40 only a 1:1 ratio of the cavitand and bipyridine. This provided co-crystal, **1a&4,4'-bipy-B**. The
41 asymmetric unit cell is comprised of one cavitand and half of a **4,4'-bipy**, adopting a self-
42 included 2:2 dimer architecture (Figure 5a). The **4,4'-bipy** acceptor shows strong ditopic Br \cdots N
43 halogen bonds ($R_{\text{BrN}} = 0.81, 2.76 \text{ \AA}$, Figure 5a, green color) with the cavitand.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60





Figure 5. (a) The self-included dimer in **1a&4,4'-bipy-B** in a CPK presentation highlighting the ditopic XB-bipy (green). The Br \cdots N XB connects the dimers along the *b*-axis, the XB Br (brown) and N (pale blue) atoms are shown as CPK models.

In this case, as there are insufficient XB acceptors present for the available donors, the Aakeröy concept⁵³ is sustained and only the strongest donor and acceptor interact as expected. This assembles the dimers through the **4,4'-bipy** (Figure 5b) into a 1-D chain along the crystallographic *b*-axis. The lack of sufficient XB acceptors present in **1a&4,4'-bipy-B**, forces the contacts between the cavitand molecules to be different from those observed for **1a&2,2'-bipy** and **1a&4,4'-bipy-A**. As in **1a&4,4'-bipy-A**, the strong self-inclusion behavior and the $\pi\cdots\pi$ interactions (motif **E**, Figure 2) between the dimers are preserved. Yet the Br \cdots O XBs observed in **1a&2,2'-bipy** and **1a&4,4'-bipy-A** are now replaced by weak CH \cdots Br bonds. Even the $\pi\cdots\pi$ stacking between the cavitand resorcinol moieties relative to the orientation of the self-included dimer is different (Figure S5A). It is noteworthy that, the **4,4'-bipy** molecule in **1a&4,4'-bipy-B** is coplanar (deviation from planarity is only 0.01 Å). This manifests that, in addition to the expected Br \cdots N halogen bonds, the **4,4'-bipy** acts as a “ π -glue” between the dimers through $\pi\cdots\pi$ interactions (Figure S5C).

The behavior of the **4,4'-bipy** in the two systems **1a&4,4'-bipy-A** and **1a&4,4'-bipy-B**, both as an XB acceptor and as an auxiliary molecule, differs considerably. To further investigate

1
2
3 this discrepancy, we probed the interaction of **4,4'-bipy** with cavitand **1b**. It has the same
4 methylene-bridged cavitand skeleton but replaces the lower rim hexyl chains of **1a** with longer
5 undecyl (C_{11}) groups. In the **1b&4,4'-bipy** crystal, the asymmetric unit contains one cavitand
6 and two **4,4'-bipy** molecules, and is defined as a self-included 2:4 dimer (Figure 6a), reminiscent
7 to the structure of **1a&4,4'-bipy-A**. However, unlike for that system, all **4,4'-bipy** molecules act
8 as two pairs of monodentate XB acceptors ($R_{BrN} = 0.83, 2.84 \text{ \AA}$ and $0.86, 2.92 \text{ \AA}$; Figure 6a).
9
10 One pair mediates inter-dimer organization using $\pi \cdots \pi$ interactions, while the other does not.
11
12 The increased length of the alkyl substituents enhances their capability to have London
13 dispersion force interactions, and these clearly play an important role in crystal packing. This
14 behavior was not observed for the shorter hexyl chains in the **1a** structures. The remaining Br
15 atoms of the dimer participate in a very weak $Br \cdots O$ XBs with the neighboring dimer ($R_{BrO} =$
16 $0.98, 3.29 \text{ \AA}$); these interactions are notable due to the unusually acute $C-Br \cdots O$ angles of 139°
17 (normally $> 170^\circ$), and link the dimers through motif **C** (Figure 2) into a 1-D chain along the
18 crystallographic a -axis, as has been previously observed both above and for the solvates.⁵²
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



1
2
3 **Figure 6.** (a) The self-included dimer in **1b**&**4,4'-bipy** presented as CPK models showing the
4 four mono-XB-bipy (green). (b) The Br \cdots O XB connects adjacent dimers along the *b*-axis. The
5 XB Br (brown) and O (red) atoms shown as CPK models for clarity.
6
7
8
9

10
11 Due to the more strongly polarized iodine atom, the tetraiodoethynyl cavitand **2** is able to
12 form stronger halogen bonds with XB acceptors than the analogous tetrabromo cavitand **1a**.
13 Cavitand **2** is expected to show the same self-included dimer behavior, but its stronger XB
14 potential donor should lead to other, maybe even capsular, structures with linearly ditopic XB
15 acceptors. To investigate this hypothesis, we challenged cavitand **2** with **4,4'-bipy** and the
16 resulting co-crystal **2**&**4,4'-bipy** is consistent with this logic, but represents a completely
17 unexpected structure. For the first time, the crystal lattice contains the solvent molecule, yet not
18 as an innocent bystander, but as an active component of the assembly. The asymmetric unit
19 contains one cavitand, one **4,4'-bipy** and three acetone molecules, thus the self-included dimer
20 has a formal composition of 2:2:7 (cavitand:**4,4'-bipy**:acetone, Figures 7a and S6). As in **1a** and
21 **1b** the self-inclusion dimer involves two of the eight available iodine atoms. The strong halogen
22 bonds ($R_{IN} = 0.77, 2.71$ and 2.72 Å) between the two **4,4'-bipy** molecules and four iodines of the
23 dimer at the 1- and 3-positions organize the dimers into a 1-D rod (Figure 7c). The two
24 remaining iodine atoms are dormant and are blocked by an acetone molecule trapped into the
25 space between the hexyl chains of the adjacent cavitand. The strong halogen bonds force the
26 cavitand dimers to orient themselves so that a cavity, reminiscent of the alternating cavity
27 structure found in bamboo, is formed. This cavity is large at
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 7. (a) The halogen-bonded “dimer of dimers” in 2&4,4’-bipy as CPK presentation showing the two di-XB-bipy in green and the acetone molecules in yellow. (b) The cavity in between the dimers, represented with a transparent yellow solid, the two located acetone molecules are provided as CPK models. (c) Schematic presentation of the formation of the bamboo-like chained cavities. The curved arrow represents the cavitand, the “fork” the two active iodoethynyl groups, the green rectangle 4,4’-bipy and the red oval the created cavity.

683 Å³, calculated using MSRoll⁵⁷ inside the X-seed^{58,59} with the probe of 1.2 Å (Figure 7b). The cavity is big enough to accommodate five acetone solvent molecules, two out of which are not disordered and can be localized (Figure S6). In addition to the strong XBs between the dimers, the crystal lattice is reinforced by $\pi \cdots \pi$ stacking of the 1-D rods through the interaction motif **E** (Figure 2).

All the methylene-bridged cavitands **1a**, **1b** and **2** form self-included dimers due to the self-complementary fit of the haloethynyl groups with the cavity; this is accentuated by the fact that the methylene hydrogens are both acidic and pointed directly towards the center of the cavity, thus enabling CH \cdots π interaction between the cavitands of the dimer (interaction motif **A**,

Figure 2).^{51,52} If the cavitand is constructed using the ethylene-bridge, -CH₂-CH₂-, instead of the methylene, the situation changes dramatically. Now the interactions driving the self-inclusion are greatly diminished and consequently self-included dimers are not formed. This leaves all the XB donor sites free to interact with the XB acceptors. Unfortunately, all attempts to obtain high quality single crystals from the tetraiodoethynyl cavitand **3** and **4,4'-bipy** were unsuccessful. However, **3** could be crystallized with **DABCO** by slow evaporation of the corresponding acetone solution to provide co-crystal **3&DABCO**. Like **4,4'-bipy**, **DABCO** is also a strong ditopic XB acceptor, yet it is a little bit shorter and has larger steric requirements due to its nearly spherical shape. As the π -systems of **4,4'-bipy** were also very important in the structures described above, these crystals are expected to adopt rather different architectures. The asymmetric unit of **3&DABCO** contains one cavitand, two **DABCO** units, and two acetone molecules, such that two of the **DABCO** molecules are ditopically halogen-bonded ($R_{IN} = 0.78$, $4 \times 2.74 \text{ \AA}$) to two adjacent cavitands, while the remaining **DABCO** interacts through a monodentate XB ($R_{IN} = 0.76$, 2.67 \AA). An acetone molecule interacts with the remaining iodine *via* a I \cdots O XB ($R_{IO} = 0.82$, 2.87 \AA , Figure 8a). An additional non-XB acetone molecule is trapped in the space between the hexyl chains as in **2&4,4'-bipy** above. As the self-inclusion does not happen in **3&DABCO**, the cavity is now open for other guests and as acetone is too polar to reside comfortably in the cavitand cavity, one of the hexyl chains of the adjacent cavitand acts as a guest and is inserted into the cavity (Figure 8b). The ditopic halogen bonds between the **DABCO** and **3** result in $(\cdots\mathbf{3}\cdots\mathbf{DABCO}\cdots\mathbf{3}\cdots\mathbf{DABCO}\cdots)_n$




Figure 8. (a) The halogen bonding in 3&DABCO (CPK model) showing the di-XB-DABCO in green, the mono-XB-DABCO in light green and the XB-acetone in yellow. (b) The “tail-including” cavitand is shown as a CPK model in orange; the “binding” cavitand is shown as a ball-and-stick representation, and the acetone molecules trapped inside the hexyl rings are depicted as yellow CPK representations. (c) The XB ribbon extending along the *c*-axis, the XB donor I (purple) and N (pale blue) are highlighted as CPK models. The hexyl chain including in the cavitand is shown in orange with the mono-XB-DABCO and the XB-acetone removed for clarity.

“wavy” ribbons which are then joined together by the hexyl chain inclusion between the 1-D chains (Figure 8c). The C-I⋯N XBs in 3&DABCO control the interactions between the components so efficiently that none of the interaction motifs that we have become accustomed to seeing **A – E** (Figure 2) are present.

Using other quite similar solvent molecules as competitive guests, when both of them can act as XB acceptors, might lead to a situation where one solvent molecule clearly displaces the other. In spite of the similarities between the polar aprotic solvents acetone ($V = 72.7 \text{ \AA}^3$) and 1,4-dioxane ($V = 94.3 \text{ \AA}^3$), both this study (*viz.* in **2&4,4'-bipy** and **3&DABCO** above) and our previous data has repeatedly demonstrated that acetone does not bind in the cavity of these halo-substituted resorcinarenes; on the other hand, earlier work on other resorcinarene systems³⁸⁻⁴⁴ has shown that 1,4-dioxane is an excellent guest for resorcinarenes and cavitands. Keeping this in mind, the slow evaporation of tetraiodoethynyl cavitand **3** in acetone in the presence of a few drops of 1,4-dioxane (**dioxane**) resulted in a co-crystal, **3&dioxane**, which crystallizes in the high symmetry orthorhombic space group *Cmcm*. Due to the high symmetry, the asymmetric unit contains 1/4 of **3** and 1.5 **dioxane** molecules. In spite of the huge excess of acetone during the crystallization, only **dioxane** is found in the crystal lattice. Taking into account the crystal symmetry, each cavitand **3** has one **dioxane** in the cavity, two acting as ditopic XB acceptors, and the remaining eight (some badly disordered) fill the voids between the cavitands. The two di-XB-**dioxane** molecules form very strong $\text{I}\cdots\text{O}$ halogen bonds ($R_{\text{IO}} = 0.77$, $4 \times 2.70 \text{ \AA}$, Figure 9a), very similarly to those observed for **3&DABCO** (Figure 8c), and connect the cavitands into similar 1D wave-like ribbons. Due to these strong $\text{I}\cdots\text{O}$ XBs, the cavitands stack “head-to-tail” on top of each other forcing the hexyl chains to point perpendicular to the stacking axis of the cavitand columns (Figures 9b and 9c) forming a herringbone-type packing (Figure 9e, only one ribbon shown). In addition to the strong cavitand-**dioxane** $\text{I}\cdots\text{O}$ halogen bonds, the **3&dioxane** manifests very weak intermolecular $\text{I}\cdots\pi$ halogen bonds ($R_{\text{I}\pi} = 0.92$, 3.40 \AA) and is the only structure in this study to do so (motif **D**, Figure 2). These are perpendicular to the $\text{I}\cdots\text{O}$ halogen

bonds joining the head-to-tail stacks together (Figure 9f). The tight head-to-tail stacking creates a cavity between the cavitands.




Figure 9. (a) The halogen bonding in **3** and **dioxane** presented as CPK models showing the di-XB-dioxanes in green. (b) The head-to-tail packing of the cavitands as CPK models with the encapsulated **dioxane** (CPK) highlighted in yellow. (c) Same view as in (b) but showing the cavitands as ball-and-stick to better illustrate the dioxanes. (d) A more detailed presentation of the inter-cavitand cavity, yellow, and the included **dioxane** as a translucent CPK model. (e) The XB ribbon extending along the *b*-axis: the XB-participating atoms are represented using CPK models, iodine (purple) and oxygen (red). (f) The independent auxiliary $I \cdots \pi$ halogen bonds, the XB atoms again are presented as CPK models, iodine (purple) and carbon (grey).

The cavity volume is 186 \AA^3 (Figure 9d) calculated using MSRoll⁵⁷ inside the X-seed^{58,59} with the probe of 1.2 \AA . As the volume of **dioxane** is 94.3 \AA^3 , this results in a packing coefficient PC

1
2
3 = 0.51, only slightly smaller than the optimum 0.55 according to Rebek and coworkers.⁶⁰ The
4 small PC allows the **dioxane** molecule to tumble inside the cavity, which is observed as a severe
5 disorder.
6
7
8
9

10 **Conclusions**

11
12
13
14 Despite the apparent geometrical and structural compatibility of these cavitand tetratopic halogen
15 bond donors (**1a**, **1b**, **2** and **3**) and ditopic halogen bond acceptors, 2,2'-bipyridine, 4,4'-
16 bipyridine, **DABCO** and 1,4-dioxane, we were unable to obtain any simple capsules. Instead,
17 these systems clearly prefer to self-assemble into self-including dimeric units (for the methylene-
18 bridged **1** and **2**) and ribbons or head-to-tail columnar stacks (for ethylene-bridged **3**). The
19 precise structure in each case is highly dependent upon the particular nature of the ditopic
20 acceptor and the identity of the solvent or additive. Despite the size complementarity, none of
21 these ditopic ligands adopt a traditional guest-like posture inside the cavity of the putative hosts.
22 However, this difficult to predict behavior results in highly exotic and intriguing architectures
23 including self-included 1-D chains, bamboo-like alternating cavity systems, and self-organized
24 parallel columns. These structures may prove useful for the design of innovative materials in the
25 future, and we are continuing our investigations into the behavior of the halo-resorcinarene
26 systems.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44

45 ASSOCIATED CONTENT

46
47
48 **Supporting Information.** Electronic Supplementary Information (ESI) available: [X-ray
49 crystallographic details. CCDC 1574168-1574174]. This material is available free of charge via
50 the Internet at <http://pubs.acs.org>.
51
52
53

54 AUTHOR INFORMATION

Corresponding Authors

*E-mail: ffpan@mail.ccnu.edu.cn; k.beyeh@uwindsor.ca; kari.t.rissanen@jyu.fi

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

The authors gratefully acknowledge financial support from the Academy of Finland (KR: grant nos. 265328, 263256 and 292746; RHAR: grant no. 272579), the University of Jyväskylä, and the University of Windsor, Canada. This work was supported by the Program of Introducing Talents of Discipline to Universities of China (111 program, B17019) and the Academy of Finland through its Centres of Excellence Programme (HYBER 2014–2019).

REFERENCES

1. Desiraju, G. R. *Crystal Engineering: the Design of Organic Solids*; Elsevier: Amsterdam, 1989.
2. Tiekink, E. R. T.; Vittal, J. J.; Zaworotko, M., Eds.; *Organic Crystal Engineering: Frontiers in Crystal Engineering*, Wiley VCH, 2010.
3. Tiekink, E. R. T.; Zukerman-Schpector, J., Eds.; *The Importance of π -Interactions in Crystal Engineering: Frontiers in Crystal Engineering*, 2^{ed}. Wiley, 2012.
4. Bond, A. D. *CrystEngComm* **2007**, *9*, 833–834.
5. Adachi, T.; Ward, M. D. *Acc. Chem. Res.* **2016**, *49*, 2669–2679.
6. Atwood, J. L.; MacGillivray, L.R. *Nature* **1997**, *389*, 469–472.
7. Gerkensmeier, T.; Iwanek, W.; Agena, C.; Fröhlich, R.; Kotila, S.; Näther, C.; Mattay, J. *Eur. J. Org. Chem.* **1999**, *9*, 2257–2262.
8. Shivanyuk, A.; Rebek, J. *Proc. Natl. Acad. Sci.* **2001**, *98*, 7662–7665.
9. Beyeh, N. K.; Rissanen, K. *Isr. J. Chem.* **2011**, *51*, 769–780.
10. Fujita, D.; Ueda, Y.; Sato, S.; Mizuno, N.; Kumasaka, T.; Fujita, M. *Nature* **2016**, *540*, 563–566.
11. Fujita, D.; Ueda, Y.; Sato, S.; Yokoyama, H.; Mizuno, N.; Kumasaka, T.; Fujita, M. *Chem* **2016**, *1*, 91–101.
12. McKinlay, R. M.; Cave, G. W. V.; Atwood, J. L. *Proc. Natl. Acad. Sci.* **2005**, *102*, 5944–5948.
13. McKinlay, R. M.; Thallapally, P. K.; Atwood, J. L. *Chem. Commun.* **2006**, 2956–2958.

14. Rathnayake, A. S.; Feaster, K. A.; White, J.; Barnes, C. L.; Teat, S. J.; Atwood, J. L. *Cryst. Growth Des.* **2016**, *16*, 3562–3564.
15. Desiraju, R.G.; Ho, S.P.; Kloo, L.; Legon, C.A.; Marquardt, R.; Metrangolo, P.; Politzer, P.; Resnati, G.; Rissanen, K. *Pure Appl. Chem.* **2013**, *85*, 1711–1713.
16. Cavallo, G.; Metrangolo, P.; Milani, R.; Pilati, T.; Priimagi, A.; Resnati, G.; Terraneo, G. *Chem. Rev.* **2016**, *116*, 2478–2601.
17. Kolář, M. H.; Hobza, P. *Chem. Rev.* **2016**, *116*, 5155–5187.
18. Wang, H.; Wang, W.; Jin, W. J. *Chem. Rev.* **2016**, *116*, 5072–5104.
19. Gilday, L.C.; Robinson, S. W.; Barendt, T. A.; Langton, M. J.; Mullaney, B. R.; Beer, P. D. *Chem. Rev.* **2015**, *115*, 7118–7195.
20. Troff, R.W.; Mäkelä, T.; Topić, F.; Valkonen, A.; Raatikainen, K.; Rissanen, K. *Eur. J. Org. Chem.* **2013**, *2013*, 1617–1637.
21. Rissanen, K. *CrystEngComm* **2008**, *10*, 1107–1113.
22. Aakeröy, C. B.; Rajbanshi, A.; Metrangolo, P.; Resnati, G.; Parisi, M. F.; Desper, J.; Pilati, T. *CrystEngComm* **2012**, *14*, 6366–6368.
23. Dumele, O.; Trapp, N.; Diederich, F. *Angew. Chem., Int. Ed.* **2015**, *54*, 12339–12344.
24. Dumele, O.; Schreiber, B.; Warzok, U.; Trapp, N.; Schalley, C. A.; Diederich, F. *Angew. Chem., Int. Ed.* **2017**, *56*, 1152–1157.
25. Beyeh, N. K.; Pan, F.; Rissanen, K. *Angew. Chem., Int. Ed.* **2015**, *54*, 7303–7307.
26. Turunen, L.; Warzok, U.; Puttreddy, R.; Beyeh, N. K.; Schalley, C. A.; Rissanen, K. *Angew. Chem., Int. Ed.* **2016**, *55*, 14033–14036.
27. Turunen, L.; Peuronen, A.; Forsblom, S.; Kalenius, E.; Lahtinen, M.; Rissanen, K. *Chem.-Eur. J.* **2017**, *23*, 11714–11718.
28. Turunen, L.; Warzok, U.; Schalley, C. A.; Rissanen, K. *Chem* **2017**, *3*, 861–869.
29. Timmeman, P.; Verboom, W.; Reinhoudt, D. N. *Tetrahedron* **1996**, *52*, 2663–2704.
30. Jasat, A.; Sherman, J. C. *Chem. Rev.* **1999**, *99*, 931–968.
31. Beyeh, N. K.; Pan, F.; Valkonen, A.; Rissanen, K. *CrystEngComm* **2015**, *17*, 1183–1188.
32. Puttreddy, R.; Beyeh, N. K.; Rissanen, K. *CrystEngComm* **2016**, *18*, 793–799.
33. Puttreddy, R.; Beyeh, N. K.; Kalenius, E.; Ras, R. H. A.; Rissanen, K. *Chem. Commun.* **2016**, *52*, 8115–8118.
34. Puttreddy, R.; Beyeh, N. K.; Rissanen, K. *CrystEngComm* **2016**, *18*, 4971–4976.
35. Puttreddy, R.; Beyeh, N. K.; Ras, R. H. A.; Rissanen, K. *ChemistryOpen* **2017**, *6*, 417–423.
36. Puttreddy, R.; Beyeh, N. K.; Ras, R. H. A.; Trant, J. F.; Rissanen, K. *CrystEngComm* **2017**, *19*, 4312–4320.
37. Kiesilä, A.; Kivijärvi, L.; Beyeh, N. K.; Moilanen, J.; Groessl, M.; Rothe, T.; Götz, S.; Topić, F.; Rissanen, K.; Lützen, A.; Kalenius, E. *Angew. Chem., Int. Ed.* **2017**, *56*, 10942–10946.


- 1
2
3 38. Beyeh, N. K.; Rissanen, K. In *Calixarenes and Beyond*; Neri, P., Sessler, J., Wang, M.
4 X. Eds.; Springer, Cham, 2016; pp 255–283.
5
6 39. Pan, F.; Beyeh, N. K.; Rissanen, K. *J. Am. Chem. Soc.* **2015**, *137*, 10406–10413.
7
8 40. Beyeh, N. K.; Pan, F.; Bhowmik, S.; Mäkelä, T.; Ras, R. H. A.; Rissanen, K. *Chem. -Eur.*
9 *J.* **2016**, *22*, 1355–1361.
10
11 41. Pan, F.; Beyeh, N. K.; Bertella, S.; Rissanen, K. *Chem. -Asian J.* **2016**, *11*, 782–788.
12
13 42. Pan, F.; Beyeh, N. K.; Ras, R. H. A.; Rissanen, K. *CrystEngComm* **2016**, *18*, 5724–5727.
14
15 43. Pan, F.; Beyeh, N. K.; Ras, R. H. A.; Rissanen, K. *Cryst. Growth Des.* **2016**, *16*, 672–
16 6733.
17
18 44. Beyeh, N. K.; Jo, H. H.; Kolesnichenko, I.; Pan, F.; Kalenius, E.; Anslyn, E. V.; Ras, R.
19 H. A.; Rissanen, K. *J. Org. Chem.* **2017**, *82*, 5198–5203.
20
21 45. Wishard, A., Gibb, B. C. In *Calixarenes and Beyond*; Neri, P., Sessler, J., Wang, M. X.
22 Eds.; Springer: Cham, 2016; pp 193–234.
23
24 46. Knighton, R. C.; Chaplin, A. B. *Tetrahedron* **2017**, *73*, 4591–4596.
25
26 47. Chavagnan, T.; Bauder, C.; Semeril, D.; Matt, D.; Toupet, L. *Eur. J. Org. Chem.* **2017**,
27 70–76.
28
29 48. Gropp, G.; Trapp, G.; Diederich, F. *Angew. Chem., Int. Ed.* **2016**, *55*, 14444–14449.
30
31 49. Elaieb, F.; Semeril, D.; Matt, D. *Eur. J. Org. Chem.* **2016**, 3103–3108.
32
33 50. Kobayashi, K.; Yamanaka, M. *Chem. Soc. Rev.* **2015**, *44*, 449–466.
34
35 51. Turunen, L.; Beyeh, N. K.; Pan, F.; Valkonen A.; Rissanen, K. *Chem. Commun.* **2014**,
36 50, 15920–15923.
37
38 52. Turunen, L.; Pan, F.; Beyeh, N. K.; Cetina, M.; Trant, J. F.; Ras, R. H. A.; Rissanen, K.
39 *CrystEngComm* **2017**, *19*, 5223–5229.
40
41 53. Aakeröy, C. B.; Wijethunga, T. K.; Desper, J.; Đaković, M. *Cryst. Growth Des.* **2016**, *16*,
42 2662–2670.
43
44 54. Lommerse, P. M.; Stone, A. J.; Taylor, R.; Allen, F. H. *J. Am. Chem. Soc.*, **1996**, *118*,
45 3108–3116.
46
47 55. Brammer, L.; Bruton, E. A.; Sherwood, P. *Cryst. Growth Des.* **2001**, *1*, 277–290.
48
49 56. Zordan, F.; Brammer, L.; Sherwood, P. *J. Am. Chem. Soc.* **2005**, *127*, 5979–5989.
50
51 57. Connolly, M. L. *J. Mol. Graphics* **1993**, *11*, 139–141.
52
53 58. Barbour, L. J. *J. Supramol. Chem.* **2001**, *1*, 189–191.
54
55 59. Atwood, J. L.; L. J. Barbour, *Cryst. Growth Des.* **2003**, *3*, 3–8.
56
57
58
59
60

1
2
3 60. Mecozzi, S.; Rebek, J. *Chem. -Eur. J.* **1998**, *4*, 1016–1022
4
5
6
7
8
9
10
11
12
13
14
15
16
17

18 For Table of Contents Use Only
19
20
21
22

23 Bamboo-like Chained Cavities and Other Halogen-Bonded Complexes from Tetrahaloethynyl
24 Cavitands with Simple Ditopic Halogen Bond Acceptors
25
26
27

28 Lotta Turunen, Fangfang Pan, Ngong Kodiah Beyeh, John F. Trant, Robin H. A. Ras and Kari
29 Rissanen
30
31
32
33



49 Resorcinarenes are bowl-shaped cavitands with excellent potential for host-guest interactions. In
50 this study, halogen bond-organized systems are reported between host haloethynyl groups and
51 ditopic halogen bond acceptors. Among numerous other reported crystal structures, including
52
53
54
55
56
57
58
59
60

1
2
3 self-included dimers and 2- and 3-dimensional networks, a one-dimensional bamboo-like rod
4
5 with large cavities was observed.
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60