

University of Groningen

## Biomarker Discovery In Chronic Obstructive Pulmonary Disease (COPD) Using Epithelial Lining Fluid

Franciosi, L.; Govorukhina, N.; Fusetti, F.; Poolman, B.; Hacken, N. ten; Postma, D.; Bischoff, R.

*Published in:*  
American Journal of Respiratory and Critical Care Medicine

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2011

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Franciosi, L., Govorukhina, N., Fusetti, F., Poolman, B., Hacken, N. T., Postma, D., & Bischoff, R. (2011). Biomarker Discovery In Chronic Obstructive Pulmonary Disease (COPD) Using Epithelial Lining Fluid: A Proteomic Approach. *American Journal of Respiratory and Critical Care Medicine*, 183(1), A1492-A1493.

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

## Biomarker Discovery In Chronic Obstructive Pulmonary Disease (COPD) Using Epithelial Lining Fluid: A Proteomic Approach

**L. Franciosi<sup>1</sup>, N. govorukhina<sup>1</sup>, F. fusetti<sup>1</sup>, B. poolman<sup>1</sup>, N. hacken ten<sup>2</sup>, D. Postma<sup>3</sup>, R. bischoff<sup>1</sup>**

<sup>1</sup>University of Groningen, groningen, Netherlands, <sup>2</sup>University Medical Center Groningen, groningen, Netherlands, <sup>3</sup>University Medical Center Groningen, Groningen, Netherlands

**Corresponding author's email: L.Franciosi@rug.nl**

### RATIONALE

Chronic Obstructive Pulmonary Disease (COPD) is the third most frequent disease worldwide with increasing mortality. Cigarette smoking is the principle risk factor and 15-20% of smokers develop COPD.

Epithelial Lining Fluid (ELF) covers the internal part of the airways and can be collected during bronchoscopy. ELF appears to be well-suited for proteomic analysis, since it contains a higher concentration of proteins (150-300 µg /mL) than other lung fluids and can be obtained from different locations of the lungs. No comprehensive proteomic analysis of human ELF has been performed to date, which makes ELF a highly interesting fluid for biomarker discovery in COPD.

### AIM

To discover proteins that change in abundance in ELF from COPD patients versus healthy controls using a quantitative proteomics approach.

### METHODS

The ELF proteome from COPD patients and healthy controls was studied by 1D polyacrylamide gel electrophoresis in the presence of SDS followed by in-gel tryptic digestion to establish the methodology and assess the feasibility of such an approach. Approximately 40 gel slices were obtained from each lane of the gel (corresponding to one patient). Digested samples were analyzed by nanoChip-LC-MS/MS using an ion trap.

We performed a quantitative pilot study of ELF from 4 COPD patients and 4 healthy controls (table 1) to test for statistically significant differences in protein levels. ELF samples were digested by trypsin, labeled with stable isotope-containing reagents (iTRAQ®, 8-plex) and processed by strong cation-exchange chromatography followed by nanoLC-MS/MS. In order to validate the results, a second quantitative analysis of an independent sample set (4 COPD vs 4 healthy) using the same methodological approach was done.

### RESULTS

The 1D electrophoretic approach resulted in more than 300 identified proteins. Most of the identified proteins were present in both COPD and healthy samples, although some proteins were only identified either in healthy control or in COPD samples.

The quantitative studies showed that a number of proteins was significantly different between ELF of COPD patients and controls, including 4 up-regulated proteins in common in both studies.

### CONCLUSIONS

This is the first study in ELF of COPD patients and healthy controls in which such a large number of proteins has been identified. The obtained results show the feasibility of this proteomic approach and the possibility to discover proteins that are differentially expressed in ELF of COPD patients and controls. We are currently validating these proteins further by western blot and immunohistochemistry.

Patient characteristics, lung function, smoking history

	Pilot study	Pilot study	Validation study	Validation study
	COPD (n=4)	Healthy (n=4)	COPD (n=4)	Healthy (n=4)
Female, number	0	2	2	2
Age, years	63.75	62.75	60	55.5
Co-morbidity, CCI	2.25	0.25	1.5	0.25
Current smoker, number	1	0	1	2
Ex smoker, number	3	0	3	1
Pack years, number	37.5	0	37.5	9.52
Inhaled corticosteroids, number	2	0	1	0
FEV <sub>1</sub> /FVC, %	58.5	76	33.25	84.25

FEV <sub>1</sub> , %predicted	71.5	116.3	40.25	111.25
-------------------------------	------	-------	-------	--------

This abstract is funded by: None

**Am J Respir Crit Care Med 183;2011:A1492**

**Internet address: [www.atsjournals.org](http://www.atsjournals.org)**

**Online Abstracts Issue**