

University of Groningen

## Real-Time Quantum Dynamics of a Single Spin 1/2 Coupled to a Spin Bath

Guerra, Marta L.; Raedt, Hans De; Novotny, M.A.

*Published in:*

Proceedings of the 24th Workshop on Computer Simulation Studies in Condensed Matter Physics (CSP2011)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

2011

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Guerra, M. L., Raedt, H. D., & Novotny, M. A. (2011). Real-Time Quantum Dynamics of a Single Spin 1/2 Coupled to a Spin Bath. In *Proceedings of the 24th Workshop on Computer Simulation Studies in Condensed Matter Physics (CSP2011)* (Vol. 15, pp. 33-36). University of Groningen, The Zernike Institute for Advanced Materials.

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

## Real-Time Quantum Dynamics of a Single Spin 1/2 Coupled to a Spin Bath

Marta L. Guerra<sup>a</sup>, Hans De Raedt<sup>b</sup>, M.A. Novotny<sup>a</sup>

<sup>a</sup>*Department of Physics and Astronomy and HPC<sup>2</sup> Center for Computational Sciences; Mississippi State University, P.O. Box 5167, Mississippi State, MS, 39762-5167, USA*

<sup>b</sup>*Department of Applied Physics, Zernike Institute for Advanced Materials, University of Groningen, Nijenborgh 4, NL-9747 AG Groningen, The Netherlands*

---

### Abstract

We performed real-time dynamics of a spin  $\frac{1}{2}$  quantum system that is coupled to a bath of  $N$  quantum spin  $\frac{1}{2}$  particles. The quantities we study are the time dependence of the magnetization and of the quantum purity,  $\mathcal{P}(t)$ . If the spins that constitute the bath have no interactions with each other, the result is compared to a recent theoretical result. We also present simulations to study the effect of interactions between the bath spins. In particular, we analyze how  $\mathcal{P}(t)$  approaches its asymptotic value for large  $t$  and  $N$ .

**Keywords:** Quantum Dynamics, Spin Bath, Quantum Purity, Decoherence

---

### 1. Introduction

Advances in quantum information processing and quantum computing have been steady for over a decade [1, 2]. However, additional theoretical understanding and improved physical implementations are necessary before a quantum computer becomes a practical device. At the simplest level a quantum computer can be viewed as a time development problem for a quantum spin system coupled to a bath that is its environment. At the computational level, the time development of the quantum system is in principle simple, since the underlying equation is just the time-dependent Schrödinger equation. However in practice there are two complications that make calculations of such quantum systems difficult. First, the precise nature of the environment must be taken into account quantum mechanically, and hence the system to be simulated is much larger than just the quantum system of the qubits of the quantum computer. Second, the development of real-time algorithms for the time-dependent Schrödinger equation are very difficult due to the imaginary phase in the propagator of the quantum system. In quantum statistical mechanics simulations this difficulty is referred to as the ‘minus sign’ problem. There has however been significant progress in simulation of quantum spin systems in real time and the physical understanding of these systems [3, 4, 5, 6].

Suppression of decoherence of spin systems is of fundamental importance for building quantum computers [2, 4]. A recent paper [7] calculated the quantum frustration of dissipation of a single spin  $\frac{1}{2}$  coupled to one or two spin baths, with no interactions between the spins of the bath. They showed that the quantum purity,  $\mathcal{P}(t)$ , decayed as an exponential for coupling to one bath but as a power law for coupling to two baths. In this paper we describe our simulation to check their expressions for a spin  $\frac{1}{2}$  coupled to one spin bath. Simulations are restricted by the largest number of spins that can be simulated on modern supercomputers, so we study the approach to the asymptotic result for a finite number of spins. Furthermore in our calculations we are not restricted to zero interactions between the bath spins, and therefore we address the question of whether a spin bath with random interactions will allow a faster approach to the asymptotic region for  $\mathcal{P}(t)$ .

## 2. Problem and Theory

We consider a quantum spin system coupled to a quantum spin bath. The Hamiltonian for the system+bath is

$$\mathcal{H} = \mathcal{H}_S + \mathcal{H}_{SB} + \mathcal{H}_B \quad (1)$$

where  $\mathcal{H}_S = \omega_0 S^z$  is the Hamiltonian of the system composed of a single spin  $\frac{1}{2}$  interacting with an external magnetic field,  $\mathcal{H}_{SB} = g S^x \sum_{k=1}^N I_k^x$  is the interactions of the system with the bath spins,  $\mathcal{H}_B$  is the Hamiltonian for interactions among the bath spins. The spin operators are  $S$  and  $I$ , for example  $S^z = \frac{1}{2} \sigma^z$  with  $\sigma^z$  the Pauli spin matrix.

For the case where  $\mathcal{H}_B = 0$  the form of the interactions with the bath ensure that the total  $x$  component of the spins of  $I$  are a conserved quantity. We consider the case where the system spin is initially in the down state. Therefore the time dependence of the  $z$  component of the single spin that makes up the system is given by [7]

$$\langle \vec{S}(t) \rangle = \text{Tr}(\rho(0) \vec{S}(t)) = \sum_{m=-\frac{N}{2}}^{\frac{N}{2}} \lambda_m \text{Tr}(\vec{S}_m(t) \rho_m(0)) \quad (2)$$

where  $\rho(0)$  is the initial quantum density matrix and

$$\lambda_m = \binom{N}{\frac{N}{2} - m} = \frac{N!}{\left(\frac{N}{2} - m\right)! \left(\frac{N}{2} + m\right)!}. \quad (3)$$

In addition  $\langle S^x(t) \rangle = \langle S^y(t) \rangle = 0$ . Define  $\Omega_m = \sqrt{(mg)^2 + \omega_0^2}$ . Then one obtains [7]

$$\langle S^z(t) \rangle = -\frac{1}{2^{N+1}} \sum_{m=-\frac{N}{2}}^{\frac{N}{2}} \frac{\lambda_m}{\Omega_m^2} \left[ m^2 g^2 \cos(\Omega_m t) + \omega_0^2 \right]. \quad (4)$$

Using the Laplace-de Moivre theorem

$$\frac{1}{2^N} \sum_m \lambda_m \approx \sqrt{\frac{2}{\pi N}} \int_{-\infty}^{\infty} dm \exp\left(-\frac{2m^2}{N}\right) \quad (5)$$

gives the asymptotic result for  $\omega_0 = 0$  and large  $N$ , valid for all times  $t$ ,

$$\langle S^z(t) \rangle = -\frac{1}{2} \exp\left(-\frac{Ng^2 t^2}{8}\right). \quad (6)$$

The quantum purity in our case is defined as

$$\mathcal{P}(t) = \text{Tr}(\rho^2) = \frac{1}{2} + 2 \langle S^z \rangle^2 = \frac{1}{2} + \frac{1}{2} \exp\left(-\frac{Ng^2 t^2}{4}\right). \quad (7)$$

Note the factor of  $\frac{1}{4}$  in the exponential, compared to Eq. (27) of Ref. [7]. A rederivation of their result [8] gives the correct factor in Eq. 7. For a pure quantum state  $\mathcal{P} = 1$ , and hence the rate of decrease of the purity is a measure of the decoherence of the spin of the system.

## 3. Results

In Fig. (1a) we show results for  $2\mathcal{P}(t) - 1$  as a function of  $N t^2$ . The quantum purity was calculated from Eq. (7) and Eq. (4), with  $g=1$ . To compare with the asymptotic form we have plotted  $2\mathcal{P}(t) - 1$  versus  $N t^2$  rather than as a function of either the number of spins  $N$  in the bath or the time  $t$ . Clearly to go to larger values of  $N t^2$  the number of spins in the bath,  $N$ , must increase. There is a limit to the number of bath spins which can be simulated, currently on

the largest supercomputers one is restricted to about  $N \approx 40$ . Our calculation results performed for  $N = 8$  and  $N = 16$  were indistinguishable from the results of Eq. (4), until numerical rounding error occurred for very small values of  $2\mathcal{P}(t) - 1$ . In Fig. (1b) we show results for the fractional error due to finite  $N$  in  $2\mathcal{P}(t) - 1$  as a function of  $N$ . The fractional error is defined as

$$\text{fractional error} = \frac{[2\mathcal{P}(t) - 1]_{\text{asymptotic}} - [2\mathcal{P}(t) - 1]_{\text{calculated}}}{[2\mathcal{P}(t) - 1]_{\text{asymptotic}}}. \quad (8)$$

As long as  $N$  is large enough the fractional error for fixed  $Nt^2$  approaches the exact result as a power law in  $N$ . The computer calculations are limited in the number of bath spins  $N$ , and hence can reach the power law approach to the asymptotic value only for small values of  $Nt^2$ .




Figure 1: (a) The decay of the quantum purity as a function of the variable  $Nt^2$ . For  $N = 8$  and  $N = 16$  the results from the simulation are the same as the results shown here from Eq. (7). (b) The fractional error in  $2\mathcal{P}(t) - 1$  as a function of  $N$  for three values of  $Nt^2$ .

The computer calculation is limited to values of  $N$  of less than about forty. However, unlike the exact calculation the computer calculations are not limited to the case  $\mathcal{H}_B = 0$ . Therefore it is natural to ask whether or not having interactions within the bath can allow a closer approach to the asymptotic limit. This is shown in Fig. (2) for  $N = 10, 20$ , and  $30$ . The exact calculation is from Eq. (7) and Eq. (4). The label ‘No BI’ is from the program for  $\mathcal{H}_B = 0$ , and is indistinguishable on this scale from the exact results. For the case of interactions within the bath the bath Hamiltonian is

$$\mathcal{H}_B = \sum_{k=1}^N \sum_{j=1, j \neq k}^N \left( g_{k,j,x} I_k^x I_j^x + g_{k,j,y} I_k^y I_j^y + g_{k,j,z} I_k^z I_j^z \right) \quad (9)$$

for the components of the spin  $\frac{1}{2}$  operators  $I_k$ . Each component of the values of  $g_{kj}$  were chosen randomly from a distribution uniformly distributed between chosen bounds  $[-g_{\max}, g_{\max}]$ . Fig. (2) shows that there exists a value of  $g_{\max}$  for a given value of  $Nt^2$  that brings the value of the purity closer to the asymptotic value. However for a fixed value of  $Nt^2$  making the random bath interactions too strong hinders the approach with  $N$  to the asymptotic limit.

#### 4. Conclusions and Discussion

We presented results for a single quantum spin  $\frac{1}{2}$  coupled to a bath of  $N$  quantum spin  $\frac{1}{2}$  particles with a coupling  $g$  between the  $x$ -components of the spin operators. The comparison of the simulation results with the exact results for no interactions between the bath spins is excellent. Furthermore, for large enough  $N$  for a fixed value of  $Nt^2$  the approach to the exact result is a power law in  $N$ . Adding random interactions among the bath spins changes the approach to the large  $N$  limit for a fixed value of  $Nt^2$ .




Figure 2:  $2\mathcal{P}(t) - 1$  is shown as a function of  $N_{\text{bath}}^2$ . The asymptotic result is shown, as is the theoretical and program result for no bath interaction (BI). Also shown are a single realization of the random bath interactions with different strengths  $g_{\text{max}}$ . The graphs are for  $N = 10, 20$ , and  $30$  from top to bottom.

A number of questions remain to be addressed. First, is there a mechanism to systematically pick the strength of the random bath interactions,  $g_{\text{max}}$ , to approach the asymptotic limit more quickly for fixed  $N$ ? Second, can adding interactions among the bath spins also allow a faster approach to the asymptotic limit when there is more than one bath? Third, how can this approach improve on current simulations of quantum computers? Answers to these questions will hopefully be found in the near future.

## Acknowledgements

This research was supported in part by the U.S. National Science Foundation through TeraGrid resources provided by TACC Lonestar under grant number DMR100059.

- [1] M. Le Bellac, A Short Introduction to Quantum Information and Quantum Computing, Cambridge University Press, Cambridge U.K., 2005.
- [2] M. Nakahara, T. Ohmi, Quantum Computing, From Linear Algebra to Physical Realizations, CRC Press, Boca Raton, FL, 2008.
- [3] V. V. Dobrovitski, H. De Raedt, An efficient scheme for numerical simulation of the spin-bath decoherence, Phys. Rev. E 67 (056702) (2003) [7 pages].
- [4] V. V. Dobrovitski, H. De Raedt, Numerical simulations of decoherence suppression in open quantum spin systems, HAIT J. of Science and Engineering 1 (2004) 586–597.
- [5] K. De Raedt, K. Michielsen, H. De Raedt, B. Trieu, G. Arnold, M. Richter, T. Lippert, H. Watanabe, N. Ito, Massive parallel quantum computer simulator, Comp. Phys. Comm. 176 (2007) 642–651.
- [6] S. Yuan, M. I. Katsnelson, H. De Raedt, Decoherence by a spin thermal bath: Role of spin-spin interactions and initial state of the bath, Phys. Rev. B 77 (2008) 184301 [8 pages].
- [7] D. D. Rao, H. Kohler, F. Sols, Quantum frustration of dissipation by a spin bath, New Journal of Physics 10 (115017) (2008) [17 pages].
- [8] M. L. Guerra, Dynamics of classical particles and quantum spin systems, Ph.D. thesis, Mississippi State University (2011).