

University of Groningen

Biophysical properties of ladderane lipids derived from anammox bacteria

Boumann, Henry; Longo, Marjorie; Stroeve, Pieter; Jetten, Mike; Poolman, Bert; Sinninghe Damsté, Jaap; Schouten, Stefan

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2007

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Boumann, H., Longo, M., Stroeve, P., Jetten, M., Poolman, B., Sinninghe Damsté, J., & Schouten, S., (2007). *Biophysical properties of ladderane lipids derived from anammox bacteria*, 1 p.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Abstracts

Oral Presentations

SO 1

Biophysical properties of ladderane lipids derived from anammox bacteria

Henry Boumann¹, Marjorie Longo², Pieter Stroeve², Mike Jetten³, Bert Poolman⁴, Jaap Sinninghe Damsté¹, Stefan Schouten¹

¹ *Department of Marine Biogeochemistry & Toxicology, Royal Netherlands Institute for Sea Research, The Netherlands;* ² *Department of Chemical Engineering and Material Science, University of California, USA;* ³ *Department of Microbiology, University of Nijmegen, The Netherlands;* ⁴ *Department of Membrane Enzymology, University of Groningen, The Netherlands*

Anammox bacteria, originally isolated from wastewater treatment systems, play an essential role in the biogeochemical nitrogen cycle by anaerobically oxidizing ammonium to dinitrogen gas. This anammox metabolism takes place in the anammoxosome, an intracytoplasmic compartment enclosed by an exceptionally dense membrane that may serve to prevent diffusion of toxic intermediates and protons. The core lipids of this membrane comprise extraordinary stair-case like 'ladderane' hydrocarbon chains that have three or five linearly concatenated cyclobutane structures. Recently, we have determined the full identity of the intact membrane lipids. Mass spectrometry analysis revealed a wide variety of ladderane lipid species with either a phosphocholine or phosphoethanolamine polar headgroup attached to the glycerol backbone. In addition, we have reconstructed the *in silico* biosynthetic pathway for the phosphoethanolamine and phosphocholine headgroups of the anammox phospholipids from the genome assembly of *K. stuttgartiensis*. Subsequently, we have isolated substantial amounts of intact ladderane lipids to exam-

ine these molecules in membrane-like environments. In agreement with molecular modeling studies, Langmuir monolayer experiments demonstrate that ladderane lipids pack very well. Similar to fluorescence depolarization results, the monolayer data also indicate that the rigid ladderane lipids are in the liquid state at physiological relevant temperatures. The purified ladderane lipids have also been reconstituted into bilayer membrane vesicles and visualized by fluorescence and cryo-electron microscopy. On-going studies include micropipette aspiration to determine the mechanical properties of the ladderane lipids assembled in giant unilamellar vesicles and monitoring the proton permeability of ladderane membranes. The combined outcome will shed further light on the function of the ladderane lipid structures in the membrane of these environmentally important organisms.

doi:10.1016/j.chemphyslip.2007.06.024

SO 2

X-ray diffraction and calorimetric study on the influence of sterols on the phase properties of phospholipids

Lin Chen¹, Jayne Lawrence², Zhi Wu Yu³, Peter Quinn¹

¹ *Departments of Biochemistry, King's College, UK;*

² *Departments of Pharmacy, King's College, UK;*

³ *Department of Chemistry, Tsinghua University, China*

The effect of cholesterol, stigmasterol and ergosterol on the formation of liquid ordered L_o phase in the equimolar mixtures of dipalmitoylphosphatidylcholine (DPPC) and dioleoylphosphocholine (DOPC) mixtures was investigated by differential scanning calorimetry and