

University of Groningen

The Structure of the CF1 Part of the ATP-Synthase Complex from Chloroplasts

Boekema, Egbert J.; Lücken, Uwe

Published in:
EPRINTS-BOOK-TITLE

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1996

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):
Boekema, E. J., & Lücken, U. (1996). The Structure of the CF1 Part of the ATP-Synthase Complex from Chloroplasts. In *EPRINTS-BOOK-TITLE*

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Chapter 26

The Structure of the CF_1 Part of the ATP-Synthase Complex from Chloroplasts

Egbert J. Boekema

*Bioson Research Institute, Biophysical Chemistry, University of Groningen,
Nijenborgh 4, 9747 AG Groningen, The Netherlands*

Uwe Lücken*

Fritz-Haber-Institute, Faradayweg 4-6, D-14195 Berlin, Germany

Summary	487
I. Introduction.....	488
II. Structural Aspects of CF_1	488
A. Overall Shape of CF_1	488
B. Structure of the Individual Subunits.....	489
1. Shape of the α - and β -Subunits.....	489
2. Shape of the Small Subunits γ , δ and ϵ	489
C. Positions and Interactions of the Subunits.....	490
D. Conformational Changes in Subunits and Subunit Positions.....	491
III. Concluding Remarks	491
Acknowledgments.....	492
References	492

Summary

The proton ATP synthase consists of a membrane integrated part, F_0 , and a hydrophilic part, F_1 . F_1 is composed of five different subunits: α , β , γ , δ and ϵ . This chapter focuses on the chloroplast F_1 (CF_1) structure and discusses the overall shape and dimensions of CF_1 , shape and size of the various subunits, subunit interactions and conformational changes in the subunit positions related to catalysis. Structural data originate mainly from X-ray diffraction and electron microscopy. Recently, the structure of F_1 from beef heart mitochondria has been determined at 2.8 Å and this structure, although not fully identical to CF_1 , can be taken as a blueprint for models of the CF_1 structure.

* Present Address: Electron Optics Laboratory, Philips, 5600 MD Eindhoven, The Netherlands

I. Introduction

The proton ATP synthase from chloroplasts, mitochondria and eubacteria consists of a membrane-integrated part, F_o , which acts as a proton channel through the membrane and a hydrophilic part, F_1 , which contains the nucleotide-binding sites. The ATP synthases from different sources have a very similar structure. This is especially the case for the F_1 part. The F_1 part from spinach chloroplasts, CF_1 , is composed of five different subunits: α (55.4 kDa), β (53.9 kDa), γ (35.7 kDa), δ (20 kDa) and ϵ (14.7 kDa). One CF_1 molecule contains three copies of the large α and β subunits, and one copy each of the smaller subunits making a total mass of nearly 400 kDa. This chapter focuses on the CF_1 structure. The lack of a 3D structure with high resolution forces us to discuss also the F_1 structure from mitochondria and eubacteria. Results obtained with related F_1 's are also relevant for CF_1 structure; at the low resolution level the mitochondrial (MF_1) and CF_1 structures have a striking similarity (Boekema et al., 1990), although all 5 subunits differ in mass and amino acid sequence, resulting in a 10% smaller mass for mitochondrial F_1 . Especially the two smallest subunits differ considerably, but the fact that the δ subunits of MF_1 and CF_1 have a very different amino acid sequence has often been overemphasized, because both δ subunits are functionally related (Engelbrecht and Junge, 1990).

There are three main methods to investigate the structure of a protein: X-ray diffraction, NMR and electron microscopy (EM). For F_1 , the first and the third method have been widely applied. By X-ray diffraction of three-dimensional crystals the structure can be determined at atomic resolution. A structure of rat liver MF_1 at the sub-atomic level has been published (Bianchet et al., 1991). More recently, based on crystals from beef heart MF_1 (Abrahams et al., 1993) a 2.8 Å resolution structure has been determined (Abrahams et al., 1994). NMR has the same potential as X-ray diffraction but has the disadvantage that the objects studied should be smaller than 40 kDa. Thus it can only be applied to the structure elucidation of the small F_1 subunits. Results of such work have not yet been reported. With EM it is very difficult to determine a high-

resolution structure, because it needs large two-dimensional (2D) crystals. With small 2D crystals a low-resolution 3D model can be obtained (Gogol et al., 1989a; Ishii et al., 1993). Low-resolution data can also be easily obtained with single-particle averaging (Boekema et al., 1986; Gogol et al., 1989b). In this chapter we will mainly discuss the outcomes of these techniques and, where necessary, extend them with results of biochemical investigations.

II. Structural Aspects of CF_1

A. Overall Shape of CF_1

If the electron density of a protein is solved by X-ray diffraction with a resolution of better than 3 Å, there is enough structural detail to define the amino acid chain backbone. If the resolution is lower, the ambiguity of fitting amino acids increases and above 5 Å only secondary elements, such as large α -helices, can be recognized. First, an X-ray study on rat liver mitochondrial F_1 was published which did not show much of the amino acid chain backbone, although a resolution of 3.5 Å resolution was claimed (Bianchet et al., 1991). But more recently, the beef heart mitochondrial ATPase structure could be solved at 2.8 Å resolution (Abrahams et al., 1994) and this enabled the determination of the amino acid chain backbone of the α - and β subunits and most of the γ subunit. We will discuss the features from both X-ray structures mainly in connection with the EM and other data on CF_1 .

The maximal diameter of CF_1 parallel to the plane of the membrane is close to 110 Å, because this is the repeating distance of CF_1 in two-dimensional crystals (Boekema and Böttcher, 1992). In crystals the repeat can be determined by EM with an error of only a few Å, even if the resolution of the image features is much lower. The diameter of bovine heart MF_1 is 100 Å (Abrahams et al., 1994). This is somewhat in discrepancy with the 120 Å diameter from the rat liver MF_1 model (Bianchet et al., 1991). The height of CF_1 vertical to the membrane plane is about 83 Å. A similar height has been found for mitochondrial F_1 (Boekema et al., 1988; Abrahams et al., 1994) and *E. coli* F_1 (90 Å; Lücken et al., 1990). Average side views of F_1 complexes in CF_oF_1 (Fig. 1) and *E. coli* F_oF_1 (Lücken et al., 1990), recorded by EM, show a trapezoid form for symmetrical views (Fig. 3) of F_1 ,

Abbreviations: CF_1 – chloroplast F_1 ; DTT – dithiothreitol; EM – electron microscopy; MF_1 – mitochondrial F_1 ; NEM – N-ethylmaleimide




Fig. 1. Cryo-electron microscopy and image analysis of unstained F_0F_1 ATPase from spinach chloroplasts. (A) EM of F_0F_1 ATPase reconstituted in lipid vesicles (B) average image of reconstituted F_0F_1 particles (C) average image of F_0F_1 particles labeled with a gold cluster, which binds specifically to the S-S disulfide in the loop of the γ subunit. Procedure: the -SH groups in CF_1 were blocked with NEM, the S-S bond was reduced with DTT and the reduced -SH groups labeled with a monomaleimidogold label. The S-S group is in the periphery of CF_1 (position marked by an arrow), in accordance with the fluorescence energy transfer data (see chapter 23). (D) structure of the bovine heart MF_1 (Abrahams et al., 1993) shown on the same scale. For representation and comparison the electron density was low-pass filtered to enhance the low-resolution features of the map. The bar represents 100 Å.

B. Structure of the Individual Subunits

1. Shape of the α - and β -Subunits

The 2.8 Å map of MF_1 (Abrahams et al., 1994) shows that the α - and β subunits are almost identically folded: each consists of N-terminal six-stranded β -barrel, a central α - β domain containing the nucleotide-binding site and a C-terminal bundle of 7 and 6 helices, respectively. The X-ray model confirms and extends all previous work on the structure of the

large subunits, such as the fact that the distribution of 21 protease-cleaved sites in the isolated α subunit is similar to that of the β subunit, thus providing experimental evidence for a similar folding topology of the two subunits (Tozawa et al., 1993).

From EM of negatively stained specimens followed by computer analysis (Boekema et al., 1992) it could be concluded that the overall shape of the isolated β subunit of spinach CF_1 is also elongated. The β subunit has dimensions of about 66–69 × 42–44 Å. This indicates that the packing of the large subunits in CF_1 and MF_1 could have a similar folding. Two independent low-resolution 3D reconstructions of bacterial F_1 from EM images also indicate that the α - and β subunits are elongated. In a model of Gogol et al. (1989a) these subunits are very elongated, e.g. 90 × 30 Å. In the model of Ishii et al., 1993 they are shown to be 70 × 30 Å.

2. Shape of the Small Subunits γ , δ and ϵ

By analytical ultracentrifugation experiments it was found that subunit γ from *E. coli* F_1 is rather elongated (Dunn, 1982). EM analysis of a CF_1 particle, from which subunits δ and ϵ were removed, showed that the diameter of the central mass was 25–28 Å, much smaller than the projected diameter of the large subunits (Fig. 2). Considering the mass of γ , 35.7 kDa, this indicates that it is an elongated subunit with a calculated length of 50 Å (Boekema et al., 1990). The X-ray map of bovine mitochondrial F_1 shows that the γ subunit is even much longer: a 90 Å long α -helix forms the center of the MF_1 structure, a second α -helix of 60 Å forms a left-handed antiparallel coiled coil with the first helix (Abrahams et al., 1994).

The chloroplast γ subunit has an interesting component, which is a flexible loop that includes a disulfide bond (McCarty and Hammes, 1987). The distance of this disulfide bond to two other sulfhydryl groups on γ , determined by fluorescence resonance energy transfer, was 44 and 47 Å, or about 38 Å, if projected in a plane parallel to the membrane. Thus this loop, which is important for ATPase activity, is really a loop protruding from the rest of the γ subunit. The position of the S-S disulfide was also investigated by cryo-EM (U. Lücken, unpublished data) using a monomaleimidogold label (Wilkins and Capaldi, 1992). Results indicate that the S-S disulfide bridge is indeed extending far from the center of CF_1 (Fig. 1).

Unfortunately, the atomic structure of δ and ϵ




Fig. 2. Electron microscopy of negatively stained CF_1 . (A) average image of holo-enzyme CF_1 in the hexagonal view (modified from Boekema et al., 1990); (B) average image of CF_1 depleted of subunits δ and ϵ (modified from Boekema et al., 1990); (C) and (D) average images of projections from CF_1 particles arranged upside-up and upside-down on the carbon support. In general, in negatively stained samples the lower half of the particle is more strongly embedded in the stain layer and this leads to a stronger contrast of the features of this part in the projection. In the predominant view (A and the similar view C) the α - and β subunits that are close to δ appear less pronounced, because the structure is slightly lifted up by δ . The α - and β subunits become more evenly embedded in the stain layer when the δ subunit is absent, as in Fig. B, or when δ is away from the carbon support in upside-down attached particles, as in D. The bar represents 20 Å.

could not be determined in the 2.8 Å density map (Abrahams et al., 1994). The chloroplast δ subunit has an elongated shape (Wagner et al., 1988). It is predicted to be an α -helical protein with only limited β -structure. Conserved amino acids are found in the N-terminal start of a putative amphipathic α -helix (Hoesche and Berzborn, 1993). The shape of the chloroplast ϵ subunit is not known, ϵ from *E. coli* has been crystallized (Codd et al., 1992), but the structure has not yet been reported.

C. Positions and Interactions of the Subunits

From thermal denaturation experiments it was found that the major forces that stabilize CF_1 must be between the α , β and γ subunits; subunits δ and ϵ have little effect on the stability (Wang et al., 1993). Thus

the three largest subunits with a total of 7 copies define the shape of CF_1 . Nevertheless, the smallest subunits are not totally unimportant for the overall shape. It was demonstrated that the positions of the α subunits in CF_1 changed when δ or δ plus ϵ were removed (Boekema and Böttcher, 1992).

Although not all interesting questions concerning the detailed structure can be answered, several conclusions concerning the arrangement of the large subunits and small subunits can be drawn:

1. The overall shape of CF_1 is asymmetric. This means that all three α - and all three β subunits 'see' a different surrounding environment. Several lines of evidence support this conclusion. For instance; the three β - β center-to-center average distances in CF_1 are 79, 77 and 73 Å (Boekema and Böttcher, 1992). Similar asymmetrical distances were found for the β subunits in *E. coli* F_1 (Wilkins and Capaldi, 1994). The 2.8 Å resolution structure of MF_1 (Abrahams et al., 1994) shows the asymmetry of the overall structure in a very detailed way. The main reason why the other X-ray map presents a symmetric F_1 structure (Bianchet et al., 1991) is the fact that a polypeptide chain can be better fitted if noisy data are 3-fold averaged. The large subunits not only 'see' a different surrounding but also have a slightly different conformation. For example: only one of the three α subunits of CF_1 reacts with Lucifer yellow which modifies lysine 378 (McCarty and Hammes, 1987). Similarly, Wilkins and Capaldi (1992) found that only one of the α subunits in *E. coli* F_1 has a cysteine group that reacts with maleimides. The inequivalent catalytic sites in CF_1 originate, of course, also from a structural difference (see chapters 24 and 25).

2. In the plane parallel to the membrane, the six copies of the large subunits alternate in position. This was first shown unambiguously by immunoelectron microscopy with monoclonal antibodies (Gogol et al., 1989b).

3. There is some evidence that the α - and β subunits in CF_1 are not exactly at the same height as shown in Fig 3. The MF_1 model from Bianchet et al. (1991) shows two slightly offset, interdigitated layers of large subunits. At the top, the β subunits protrude about 15 Å higher than the α subunits. The CF_1 data indicate a similar arrangement

(Boekema et al., 1992). However, the 2.8 Å resolution structure of MF₁ shows that only one β subunit (β_{DP}) is at a substantially higher level in the structure, whereas two α subunits are extending slightly more to the base of F₁.

4. A seventh mass seen in the center of the α₃β₃ configuration (Fig. 2) is formed by the small subunits and consists mainly of subunit γ (Boekema et al., 1990; Wilkens and Capaldi, 1992).

5. The small subunits are positioned in the lower half of F₁, which is the part of F₁ closest to the membrane. This was already inferred based on biochemical evidence, but recently Ishii et al (1993) directly showed that the central mass is in the lower half of F₁.

6. Subunit α interacts with an α subunit and with subunit I, a F₀ subunit, which contributes to the stalk connecting F₁ to F₀ (Beckers et al., 1992). Although subunit δ is partly hidden within CF₁, it must also be involved in the stalk, because it forms a protrusion on F₁. In original EM images the protrusion was barely visible, but the large subunits around δ look differently embedded in projections of negatively stained specimens (Fig. 2). The protrusion is clearly visible in the structure of bovine heart MF₁ (Abrahams et al., 1993), where it consists of two α-helices in a coiled-coil making a 40 Å stem (Fig. 1D). But it needs to be confirmed if the entire protrusion consists of only the δ subunit.

7. In CF₁ the subunits γ and ε are close and interact with each other (McCarty and Hammes, 1987). Interestingly, a complex of γ and ε was purified and crystallized from *E. coli* (Cox et al., 1993).

8. According to Fig. 3, subunit γ is the central CF₁ subunit that interacts with all other F₁ subunits and with the stalk formed by F₀. Because it protrudes from MF₁, it is very likely that conformational changes occurring in the stalk of F₀F₁ can be of catalytic relevance if mediated by subunit γ. But also other CF₁ subunits may interact with the stalk and be directly influenced by changes occurring in the stalk. In *E. coli* F₁, a β subunit, different from the one which interacts with the ε subunit, binds to the b subunit of the F₀ part (Wilkens et al., 1994).




Fig. 3. Views of a model for CF₁ ATP synthase. A. The model in the hexagonal projection, parallel to the plane of the membrane, seen from the membrane. B. The model seen from aside in a 'symmetrical view'. C. The model seen from aside in an 'asymmetrical view'. The positions of 3B and 3C differ by 30°. The dimensions of the F₀ part and the stalk, symbolized as cylinders, have been determined, but the positions of the subunits within the membrane are still not fully clear.

D. Conformational Changes in Subunits and Subunit Positions

It has been shown that there is evidence for movements of subunit ε that could play a role in the catalytic mechanism and regulation of ATP synthesis/hydrolysis (Gogol et al., 1990; Boekema and Böttcher, 1992). Wilkens and Capaldi (1994) showed that both γ and ε subunits are mobile, probably by moving as a single domain through distances as great as 10–20 Å.

III. Concluding Remarks

Further structural data are in demand, even after the determination of the MF₁ structure from beef heart mitochondria, which is a breakthrough for the field of bioenergetics. For the precise mechanism of ATP synthesis in chloroplasts it will also be necessary to solve the CF₁ structure at atomic resolution by X-ray diffraction. Only based on such a structure will all the biochemical data attain their full significance. But F₁ is not only a fixed structure, as seen by X-rays

in a 3D crystal. Large rearrangements, such as the movement of the γ and ϵ subunit cannot be easily studied by X-ray diffraction if the conformation is unstable. Here, cryo-EM, which usually results in a much lower resolution than X-ray diffraction, will continue to be of value.

Acknowledgments

Support by the Netherlands Foundation for Chemical Research (SON) and by a grant of the European Union BIO-CT93-0078 is gratefully acknowledged.

References

- Abrahams JP, Lutter R, Todd RJ, van Raaij MJ, Leslie AGW and Walker JE (1993) Inherent asymmetry of the structure of F_1 -ATPase from bovine heart mitochondria at 6.5 Å resolution. *EMBO J* 12: 1775–1780
- Abrahams JP, Leslie AGW, Lutter R and Walker JE (1994) Structure at 2.8 Å resolution of F_1 -ATPase from bovine heart mitochondria. *Nature* 370: 621–628
- Beckers G, Berzborn RJ and Strotmann H (1992) Zero-length crosslinking between subunits δ and I of the H^+ -translocating ATPase of chloroplasts. *Biochim Biophys Acta* 1101: 97–104
- Bianchet M, Ysern X, Hullihien J, Pedersen PL and Amzel, LM (1991) Mitochondrial ATP synthase. Quaternary structure of the F_1 moiety at 3.6 Å determined by X-ray diffraction. *J Biol Chem* 266: 21197–21201
- Boekema EJ and Böttcher B (1992) The structure of ATP synthase from chloroplasts. Conformational changes of CF_1 studied by electron microscopy. *Biochim Biophys Acta* 1098: 131–143
- Boekema EJ, Berden JA, and van Heel MG (1986) Structure of mitochondrial F_1 -ATPase studied by electron microscopy and image processing. *Biochim Biophys Acta* 851: 353–360
- Boekema EJ, Schmidt G, Gräber P and Berden JA (1988) Structure of the ATP-synthase from chloroplasts and mitochondria by electron microscopy. *Zeitschrift Naturforsch.* 43c: 219–225
- Boekema EJ, Xiao J, and McCarty RE (1990) Structure of the ATP synthase from chloroplasts studied by electron microscopy. Localization of the small subunits. *Biochim Biophys Acta* 1020:49–56
- Boekema EJ, Harris D, Böttcher B and Gräber P (1992) The structure of the ATP-synthase from chloroplasts. In: Murata N (ed) *Research in Photosynthesis*, Vol 2, pp 645–652. Kluwer Academic Publishers, Dordrecht
- Codd R, Cox G, Guss JM, Solomon RG and Webb D (1992) The expression, purification and crystallization of the ϵ subunit of the F_1 portion of the ATPase of *Escherichia coli*. *J Mol Biol* 228: 306–309
- Cox GB, Cromer BA, Guss JM, Harvey, I, Jeffrey PD, Solomon RG and Webb DC (1993) Formation in vivo, purification and crystallization of a complex of the γ and ϵ subunits of the F_0F_1 -ATPase of *Escherichia coli*. *J Mol Biol* 229: 1159–1162
- Dunn SD (1982) The isolated γ subunit of *Escherichia coli* F_1 ATPase binds the ϵ subunit. *J Biol Chem* 257: 7354–7359
- Engelbrecht S and Junge W (1990) Subunit δ of H^+ -ATPases At the interface between proton flow and ATP synthesis. *Biochim Biophys Acta* 1015: 379–390
- Gogol EP, Lücken U, Bork T and Capaldi RA (1989a) Molecular architecture of *Escherichia coli* F_1 adenosinetriphosphatase. *Biochemistry* 28: 4709–4716
- Gogol EP, Aggeler R, Sagermann M and Capaldi RA (1989b) Cryoelectron microscopy of *Escherichia coli* F_1 adenosinetriphosphatase decorated with monoclonal antibodies to individual subunits of the complex. *Biochemistry* 28: 4717–4724
- Gogol EP, Johnston E, Aggeler R and Capaldi RA (1990) Ligand-dependent structural variations in *Escherichia coli* F_1 ATPase revealed by cryoelectron microscopy. *Proc Natl Acad Sci USA* 87: 9585–9589
- Hoesche JA and Berzborn R (1993) Primary structure, deduced from cDNA, secondary structure analysis and conclusions concerning interaction surfaces of the δ subunit of the photosynthetic ATP-synthase (E.C. 3.6.1.34) from millet (*Sorghum bicolor*) and maize (*Zea mays*). *Biochim Biophys Acta* 1142: 293–305
- Ishii N, Yoshimura H, Nagayama K, Kagawa Y and Yoshida M (1993) Three-dimensional structure of F_1 -ATPase of thermophilic bacterium PS3 obtained by electron crystallography. *J Biochem* 113: 245–250
- Lücken U, Gogol EP and Capaldi RA (1990) Structure of the ATP synthase complex (ECF_1F_0) of *Escherichia coli* from cryoelectron microscopy. *Biochemistry* 29: 5339–5343
- McCarty, RE and Hammes GG (1987) Molecular architecture of chloroplast coupling factor 1. *Trends Biol Sci* 12: 234–237
- Tozawa K, Miyauchi M and Yoshida M (1993) Structure of the α subunit of F_1 -ATPase probed by limited proteolysis. *J Biol Chem* 268: 19044–19054
- Wagner R, Apley EC, Engelbrecht S and Junge W (1988) The binding of eosin-labeled subunit δ to the isolated chloroplast ATPase, CF_1 , as revealed by rotational diffusion in solution. *FEBS Lett* 230: 109–115
- Wang Z-Y, Freire E and McCarty RE (1993) Influence of nucleotide binding site occupancy on the thermal stability of the F_1 portion of the chloroplast ATP synthase. *J Biol Chem* 268: 20785–20790
- Wilkens S and Capaldi RA (1992) Monomaleimidogold labeling of the γ subunit of the *Escherichia coli* F_1 ATPase examined by cryoelectron microscopy. *Arch Biochem Biophys* 299: 105–109
- Wilkens S and Capaldi RA (1994) Asymmetry and structural changes in ECF_1 examined by cryoelectron microscopy. *Biol Chem Hoppe-Seyler* 375: 43–51
- Wilkens S, Dunn SD and Capaldi RA (1994) A cryoelectron microscopy study of the interaction of the *Escherichia coli* F_1 -ATPase with subunit β dimer. *FEBS Lett.* 354: 37–40