

## University of Groningen

### Nodal metasases and biomarkers in melanoma

Kruijff, Schelto

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

2011

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Kruijff, S. (2011). *Nodal metasases and biomarkers in melanoma*. [s.n.].

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

# Nodal metastases and biomarkers in melanoma

Schelto Kruijff

2011

The printing of this thesis was financially supported by:

## MediRisk

Medisch Spectrum Twente (MST)

Maatschap chirurgie, MST

Eurocept B.V.

Novartis Pharma B.V.

Johnson & Johnson Medical B.V.

Roche Nederland B.V.

Mundipharma Pharmaceuticals B.V.

Stichting Melanoom

DiaSorin SA/NV

Covidien Nederland B.V.

Kruijff, S

Nodal metastases and biomarkers in melanoma

Thesis, University of Groningen, The Netherlands

ISBN: 978-90-367-5180-3

ISBN: 978-90-367-5179-7

© Copyright 2011 Schelto Kruijff, The Netherlands

All rights reserved. No part of this book may be reproduced, stored in a retrieval system or transmitted in any form or by any means, without prior permission of the author.

Lay out : Chris Bor, Medische fotografie en illustratie, AMC, Amsterdam

Printed by : Buijten en Schipperheijn, Amsterdam

Foto omslag : Duitse kinderen op Nederlands strand in 1930, fotograaf onbekend

RIJKSUNIVERSITEIT GRONINGEN

# Nodal metastases and biomarkers in melanoma

Proefschrift

ter verkrijging van het doctoraat in de  
Medische Wetenschappen  
aan de Rijksuniversiteit Groningen  
op gezag van de  
Rector Magnificus, dr. E. Sterken,  
in het openbaar te verdedigen op  
maandag 12 december 2011  
om 12.45 uur

door

Schelto Kruijff  
geboren op 18 mei 1977  
te Rijpwetering

Promotor : Prof. dr. H.J. Hoekstra

Beoordelingscommissie: Prof. dr. E. Heineman  
Prof. dr. I.H.M. Borel Rinke  
Prof. dr. G.A.P. Hospers

Paranimfen:

Philip de Reuver

Herman Frima

Jelle Jansen

Voor Annelies



## TABLE OF CONTENTS

Chapter 1	General introduction and outline of this thesis	9
<b>Part I</b>	<b>Detection and treatment of nodal melanoma metastases</b>	
Chapter 2	Detection of melanoma nodal metastases; differences in detection between elderly and younger patients do not affect survival	27
Chapter 3	Therapeutic groin dissection for melanoma: risk factors for short term morbidity	41
<b>Part II</b>	<b>Prognostic markers in melanoma</b>	
Chapter 4	Breslow thickness in the netherlands: a population-based study of 40,880 patients comparing young and elderly patients	57
Chapter 5	S-100B concentrations predict disease-free survival in stage III melanoma patients	69
Chapter 6	The value of pre-operative S-100B and SUV in clinically Stage III melanoma patients undergoing therapeutic lymph node dissection	83
Chapter 7	The use of S-100B to evaluate therapy effects during Bevacizumab induction treatment in AJCC stage III melanoma	99
Chapter 8	The current status of S-100B as a biomarker in melanoma	111
Chapter 9	General conclusion and future perspectives	121
Chapter 10	Summary	133
	Samenvatting	141
	List of contributing authors	147
	List of publications	149
	Dankwoord/Acknowledgement	151
	Curriculum Vitae	152



# 1



## General introduction and outline of thesis

---

### GENERAL

Melanoma is the most malignant of all skin cancer types. It causes more than 75% of all skin cancer related mortality.<sup>1</sup> Cutaneous melanoma incidence has increased rapidly in the last years and estimates predict a continuing increase.<sup>2</sup> In the Netherlands, the total number of melanoma patients is expected to increase from 2,400 patients in 2000 to 4,800 patients in 2015 and already reached 3,500 patients in 2005.<sup>2</sup> The incidence and mortality of populations have reached either a plateau or maybe a decline in younger age groups while there is an increase in older age groups.<sup>3-7</sup> In Australia, mortality rates seem to have reached a plateau, even though incidence rates are increasing. This may be due to the fact that most new melanomas are thinner which usually do not lead to short term death.<sup>8</sup>

#### *AJCC Staging*

Patients with melanoma are staged according to the American Joint Committee on Cancer TNM-staging manual. In 2009 the 7<sup>th</sup> version was published and a summary is given in table 1 and 2.<sup>9</sup> This updated manual has an evidence-based approach and reflects the improved understanding of the disease.<sup>9</sup> The main histological biomarkers influencing T- staging are Breslow thickness and ulceration. N-staging is influenced by the involvement of regional lymph nodes. In the latest version of the AJCC staging system, micro-metastases were also included which are nowadays detected as a consequence of the routine use of the sentinel node biopsy procedure in the treatment of melanoma. The M-staging describes metastatic disease and differentiates between subcutaneous or extra regional involved lymph nodes, pulmonary metastases and other metastatic disease. For the first time a serum biomarker LDH was added to the AJCC staging system.

**TABLE 1** TNM Staging categories for melanoma (American Joint Committee on Cancer 2009)<sup>(9)</sup>

<b>Tumour</b>	<b>Breslow thickness</b>	<b>Ulceration status/mitoses</b>
<b>Tis</b>	NA	NA
<b>T1</b>	≤ 1,0 mm	T1a: without ulceration and mitoses < 1/mm <sup>2</sup> T1b: with ulceration and/or mitoses ≥ 1/mm <sup>2</sup>
<b>T2</b>	1,01-2,0 mm	T2a: without ulceration T2b: with ulceration
<b>T3</b>	2,01-4,0 mm	T3a: without ulceration T3b: with ulceration
<b>T4</b>	> 4,0 mm	T4a: without ulceration T4b: with ulceration
<b>Node</b>	<b>No. of metastatic nodes</b>	<b>Nodal metastatic burden</b>
<b>N0</b>	0	NA
<b>N1</b>	1	N1a: micrometastasis* N1b: macrometastasis**
<b>N2</b>	2-3	N2a: micrometastasis* N2b: macrometastasis** N2c: In-transitmetastases/satellites without metastatic nodes.
<b>N3</b>	4+ metastatic nodes, or matted nodes, or in-transitmetastases/satellites with metastatic nodes	
<b>Metastases Site</b>	<b>Serum LDH</b>	
<b>M0</b>	No distant metastasis	NA
<b>M1a</b>	Distant skin, subcutaneous or nodal metastases	Normal
<b>M1b</b>	Lung metastasis	Normal
<b>M1c</b>	All other visceral metastasis	Normal
	Any distant metastasis	Elevated

\*Micrometastases are diagnosed after sentinel lymph node biopsy. \*\*Macrometastases are defined as clinically detectable nodal metastases confirmed pathologically. LDH Lactate dehydrogenase. NA Not applicable

### *Treatment and staging of primary localized melanoma; AJCC stage I/II*

Local control is the main component of primary cutaneous melanoma treatment. Therefore, surgical resection is the first choice of treatment. The treatment of a suspect nevus or skin lesion is a diagnostic excision with a 2 to 3 millimetre margin of normal skin. Most important is a disease free margin. According to the Dutch melanoma guidelines and most other international guidelines the therapeutic, surgical excision margins are defined as 0,5 cm for in-situ melanoma, 1 cm for T 1-2 melanoma and 2 cm for T 3-4 melanoma (Table 3).<sup>10-28</sup> In case of unfavourable melanoma characteristics, excision margins of more than 2 centimetres, if not harmful for the patient, are advised in the majority of guidelines.<sup>19,30</sup>

### *Nodal staging: AJCC stage IIIa en Sentinel node biopsy (SLNB)*

The value of sentinel node biopsy (SLNB) in Stage I and II patients has been controversial.<sup>31</sup> In most countries SLNB is performed as a staging procedure in patients with melanomas

**TABLE 2** Anatomic stage groupings for melanoma (AJCC 2009)<sup>(9)</sup>

Clinical staging*				Pathological staging**			
<b>O</b>	Tis	N0	M0	<b>O</b>	Tis	N0	M0
<b>IA</b>	T1a	N0	M0	<b>IA</b>	T1a	N0	M0
<b>IB</b>	T1b	N0	M0	<b>IB</b>	T1b	N0	M0
	T2a	N0	M0		T2a	N0	M0
<b>IIA</b>	T2b	N0	M0	<b>IIA</b>	T2b	N0	M0
	T3a	N0	M0		T3a	N0	M0
<b>IIB</b>	T3b	N0	M0	<b>IIB</b>	T3b	N0	M0
	T4a	N0	M0		T4a	N0	M0
<b>IIC</b>	T4b	N0	M0	<b>IIC</b>	T4b	N0	M0
<b>III</b>	Any T	N1-3	M0	<b>IIIA</b>	T1-4a	N1a	M0
					T1-4a	N2a	M0
				<b>IIIB</b>	T1-4b	N1a	M0
					T1-4b	N2a	M0
					T1-4a	N1b	M0
					T1-4a	N2b	M0
					T1-4a/b	N2c	M0
				<b>IIIC</b>	T 1-4b	N1b	M0
					T 1-4b	N2b	M0
					Any T	N3	M0
<b>IV</b>	Any T	Any N	M1	<b>IV</b>	Any T	Any N	M1

\* Clinical staging includes microstaging of the primary melanoma and clinical/radiologic evaluation for metastases. By convention, it should be used after complete excision of the primary melanoma with clinical assessment for regional and distant metastases. \*\*Pathologic staging includes microstaging of the primary melanoma and pathologic information about the regional lymph nodes after partial (ie, sentinel node biopsy) or complete lymphadenectomy. Pathologic stage 0 or stage IA patients are the exception; they do not require pathologic evaluation of their lymph nodes.

over 1 mm Breslow thickness or under 1 mm in case of prognostic unfavourable tumour characteristics. In approximately half of the guidelines SLNB is stated to be a standard, routine procedure which all patients meeting criteria should undergo (Table 3).

The results of the Multicenter Selective Lymph Node Trial (MSLT), the 3rd interim analysis, showed that the SLNB procedure reflects mainly staging information and patients who underwent a SLNB were associated with a longer disease free interval.<sup>31,32</sup> According to the Dutch melanoma guidelines, SLNB is only advised combined with inclusion in clinical trials or for patients who wish to be optimal informed about the stage of their disease.<sup>19</sup>

However, today the technique seems to have undisputedly proven to be a reliable staging method and the fourth interim analysis of the MSLT I study reveals that SLNB indeed leads to improved melanoma specific survival.<sup>33</sup>

**TABLE 3** Guidelines for primary treatment and staging primary stage I-II melanoma.

Country	Excision margins (cm)				Initial staging				
	Primary	Tis	T	T	SLNB *		Imaging		Blood tests
			≤2mm	>2mm	Tumour	Routine	US	CXR	
Australia & NZ <sup>(10)</sup>	0.2	0.5	1-2	2	>1mm	-	-	-	-
Austria <sup>(11)</sup>	0.3-1.0	0.5	1-2	3	>1mm	+	+	+	LDH,CBC
Belgium <sup>(12)</sup>	0.2	0.5	1***	2***	≥pT1b	-	+	+	-
Canada (BC) <sup>(13)</sup>	NR	0.5-1	1	2	1-4mm	+	-	0	-
Denmark <sup>(24)</sup>	0.5	0.5	1-2	2-4	pT1b	+	-	-	-
France <sup>(15)</sup>	NR	0.5	1-2	2-3	≥pT1b	-	-	-	-
Germany <sup>(16)</sup>	NR	0.5	1	2	>1mm	+	+	+	LDH,S100,CBC
Ireland <sup>(17)</sup>	0.1-0.3	0	1-2	2	≥pT1b	+	***	***	LDH**
Italy <sup>(18)</sup>	1.0	0	0	1-2	>1mm	-	-	-	-
Netherlands <sup>(19)</sup>	0.2	0.5	1	2	>1mm	-	-	-	-
Norway <sup>(20)</sup>	0.2-0.5	0.5	1	2-3	Trial	-	-	-	-
Poland <sup>(21)</sup>	0.1-0.2	0.5	1	2	≥1 mm	+	-	-	-
Scotland <sup>(22)</sup>	0.2	0.2-0.5	1-2	2	≥pT1b	-	-	-	-
South Africa <sup>(23)</sup>	0.2	0.5	1-2	2	NR	-	-	-	-
Spain <sup>(24)</sup>	0.1-0.2	0.5	1-2	2-3	≥pT1b	+	+	+	LDH
Sweden <sup>(25)</sup>	0.2	0.5	1-2	2	≥pT1b	+	-	-	-
Switzerland <sup>(16)</sup>	NR	0.5	1	2	>1mm	+	-	-	-
United Kingdom <sup>(27)</sup>	0.2-0.5	0.2-0.5	1-2	2-3	Stage II	-	***	***	LDH, CBC**
United States <sup>(28)</sup>	0.1-0.3	0.5	1-2	2	≥pT1b	+	-	-	-

\*All guidelines recommend that completion lymph node dissection (CLND) should follow positive SNLB or otherwise in clinical trial setting. \*\* Only for stage IIB and higher. \*\*\* T≤1 mm and T>1mm. CBC Complete blood cell count CXR Chest X-ray. NR Not reported in guideline. LDH Lactate dehydrogenase. LNB Sentinel lymph node biopsy. US Ultrasound of draining lymph node basins with/without abdomen

### *Palpable nodes : AJCC Stage IIIb*

In AJCC stage III melanoma patients, a therapeutic lymph node dissection is the first choice of treatment. However, this is a group of patients with a remarkable heterogeneity in survival. Recently, Charles Balch published a multivariate analysis of prognostic factors among 2,313 patients with stage III melanoma. The five year overall survival was 63% varying from 67% for patients with nodal micro-metastases to 43% for nodal macrometastases.<sup>34</sup>

Staging studies analyzing patients with palpable lymph nodes, whole-body FDG-PET and/or spiral CT have proved their value with 27% upstaging to stage IV disease. A sensitivity of 79-92% and a specificity of 86-90% to detect distant metastases was found.<sup>35,36</sup> The combined FDG-PET and spiral CT scan leads to a change in the planned dissection in 37% of the patients.<sup>37</sup> Analyzing the heterogeneity of stage III melanoma, the suspicion rises that numerous patients have a form of subclinical dissemination, which can even remain undetected

by standard FDG-PET and CT imaging.<sup>38</sup> Biomarkers could identify these high risk melanoma patients with the goal to treat this group more aggressively and to downsize follow up frequency in low risk melanoma patients.

### *Metastatic disease (AJCC stage IV)*

For stage IV melanoma treatment results remain unsatisfactory. No systemic therapy has demonstrated to affect overall survival, although the recent immunotherapy with Ipilimumab and the introduction of the BRAF pathway inhibitors have shown promising results.<sup>39,40</sup> Treatment of metastatic disease is mainly based on the individual situation of the patient. If disseminated melanoma disease presents itself as a single isolated metastasis, metastasectomy might be the only treatment.<sup>41,42</sup> The failed phase III Canvaxin trial enrolled 496 patients at 80 centers between June 1998 and March 2005. Canvaxin did as well as expected, but the placebo arm did better than predicted for surgery in stage IV melanoma. At 5 years, 40% of patients in both arms of the trial were still alive (unpublished data SSO conference 2011). All had received surgery for stage IV disease before entering the trial, which was testing postsurgical adjuvant use of the vaccine. Although the trial was not designed to test the efficacy of surgery in stage IV disease, the largest multicenter clinical trial undertaken using surgical resection as initial therapy for melanoma metastatic to distant sites prolonged survival and suggested that the more widespread use of surgery is indicated in stage IV melanoma. The role of surgery versus the 'best systemic treatment' for limited stage IV disease is now studied.<sup>43</sup> Radiotherapy can be indicated for local control and reduction of pain in metastatic disease.<sup>44</sup>

### *Follow-up*

The overall prevalence of melanoma is increasing as a result of the rising worldwide incidence of melanoma and the stabilizing mortality rates.<sup>45</sup> This has resulted in a growing number of melanoma patients needing follow-up and consequently more doctors are required to deliver this service. International consensus on the follow-up of melanoma patients does not exist and high-frequency follow-up after treatment for melanoma is still standard practice in many countries around the world.<sup>46</sup> The main purpose of follow-up is early detection of recurrent disease and second primary melanomas that could be treated successfully by surgery or other treatment modalities.<sup>47</sup> However, it is still difficult to predict the behavior of melanoma in the individual patient and overall treatment results of metastatic or recurrent disease are disappointing.<sup>47</sup> No evidence-based optimal follow-up schedule for melanoma patients exists, although most guidelines for management advise a more vigorous follow-up strategy for patients in more advanced stages. Future investigations should be focused on determining patient-tailored follow-up regimens. Therefore, in our centre a prospective, randomized, high quality methodological research has been started in order to develop meaningful applicable guidelines (MELFO).<sup>48</sup>

**TABLE 4** Follow-up recommendations per guideline for follow-up in stage I-III primary melanoma

Country	Year	Follow-up schedule	Number of visits per year after diagnosis					Routine additional testing Type and number per year				
			AJCC or T-class	1	2	3	4	5	6-10	USLN	CXR	PET/CT
Australia & New Zealand <sup>(10)</sup>	2008	I	2	2	2	2	2	1	-	-	-	-
		II	3-4	3-4	3-4	3-4	3-4	1	-	-	-	-
		III	3-4	3-4	3-4	3-4	3-4	1	-	-	-	-
Austria <sup>(11)</sup>	2010	Tis	1	1	1	1	1	1	-	-	-	-
		T1	2	2	2	2	2	2	1	-	-	-
		T2-3a	2-4	2-4	2-4	2-4	2-4	2-4	2	1-2	1-2	-
		T3b-4b	4	4	4	4	4	4	2	1-2	1-2	-
		III	4-12	4-12	4-12	4-12	4-12	4-12	4	4	4	-
Belgium <sup>(12)</sup>	2007	I	4	4	2	2	2	1	-	-	-	-
		II	4	4	2	2	2	1	-	-	-	-
		III	4	4	2	2	2	1	-	-	-	-
Canada (British Columbia) <sup>(13)</sup>	2009	I	2	2	1	1	1	1 <sup>a</sup>	-	-	-	-
		II	2-4	2-4	2	2	1	1 <sup>a</sup>	-	-	-	-
		III	2-4	2-4	2	2	1	1 <sup>a</sup>	-	-	-	-
Denmark <sup>(14)</sup>	2003	Ia	1	1	1	1	1	1	-	-	-	-
		Ib	4	4	2	2	2	1	-	-	-	-
		II	4	4	2	2	2	1	-	-	-	-
		III	4	4	2	2	2	1	-	-	-	-
France <sup>(15)</sup>	2005	I	2	2	2	2	2	1 <sup>a</sup>	-	-	-	-
		IIA-IIIB	4	4	4	4	4	1 <sup>a</sup>	2-4	-	-	-
		IIC-III	4	4	4	4	4	1 <sup>a</sup>	2-4	-	-	-
Germany <sup>(16)</sup>	2007	I (≤1mm)	2	2	2	2	2	1-2	-	-	-	-
		I-IB	4	4	4	4	4	1-2	2	-	-	2-4
		IIC-III	4	4	4	4	4	2	2-4	2	-	2-4
Ireland <sup>(17)</sup>	2006	I (≤1mm)	2	2	-	-	-	-	-	-	-	-
		I-II	2	2	2	2	2	-	-	-	-	-
		III	2	2	2	2	2	1 <sup>a</sup>	-	-	-	-
Italy <sup>(18)b</sup>	2009	O	1	1	1	1	1	1	-	-	-	-
		I	2	2	2	1	1	1	-	-	-	-
		IIAB-IIIA	3	3	2	2	1	1	1-3	1-2	-	-
		IIC-IIIB	3	3	3	1	1	1	1-3	1	1	-
		IIIC	3	3	3	2	2	1	2-3	2	0-3	2-3
Netherlands <sup>(19)</sup>	2005	Id	4	3	2	2	2	-	-	-	-	-
		II	4	3	2	2	2	1	-	-	-	-
		III	4	3	2	2	2	1	-	-	-	-
Norway <sup>(20)</sup>	2007	Tis	1	-	-	-	-	-	-	-	-	-
		<1 mm	4	4	4	-	-	-	-	-	-	-
		≥1 mm	4	4	4	2	2	-	-	-	-	-

General contents follow-up visits						Follow-up by which doctor?
History	Scar Exam	ITM Exam	LRLN	ComplSkin	Self-exam	
+	+	+	+	-	+	Doctor preferred by patient
+	+	+	+	-	-	Surgeon in specialized center
+	+	+	+	+	-	Dermatologist
+	+	+	+	+	-	Not specified
+	+	+	+	+	-	Yr 1-5: DMG Member Yr 5-10: GP
+	+	+	+	+	+	Not specified
+	+	+	+	+	-	Dermatologist
+	+	+	+	+	+	Pigmented skin lesion specialist
+	+	+	+	+	-	Surgeon or Dermato-logist
+	+	+	+	-	-	Not specified
+	+	+	+	+	-	Surgeon or Dermato-logist



TABLE 4 cont.

Country	Year	Follow-up schedule AJCC or T-class	Number of visits per year after diagnosis						Routine additional testing Type and number per year			
			1	2	3	4	5	6-10	USLN	CXR	PET/CT	S-100
Poland <sup>(21)</sup>	2009	All stages	3-4	3-4	2	2	2	1	-	-	-	-
Scotland <sup>(22)</sup>	2007	All stages	Frequency and duration individually based on tumor and patient characteristics.						-	-	-	-
South Africa <sup>(23)</sup>	2004	All stages	Frequency and duration individually based on tumor characteristics.						-	-	-	-
Spain <sup>(24)</sup>	2006	O	1	1	1	1	1	1 <sup>a</sup>	-	-	-	-
		IA	2	2	2	1	1	1 <sup>a</sup>	1	1	-	-
		IB-IIA	2-4	2-4	2-4	1-2	1-2	1 <sup>a</sup>	2	2	-	-
		IIB/C-III	2-4	2-4	2-4	1-3	1-3	1 <sup>a</sup>	-	-	1-2	-
Sweden <sup>(25)</sup>	2007	All stages	No follow-up schedule provided by national guideline. <sup>c</sup>						-	-	-	-
Switzerland <sup>(26)</sup>	2005	<1mm	2	2	2	2	2	1 <sup>a</sup>	-	-	-	-
		IB-IIB	4	4	4	2	2	1 <sup>a</sup>	1	1	-	-
		IIC-III	4	4	4	2	2	1/2 <sup>a</sup>	1	-	1	-
United Kingdom <sup>(27)</sup>	2002	<1 mm	4	4	4	-	-	-	-	-	-	-
		≥1 mm	4	4	4	2	2	-	-	-	-	-
United States of America <sup>(28)</sup>	2009	O	1	1	1	1	1	1 <sup>d</sup>	-	-	-	-
		IA-IIA	1-4	1-4	1-4	1-4	1-4	1 <sup>d</sup>	-	-	-	-
		IIB-IV	2-4	2-4	1-4	1-4	1-4	1 <sup>d</sup>	-	-	-	-

<sup>a</sup> Lifelong follow-up

<sup>b</sup> Personal communication from dr. C.R. Rossi, University of Padova, Italy

<sup>c</sup> No follow-up for Breslow <1 mm and melanoma in situ

<sup>d</sup> At least annual skin exam for life

Compl skin Complete skin examination

CXR Chest X-ray

## SUMMARY

The incidence of melanoma has increased drastically although mortality rates have stabilized which has resulted in a growing number of melanoma patients needing follow-up. Research should be continued to search for an adequate and optimal follow up regimen. Survival rates of stadium III and IV are still disappointing. Up to today, no systemic therapy has proven survival benefit. New promising treatment strategies with monoclonal antibodies and inhibitors are currently tested and new phase II and III trials designed.<sup>39,40</sup> Biomarkers should be evaluated to optimize the staging system as to strive to a more tailored systemic therapy regimen.

General contents follow-up visits						Follow-up by which doctor?
History	Scar Exam	ITM Exam	LRLN	ComplSkin	Self-exam	
+	+	+	+	+	-	Specialist in oncology center
+	+	+	+	+	+	Appropriate local specialist
+	+	+	+	+	-	Not specified
+	+	+	+	+	-	Dermatologist
+	+	+	+	+	+	Not specified
+	+	+	+	+	-	Not specified
+	+	+	+	+	-	Not specified
+	+	+	-	-	+	Not specified

ITM Exam Examination of in-transit metastasis pathway  
LRLN Palpation of loco-regional lymph nodes  
PET/CT Positron emission tomography with/without CT scanning  
USLN Ultrasonography of draining lymph node basin(s)  
DMG Danish Melanoma Group  
GP General practitioner

## OUTLINE OF THIS THESIS

Improvement in staging allows more precise classification and more accuracy of predicting the likely prognosis and outcomes for individuals with a disease. However, cutaneous melanoma still represents a paradox among all solid tumors. It is the cancer for which the best prognostic markers are available, yet there is very little understanding of their biological significance. It can be expected that the strongest biological markers, such as the ones that were included in the last AJCC staging system, are surrogates of key biological events. The studies in this thesis discuss the use of prognostic markers in the treatment of melanoma and try to create understanding about the correlation between melanoma biomarkers and tumor biology. Also, the potential use of various prognostic markers as LDH, S-100B, SUV and Breslow thickness in melanoma disease is evaluated.

**TABLE 5** Proposed optimal follow-up schedule and experimental schedule MELFO study <sup>(48)</sup>

AJCC Stage	Number of visits per year after diagnosis					
	1	2	3	4	5	6-10
IA	1					
IB	1	1	1	1	1	
IIA	2	2	1	1	1	1
IIB	3	3	2	1	1	1
IIC	3	3	2	1	1	1
III	4	3	2	2	2	1
IV	Individually tailored to patient's needs					

## PART I: DETECTION AND TREATMENT OF NODAL MELANOMA METASTASES

The overall prevalence of melanoma is increasing and therefore patients needing follow-up are increasing. International consensus on the follow-up of melanoma patients has not been achieved and high-frequency follow-up still practiced in many countries around the world. In chapter two the reader is informed about the influence of lymph nodes metastases detected by patients or by physicians at the outpatient department and the influence on survival. We evaluated the role of the method of detection in nodal disease in the prognosis of melanoma patients who underwent therapeutic lymph node dissection (TLND). Understanding the outcomes of self-detection or physician detection is of vital importance for the design of follow-up studies. Numerous studies have also demonstrated that older melanoma patients have a lower survival rate, especially those over 60 years of age.<sup>49-54</sup> The older patient might not detect positive lymph nodes because of a certain physical negligence that occurs with aging. Therefore we also evaluated the difference in self detection between elderly en younger patients and influences on survival.

Ilio-inguinal lymph node dissection for stage III melanoma is often complicated by wound healing disturbances. In the third chapter we conducted an extensive retrospective study to assess the present morbidity of therapeutic or completion lymph node dissection to determine risk factors for short term morbidity and to evaluate recommendations on peri-operative treatment. A retrospective study was performed to investigate the wound healing disturbances after therapeutic ilio-inguinal lymph node dissection.

## PART II : PROGNOSTIC MARKERS IN MELANOMA

When compared with other markers and tumor characteristics as ulceration, Clark level and Mitosis index, Breslow thickness is the most important biomarker and predictor for mortality.<sup>9,55</sup> Because of the increasing melanoma incidence in the last decades and predictions

showing a continuing increase in the next years the aim of the study described in the fourth chapter, was to assess trends in melanoma incidence, Breslow thickness (BT), and melanoma survival among young and elderly patients in the Netherlands. Patients diagnosed with melanoma in the Netherlands 1994-2008 were selected from the Netherlands Cancer Registry (NCR) and overall 40,880 patients were included and analyzed.

For stadium IV melanoma it was already known that the tumor marker S-100B in melanoma patients could be used for the detection of recurrences.<sup>56</sup> However, the role of S-100B for patients with palpable lymph node metastases before undergoing therapeutic lymph node dissection, was still unknown. In chapter five the role of S-100B for disease-free survival (DFS) was evaluated and compared with the tumor marker LDH.

In chapter six, S-100B values and Standardized Uptake Value (SUV) in FDG-PET for clinically stage III melanoma patients were analyzed as indicators of early recurrence in stage III disease. Aim was to assess the association between these markers and their relation with survival.

In chapter seven the effect of neoadjuvant treatment with bevacizumab in stage III melanoma is discussed. For patients with palpable nodes, a lack of resources in healthcare has resulted in a waiting list of at least 5 weeks before a lymph node dissection can be performed. A feasibility study was performed to investigate the effect of induction treatment with bevacizumab for Stage III melanoma, while waiting for lymph node dissection.

In chapter eight a review was composed describing the role of S-100B and how it can be used for the different tumor stages in melanoma.

A summary of the work undertaken is given in English and Dutch at the end of this thesis. Finally, in the 'future perspectives' section, new research developments in the field of melanoma markers are discussed.

## REFERENCES


1. Rinne D, Baum RP, Hor G et al. Primary staging and follow-up of high risk melanoma patients with whole-body 18F-fluorodeoxyglucose positron emission tomography: results of a prospective study of 100 patients. *Cancer* 1998;82:1664-71.
2. De Vries E, van de Poll-Franse LV, Louwman WJ, de Gruijl FR, Coebergh JW. Predictions of skin cancer incidence in the Netherlands up to 2015. *Br J Dermatol*. 2005;152:481-8.
3. Ulmer MJ, Tonita JM, Hull PR. Trends in invasive cutaneous melanoma in Saskatchewan 1970-1999. *J Cutan Med Surg*. 2003;7:433-42.
4. Venzon DJ, Moolgavkar SH. Cohort analysis of malignant melanoma in five countries. *Am J Epidemiol*. 1984;119:62-70.
5. Kricger A, Armstrong BK, Goumas C, et al. Personal sun exposure and risk of multiple primary melanomas. *Cancer Causes Control* 2007;18:295-304.
6. Armstrong BK, Kricger A. Cutaneous melanoma. *Cancer Surv*. 1994; 19-20:219-40.
7. Gaudette LA, Gao RN. Changing trends in melanoma incidence and mortality. *Health Rep*. 1998;10:29-41.
8. Giles GG, Armstrong BK, Burton RC, Staples MP, Thursfield VJ. Has mortality from melanoma stopped rising in Australia? Analysis of trends between 1931 and 1994. *BMJ* 1996;312:1121-5.
9. Balch CM, Greshenwald JE, Soong SJ, et al. Final version of 2009 AJC melanoma staging and classification. *J Clin Oncol* 2009;27:6199-206.
10. Australian Cancer Network Melanoma Guidelines Revision Working Party Clinical Practice Guidelines for the Management of Melanoma in Australia and New Zealand.
11. Österreichische Gesellschaft für Chirurgische Onkologie. ACO-Manual, Malignes Melanom der Haut.
12. College voor Oncologie. Nationale Richtlijnen voor de Klinische Praktijk, Melanoma Versie 1.2007.
13. British Columbia Cancer Agency, Cancer Management Guidelines, Melanoma, (2008).
14. Dansk Melanom Gruppe. Protokoller: klini og pathologi.
15. Négrier S, Saiag P, Guillot B, and et al. Recommendations pour la Pratique Clinique: Standards, Options et Recommendations 2005 pour la prise en charge des patients adultes atteints d'un mélanome cutané MO. *Ann Dermatol Venereol* 2005.
16. Garbe C, Hauschild A, Volkenandt M, and et al. Evidence and interdisciplinary consensus-based German guidelines: diagnosis and surveillance of melanoma. *Melanoma Res* 2007;17:393-9.
17. Royal College of Surgeons in Ireland, Guidelines Committee. Management of Cutaneous Melanoma, Clinical Guidelines.
18. Consiglio Nazionale dello Ricerche. Basi scientifiche per linee guida, il melanoma cutaneo.
19. Kwaliteitsinstituut voor de Gezondheidszorg CBO. Richtlijn Melanoom van de huid (2005).
20. Norsk Melanom Gruppe. Veileider 2007.
21. Ruka W, Krzakowski M, Placek W, and et al. Czerniaki skóry - zasady postępowania diagnostyczno-terapeutycznego. Cutaneous melanoma - diagnostic and therapeutic guidelines. *Onkologia w praktyce klinicznej*5, 20-32 (2009)
22. Scottish Intercollegiate Guidelines Network (SIGN). Guideline 72: Cutaneous melanoma, a national clinical guideline (2003).
23. Whitaker DK and Sinclair W. Melanoma Advisory Board. Guideline on the management of melanoma. *S Afr Med J*94, 697-708 (2004).
24. Generalitat Valenciana, Consellaria de Sanitat. Guía de Prevención y Tratamiento del Melanoma. (2006).
25. Svenska Melanomstudiegruppen. Nationellt Vardprogram Malignt hudmelanom, (2007).
26. Dummer R, Meier S, Beyeler M, and et al. Die aktualisierten schweizerischen Richtlinien zur Behandlung des kutanen Melanoms – wo stehen wir heute? *Schweiz Med Forum* 2005;6:196-201.
27. Roberts DLL, Anstey AV, Barlow RJ, and et al. U.K. guidelines for the management of cutaneous melanoma. *Br J Dermatol* 2002;146, 7-17.
28. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology, Melanoma V.I. 2010.
29. MJ Speijers, AB Francken, JEHM Hoekstra-Weebers, E Bastiaannet, S Kruijff, HJ Hoekstra. Optimal Follow-up for Melanoma. *Expert Rev Dermatol*. 2010;5:461-478.


30. Lens MB, Dawes M, Goodacre T et al. Excision margins in the treatment of primary cutaneous melanoma: a systematic review of randomized controlled trials comparing narrow vs wide excision. *Arch Surg* 2002; 137:1101-1105.
31. Morton DL, Cochran AJ, Thompson JF, et al. Sentinel node biopsy for early-stage melanoma: accuracy and morbidity in MSLT-I, an international multicenter trial. *Ann Surg.* 2005;242:302-11.
32. Morton DL, Thompson JE, Cochran AJ, et al. Sentinel-node biopsy or nodal observation in melanoma. *N Engl J Med.* 2006;355:1307-17.
33. Morton DL, Thompson JE, Cochran AJ, et al. Sentinel node biopsy and immediate lymphadenectomy for occult metastases versus nodal observation and delayed lymphadenectomy for nodal recurrence. *Ann Surg Oncol* 2010, 17 Suppl 1, S22-S23 (Abstract 53).
34. Balch CM, Gershenwald JE, Soong S, et al. Multivariate Analysis of Prognostic Factors Among 2,313 Patients With Stage III Melanoma: Comparison of Nodal Micrometastases Versus Macrometastases. *J Clin Oncol.* 2010;28:2452-9.
35. Bastiaannet E, Wobbes T, Hoekstra OS, et al. Prospective comparison of [18F] fluorodeoxyglucose positron emission tomography and computed tomography in patients with melanoma with palpable lymph node metastases: diagnostic accuracy and impact on treatment. *J Clin Oncol* 2009;27:4774-80.
36. Mijnhout GS, Hoekstra OS, van Tulder MW, Teule GJ, Deville WL. Systematic review of the diagnostic accuracy of (18)F-fluorodeoxyglucose positron emission tomography in melanoma patients. *Cancer* 2001;91:1530-42.
37. Aukema TS, Valdés Olmos RA, Wouters MW, et al. Utility of Preoperative 18F-FDG PET/CT and Brain MRI in Melanoma Patients with Palpable Lymph Node Metastases. *Ann Surg oncol* 2010;17:2773-8.
38. Kruijff S, Bastiaannet E, Muller-Kobold AC, et al. S-100B concentrations predict disease-free survival in stage III melanoma patients. *Ann Surg Oncol* 2009;16:3455-62.
39. Hodi FS, O'Day SJ, McDermott DF, et al. Improved survival with ipilimumab in patients with metastatic melanoma. *New Engl J Med* 2010; 363:711-23.
40. Flaherty KT, Puzanov I, Kim KB, et al. Inhibition of mutated, activated BRAF in metastatic melanoma. *N Engl J Med.* 2010;363:809-19.
41. Ollila DW. Complete metastasectomy in patients with stage IV metastatic melanoma. *Lancet Oncol* 2006;7:919-924.
42. Andrews S, Robinson L, Cantor A et al. Survival after surgical resection of isolated pulmonary metastases from malignant melanoma. *Cancer Control* 2006; 13:218-223.
43. <http://clinicaltrials.gov/show/NCT01013623>.
44. Schmidt-Ullrich RK, Johnson CR. Role of radiotherapy and hyperthermia in the management of malignant melanoma. *Semin Surg Oncol* 1996; 12:407-415
45. Karim-Kos HE, de Vries E, Soerjomataram I, Lemmens V, Siesling S, Coebergh JW. Recent trends of cancer in Europe: a combined approach of incidence, survival and mortality for 17 cancer sites since the 1990s. *Eur. J. Cancer* 2008;44:1345-1389.
46. Francken AB, Bastiaannet E, Hoekstra HJ. Follow-up in patients with localized primary cutaneous melanoma. *Lancet Oncol.*2005;6: 608-621.
47. Brada M. Is there a need to follow-up cancer patients? *Eur. J. Cancer* 1995;5, 655-657.
48. <http://dissertations.ub.rug.nl/FILES/faculties/medicine/2007/a.b.francken/melfo.pdf>.
49. Austin PF, Cruse CW, Lyman G, Schroer K, Glass F, Reintgen DS. Age as a prognostic factor in the malignant melanoma population. *Ann Surg Oncol.* 1994;1:487-94.
50. Averbook BJ, Russo LJ, Mansour EG. A long-term analysis of 620 patients with malignant melanoma at a major referral center. *Surgery.* 1998;124:746-55.
51. Balch CM, Soong SJ, Bartolucci AA, et al. Efficacy of an elective regional lymph node dissection of 1 to 4 mm thick melanomas for patients 60 years of age and younger. *Ann Surg.* 1996;224:255-63.
52. Cohen HJ, Cox E, Manton K, Woodbury M. Malignant melanoma in the elderly. *J Clin Oncol.* 1987;5:100-6.
53. Loggie B, Ronan SG, Bean J, Das Gupta TK. Invasive cutaneous melanoma in elderly patients. *Arch Dermatol.* 1991;127:1188-93.

54. Sahin S, Rao B, Kopf AW, et al. Predicting ten-year survival of patients with primary cutaneous melanoma: corroboration of a prognostic model. *Cancer*.1997;80:1426–31.
55. Francken AB, Shaw HM, Thompson JE, et al. The prognostic importance of tumor mitotic rate confirmed in 1317 patients with primary cutaneous melanoma and long follow-up. *Ann Surg.Oncol* 2004;11:426-33.
56. Smit LH, Korse CM, Hart AA, et al. Normal values of serum S-100B predict prolonged survival for stage IV melanoma patients. *Eur J Cancer* 2005; 41:386-392.









# Part I

Detection and treatment of  
nodal melanoma metastases

# 2

---





# Detection of melanoma nodal metastases; differences in detection between elderly and younger patients do not affect survival

| S Kruijff, E Bastiaannet, AJH Suurmeijer, HJ Hoekstra | Ann Surg

Oncol. 2010;17:3008-14

---

## ABSTRACT

**Background:** Melanoma lymph nodes metastases may be detected by patients or by physicians. Understanding the outcomes of self- or physician detection is essential for the design of follow-up studies. We evaluated the role of the method of detection in nodal disease in the prognosis of melanoma patients who underwent therapeutic lymph node dissection (TLND).

**Methods:** All melanoma patients with palpable lymph nodes were included in a prospective database (n=98) and the method of detection was recorded. Detection of lymph node metastases compared with pathological findings in the TLND was assessed by multivariate logistic regression. Disease free survival (DFS) and disease specific survival (DSS) were assessed by univariate and multivariate Cox proportional hazard analysis.

**Results:** Nodal metastases were detected by physicians in 45% and by patients in 55% (p<0.001). Age was significantly associated with method of detection. Patients ≤ 60 years detected 69% their lymph node metastases as opposed to 32% of patients > 60 years (OR 0.3; p= 0.007). However, this was not associated with prognostic findings in TLND, number of positive nodes, tumor size, or extranodal spread. Method of detection or age at the time of nodal metastases was not significantly associated with two year DFS or DSS.

**Conclusion:** 45% of lymph node metastases in Stage I-II melanoma patients are physician detected. Younger patients detect their own lymph node metastases significantly more often than elderly patients. However, neither the method of detection nor age correlates with DSS. More frequent follow-up would not alter DFS and DSS significantly.

## INTRODUCTION

The incidence of melanoma is increasing worldwide. In the Netherlands, the incidence increased from 9.5 to 13.7 per 100,000 in men and from 13.4 to 18.5 per 100,000 in women between 1989-2003. Approximately 90% of patients have stage I or II melanoma at diagnosis and the incidence of patients with a thin melanoma has increased.<sup>1-5</sup> Although the early diagnosis of cutaneous melanoma with small Breslow's tumour thickness has been responsible for the levelling off of overall melanoma mortality, the incidence at the same time has continued to rise.<sup>6</sup> The number of melanoma patients requiring follow-up surveillance has doubled in the Netherlands.<sup>1</sup> Despite improved public awareness and earlier diagnosis and treatment, mortality from melanoma increased from 2.5 to 3.6/100,000 in men and from 2.0 to 2.4/100,000 in women in the Netherlands between 1989-2003.<sup>1</sup>

Several studies have been performed concerning survival in patient versus physician detected recurrences. Since most of first melanoma recurrences are detected by patients or their partners, the value of high frequency follow-up has often been questioned.<sup>7,8</sup> Over half of patients (55%) with early stage melanoma detect their own recurrence(s).<sup>9-11,14,16-23</sup> In recent decades, several follow-up schedules have been suggested with large variations and without international consensus.<sup>3</sup> Proposals for these schedules are based on risk calculations concerning melanoma recurrence and cost effectiveness. Several authors have recommended a reduction in the intensity of follow-up regimens based on these factors.<sup>9-12</sup> Others claim that a more intensive follow-up will improve outcome because of earlier detection.<sup>13-15</sup>

Of the various follow-up methods performed by physicians, only medical history and physical examination seem to be cost effective.<sup>23</sup> Despite these findings, most melanoma patients are still followed up on a frequent basis, mostly for education, reassurance or inclusion in clinical trials.

Studies concerning the method of nodal metastasis detection have not shown a difference in survival comparing physician- versus patient detected recurrences. Intense follow-up does not seem to contribute to disease free interval or overall survival, as physician detected lesions do not have a better prognosis than those detected by patients themselves.<sup>3,19</sup>

Patients with clinical stage III b melanoma (palpable lymph node metastases) have a 5-year survival of 59%.<sup>24</sup> A better understanding of the method of detection of palpable melanoma metastases (i.e. by patient or physician) and its influence on survival is essential to resolve the above mentioned follow up controversy.

The aim of this study was to evaluate the role of the method of detection, by patient or physician, in melanoma patients with palpable lymph nodes and to analyse the association with pathological findings in therapeutic lymph node dissection (TLND) and the impact of the detection method on disease free and disease specific survival. The method of detection of positive nodes (patient versus physician) might not only have impact on survival but might also influence the number of positive nodes, the tumor size or the presence of extranodal growth in the TLND.

## MATERIAL AND METHODS

Patients with primary melanoma were treated with wide local resection (1 or 2 cm) with or without sentinel lymph node biopsy and, if indicated, therapeutic lymph node dissection. Stage I and II melanoma patients were followed after treatment in accordance with the Dutch National Guideline Treatment of Melanoma ([www.oncoline.nl](http://www.oncoline.nl)). Postoperative follow-up included physical examination every 3 months for the first year, every 4 months for the second year, and every 6 months thereafter. Standard radiographic or serum investigations were not performed.

All patients with clinically and cytological proven lymph node metastases of melanoma (AJCC stage III b) were included in this study. If palpable positive nodes were found during regular follow-up this was recorded as “physician detected nodal metastases”. When positive palpable nodes were recorded after a patient-initiated visit, this was recorded as “patient detected nodal metastases”. Patients without distant metastases on FDG-PET and CT received a TLND with curative intent and were entered into a single prospective institutional database. The Stage IV upstaged patients were referred for further palliative treatment to their primary physician, medical oncologist or radiation oncologist and excluded for further follow up. Patients, who presented with local recurrence or in transit disease, were not included.

Patients with nodal metastases of  $\geq 3$  cm and/or  $\geq 3$  positive lymph nodes and/or extra nodal (EN) disease received adjuvant radiotherapy (20 x 2.4 Gy). Data examined include patient demographics, clinical and histopathologic characteristics of the tumor, date and type of operation, site and date of first (nodal) and second recurrence, method of detection of recurrence (i.e. patient or physician during standard follow-up), and status at last follow-up.

The study was approved by the Medical Ethical Committee of the University Medical Centre Groningen (UMCG).

### *Statistics*

For retrospective analysis, patients were divided by age (less than or equal to 60 versus greater than 60 years of age). Factors associated with method of detection were analysed by univariate and multivariate logistic regression analysis. Four subgroups were generated based on the combinations of young versus elderly and physician versus patient detection to assess for associations with pathological variables. Factors associated with disease free survival (DFS) and disease specific survival (DSS) were assessed by univariate and multivariate Cox proportional hazard analysis. For DFS, any recurrence was recorded as an event; for DSS, death due to melanoma was considered an event. Follow-up was truncated at 4 years.

## RESULTS

### *Patients*

In the period 2003-2008, 98 patients 54 males (55.1%) and 44 females (44.9%), median age 57.7 years (range 28.5-86.7) with stage III b melanoma, based on PET and spiral CT, underwent a TLND (Table 1). Median time from primary melanoma treatment to nodal recurrence or nodal recurrence of an unknown primary melanoma was 22.0 (0-315.3) months.

**TABLE 1** Patient and tumour characteristics of 98 patients with clinically and cytological proven lymph node metastases of melanoma (AJCC stage IIIb)

		Number	Percentage
Sex	Male	54	55.1
	Female	44	44.9
Age (years)	≤60	61	62.2
	>60	37	37.8
<b>Primary melanoma</b>			
Localisation	Upper extremities	11	11.2
	Head & Neck	8	8.2
	Lower extremities	39	39.8
	Trunk	34	34.7
	Unknown primary	6	6.1
Breslow thickness	≤2.0	43	43.9
	>2.0	49	50.0
	Unknown	6	6.1
Clark level	I-III	33	33.7
	IV-VI	59	60.2
	Unknown	6	6.1
Ulceration	Yes	22	22.5
	No	76	77.5
<b>Lymph node metastases</b>			
Lymph node dissection	Axilla	34	34.7
	Groin	50	51.0
	Neck	14	14.3
Number of nodes removed	<15	48	49.0
	15 or more	50	51.0
Nr of nodes positive	2 or less	53	54.1
	>2	45	45.9
Tumorsize (cm)	≤ 3.0	62	63.3
	>3.0	36	36.7
Extranodal growth	Yes	27	27.5
	No	71	72.5

## Detection

Physicians detected 45% of the nodal metastases and patients detected 55% ( $p=0.001$ ) (Table 2). Of the 61 patients in the younger age group ( $\leq 60$  years), 19 nodal metastases were physician detected (31%) whereas 42 nodal metastases were patient detected (69%). Of the 37 elderly patients ( $> 60$  years), 12 nodal metastases were patient detected (32%) and 25 nodal metastases physician detected (68%) ( $p=0.001$ ).

Table 2 also shows the results of univariate and multivariate logistic regression analysis of the association between method of detection and pathological information from the lymph node dissection, sex, age, and nodal bearing area (axilla, groin, neck). Age was significantly associated with method of detection: 69% of patients  $\leq 60$  yrs detected their nodal metastases versus 32% of the patients  $>60$  yrs (OR 0.3;  $p=0.007$  in multivariate analysis).

**TABLE 2** Detection of lymph node metastases in association with pathological results of lymph node dissection

		% patient-detected	Univariate analysis		Multivariate analysis	
			OR (95%CI)	p-value	OR (95%CI)	p-value
Overall: 44.9% detected by physician and 55.1% by patients ( $p<0.001$ )						
Sex	Male	48.2	1.0		1.0	
	Female	63.6	1.9 (0.8-4.3)	0.13	1.8 (0.7-4.6)	0.20
Age	$\leq 60$	68.9	1.0		1.0	
	$>60$	32.4	0.2	0.001	0.3 (0.1-0.7)	0.007
Breslow thickness	$\leq 2.0$	67.4	1.0		1.0	
	$>2.0$	42.9	0.4 (0.2-0.8)		0.5 (0.2-1.3)	
	Unknown primary	66.7	0.9 (0.2-5.9)	0.05	2.1 (0.3-15)	0.22
Lymph node region	Axilla	61.8	1.0			
	Groin	50.0	0.6 (0.3-1.5)			
	Neck	57.1	0.8 (0.2-2.9)	0.56		
Nodes removed (cat)*	$<15$	58.3	1.0			
	$>15$	52.0	0.7 (0.3-1.7)	0.53		
Nodes positive*	2 or less	47.2	1.0		1.0	
	$>2$	64.4	2.0 (0.9-4.6)	0.09	1.9 (0.7-4.9)	0.17
Tumorsize (cm)	cont		1.2 (0.9-1.6)	0.09	1.2 (0.9-1.6)	0.17
Extranodal growth	Yes	66.7	1.0		1.0	
	No	50.7	0.5 (0.2-1.3)	0.16	0.7 (0.2-2.0)	0.48

Patients with a thin primary melanoma had a higher ratio of patient-detected positive lymph nodes ( $p=0.05$ ). This was, however, not statistically significant in multivariate analysis.

Sex, regional nodal basin site, number of nodes removed, number of positive nodes, tumor size and extranodal growth were not associated with method of detection in multivariate analysis.

There were no significant differences in pathological findings when the groups were divided in four groups: patient detected younger age ( $\leq 60$  yrs), patient detected elderly age ( $>60$  yrs), physician detected younger age ( $\leq 60$  yrs), and physician detected elderly age ( $>60$  yrs).




### Survival

Multivariate analysis showed no significant difference in 2-year DFS for physician versus patient detected nodal recurrences (Table 3). Univariate analyses showed that DFS was associated with the lymph node bearing area, the number of positive nodes, node size, and extranodal growth pattern.


The 2-year DSS in males (53%) was significantly shorter than that in females (75%) (HR 0.3;  $p=0.004$ ). DSS was significantly reduced in patients with  $\geq 2$  positive nodes (51%) compared to patients with  $< 2$  positive nodes (72%) (HR 2.2;  $p=0.03$ ). Multivariate analysis showed no significant differences for DSS related to age, Breslow thickness, ulceration, lymph node bearing area (axilla, groin, neck), number of removed nodes ( $\leq 15$  nodes or  $>15$  nodes), tumor size ( $\leq 3\text{cm}$  or  $>3\text{cm}$ ), or extranodal growth (Table 3). Although in univariate analysis a trend in DSS was noted ( $p=0.08$ ) in favour of the physician detected group, in multivariate analysis, when compared to detection by patients, no significant difference could be found.

**TABLE 3** Disease Specific Survival (DSS) and Disease Free survival (DFS) in melanoma patients AJCC stage IIIB in correlation to detection by physician versus detection by patient

		DSS	HR	p-value	HR	p-value	DFS	HR	p-value	HR	p-value
Sex	Male	53.2	1.0	0.02	1.0	0.004	35.2	1.0	0.62		
	Female	75.4	0.4		0.3		36.8	0.9			
Age	$\leq 60$	59.7	1.0	0.28			29.7	1.0	0.17		
	$>60$	66.6	0.7				45.2	0.7			
Breslow	$\leq 2.0$	65.8	1.0	0.81			45.4	1.0	0.83		
	$>2.0$	64.5	0.9				30.4	1.06			
Ulceration	Yes	61.5	1.0	0.61			22.6	1.0	0.21		
	No	62.8	0.8				40.5				
Loc lymph	Axilla	57.2	1.0	0.37			38.8	1.0	0.07	1.0	0.2
	Groin	60.5	0.8				26.0	1.3		1.2	
	Neck	78.6	0.5				66.7	0.4		0.4	
Removed	$<15$	63.2	1.0	0.9			39.2	1.0	0.78		
	15 or more	61.8	1.1				32.9	1.1			
Positive	2 or less	72.5	1.0	0.02	1.0	0.03	46.7	1.0	0.009	1.0	0.1
	$>2$	51.4	2.2		2.2		23.8	2.0		1.6	
Tumorsize	$\leq 3.0$	69.6	1.0	0.06	1.0	0.39	45.3	1.0	0.01	1.0	0.2
	$>3.0$	47.9	1.9		1.4		17.3	2.0		1.5	
Extranod	Yes	56.8	1.0	0.52			27.6	1.0	0.03	1.0	0.2
	No	64.8	0.8				39.4	0.6		0.7	
Detection	Physician	67.6	1.0	0.08	1.0	0.14	36.2	1.0	0.2	1.0	0.5
	Patient	58.4	1.8		1.7		36.6	1.4		1.2	



**FIGURE 1** KM curve DSS melanoma patients with positive lymph nodes detected by physician versus detected by patient



**FIGURE 2** KM-curve DFS melanoma patients with positive lymph nodes detected by physician versus detected by patient

## DISCUSSION

The most efficacious follow-up scheme for melanoma patients is uncertain. Understanding the impact of the method of detection of nodal recurrence on prognosis is important in determining the most appropriate follow-up. Despite controversy about follow-up in melanoma, our results show that nodal metastases are still detected by physicians in almost 45% of the patients. Young patients detect their own lymph node metastases significantly more often than older patients (68% versus 32%) ( $p=0.007$ ). However, in patients undergoing close clinical surveillance neither the method of detection (patient versus physician) nor age seems to have a significant influence on two-year DSS or DFS. Furthermore, no correlation could be found between the methods of detection and the number of positive nodes, lymph node size, or extra nodal growth.

Despite this encouraging percentage of almost 45% detected nodes by physicians, this rate does not seem to result in any survival advantage. Earlier follow up studies describe an increasing controversy about the efficacy of an intensive follow-up scheme for melanoma patients as is still organised in a lot of hospitals today.

This is the first study evaluating the method of detection of nodal recurrence in stage III b melanoma and the impact of patient or physician detected nodal metastases on survival.

Patients with clinical stage III b melanoma (palpable lymph node metastases) have a 5-year survival of 59%.<sup>24</sup> The most well recognized causes for this low survival rate is the variation in the number of positive nodes, tumor size, and presence of extranodal growth.

Another cause for low survival rates in this patient category might be under staging. Bastiaannet et al. revealed that staging with FDG-PET of clinically stage III b patients resulted in upstaging of 22% to stage IV. In 19% treatment was changed, usually from surgery to systemic treatment.<sup>26</sup> Patients in our current study were all staged with FDG-PET and CT and therefore can be considered as “true” AJCC stage III b patients.

Survival might also be influenced by follow up and method of detection. We expected physician detected positive lymph node status to correlate with a lower number of positive nodes, smaller tumor size, and absence of extranodal growth in the TLND. This association was not found despite a positive trend in DSS ( $p= 0.08$ ) in the advantage of physician detected nodes (Table 3).

Follow-up of melanoma has different purposes: to detect first melanoma recurrence, to assess treatment efficacy, and to detect a second primary melanoma (2-6%).<sup>28-30</sup> Also of great importance are patient reassurance and documentation or inclusion for clinical trials. Follow-up schemes are often of high intensity; however, resources in many centres are insufficient to deal with the increasing patient load. Therefore, identifying and integrating evidence concerning the efficacy of follow-up strategies is important.

In 2003 Garbe et al performed a prospective follow-up study in melanoma patients following an intensive follow-up scheme (every 3 months for the first five years and every six months during the sixth to tenth years).<sup>14</sup> The authors claimed that 83% of all recurrences were detected by a physical examination during regular follow-up and only 17% were detected by patients

themselves. However, various AJCC melanoma stages were included in this study. Francken et al performed a large retrospective study and found that only 62% of all types of recurrences were identified by patients themselves; however, three quarters of first melanoma recurrences (FMR) were detected by patients or their partners, of which 11% were due to self examination.<sup>3</sup> In the present study no significant survival difference was found comparing patient - versus physician-detected nodal metastasis.

Additionally, patient age did not seem to have a significant influence on two-year DSS or DFS. In patients older than 60 years, almost 70% of the detected nodes were found by a physician; in contrast, patients at or below 60 years of age detected their own nodes 70% of the time. Earlier literature reports age as an independent prognostic factor, perhaps presenting a surrogate for declining host defence mechanism associated with advancing age.<sup>2</sup> Numerous studies have demonstrated that older patients have a lower survival rate, especially those over 60 years of age.<sup>31-36</sup> The older patient might not detect positive lymph nodes because of a certain physical negligence that occurs with aging. However, our results surprisingly did not identify any survival disadvantage associated with delayed detection of nodal metastases. This might be explained by a small patient population or a follow-up period that is too short. Another explanation could be related to the recent report by Conway et al, who found that lymphatic function declines with age.<sup>37</sup> Hypothetically, in older patients, declining lymphatic function might modify metastatic patterns and slow the process of dissemination.

Overall, it can be concluded that no study to date has proven any disease free survival or overall survival benefit related to intense follow-up surveillance as there are no prospective studies that compare high frequency to no or low frequency follow up.

With the frequency of follow-up which was used in this study, the DFS and DSS for patients capable themselves of detecting a palpable node in the regional nodal basin and those unable to do so, were not significantly different. Otherwise stated, it could be concluded that the above frequency of follow-up is sufficiently close for the detection and treatment of nodal disease. More frequent follow-up probably would not alter the DFS and DSS significantly. Therefore, arguments for more frequent follow-up than the above described scheme are hard to justify as physician detected lesions do not have a better prognosis than those detected by patients.<sup>3,7,8</sup> Whether follow-up with sparser intervals could be designed in a responsible way cannot be answered by this study.

Prospective, randomized, high quality methodological research is required in order to develop meaningful applicable guidelines. Currently no international consensus has been reached concerning the optimal frequency of follow-up visits for melanoma patients. Follow-up should be based on individual patient characteristics; multiple patient factors should be used to design the most appropriate follow-up.<sup>39</sup> A clinical randomized trial (MELFO) is currently underway at the University Medical Centre Groningen to evaluate the safety and cost effectiveness of reduced follow-up surveillance.<sup>40</sup> In this RCT a high frequent follow up scheme will be compared with a less frequent and better differentiated follow up scheme adjusted to melanoma AJCC stage.

In conclusion, more than 55% of lymph node metastases in melanoma are detected by the patient. Younger patients detect their own lymph node metastases significantly more often than elderly patients without any impact on DSS and DFS. The data of our study will add to the controversy about the value of high frequency follow-up regimens in melanoma. A prospective, randomized, high quality methodological research has been started in order to develop meaningful applicable guidelines (MELFO).

## REFERENCES

1. [Http://www.ikcnet.nl](http://www.ikcnet.nl)
2. Balch CM, Buzaid AC, Soong SJ et al. Final version of the American Joint Committee on Cancer staging system for cutaneous melanoma. *J Clin Oncol* 2001; 19:3635-3648.
3. Francken AB, Bastiaannet E, Hoekstra HJ. Follow-up in patients with localised primary cutaneous melanoma. *Lancet Oncol* 2005; 6:608-621.
4. De Vries E, Bray FI, Coebergh JW et al. Changing epidemiology of malignant cutaneous melanoma in Europe 1953-1997: rising trends in incidence and mortality but recent stabilizations in western Europe and decreases in Scandinavia. *Int J Cancer* 2003; 107:119-126.
5. Goldsmith LA, Askin FB, Chang AE et al. Diagnosis and Treatment of Early Melanoma: NIH Consensus Development Panel on Early Melanoma. *JAMA* 1992; 268: 1314-19
6. Garbe C, Eigentler TK. Diagnosis and treatment of cutaneous melanoma: state of the art 2006. *Melanoma Res* 2007; 17:117-127.
7. Francken AB, Shaw HM, Accortt NA et al. Detection of first relapse in cutaneous melanoma patients: implications for the formulation of evidence-based follow-up guidelines. *Ann Surg Oncol* 2007; 14:1924-1933.
8. Francken AB, Thompson JF, Bastiaannet E et al. [Detection of the first recurrence in patients with melanoma: three quarters by the patient, one quarter during outpatient follow-up]. *Ned Tijdschr Geneesk* 2008; 152:557-562.
9. Hofmann U, Szedlak M, Rittgen W et al. Primary staging and follow-up in melanoma patients- monocenter evaluation of methods, costs and patient survival. *Br J Cancer* 2002; 87:151-157.
10. Mooney MM, Kulas M, McKinley B et al. Impact on survival by method of recurrence detection in stage I and II cutaneous melanoma. *Ann Surg Oncol* 1998; 5:54-63.
11. Shumate CR, Urist MM, Maddox WA. Melanoma recurrence surveillance. Patient or physician based? *Ann Surg* 1995; 221:566-569.
12. Moloney DM, Gordon DJ, Briggs JC et al. Recurrence of thin melanoma: how effective is follow-up? *Br J Plast Surg* 1996; 49:409-413.
13. Kittler H, Weitzdorfer R, Pehamberger H et al. Compliance with follow-up and prognosis among patients with thin melanomas. *Eur J Cancer* 2001; 37:1504-1509.
14. Garbe C, Paul A, Kohler-Spath H et al. Prospective evaluation of a follow-up schedule in cutaneous melanoma patients: recommendations for an effective follow-up strategy. *J Clin Oncol* 2003; 21:520-529.
15. Sylaidis P, Gordon D, Rigby H et al. Follow-up requirements for thick cutaneous melanoma. *Br J Plast Surg* 1997; 50:349-353.
16. Weiss M, Loprinzi CL, Creagan ET et al. Utility of follow-up tests for detecting recurrent disease in patients with malignant melanomas. *JAMA* 1995; 274:1703-1705.
17. Regan MW, Reid CD, Griffiths RW et al. Malignant melanoma, evaluation of clinical follow up by questionnaire survey. *Br J Plast Surg* 1985; 38:11-14.
18. Poo-Hwu WJ, Ariyan S, Lamb L et al. Follow-up recommendations for patients with American Joint Committee on Cancer Stages I-III malignant melanoma. *Cancer* 1999; 86:2252-2258.
19. Moore DK, Zhou Q, Panageas KS et al. Methods of detection of first recurrence in patients with stage I/II primary cutaneous melanoma after sentinel lymph node biopsy. *Ann Surg Oncol* 2008; 15:2206-2214.
20. Kersey PA, Iscoe NA, Gapski JA et al. The value of staging and serial follow-up investigations in patients with completely resected, primary, cutaneous malignant melanoma. *Br J Surg* 1985; 72:614-617.
21. Dicker TJ, Kavanagh GM, Herd RM et al. A rational approach to melanoma follow-up in patients with primary cutaneous melanoma. *Scottish Melanoma Group. Br J Dermatol* 1999; 140:249-254.
22. Baughan CA, Hall VL, Leppard BJ et al. Follow-up in stage I cutaneous malignant melanoma: an audit. *Clin Oncol (R Coll Radiol)* 1993; 5:174-180.
23. Basseres N, Grob JJ, Richard MA et al. Cost-effectiveness of surveillance of stage I melanoma. A retrospective appraisal based on a 10-year experience in a dermatology department in France. *Dermatology* 1995; 191:199-203.
24. Balch CM et al ; Final Version of 2009 AJCC Melanoma Staging and Classification . *Journal of Clinical Oncology* 2009; 27 (36) 6199


25. Balch CM, Soong SJ, Gershenwald JE et al. Prognostic factors analysis of 17,600 melanoma patients: validation of the American Joint Committee on Cancer melanoma staging system. *J Clin Oncol* 2001; 19:3622-3634.
26. Bastiaannet E, Oyen WJ, Meijer S et al. Impact of [18F] fluorodeoxyglucose positron emission tomography on surgical management of melanoma patients. *Br J Surg* 2006; 93:243-249.
27. Kruijff S, Bastiaannet E, Muller Kobold AC et al. S-100B Concentrations Predict Disease-Free Survival in Stage III Melanoma Patients. *Ann Surg Oncol* 2009;16:3455-62.
28. Ariyan S, Poo WJ, Bologna J et al. Multiple primary melanomas: data and significance. *Plast Reconstr Surg* 1995; 96:1384-1389.
29. Brobeil A, Rapaport D, Wells K et al. Multiple primary melanomas: implications for screening and follow-up programs for melanoma. *Ann Surg Oncol* 1997; 4:19-23.
30. Johnson TM, Hamilton T, Lowe L. Multiple primary melanomas. *J Am Acad Dermatol* 1998; 39:422-427.
31. Austin PF, Cruse CW, Lyman G et al. Age as a prognostic factor in the malignant melanoma population. *Ann Surg Oncol* 1994; 1:487-494.
32. Averbook BJ, Russo LJ, Mansour EG. A long-term analysis of 620 patients with malignant melanoma at a major referral center. *Surgery* 1998; 124:746-755.
33. Balch CM, Soong SJ, Bartolucci AA et al. Efficacy of an elective regional lymph node dissection of 1 to 4 mm thick melanomas for patients 60 years of age and younger. *Ann Surg* 1996; 224:255-263.
34. Cohen HJ, Cox E, Manton K et al. Malignant melanoma in the elderly. *J Clin Oncol* 1987; 5:100-106.
35. Loggie B, Ronan SG, Bean J et al. Invasive cutaneous melanoma in elderly patients. *Arch Dermatol* 1991; 127:1188-1193.
36. Sahin S, Rao B, Kopf AW et al. Predicting ten-year survival of patients with primary cutaneous melanoma: corroboration of a prognostic model. *Cancer* 1997; 80:1426-1431.
37. Conway WC, Faries MB, Nicholl MB et al. Age-related lymphatic dysfunction in melanoma patients. *Ann Surg Oncol* 2009; 16:1548-1552.
38. Nieweg OE, Kroon BB. The conundrum of follow-up: should it be abandoned? *Surg Oncol Clin N Am* 2006; 15:319-330.
39. Francken AB, Hoekstra HJ. Follow-up of melanoma patients: the need for evidence-based protocols. *Ann Surg Oncol* 2009; 16:804-805.
40. [Http://dissertations.ub.rug.nl/FILES/faculties/medicine/2007/a.b.francken/melfo.pdf](http://dissertations.ub.rug.nl/FILES/faculties/medicine/2007/a.b.francken/melfo.pdf)





# 3

---



# Therapeutic groin dissection for melanoma; Risk factors for short term morbidity | HPAM Poos, S Kruijff, E Bastiaannet,

RJ van Ginkel, HJ Hoekstra | Eur J Surg Oncol. 2009;35:877-83

---

## ABSTRACT

**Aims:** Ilio-inguinal lymph node dissection for stage III melanoma is often complicated by wound healing disturbances. A retrospective study was performed to investigate the wound healing disturbances after therapeutic ilio-inguinal lymph node dissection.

**Patients and methods:** Between 1989 and 2007, 139 consecutive patients, 73 females (53%) and 66 males (47%), median age 55 (range 20-86) years underwent a therapeutic ilio-inguinal lymph node dissection. Data were recorded on early complications: haematoma, wound infection, wound necrosis and seroma. Univariate and multivariate logistic regression analyses were used to evaluate the influence of a wide range of variables on postoperative complications.

**Results:** Seventy-two patients had one or more early wound complications (49.7%). These complications comprised haematoma (n=3, 2.1%), wound infection (n=30, 20.7%), wound necrosis (n=25, 17.5%) and seroma (n=31, 21.8%). Wound infections were significantly more common in patients with a body mass index (BMI) of >25 (p=0.019). Wound necrosis developed significantly more often if the Bohler Braun splint was not used postoperatively (p=0.002). The occurrence of one or more early complications was significantly associated with the non-use of a Bohler Braun splint (p=0.026) and age of >55 years (p=0.015).

**Conclusions:** High BMI was significantly correlated with the occurrence of wound infections. Bed rest with the knee and hip in flexion using a Bohler splint improved wound healing after therapeutic ilio-inguinal lymph node dissection.

## INTRODUCTION

The incidence of melanoma is still increasing worldwide. In the Netherlands, the incidence currently amounts to 16.1/100,000, with a mortality-rate of 3.0/100,000. The anatomical localisation of preference in males is the trunk, and in females the leg.<sup>1</sup> In about 90% of the cases, melanoma is diagnosed at tumour stage I or II.<sup>2</sup> Approximately 16-28% of patients with a primary melanoma will develop a tumour recurrence, 20-28% will develop a local recurrence or intransit metastases, 26-60% will develop regional lymph node metastases and 15-50% will develop distant metastases.<sup>3</sup> The value of elective lymph node dissection has been studied to improve disease-free survival and overall survival. Four studies concluded that elective lymph node dissection did not influence the survival of melanoma patients.<sup>4-7</sup> Morton et al. developed the concept of sentinel lymph node biopsy (SLNB) to detect regional lymph node metastases at an early stage.<sup>8</sup> In patients with a primary melanoma of intermediate tumour thickness (1.2-3.5 mm) and a positive SLNB, completion lymph node dissection led to significant improved disease-free survival. An improved overall survival was not found, however.<sup>9</sup> Sentinel lymph node biopsy is a diagnostic procedure that involves a certain degree of morbidity.<sup>10-11</sup> Therefore, studies on the value of preoperative ultrasound examination of the regional lymph nodes are currently being performed.<sup>12</sup> Morbidity after elective and therapeutic ilio-inguinal lymph node dissection can comprise: wound necrosis, wound infection, seroma and/or lymphoedema.<sup>13-23</sup> Efforts have been made to improve the surgical technique and reduce the risk of complications. An ilio-inguinal lymph node dissection can be conducted via one vertical incision crossing the inguinal fold, that removes a skin ellipse (Fig. 1A), or via two separate incisions above and below the inguinal fold (Fig. 1D).<sup>13,24-26</sup> At the University Medical Centre Groningen (UMCG), the first technique, with the excision of a skin ellipse, is used.<sup>13,26</sup>

Previously, an extensive analysis on the complications that can arise after ilio-inguinal lymph node dissection was performed at the UMCG. Morbidity was present, but limited, while the risk of lymphoedema was small.<sup>13</sup> Recent research into the morbidity of sentinel lymph node biopsy in the ilio-inguinal region has shown that nowadays, more complications occur after completion lymph node dissections.<sup>10</sup> This prompted us to conduct a more extensive retrospective study to assess the present morbidity of therapeutic or completion lymph node dissection to determine the risk factors for short term morbidity and to evaluate recommendations on perioperative treatment.

## METHODS

### *Patients*

From 1989 to 2007, 139 consecutive patients, 73 females (53%) and 66 males (47%), with a median age of 55 (range 20-86) years underwent 143 ilio-inguinal lymph node dissections for locoregional metastasized melanoma. The melanomas were localised on the trunk (n=11,

7.6%), penis or scrotum (n=4, 2.8%), thigh (n=46, 31.7%), lower leg (n=39, 26.9%) and foot (n=30, 20.7%). Thirteen patients had an unknown primary tumour (9%). The median Breslow thickness was 2.5 (range 0.70-20.0) mm. Four patients underwent a bilateral ilio-inguinal lymph node dissection in two separate surgical procedures. Indications for ilio-inguinal lymph node dissection were histologically or cytologically proven ilio-inguinal lymph node metastases (n=112) or positive sentinel lymph node biopsy (n=31). The clinical data of the patients who underwent these 143 therapeutic lymph node dissections are shown in Table 1.


**TABLE 1:** Clinical data on 143 therapeutic ilio-inguinal lymph node dissections for stage III melanoma

Variable	No. of patients (%)
Gender	
M	69 (48.3)
F	74 (51.7)
Median age (years)	55 (20-86)
Localisation primary tumor	
Trunk	11 (7.6)
Penis / Scrotum	4 (2.8)
Thigh	46 (31.7)
Lower leg	39 (26.9)
Foot	30 (20.7)
Unknown	13 (9.0)
Breslow thickness	
T1 (<1.00 mm)	11 (7.7)
T2 (1.00-2.00 mm)	33 (23.1)
T3 (2.00-4.00 mm)	42 (29.4)
T4 (>4.00 mm)	39 (27.3)
Unknown	18 (12.6)
Median (mm)	2.5 (0.70 – 20.0)
Indication ilio-inguinal lymph node dissection	
Macrometastases	112 (78.3)
positive SLNB	31 (21.7)
Risk factors	
BMI > 25	76 (53.1)
Smoking	41 (28.7)
Cardiovascular disease	37 (25.9)
Pulmonary disease	8 (5.6)
Diabetes mellitus	5 (3.5)
Comorbidity*	43 (30.1)

\* Comorbidity: presence of cardiovascular disease and/or pulmonary disease and/or diabetes mellitus


### *Surgical technique*

First an ellipse shaped incision of the skin cranially, 2 cm medial from the superior anterior iliac spine, to 15 cm below Poupart's ligament was performed. The ellipse was approximately 4-6 cm at its broadest point. Any scar tissue from previous lymph node excisions was included in the skin ellipse (Figure 1A). The superficial dissection contained the subcutaneous fat and lymphatic tissue underneath the skin ellipse, limited by the long adductor muscle medially and the sartorius muscle laterally. The saphenous magna vein was taken along. The femoral artery and the femoral nerve became visible, as well as the femoral vein on the dorsal side of the femoral artery. The lymphatic tissue located at the superior aspect of the femoral canal, medially from the femoral vein is called the Cloquet node. It is thought that this tissue represents the leading lymph node draining into the pelvis from the inguinal basin.<sup>27</sup> Poupart's ligament was transected longitudinally 2 cm lateral from the neurovascular bundle, which exposed the iliac and obturator lymph nodes. Then the iliac lymph node dissection was performed. All the lymph nodes were excised up to the bifurcation of the communicating iliac artery, as well as the lymph nodes in the obturator fossa. Poupart's ligament was closed. The origin of the sartorius muscle was transected at the level of the anterior iliac spine and preserved over a length of about 10 cm (Figure 1B). Then it was reverted medially to cover the neurovascular femoral bundle and fixed to Poupart's ligament (Figure 1C).



**FIGURE 1.**

- A.** Ellipse shaped incision
- B.** The origin of the sartorius muscle transected and preserved over a length of about 10 cm
- C.** The sartorius muscle reverted medially to cover the neurovascular femoral bundle and fixed to Poupart's ligament
- D.** Separate incision above and below the groin



**FIGURE 2.** Bed rest with the hip and the knee in flexion on a Bohler Braun splint.

### *Postoperative treatment*

During the first 10 days postoperatively, the patient had strict bed rest. From 1989 to 2000, in the majority of patients (n=58, 70.7%) a Bohler Braun splint was used for the first 5 or 10 days to place the patient's hip and knee in flexion in order to reduce the tension along the wound. This concept was thought to support wound healing (Figure 2). A Bohler Braun Splint is often used in orthopaedics to provide stabilization with or without traction for a variety of angulated lower extremity fractures. Since 2000 the distal end of the bed was elevated for the first 5 days with the hip and knee in slight flexion and in only 5 patients (8.2%) a Bohler Braun splint was used after 2000. The use of the Bohler Braun splint was gradually abandoned in the late nineties because the 'old hospitals beds' were then replaced by 'modern hospital beds' with the option to place the hip and the knee in flexion. The wound drains were removed after a minimum of 7 days, and if the production was less than 20 ml per 24 hours. For the first 6 months postoperatively, patients were prescribed a tailor made support stocking that had been made for the affected leg prior to surgery. Perioperatively and during the period of bed rest, all patients received prophylactic low molecular weight heparin (fraxiparine). Antibiotics were not administered routinely during surgery.

### *Complications*

Four types of complications were scored: 1, haematoma that required surgical intervention; 2, wound infection if the wound was culture-positive and/or therapeutic antibiotics were indicated; 3, wound necrosis if the edges of the wound were necrotic, irrespective of whether necrotomy had been performed, and if there was secondary wound healing; 4, seroma that required puncture.

### *Variables*

The following variables were analysed for their possible influence on postoperative morbidity: age ( $\leq 55$  years vs  $> 55$  years), gender, smoking, body mass index (BMI,  $\leq 25$  vs  $> 25$ ), comorbidity (diabetes mellitus (DM), cardiovascular disease and chronic obstructive pulmonary disease), surgeon (fellow vs staff surgeon), operative time (150 min vs  $> 150$  min), bed rest (Bohler Braun splint vs no Bohler Braun splint), operative period (1989-2000 vs 2001-2007) and surgical indication (macrometastases vs positive SLNB).

**TABLE 2:** Univariate and multivariate analysis on Infection, Wound necrosis and 1 or more complications

Variable	Wound infection n=30 (%)	Univariate Analysis p-value	Multivariate analysis OR (95%CI) p-value	Wound necrosis n=25 (%)
Age (years)				
≤ 55	12/74 (16.2)		1.57 (0.62-3.94)	11/74 (14.9)
> 55	18/69 (26.1)	0.147	0.339	14/69 (20.3)
Gender				
M	16/69 (23.2)			14/69 (20.3)
F	14/74 (18.9)	0.531		11/74 (14.9)
Smoking				
+	7/41 (17.1)			9/41 (22.0)
-	23/102 (22.6)	0.467		16/102 (15.7)
BMI				
> 25	22/76 (29.0)		3.28 (1.21-8.88)	16/76 (21.1)
≤ 25	6/58 (10.3)	0.009	0.019	7/58 (12.1)
Comorbidity				
+	13/43 (30.2)		1.39 (0.53-3.65)	12/43 (27.9)
-	17/100 (17.0)	0.075	0.500	13/100 (13.0)
Surgeon				
Staff	9/54 (16.7)			8/54 (14.8)
Fellow	21/89 (23.6)	0.324		17/89 (19.1)
Operative time				
≤ 150 min	14/63 (22.2)			12/63 (19.1)
> 150 min	16/80 (20.0)	0.746		13/80 (16.3)
Bed rest				
Bohler Braun splint +	12/63 (19.0)			5/63 (7.9)
Bohler Braun splint –	18/80 (22.5)	0.615		20/80 (25.0)
Operation year				
89-00	14/82 (17.1)		1.51 (0.63-3.62)	13/82 (15.9)
01-07	16/61 (26.2)	0.184	0.353	12/61 (19.7)
Indication				
Macrometastasis	23/112 (20.5)			17/112 (15.2)
Positive SLNB	7/31 (22.6)	0.805		8/31 (25.8)

### *Statistical analysis*

The statistical analyses were performed with SPSS version 14.0.2. Early complications were analysed separately and also as combined variables. Chi square tests and multivariate logistic regression were used to assess factors associated with the early complications. Factors with  $p < 0.20$  in the univariate analysis were entered into the multivariate model.

Univariate Analysis p-value	Multivariate analysis OR (95%CI) p-value	1 or more complications n=72 (%)	Univariate Analysis p-value	Multivariate analysis OR (95%CI) p-value
		27/74 (36.5)		2.71 (1.21-6.05)
0.393		45/69 (65.2)	0.001	0.015
		38/69 (55.1)		
0.393		34/74 (45.9)	0.275	
		21/41 (51.2)		
0.372		51/102 (50.0)	0.895	
	1.89 (0.67-5.35)	44/76 (57.9)		2.05 (0.94-4.50)
0.172	0.229	23/58 (39.7)	0.036	0.071
	2.72 (0.99-7.50)	30/43 (69.8)		2.13 (0.86-5.29)
0.031	0.053	42/100 (42.0)	0.002	0.101
		26/54 (48.1)		
0.513		46/89 (51.7)	0.682	
		35/63 (55.6)		
0.662		37/80 (46.3)	0.269	
	6.62 (1.94-22.53)	24/63 (38.1)		3.14 (1.14-8.57)
0.008	0.002	48/80 (60.0)	0.009	0.026
		35/82 (42.7)		1.30 (0.48-3.54)
0.552		37/61 (60.7)	0.034	0.608
	0.43 (0.13-1.38)	57/112 (50.9)		
0.168	0.155	15/31 (48.4)	0.805	

## RESULTS

### *General*

The median duration of hospitalisation was 14 (range 7-44) days. There was no mortality due to surgery. Overall perioperative morbidity was 55.2%. General complications were observed in eight cases: two delirium (1%), four urinary tract infections (3%) and two bladder retentions (1%); there were no pulmonary and thrombo-embolic complications.



Early wound complications comprised: haematoma (n=3, 2.1%), wound infection (n=30, 20.7%), wound necrosis (n=25, 17.5%) and seroma (n=31, 21.8%). Seventy-two patients had one or more early wound complications (49.7%).

### *Multivariate analysis*

No significant associations were found between the different variables and the complications comprised haematoma and seroma. Wound infections occurred significantly more frequently in patients with a BMI of >25. Wound necrosis was more common if the Bohler Braun splint was not used postoperatively. The occurrence of one or more postoperative complications was significantly correlated with not applying a Bohler Braun splint and with more advanced age.

Comorbidity was related to a higher risk of wound necrosis, although not significantly proven. An overview of the univariate and multivariate analyses of local complications is presented in Table 2.

## DISCUSSION

### *General*

In the present study, a BMI of >25 was correlated with a higher risk of wound infection. Additionally, advanced age (>55 years) was associated with a higher incidence of one or more early complications. And, applying a Bohler Braun splint in the postoperative phase of strict bed rest after ilio-inguinal lymph node dissection reduced the risk of wound necrosis and significantly prevented one or more early wound complications. At least 10 studies have been

**TABLE 3:** Literature overview of wound complications in melanoma patients who underwent therapeutic or elective groin dissection

Author	Year	No. of patients	Inguinal / Ilio-inguinal	Thera-peutic / Elective	Wound infection %	Wound necrosis %
Baas et al. <sup>13</sup>	1992	151	8 / 143	138/31	9	3**
Beitsch et al. <sup>14</sup>	1992	168	168 / 0	132/45	29	26
Karakousis et al. <sup>17</sup>	1994	205	94 / 111	90/115	16	8
Pearlman et al. <sup>19</sup>	1995	19	0 / 19	19/0	26*	?
Strobbe et al. <sup>22</sup>	1999	52	0 / 52	52/0	17	15
Hughes et al. <sup>16</sup>	2000	132	60 / 72	132/0	13	8
Lawton et al. <sup>18</sup>	2002	56	0 / 56	12 / 44	30	4
Serpell et al. <sup>21</sup>	2003	27	25 / 2	27/0	25	7
Van Akkooi et al. <sup>23</sup>	2007	129	0 / 129	129/0	29	?
Sabel et al. <sup>20</sup>	2007	212	181 / 31	212/0	19*	?
Poos et al.	2008	139	0 / 143	143/0	21	18/7**

\* Wound infection with or without partial flap necrosis/wound dehiscence

\*\* Requiring surgical treatment

published about wound healing disturbances after lymph node dissection since the early 1990s.<sup>13-23</sup> Unfortunately, it was not possible to perform a subgroup analysis on the series that compares therapeutic lymph node dissection with elective procedures.<sup>13,14,17,18</sup> There was a wide variation in the incidence of complications: haematoma 0-5%, wound infection 9-30%, wound necrosis 3-26% and seroma 5-46% (Table 3). The rates found in the present study were very similar.

### *Our institute*

However, the results of the present study were much worse than those published by our research group over the period 1970-1984.<sup>13</sup> In the 1970s and the 1980s, our ilio-inguinal lymph node dissection patients were given 10 days of bed rest with a Bohler Braun splint that held their hip and knee in flexion (Figure 2). This might explain the lower number of postoperative complications: wound infection 9%, wound necrosis 3% and seroma 17%. From the 1990s, we reduced the period of bed rest with a Bohler Braun splint to 5 days and in the year 2000, we abandoned the Bohler Braun splint in the postoperative phase. In our study, three factors appeared to significantly influence early complications: a BMI of >25, an age of >55 years and abandoning the use of a Bohler Braun splint. The only factor that changed between 1989 and 2007 was the duration/non-use of the Bohler Braun splint. In the study by Baas et al., in which it was standard practice to use a Bohler Braun splint for the first 10 days postoperatively, the median age was only 44 years compared to 55 years in the present study and only 17.2% had severe obesity (BMI >30) compared to the 53% moderate obesity (BMI >25) in the present study. These findings most likely explain the differences in the complication rates between the two retrospective studies performed at our institute.

Seroma %	Incision type	Bed rest days
17	Vertical	10
14	Vertical	?
5	Vertical	0
?	Separate	1
15	Vertical /Separate	0
23	Separate	?
21	Separate	0
46	Separate	7
20	Separate	3
?	Vertical /Separate	?
22	Vertical	10

### *Bed rest*

The use of a Bohler Braun splint to nurse the hip and the knee in flexion during the period of strict bed rest has not been described in any of the other publications. In the studies by Serpell and van Akkooi, bed rest was prescribed for 7 and 3 days, respectively.<sup>21,23</sup> Karakousis et al. reported to prescribe flexion of the leg during bed rest and at night, but contrastingly, they mobilised patients directly after the operation and this resulted in a low risk of complications.<sup>17</sup> In the other studies, the patients were mobilised as soon as possible with a support stocking.<sup>18,19,22</sup> It has to be noticed that nowadays, it is also possible to elevate the legs with the hip and the knee in flexion by adjusting the position of modern hospital beds.

### *Patient characteristics*

Advanced age (>50 years), obesity and smoking are known to increase the risk of wound complications.<sup>14,28</sup> Age over 55 years was only correlated with the combined variable (one or more early complications), which is probably the result of smaller numbers in the separate analysis. A longer duration of surgery might also contribute to a higher risk of complications.<sup>28</sup> We were unable to confirm the latter in the present study.

### *Surgical techniques*

Different surgical techniques for performing an ilio-inguinal nodal dissection are known. Firstly, the ilio-inguinal node dissection can be performed in two different ways: either one straight incision with a skin ellipse and vertical transection of the inguinal ligament or two separate incisions above and below the groin, which was introduced by Baronofsky (Figure 1D).<sup>13,24,26</sup> The vertical incision is thought to cause minimal damage to the lymphatic vessels and therefore less risk of lymphoedema.<sup>13</sup> In contrast two separate incisions are thought to leave the groin free, which should reduce the risk of wound infection.<sup>24,25</sup> Sabel et al. did not find a difference in complication rates between the two different techniques. However, in their study, two incisions were only used in 15% of the ilio-inguinal node dissections.<sup>20</sup> Pearlman et al. and Sabel et al. did not report wound necrosis as a separate wound complication; van Akkooi et al. did not mention wound necrosis at all. The low incidences of wound necrosis in the studies mentioned before (Table 3) could suggest that using separate incisions may be less invasive.

### *Lymph node dissection after SLNB*

Completion lymph node dissection after a positive SLNB may cause an increased risk of wound infection and postoperative oedema, possibly because two operations are performed on the groin within a very short period.<sup>10</sup>

The study by Sabel et al. described that patients who underwent a completion lymph node dissection after positive SLNB (n=132) had significantly less wound complications and less often lymphoedema than patients who underwent a therapeutic lymph node dissection for palpable lymph node metastases (n=80).<sup>20</sup> However, in this series 181 superficial nodal

dissections were performed and only 31 ilio-inguinal lymph node dissections were carried out. The indication for a deep nodal dissection was based on a positive Cloquet's node or intraoperative suspicion for iliac nodal disease. This might explain the differences with the current study.

### *Surgical volume*

It has been suggested in some studies that experience and/or surgical volume are correlated with decreased morbidity; we were not able to confirm this in the present study.<sup>29</sup> It should be noted that surgical oncology fellows generally performed the operations under the supervision of an experienced surgical oncologist.

### *Consideration and conclusion*

Therapeutic groin dissection for melanoma remains a surgical procedure with a high morbidity. Presently, more complications occur after therapeutic ilio-inguinal lymph node dissection than 20 years ago. To comply with our data there are two explanations for this difference: (1) nowadays patients have a higher BMI and (2) a Bohler Braun splint was not used anymore during strict bed rest for 10 days. The majority of patients had a BMI of >25, taking along a higher risk of pulmonary and thrombo-embolic complications.


For these patients early mobilisation is indicated. It has not been investigated so far whether long term immobilisation after an ilio-inguinal lymph node dissection to reduce the risk of wound healing complications is cost-effective. This study was performed retrospectively and therefore it is possible that there were differences in the reported complications in the medical records. Lymphoedema, as a complication of groin dissection was not included in this study, as the extent of this complication cannot be obtained from medical records, because lymphoedema should be identified clinically by measuring the volume of the leg. In a recent study the limb volume was measured, and slight oedema was found after a performed groin dissection.<sup>10</sup> Considering our results, we advise performing a groin dissection with one vertical ellipse shaped incision and strict postoperative bed rest of 5 days with the hip and the knee in flexion, to reduce tension on the wound and to facilitate healing of the transferred sartorius muscle. This regimen is followed by mobilisation with a tailor made support stocking. However, we are aware that only a randomized controlled trial can determine which technique (vertical vs separate incisions) and which postoperative treatment cause the least morbidity. If the patient has recently undergone SLNB and required a completion lymph node dissection, prophylactic antibiotics are indicated during surgery to reduce the risk of wound infection. Whether perioperative antibiotics in therapeutic groin dissections are valuable, can also only be evaluated in a well designed prospective trial.

## REFERENCES

1. De Vries E, Coebergh JW. Cutaneous malignant melanoma in Europe. *Eur.J.Cancer* 2004; 40:2355-2366.
2. Garbe C, Eigentler TK. Diagnosis and treatment of cutaneous melanoma: state of the art 2006. *Melanoma Res.* 2007; 17:117-127.
3. Francken AB, Bastiaannet E, Hoekstra HJ. Follow-up in patients with localised primary cutaneous melanoma. *Lancet Oncol.* 2005; 6:608-621.
4. Balch CM, Soong SJ, Bartolucci AA, et al. Efficacy of an elective regional lymph node dissection of 1 to 4 mm thick melanomas for patients 60 years of age and younger. *Ann.Surg.* 1996; 224:255-263.
5. Cascinelli N, Morabito A, Santinami M, et al. Immediate or delayed dissection of regional nodes in patients with melanoma of the trunk: a randomised trial. *WHO Melanoma Programme. Lancet* 1998; 351:793-796.
6. Sim FH, Taylor WF, Pritchard DJ, Soule EH. Lymphadenectomy in the management of stage I malignant melanoma: a prospective randomized study. *Mayo Clin.Proc.* 1986; 61:697-705.
7. Veronesi U, Adamus J, Bandiera DC, et al. Inefficacy of immediate node dissection in stage 1 melanoma of the limbs. *N.Engl.J.Med.* 1977; 297:627-630.
8. Morton DL, Wen DR, Wong JH, et al. Technical details of intraoperative lymphatic mapping for early stage melanoma. *Arch. Surg.* 1992; 127:392-399.
9. Morton DL, Thompson JE, Cochran AJ, et al. Sentinel-node biopsy or nodal observation in melanoma. *N.Engl.J.Med.* 2006; 355:1307-1317.
10. de Vries M, Vonkeman WG, van Ginkel RJ, et al. Morbidity after inguinal sentinel lymph node biopsy and completion lymph node dissection in patients with cutaneous melanoma. *Eur.J.Surg. Oncol.* 2006; 32:785-789.
11. Morton DL, Cochran AJ, Thompson JE, et al. Sentinel node biopsy for early-stage melanoma: accuracy and morbidity in MSLT-I, an international multicenter trial. *Ann.Surg.* 2005; 242:302-11.
12. Voit C, Kron M, Schafer G, et al. Ultrasound-guided fine needle aspiration cytology prior to sentinel lymph node biopsy in melanoma patients. *Ann.Surg. Oncol.* 2006; 13:1682-1689.
13. Baas PC, Schraffordt Koops H, Hoekstra HJ, et al. Groin dissection in the treatment of lower-extremity melanoma. Short-term and long-term morbidity. *Arch.Surg.* 1992; 127:281-286.
14. Beitsch P, Balch C. Operative morbidity and risk factor assessment in melanoma patients undergoing inguinal lymph node dissection. *Am.J.Surg.* 1992; 164:462-465.
15. Fife K, Thompson JE. Lymph-node metastases in patients with melanoma: what is the optimum management? *Lancet Oncol.* 2001; 2:614-621.
16. Hughes TM, A'Hern RP, Thomas JM. Prognosis and surgical management of patients with palpable inguinal lymph node metastases from melanoma. *Br.J.Surg.* 2000; 87:892-901.
17. Karakousis CP, Driscoll DL. Groin dissection in malignant melanoma. *Br.J.Surg.* 1994; 81:1771-1774.
18. Lawton G, Rasque H, Ariyan S. Preservation of muscle fascia to decrease lymphedema after complete axillary and ilioinguinofemoral lymphadenectomy for melanoma. *J.Am.Coll.Surg.* 2002; 195:339-351.
19. Pearlman NW, Robinson WA, Dreiling LK, et al. Modified ilioinguinal node dissection for metastatic melanoma. *Am.J.Surg.* 1995; 170:647-649.
20. Sabel MS, Griffith KA, Arora A, et al. Inguinal node dissection for melanoma in the era of sentinel lymph node biopsy. *Surgery* 2007; 141:728-735.
21. Serpell JW, Carne PW, Bailey M. Radical lymph node dissection for melanoma. *ANZ.J.Surg.* 2003; 73:294-299.
22. Strobbe LJ, Jonk A, Hart AA, et al. Positive iliac and obturator nodes in melanoma: survival and prognostic factors. *Ann.Surg. Oncol.* 1999; 6:255-262.
23. Van Akkooi AC, Bouwhuis MG, van Geel AN, et al. Morbidity and prognosis after therapeutic lymph node dissections for malignant melanoma. *Eur.J.Surg.Oncol.* 2007; 33:102108.
24. Baronofsky, ID. Technique of inguinal node dissection. *Surgery* 1948; 24:555-567.
25. Fraley EE, Hutchens HC. Radical ilio-inguinal node dissection: the skin bridge technique. A new procedure. *J.Urol.* 1972; 108:279-281.
26. Van der Ploeg E, Schraffordt Koops H. A modified technique of groin dissection. *Arch.Chir Neerl.* 1972; 24:31-36.

27. Essner R, Scheri R, Kavanagh M, et al. Surgical management of the groin lymph nodes in melanoma in the era of sentinel lymph node dissection. *Arch.Surg.* 2006; 141:877-882.
28. Garibaldi RA, Cushing D, Lerer T. Risk factors for postoperative infection. *Am.J.Med.* 1991; 91:1585-1635.
29. Birkmeyer JD, Stukel TA, Siewers AE, et al. Surgeon volume and operative mortality in the United States. *N.Engl.J.Med.* 2003; 349:2117-2127.
30. Hill JO, Wyatt HR, Reed GW, et al. Obesity and the Environment: Where Do We Go From Here? *Science* 2003; 299: 853-5.








# Part II

Prognostic markers in melanoma



# 4

---



# Breslow thickness in the netherlands: a population-based study of 40,880 patients comparing young and elderly patients

| S Kruijff, E Bastiaannet, AB Francken, M Schaapveld, M van der Aa, HJ Hoekstra | Submitted

---

## ABSTRACT

**Background:** Melanoma incidence has increased rapidly in the last decades, and predictions show a continuing increase in the years to come. The aim of this study was to assess trends in melanoma incidence, Breslow thickness (BT), and melanoma survival among young and elderly patients in the Netherlands.

**Methods:** Patients diagnosed with invasive melanoma between 1994 and 2008 were selected from the Netherlands Cancer Registry (NCR). Incidence (per 100,000) over time was calculated for young (<65 years) and elderly patients ( $\geq 65$  years). Distribution of BT for young and elderly males and females was assessed. Regression analysis of the log-transformed BT was used to assess changes over time. Relative survival was calculated as the ratio of observed survival to expected survival.

**Findings:** Overall 40,880 patients were included (42.3% male, 57.7% female). Melanoma incidence increased more rapidly among the elderly (5.4% Estimated Annual Percent Change (EAPC),  $p < 0.0001$ ) than among younger patients (3.9% EAPC,  $p < 0.0001$ ). The overall BT declined significantly over time ( $p < 0.001$ ). Among younger patients, BT decreased for almost all locations. Among elderly males, BT decreased for melanomas in the head and neck region ( $p = 0.001$ ) and trunk ( $p < 0.001$ ), but did not decrease significantly for the other regions. Among elderly females, BT only decreased for melanomas at the trunk ( $p = 0.01$ ). The relative survival of elderly patients was worse compared with that of younger patients ( $p < 0.001$ ).

**Interpretation:** Melanoma incidence increases more rapidly for elderly than for younger patients and the decline in Breslow thickness is less prominent among elderly patients than among young patients. Campaigns in the Netherlands must focus more on early melanoma detection in the elderly.

## INTRODUCTION

The incidence of cutaneous melanoma has increased in the last decades and estimates predict a continuing increase in the coming years.<sup>1</sup> High incidence rates are found in populations of predominantly European origin, with white populations in Australia and New Zealand having the highest incidence rates and Asian and black populations having the lowest.<sup>2</sup>

In Australia, despite rising incidence rates, mortality rates seem to have reached a plateau, possibly due to the fact that most newly diagnosed melanomas are thinner melanomas, which usually do not lead to death.<sup>3</sup> In the Netherlands, the total number of melanoma patients is expected to increase from 2,400 patients in 2000 to 4,800 patients in 2015, and has already reached 3,500 patients in 2005.<sup>1</sup> In contrast with Australia, in the Netherlands this increase in incidence rate is accompanied by a rising mortality rate. This finding seems to contradict the earlier hypothesis that rising incidence in the Netherlands might be the result of increased awareness only.<sup>4</sup>

Increased exposure to the sun and sunburns during childhood are still the most important risk factors for melanoma. A recent meta-analysis has also shown that significantly increased melanoma risk is associated with sun bed and sun lamp exposure.<sup>5,6</sup>

Various studies have already shown that the melanoma incidence and mortality among elderly individuals (>65 years) is growing rapidly.<sup>7,8</sup> As the geriatric population increases in most industrialized nations (life expectancy almost doubled during the last century), melanoma will become an important health issue for the elderly age group in this century.<sup>9-11</sup>

The most important predictors of mortality, included in the American Joint Committee on Cancer (AJCC) staging, are tumor characteristics such as Breslow thickness (BT), ulceration, mitotic rate, and the presence of metastases.<sup>12,13</sup> However, BT has been shown to be the single most important prognostic factor for survival.<sup>12-14</sup> Therefore, the aim of this study was to assess incidence rates and differences in BT and survival between young and elderly patients with invasive melanoma in the Netherlands between 1994 and 2008.

## METHODS

### *Patients*

Patients diagnosed with invasive melanoma between 1994 and 2008 were selected from the Netherlands Cancer Registry (NCR), which covers all patients in the Netherlands. The nationwide Dutch network and registry of histopathology and cytopathology (PALGA) regularly submits reports of all diagnosed malignancies to the cancer registries. The national hospital discharge databank, which receives discharge diagnoses of admitted patients from all Dutch hospitals, completes case ascertainment. After notification, well-trained registry personnel collect data on diagnosis, staging, and treatment from the medical records, including pathology and surgery reports, using the registration and coding manual of the Dutch Association

of Comprehensive Cancer Centers. For the present study, patients with their first primary melanoma were selected. Stage was defined according to the AJCC staging system. The study was approved by the local medical ethics committee.

### *Statistical analysis*

Incidence of invasive melanoma per 100,000 Dutch individuals was calculated for young (younger than 65 years) and elderly (65 years and older) patients in the Netherlands. Because BT measurement was missing for 9.5% of patients, we used multiple imputation (5 imputations) to generate a complete dataset. A model was built that included sex, location, stage, year of diagnosis, age, and status as predictors to assess the patients with missing BT measurements. Median BT was calculated according to sex, age, and localisation, as well as Breslow distribution over time. Because the TNM Classification of Malignant Tumors changed in 2003, stage distribution before and after 2003 were not comparable and therefore a composite stage variable was not used in the analysis. BT was divided according to the AJCC staging system into the following size categories:  $\leq 1.0$  mm, 1.0-2.0 mm, 2.0-4.0 mm, and  $>4.0$  mm. Patients with an unknown primary were excluded from the analysis.

BT distributions for young and elderly males and females were assessed over time. Because BT is skewed toward smaller tumours (most melanomas are  $\leq 1.0$  mm), BT was log-transformed. A regression analysis was modeled with  $\log(\text{Breslow})$  to assess the changes over the years of incidence according to age, sex, and location.

Vital status and date of last follow-up were established either directly from the patient's medical record or through linkage of cancer registry data with municipal population registries (follow-up until January 1, 2009), which record information on vital status. Relative survival is the preferred way to describe the prognosis of (elderly) patients with melanoma, as it takes into account the risk of dying from causes other than melanoma. Relative survival was calculated as the ratio of the observed survival among cancer patients to the survival that would have been expected based on the corresponding general population (with respect to age, sex, and year of diagnosis). National lifespan tables were used to estimate expected survival (Ederer II method). Relative Excess Risks of death (RERs) for year of diagnosis were estimated using a Poisson regression model.

## **RESULTS**

This study includes 40,880 patients diagnosed with melanoma between 1994 and 2008. Table 1 shows the characteristics of the melanoma patients in the Netherlands between 1994 and 2008 of which 42% were male and 58% were female. Median age was 54 years (range 0-105 years). The trunk was the most frequently involved melanoma site (36%). From 1994 to 1996, 5,907 new melanoma patients were diagnosed (15%), increasing to 11,023 (27%) in the period from

2006 to 2008. Almost one-half (49%) of newly diagnosed melanoma patients had a BT  $\leq 1.0$  mm, while for 12% the BT was  $>4.0$  mm for thick melanomas (Table 1).


**TABLE 1** Characteristics of melanoma patients in the Netherlands, 1994-2008

Characteristic		Original data		Multiple imputation
		N	%	%
Sex	Male	17 305	42.3	42.3
	Female	23 575	57.7	57.7
Age	$\leq 40$ years	9378	23.0	23.0
	41-54 years	11 506	28.1	28.1
	55-64 years	8217	20.1	20.1
	$\geq 65$ years	11 779	28.8	28.8
Location	Head & neck	5612	13.7	13.7
	Trunk	14 693	35.9	35.9
	Upper extremities	8163	20.0	20.0
	Lower extremities	12 246	30.0	30.0
	Other	166	0.4	0.4
Year	1994-1996	5907	14.5	14.5
	1997-1999	6720	16.4	16.4
	2000-2002	7779	19.0	19.0
	2003-2005	9451	23.1	23.1
	2006-2008	11 023	27.0	27.0
Breslow category	$\leq 1.0$ mm	19 325	47.3	48.7
	1.0-2.0 mm	8622	21.1	22.7
	2.0-4.0 mm	5598	13.7	16.7
	$>4.0$ mm	3451	8.4	11.9
	Unknown	3884	9.5	


Melanoma incidence increased more rapidly among elderly patients ( $\geq 65$  years) than among younger patients (Figure 1). In 1994, melanoma incidence was 189 among elderly patients and 144 among younger patients, and in 2008 these numbers increased to 362 and 241, respectively. The Estimated Annual Percentage Change (EAPC) for young patients was 3.9% ( $p < 0.0001$ ) and for the elderly the EAPC was 5.4% ( $p < 0.0001$ ). Especially since 2002, melanoma incidence appears to be increasing at a faster rate among the elderly than among their younger counterparts.

Figure 2 shows the proportion of patients that were diagnosed with thin, median, and thick melanomas over time. The percentage of young patients with a thick melanoma (Breslow  $>4.0$  mm) has declined over time. In 1994, 16% of males and 10% of females had a thick melanoma; by 2008, these incidences had declined to 9% and 5%, respectively ( $p < 0.001$ ).

A decline was also observed among the elderly population, and especially among elderly females, although it was not as precipitous as among young patients. In 1994, 25% of elderly



**FIGURE 1** Incidence of invasive melanoma among young and elderly patients




**FIGURE 2** Distribution of Breslow thickness over time among young and elderly patients

males and 20% of elderly females had a thick melanoma, compared to, respectively, 19% and 18% in 2008 ( $p < 0.001$ ). Although younger patients generally had thinner melanomas than elderly patients, the proportion of thin melanomas increased in both age groups. In 1994, 43% of young males and 53% of young females had a thin melanoma (Breslow  $\leq 1.0$  mm); these numbers

**TABLE 2** Changes in Breslow thickness (log transformation) over time according to age, sex, and lesion location

			Regression coefficient (95% CI)		p-value
Young	Males	Head & neck	-0.43	-0.66 to -0.20	<0.001
		Trunk	-0.26	-0.37 to -0.14	<0.001
		Upper extremities	-0.36	-0.56 to -0.16	<0.001
		Lower extremities	-0.16	-0.41 to -0.1	0.2
	Females	Head & neck	-0.21	-0.51 to 0.1	0.2
		Trunk	-0.31	-0.45 to 0.17	<0.001
		Upper extremities	-0.51	-0.70 to -0.32	<0.001
		Lower extremities	-0.33	-0.46 to -0.19	<0.001
Elderly	Males	Head & neck	-0.35	-0.56 to -0.14	0.001
		Trunk	-0.42	-0.60 to -0.25	<0.001
		Upper extremities	-0.20	-0.47 to 0.1	0.2
		Lower extremities	-0.28	-0.61 to 0.1	0.1
	Females	Head & neck	-0.18	-0.44 to 0.1	0.2
		Trunk	-0.36	-0.62 to -0.10	0.01
		Upper extremities	-0.15	-0.37 to 0.06	0.2
		Lower extremities	-0.13	-0.32 to 0.1	0.2
Overall	Males & females	All locations	-0.25	-0.30 to -0.20	<0.001



**FIGURE 3** Cumulative relative survival among young and elderly melanoma patients

**TABLE 3** Relative Excess Risk (RER) for elderly (65 years and older) versus young patients, stratified according to Breslow category and lesion location

Stratification		Adjusted RER* elderly vs. young	p-value (adjusted*)
Breslow thickness	≤1.0 mm	1.9 (1.4-2.6)	<0.001
	1.0-2.0 mm	1.6 (1.3-2.0)	<0.001
	2.0-4.0 mm	1.6 (1.4-1.9)	<0.001
	>4.0 mm	1.7 (1.5-1.9)	<0.001
Location	Head & neck	1.7 (1.5-2.0)	<0.001
	Trunk	1.7 (1.5-1.8)	<0.001
	Upper extremities	1.5 (1.3-1.8)	<0.001
	Lower extremities	2.0 (1.8-2.4)	<0.001

\*adjusted for sex, year, N, M, Breslow thickness, and location (when not stratified for that factor)

increased to 51% and 62%, respectively, in 2008. In 1994, 31% of elderly males and 34% of elderly females had a thin melanoma, compared with 40% and 42%, respectively, in 2008 ( $p < 0.001$ ).

Table 2 depicts the changes in BT over time according to age, sex, and location. BT significantly decreased over time for head and neck, trunk, and upper extremities among young males ( $p < 0.001$ ), and for trunk, upper extremities, and lower extremities among young females ( $p < 0.001$ ). However, among elderly males a significant decrease in BT over the years was only observed for melanomas in the head and neck ( $p = 0.001$ ) and trunk ( $p < 0.001$ ), and for elderly females a decrease in BT is noted only on the trunk ( $p = 0.01$ ).

Relative survival is worse among elderly patients than among patients younger than 65 years (RER 2.1 for elderly patients; 95% CI 2.0–203;  $p < 0.001$ ). Adjusted for BT, location, sex, year, lymph node status, and distant metastases, RER remained higher among the elderly (RER 1.7; 95% CI 1.6–1.8);  $p < 0.001$ ). The RER for elderly (65 years and older) versus young patients was significantly worse for the elderly in all BT categories ( $\leq 1.0$  mm, 1.0–2.0 mm, 2.0–4.0 mm, > 4.0 mm) and locations (head and neck, trunk, upper extremities, and lower extremities).

## DISCUSSION

Melanoma incidence increases more rapidly among the elderly than among younger individuals. Additionally, rising incidence is accompanied by decreasing Breslow thickness (BT) mainly in younger individuals. The decrease in melanoma Breslow thickness was less prominent among elderly individuals than younger individuals, and elderly individuals are still frequently diagnosed with thick melanomas. For most melanoma locations, BT in the elderly has not been declining. Especially in elderly men, the proportion of thick melanomas has declined only minimally. The larger proportion of thick melanomas only partly explains the worse survival among elderly melanoma patients when compared with the younger



population; even within several strata of thickness, survival remains worse among the elderly than among younger patients.

In general, for males and females young and old, the percentage of thin melanomas increased over the period 1994-2008. This is in accordance with trends in several population-based studies in various other industrialized countries. Lashithiotakis et al, who studied 1980 patients diagnosed with melanoma in southern Germany in the period 1976-2003, observed that median BT decreased steadily during that period ( $p < 0.01$ ).<sup>15</sup> Buettner et al studied all 45,483 melanoma patients diagnosed between 1976 and 2000, and also noted a significant decrease in median BT ( $p < 0.0001$ ).<sup>16</sup>

In our study, the highest proportion of thin melanomas was observed in young females, a finding that is shared by many other studies.<sup>17</sup> Women are likely more alert to skin changes, resulting in thinner melanomas at diagnosis. Various studies have also shown that female patients were more likely to discover their melanoma or melanoma recurrence by themselves, even when adjusted for localisation.<sup>18,19</sup>

A trend towards prognostically more favorable melanomas has been described in the United States, Australia, and Europe, and can most likely be attributed to increased awareness in the population leading to earlier detection and treatment.<sup>20,21</sup>

The most prominent finding in our data was that elderly patients present with significantly thicker melanomas, a phenomenon described in both Europe and Australia.<sup>9-11, 22-25</sup> The question arises: why do elderly individuals fail so comprehensively to follow the trend towards early detection of melanoma that has been observed for the younger age group? Elderly patients have more difficulty recognizing or detecting changes in their skin, and therefore delay visits to their general practitioner.<sup>22</sup> Difficulty in detecting melanoma could be related to deteriorating vision, increased isolation because of loss of a partner, decreased flexibility, development of skin lesions in locations that may be difficult to see, and reduced awareness. Another difficulty for elderly patients is often the development of multiple seborrheic keratoses that may appear similar to melanoma. Additionally, melanomas are found on the back in 48% of cases in elderly men, which does not facilitate detection.<sup>11</sup>

Despite physical deterioration, it has been demonstrated that older people should be very well able to detect skin changes associated with early melanoma.<sup>25</sup> This finding does suggest that public education campaigns and increased alertness in the context of elderly care might be useful to encourage the detection of skin changes. Although previous educational campaigns have been effective at promoting awareness, they have generally been focused on younger individuals, with a continuous lack of attention directed toward elderly individuals.<sup>11</sup>

Various studies report age as an independent prognostic factor, perhaps presenting a surrogate for declining host defence mechanism associated with advancing age.<sup>9</sup> Treatment outcome is also influenced negatively, because older patients often receive a delayed diagnosis, incorrect staging, suboptimal surgery, and lower standard radio- or chemotherapy, possibly due to a reluctance to treat aggressively because of patients' co-morbidities or disabilities.<sup>9,10</sup> Numerous studies have already demonstrated that elderly patients, especially those over 60 years of age, have a lower disease-specific survival rate.<sup>26-28</sup>

In Australia, skin cancer has been identified as a national health priority, and awareness in the population is at a high level thanks to health-promotion activities.<sup>29</sup> In contrast, some studies demonstrate that melanoma awareness in Europe is not at such a high level, and focus has been mainly on young individuals.<sup>7,22,30</sup> This hypothesis might explain the high percentage of thick melanomas in the elderly, which (given the rising incidence) could have been caused by sun exposure in the era before melanoma became a health priority.

In conclusion, melanoma incidence has increased much more rapidly among the elderly than among younger individuals. Additionally, in contrast to younger patients, in which rising incidence is accompanied by a decline in BT, elderly patients still have the highest proportion of thick melanomas and the decline in BT among the elderly is not present at all melanoma locations. Campaigns in the Netherlands should focus more on early detection of melanoma in the elderly.


## REFERENCES

1. De Vries E, van de Poll-Franse LV, Louwman WJ, de Gruijl FR, Coebergh JW. Predictions of skin cancer incidence in the Netherlands up to 2015. *Br J Dermatol*. 2005;152:481-8.
2. Parkin D, Whelan S, Ferlay J. Cancer incidence in five continents VII. IACR Sci. Publ. No. 143. 1997. Lyon, International Agency for Research on Cancer.
3. Giles GG, Armstrong BK, Burton RC, Staples MP, Thursfield VJ. Has mortality from melanoma stopped rising in Australia? Analysis of trends between 1931 and 1994. *BMJ* 1996;312:1121-5.
4. Hollestein LM, van den Akker SA, Nijsten T, et al. Trends of cutaneous melanoma in the Netherlands: increasing incidence rates among all Breslow thickness categories and rising mortality rates since 1989. *Ann Oncol*. 2011 May 4, Epub ahead of print.
5. Bulliard JL. Site-specific risk of cutaneous malignant melanoma and pattern of sun exposure in New Zealand. *Int J Cancer* 2000;85:627-32.
6. Gallagher RP, Spinelli JJ, Lee TK. Tanning beds, sunlamps, and risk of cutaneous malignant melanoma. *Cancer Epidemiol Biomarkers Prev*. 2005;14:562-6.
7. Armstrong BK, Kricger A. Cutaneous melanoma. *Cancer Surv*. 1994;19-20:219-40.
8. Gaudette LA, Gao RN. Changing trends in melanoma incidence and mortality. *Health Rep*. 1998;10:29-41.
9. Testori A, Soteldo J, Sances D, et al. Cutaneous melanoma in the elderly. *Melanoma Res*. 2009;19:125-34.
10. Chang CK, Jacobs IA, Vizgirda VM, Salti GI. Melanoma in the elderly patient. *Arch Surg*. 2003 Oct;138:1135-8.
11. Kelly JW. Melanoma in the elderly- a neglected public health challenge. *Med J Aust*. 1998;169:403-4.
12. Balch CM, Buzaid AC, Soong SJ, et al. Final version of the American Joint Committee on Cancer staging system for cutaneous melanoma. *J Clin Oncol*. 2001;19:3635-48.
13. Francken AB, Shaw HM, Thompson JF, et al. The prognostic importance of tumor mitotic rate confirmed in 1317 patients with primary cutaneous melanoma and long follow-up. *Ann Surg Oncol* 2004;11:426-33.
14. De Vries E, Houterman S, Janssen-Heijnen ML, et al. Up-to-date survival estimates and historical trends of cutaneous malignant melanoma in the south-east of The Netherlands. *Ann Oncol* 2007;18:1110-6.
15. Lasithiotakis KG, Leiter U, Gorkievicz R, et al. The incidence and mortality of cutaneous melanoma in Southern Germany: trends by anatomic site and pathologic characteristics, 1976 to 2003. *Cancer* 2006;107:1331-9.
16. Buettner PG, Leiter U, Eigentler TK, Garbe C. Development of prognostic factors and survival in cutaneous melanoma over 25 years: an analysis of the Central Malignant Melanoma Registry of the German Dermatological Society. *Cancer* 2005;103:616-24.
17. De Vries E, Nijsten TE, Visser O, et al. Superior survival of females among 10,538 Dutch melanoma patients is independent of Breslow thickness, histologic type and tumor site. *Ann Oncol* 2008;19:583-9.
18. Koh HK, Miller DR, Geller AC, et al. Who discovers melanoma? Patterns from a population-based survey. *J Am Acad Dermatol*. 1992;26:914-9.
19. Francken AB, Shaw HM, Accortt NA, et al. Detection of first relapse in cutaneous melanoma patients: implications for the formulation of evidence-based follow-up guidelines. *Ann Surg Oncol* 2007;14:1924-33.
20. Shafir R, Hiss J, Tsur H, Bubis JJ. The thin malignant melanoma: changing patterns of epidemiology and treatment. *Cancer* 1982;50:817-9.
21. MacKie R, Hunter JA, Aitchison TC, et al. Cutaneous malignant melanoma, Scotland, 1979-89. The Scottish Melanoma Group. *Lancet* 1992;339:971-5.
22. Criscione VD, Weinstock MA. Melanoma thickness trends in the United States, 1988-2006. *J Invest Dermatol*. 2010;130:793-7.
23. Melia J, Cooper EJ, Frost T, et al. Cancer Research Campaign health education programme to promote the early detection of cutaneous malignant melanoma. *Br J Dermatol*. 1995;132:414-21.
24. Hersey P, Sillar RW, Howe CG, et al. Factors related to the presentation of patients with thick primary melanomas. *Med J Aust*. 1991;154:583-7.
25. Hanrahan PF, Hersey P, Menzies SW, et al. Examination of the ability of people to identify early changes of melanoma in computer-altered pigmented skin lesions. *Arch Dermatol* 1997; 133: 301-311
26. Austin PF, Cruse CW, Lyman G, et al. Age as a prognostic factor in the malignant melanoma population. *Ann Surg Oncol* 1994; 1:487-494.

27. Cohen HJ, Cox E, Manton K, et al. Malignant melanoma in the elderly. *J Clin Oncol* 1987; 5:100-106.
28. Loggie B, Ronan SG, Bean J, et al. Invasive cutaneous melanoma in elderly patients. *Arch Dermatol* 1991; 127:1188-1193.
29. Rassaby J, Larcombe I, Hill D, Walker M. Slip, Slop, Slap! Health education about skin cancer. *Cancer Forum* 1983;7:63-9.
30. Levi F, Franceschi S, Te VC, Randimbison L, La VC. Trends of skin cancer in the Canton of Vaud, 1976-92. *Br J Cancer* 1995;72:1047-53.

# 5

---



## S-100B concentrations predict disease-free survival in stage III melanoma patients | S Kruijff, E Bastiaannet, AC Muller Kobold,

RJ van Ginkel, AJH Suurmeijer, HJ Hoekstra | Ann Surg Oncol. 2009;16:3455-62.

---

### ABSTRACT

**Background:** Elevation of the tumor marker S-100B in melanoma patients is a highly specific indicator of recurrence.

**Materials and methods:** The role of S-100B in disease free survival (DFS) was evaluated in stage III melanoma patients (staged with fluorodeoxyglucose positron emission tomography [FDG-PET] and computed tomography [CT]) with palpable lymph node metastases who underwent therapeutic lymph node dissection. S-100B and LDH were measured on the day before surgery (d= -1) and on days 1, 2, and 7 postoperatively. Multivariate logistic regression was used to study factors associated with preoperative elevation of S-100B. Univariate (log-rank test) and multivariate (Cox regression) survival analyses were performed to identify factors associated with DFS.

**Results:** Between 2004 and 2008, 56 patients (median age 57, range 24-93) years, 27 males (48%) and 29 females (52%) entered the study. Preoperative S-100B elevation was found in 27 patients (48%) and elevated LDH in 20 patients (36%). No association was found between these two markers at any time. Multivariate analysis showed that elevated S-100B preoperatively (hazard ratio [HR] 2.7, p=0.03) was associated with DFS. S-100B elevation was associated with increased tumor size (odds ratio [OR] 3.40; p=0.03).

**Conclusion:** Elevated S-100B preoperatively in patients with optimally staged clinical stage III melanoma is associated with decreased disease-free survival. S100-B could be used as a prognostic marker in the stratification of new adjuvant trials to select stage III melanoma patients for adjuvant systematic treatment.

## INTRODUCTION

The incidence of melanoma is still increasing in the Netherlands. Between 1989 and 2003, the incidence increased from 9.5 to 13.7/100,000 in men and from 13.4 to 18.5/100,000 in women.<sup>1</sup> About 90% of the patients had stage I or II melanoma.<sup>2</sup> After surgical treatment, 20-28% of melanoma patients present with locoregional recurrence or regional dermal metastases, 26-60% with regional recurrences, and 15-50% with distant metastases.<sup>3</sup> Despite improved public awareness as well as early diagnosis and treatment, mortality from melanoma increased from 2.5 to 3.6/100,000 in men and from 2.0 to 2.4/100,000 in women in the Netherlands between 1989 and 2003.<sup>4</sup> It is impossible to predict the metastatic behavior of melanoma in individual patients. At present, the Multicentre Selective Lymphadenectomy Trial (MSLT I) is investigating whether sentinel lymph node biopsy reduces the risk of locoregional metastases and improves survival.<sup>4</sup> In patients with clinical stage III melanoma (palpable lymph node metastases), 5-year survival varied between 26 and 67%.<sup>5</sup> Recent staging studies with whole-body fluorodeoxyglucose positron emission tomography (FDG-PET) and/ or spiral computed tomography (CT) in patients with clinical stage III melanoma have shown that about one quarter of these patients had distant metastases.<sup>6,7</sup> Furthermore, the uptake of FDG in the lymph node metastases in patients with clinical stage III melanoma proved to be of prognostic value.<sup>8</sup> FDG-PET or spiral CT improved the staging of clinical stage III melanoma and led to better selection of patients for therapeutic lymph node dissection.<sup>6,7</sup> Serum lactate dehydrogenase (LDH) had high specificity for melanoma, but low sensitivity. It was an independent prognostic factor in stage IV melanoma.<sup>9-12</sup> Increased S-100B concentrations were first detected in melanoma patients in 1980.<sup>13</sup> Later, S-100B was found to be a serological tumor marker for melanoma and is mostly increased in stage III and IV.<sup>14,15</sup> S-100B is a 21 kilodalton (kDa) protein that was first isolated from the central nervous system in vertebrates. The name is derived from its 100% solvency in saturated ammonium sulfate with neutral pH. It has calcium-binding properties and as a dimer, consists of two isomers a and b. All possible combinations can occur (S-100aa, S-100ab, S-100bb).<sup>15</sup> The S-100B protein is of neuroectodermal and mesodermal origin and is expressed in various parts of the body. S-100B is chiefly found in glial cells and Schwann cells.<sup>16</sup> Serum S-100B can be measured with an immunoradiometric assay (Sangtec 100 IRMA/RIA assay) or with an immunoluminometric assay (LIA Mat Sangtec 100). This study addressed the question whether the perioperative measurement of the tumor markers S-100B has prognostic value in FDG-PET and spiral CT staged patients with stage III melanoma who are selected for therapeutic lymph node dissection.

## MATERIALS EN METHODS

### *Patients*

Melanoma patients, earlier staged AJCC I and II, now presenting with clinically and cytologically proven regional nodal metastases of melanoma (AJCC stage IIIb and IIIc) were staged with whole-body FDG-PET (Siemens ECAT EXACT HR + scanner) and with 64-slice spiral CT (Siemens Somatom Sensation). Sentinel node positive patients and patients with regional dermal metastases were excluded for this study. The patients whose test results were negative for distant metastases were eligible for therapeutic lymph node dissection with curative intent. Nodal metastases of  $\geq 3$  cm and/or C 3 positive lymph nodes and/or extra nodal growth received adjuvant radiotherapy (20 9 2.4 Gy).<sup>17,18</sup> LDH and S-100B were measured preoperative (day -1) and postoperative days 1, 2, and 7. The study was approved by the Medical Ethical Committee of the UMCG.

### *Methods*

LDH was analyzed routinely by means of Roche Modular (Hitachi) with an enzymatic activity measurement; normal values of LDH were considered to be  $> 250$  U/L.

Concentrations of S-100B were measured using Sangtec 100 immunoassay on the Advantage (Nichols). The S-100B half-life is estimated to be 30 min.<sup>19</sup> Approximately 10 times the half-life after lymph node dissection, S-100B values should reduce to the minimum possible. When tumor cells are still present the release of S-100B will continue or may even increase. Therefore, S-100B levels were measured on days -1, 1, 2, and 7. Levels were calculated on the basis of a calibration curve and checked against internal standards with a known concentration of S-100B. In this assay, a cut-off point of 0.15  $\mu\text{g/L}$  was set. The reference values for the S-100B assay (Liason Sangtec 100) were established by analysis of S-100B values of 120 healthy men and women according to the CLSI C28A2 guideline. Concentrations within this range were considered to be normal, whereas concentration above this range were considered to be elevated. A postoperative increase of S-100B was defined as any increase on the postoperative days 1, 2, or 7 (day 1 as baseline value, elevation on day 2 compared with day 1, and/ or elevation on day 7 compared with day 2).

### *Statistics*

Data were analyzed with STATA version 10.0. Logistic regression analysis was used to assess factors associated with elevated S-100B concentrations. Factors associated with disease-free survival (DFS) were analyzed with standard univariate (log-rank test) and multivariate (Cox proportions hazard) survival methods.



**TABLE 1: Characteristics of 56 AJCC stage III melanoma patients**

<b>Variable</b>	<b>No. of patients (%)</b>
Gender	
M	27 (48.2)
F	29 (51.8)
Median age (years)	57.0
Localisation primary tumour	
Trunk	24 (42.8%)
Head & neck	3 (5.3%)
Upper extremities	7 (12.5%)
Lower extremities	22 (55.8%)
Unknown primary	0
Breslow thickness primary tumour	
T1 (<1.00 mm)	7 (12.5%)
T2 (1.00-2.00 mm)	20 (35.7%)
T3 (2.00-4.00 mm)	21 (37.5%)
T4 (>4.00 mm)	8 (14.2%)
Clark level primary tumour	
1-3	24 (42.6%)
4-5	31 (55.4%)
unknown	1 (1.8%)
Localisation lymph nodes	
neck	6 (10.7%)
axilla	20 (35.7%)
groin	30 (53.5%)
No. of nodes removed	
< 15	27 (48.2%)
≥ 15	29 (52.8%)
No. of positive nodes	
< 2	36 (64.3%)
≥ 2	20 (35.7%)
Extranodal growth	
Yes	24 (42.6%)
No	42 (56.4%)
Tumour size	
< 3 cm	28 (50%)
≥ 3 cm	28 (50%)
Ulceration primary tumour	
Yes	13 (23.2%)
No	43 (76.8%)
Comorbidity	
Yes	16 (28.6%)
No	40 (71.4%)

## RESULTS

### *Patients*

In the period from January 2004 to January 2008, 56 patients, 27 males (48%) and 29 females (52%), median age 57 (range 24-93) years underwent therapeutic lymph node dissection (Table 1). Preoperatively, 27 patients had elevated S-100B concentrations (48%), while 20 patients had increased LDH levels (36%). The range on day -1, 1, 2, and 7 for S-100B was 0.03–23.3, 0.02–1.5, 0.03–0.82, 0.01–0.22 and for LDH 109–428, 106–816, 108–335, 115–802. No associations were found between S-100B and LDH levels on days -1, 1, 2, and 7 ( $p=0.45$ ,  $p=0.87$ ,  $p=0.84$ , and  $p=0.90$ , respectively). The S-100B concentration normalized postoperatively in 22 of the 27 patients (81.4%) and the LDH in 18 of the 20 patients (90%).

Preoperative S-100B elevation was associated with increased lymph node size (odds ratio [OR] 3.4,  $p=0.03$ ), however not significantly with extranodal growth ( $p=0.8$ ), patients (43%), of which 17 patients were diagnosed with distant metastasis and 7 patients with a regional recurrence. As shown in Table 3, DFS was not associated with sex, localization of the lymph node, or the size of the lymph nodes. Two-year DFS in patients with elevated preoperative S-100B concentrations was 34% versus 61% for patients without elevated preoperative S-100B concentrations (hazard ratio [HR] 2.6,  $p=0.03$ ). In the patients whose S-100B increased postoperatively, 2-year DFS was 30% versus 51% for patients without a postoperative S-100B increase (HR 2.0,  $p=0.1$ ) (Figure 1). In multivariate analysis (model with positive lymph nodes, extranodal growth, preoperative and postoperative S-100B and substage IIb or IIc), extranodal growth (HR 0.4,  $p=0.05$ ), and elevated preoperative S-100B concentrations (HR 2.6,  $p=0.03$ ) were significantly associated with decreased DFS. Patients with increasing postoperative S-100B serum levels showed a trend toward statistical significance when correlated to DFS (HR 2.6,  $p=0.07$ ).

**TABLE 2:** Logistic regression analysis on elevated S100-b in stage III melanoma patients

Variable	Elevated S-100b %	OR	p-value
Sex			
Male	51.9	Ref	
Female	44.8	0.75 (0.26-2.16)	0.599
Age			
< 55 years	44.0	Ref	
≥ 55 years	51.6	1.36 (0.47-3.91)	0.571
Positive lymph nodes			
≤ 2	44.4	Ref	
> 2	55.0	1.52 (0.51-4.59)	0.450
Extranodal growth			
Yes	52.2	Ref	
No	48.4	0.86 (0.29-2.53)	0.783
Tumour size			
< 3	35.7	Ref	
≥ 3	65.4	3.40 (1.11-10.4)	0.032


**TABLE 3:** Disease-free survival (DFS) of AJCC stage III melanoma patients

Variable		%	2 year DFS (%)	Univariate analysis		Multivariate analysis	
				HR (95% CI)	p-value	HR (95% CI)	p-value
Sex	Male	48.2	45.1	Ref			
	Female	51.8	47.2	1.16 (0.52-2.59)	0.713		
Age (yrs)	< 55	44.6	46.7	Ref			
	≥ 55	55.4	46.7	1.10 (0.49-2.48)	0.821		
Co-morbidity	Yes	28.6	32.3	Ref			
	No	71.4	54.9	0.82 (0.36-1.89)	0.649		
Clark level	1-3	44.6	46.1	Ref			
	4-5	55.4	45.9	0.81 (0.36-1.81)	0.608		
Breslow thickness	< 2	48.2	52.7	Ref			
	≥ 2	51.8	40.7	1.30 (0.58-2.91)	0.519		
Localisation LN	Neck	10.7	83.3	Ref			
	Axilla	35.7	50.9	3.29 (0.41-26.4)			
	Groin	53.6	35.8	4.94 (0.65-37.5)		0.237	
No. of LNs	< 15	48.2	50.8	Ref			
	≥ 15	51.8	43.8	0.93 (0.41-2.11)	0.869		
Positive LNs	≤ 2	64.3	63.5	Ref		Ref	
	> 2	35.7	22.5	3.32 (1.47-7.50)	0.004	1.98 (0.81-4.87)	0.135
Ulceration	Yes	76.8	30.8	Ref			
	No	23.2	50.8	1.59 (0.66-3.85)	0.303		
Extranodal growth	Yes	42.6	28.4	Ref		Ref	
	No	57.4	57.9	0.37 (0.16-0.84)	0.018	0.32 (0.13-0.81)	0.015
LN size (cm)	< 3	49.0	53.2	Ref			
	≥ 3	51.0	34.5	1.45 (0.65-3.25)	0.365		
Pre-operative S100-b	Normal	51.8	60.6	Ref		Ref	
	Elevated	48.2	34.3	2.56 (1.09-6.00)	0.030	2.69 (1.13-6.44)	0.026
Peri-operative S100-b	No increase	82.1	51.3	Ref		Ref	
	Increase	17.9	30.0	2.00 (0.83-4.81)	0.124	3.04 (1.14-8.13)	0.027

LN=Lymph node

## DISCUSSION

This is the first study that demonstrates a potential prognostic value of preoperative S-100B concentrations in patients who underwent a therapeutic lymph node dissection after preoperative staging with FDG-PET and CT. According to the literature, the proportions of patients with elevated S-100B concentrations were 0-9% in stage I/II, 5-98% in stage III, and 40-100% in stage IV melanoma (Table 4). The highest S-100B concentrations were found in patients with liver and/or skeletal metastases.<sup>20,21</sup> Serum S-100B concentrations are therefore correlated with the clinical stage of the disease. The aim of this study was to evaluate the



**FIGURE 1** Kaplan-Meier curve for preoperative S-100B and disease-free survival of AJCC stage III melanoma patients. **A** Elevated preoperative S-100B (34% 2-year DFS) versus not elevated (61% 2-year DFS) levels are associated with a decreased disease-free survival ( $p=0.03$ ). **B** Increasing postoperative S-100B levels (30% 2-year DFS) versus not increasing (51% 2-year DFS) ( $p=0.1$ )

**TABLE 4:** Review literature S100-B values above cut-off point in melanoma patients

Reference	N (patients)	Assay	Cut- off Value	Stage I/II	Stage III	Stage IV
Guo et al. 1995 <sup>12</sup>	126	IRMA	0.15	1%	9%	74%
Miliotes et al. 1996 <sup>34</sup>	67	IRMA	0.05	47.8%	-	-
Abraha et al. 1997 <sup>35</sup>	97	IRMA	0.2	9%	82%	-
Bosserhof et al. 1997 <sup>36</sup>	112	IRMA	0.15	0%	62%	-
Buer et al 1997 <sup>21</sup>	99	IRMA	3.00	-	-	22.2%
Henze et al. 1997 <sup>26</sup>	73	IRMA	0.3	4%	21%	79%
Tofani et al. 1997 <sup>37</sup>	53	IRMA	0.5	0%	55%	-
Schultz et al. 1998 <sup>38</sup>	84	IRMA	0.3	0%	31%	69%
Bonfrer et al 1998 <sup>39</sup>	251	LIA	0.16	I/II:1%	-	III/IV: 79%
Seregni et al. 1998 <sup>40</sup>	438	IRMA	0.2	I:4.2%,II:5.3%	38.5%	-
Hauschild et al. 1999 <sup>41</sup>	412	IRMA	0.2	2%	19%	68%
Curry et al 1999 <sup>19</sup>	147	IRMA/LIA	0.2 / 0.12	-	23%/47.5%	-
Berking et al. 1999 <sup>42</sup>	352	LIA	0.2	7%	8%	48%
Jackel et al. 1999 <sup>14</sup>	276	LIA	0.12	6%	5%	48%
Kaskel et al. 1999 <sup>43</sup>	570	LIA	0.11	9%	94%	-
Brouard et al. 2000 <sup>44</sup>	122	LIA	0.09	0%	54%	84%
Vuoristo et al 2000 <sup>45</sup>	50	LIA	0.12	-	-	64.0%
Jury et al. 2000 <sup>46</sup>	214	LIA	0.2	7%	29%	92%
Mohammed et al. 2001 <sup>18</sup>	68	LIA	0.2	-	-	73.5%
Martenson et al. 2001 <sup>17</sup>	727	LIA	0.10	12%	-	III/IV: 72%
Juergenson et al 2001 <sup>47</sup>	50	LIA	0.12	-	-	I/II/III/IV:50%
Rebmann et al 2002 <sup>48</sup>	183	LIA	0.12	-	-	I/II/III/IV: 21%
Banfalvi et al. 2002 <sup>27</sup>	59	LIA	0.12	40%	25%	59%
Garbe et al 2003 <sup>29</sup>	296	LIA	0.12			
Banfalvi et al. 2003 <sup>28</sup>	478	LIA	0.18	I:29%, II: 19%	32%	48%
Andres et al 2004 <sup>49</sup>	85	LIA	0.15	-	I/II/III:15%	89%
Ugurel et al 2005 <sup>50</sup>	300	LIA	0.12	-	-	I/II/III/IV: 20%
Smitt et al 2005 <sup>13</sup>	145	LIA	0.16	-	-	78%
Schmidt et al 2005 <sup>51</sup>	85	LIA	0.15	-	-	59%
Domingo-et al 2007 <sup>52</sup>	97	LIA	0.15	-	II/III: 21%	-
Cao et al 2007 <sup>53</sup>	42	LIA	0.20	-	-	50%
Tarhini et al 2008 <sup>33</sup>	670	LIA	0.15	-	IIb/III: 13%	-
Kruijff et al 2009	56	LIA	0.15	-	48%	-

(LIA: Lumino imunometric Assay, IRMA: Immuno Radiometric Assay)

prognostic value of preoperative and postoperative concentrations of the tumor marker S-100B in clinically stage III melanoma patients who underwent a so-called “curative lymph node dissection” after preoperative staging with whole-body FDG-PET and spiral CT. Our results showed that preoperative elevated concentrations of S-100B was statistically significantly associated with a decreased DFS. It is difficult to compare the studies investigating the

value of S-100B in patients with melanoma, because various assays (IRMA or LIA-mat) and cut-off points and groups of patients with different stages were used (Table 4).<sup>8</sup> Garbe et al. also found a relation between S-100B and disease-free survival in stage II and III melanoma patients. Postoperative elevated S-100B preceded detection of developing metastases by imaging techniques.<sup>2</sup> According to Martenson et al. and Curry et al., elevated S-100B in stage III melanoma after lymph node dissection may form an independent prognostic factor.<sup>20,22</sup> Four studies demonstrated that elevated S-100B was associated with shortened (disease-free) survival.<sup>20,23-25</sup> In stage IV patients with normal S-100B levels, survival was better when compared to those with elevated S-100B levels.<sup>15,26</sup> Increasing concentrations of S-100B in stage IV patients who were receiving systemic treatment were associated with disease progression.<sup>14,27-29</sup> In a study on 1007 melanoma patients, S-100B was not only correlated with the clinical stage of the disease, but it also appeared to be an independent prognostic marker in stages II and III. The 5-year survival of patients with S-100B values of  $< 0.10 \mu\text{g/L}$  was 91%, compared with 51% when the values were  $\geq 0.10 \mu\text{g/L}$ .<sup>20</sup> In two other studies, similar patterns were found.<sup>30,31</sup>

Two studies comparing 5-S-cysteinyl-dopa, LDH and S-100B as prognostic markers found that S-100B had the highest specificity.<sup>30,31</sup> Several studies have therefore recommended the use of S-100B as a tumor marker to monitor the course of disease in stage III melanoma patients and to evaluate the effect of therapy in stage IV patients. Successful treatment with lymph node dissection, chemotherapy, or immunotherapy was associated with decreased S-100B concentrations, whereas increased concentrations were an expression of disease progression.<sup>14,27,28</sup> The prognostic value of the preoperative and postoperative course of S-100B concentrations in "true stage III" melanoma patients has not been studied before. In advance, we hypothesized that a significant decrease in S-100B concentration after therapeutic lymph node dissection would correlate with better DFS. In this way we assumed S-100B could be used as a quality-control marker for dissection. In our study, 27 patients (48.2%) had elevated S-100B preoperatively, and indeed in 22 of them (81.4%) the concentrations normalized postoperatively. However, 12 patients of these 22 (55%) developed tumor recurrence during a median follow-up of 1.18 years. Multivariate analyses showed that preoperative S-100B elevation was an important predictor of decreased DFS ( $p=0.03$ ).

Although the results indicated that therapeutic lymph node dissection did have a temporary influence on the postoperative course of S-100B, a more important predictor for shortened DFS seemed preoperative S-100B elevation. For AJCC stage III, postoperative increasing S-100B serum were associated with a decreased DFS. However, when adjusted for stage III substages, probably because of small numbers, only a trend toward statistical significance was observed. S-100B is located in the cytoplasm of melanoma cells. Elevated serum S100-B occurs as the result of loss of melanoma cell integrity.<sup>19</sup> Therefore, the hypothesis that elevation of S-100B should be interpreted as a process of subclinical microscopic metastatic disease seems to be suggested by these data. Preoperative elevated S-100B could be an expression of early dissemination not detected by standard imaging tests.<sup>33</sup> However, the tumor marker is not suitable for screening purposes or for the early detection of recurrence in the follow-up of

melanoma patients.<sup>34</sup> The prognostic biochemical marker S-100B may be of value in the design of future trials on adjuvant systemic chemotherapy or immunotherapy. At present, LDH is the most prominent serum parameter in stage IV melanoma, and it has been included in the new AJCC staging system.<sup>5</sup> Our study on optimally staged clinical stage III melanoma patients has now shown that preoperative S-100B elevation is associated with significantly poorer survival. These results are supported by studies from M.D. Anderson Cancer Center, Barcelona and a recent meta-analysis and a pooled analysis of ECOG studies in high-risk surgically resected melanoma by Kirkwood.<sup>33,35-37</sup> These studies showed that direct postoperative (baseline) S-100B is indeed a prognostic marker for disease-free survival (DFS) and overall survival (OS). Our results seem in accordance with these data, but more importantly reveal that S-100B is a prognostic marker for DFS when determined before dissection. In our study, the 2-year DFS is slightly better than in the recently published ECOG, which is explained by the fact that the patients were staged with FDG-PET and spiral CT.<sup>37</sup> A more refined individualized risk assessment for adjuvant treatment is possible with serum marker S-100B. Therefore, S-100B could be included in future stratification research for stage III melanoma. In summary, preoperative elevation of S-100B levels in patients with true (clinical) stage III melanoma is associated with decreased disease-free survival and might have a prognostic clinical value in high-risk stage III melanoma patients. Therefore S-100B could be used to select patients for adjuvant systematic treatment in the stratification for new adjuvant therapeutic trials as well as to provide information for stage III melanoma patients who want to be informed about disease prognosis.

## REFERENCES

- Siesling S, Visser O, van Dijk JA, Coebergh JW. Trends in the incidence and death from cancer from 1989-2003 in The Netherlands. *Ned Tijdschr Geneesk.* 2006;150:2490-6.
- Garbe C, Eigentler TK. Diagnosis and treatment of cutaneous melanoma: state of the art 2006. *Melanoma Res.* 2007;17:117-27.
- Francken AB, Bastiaannet E, Hoekstra HJ. Follow-up in patients with localised primary cutaneous melanoma. *Lancet Oncol.* 2005;6:608-21.
- Morton DL, Thompson JE, Cochran AJ, Mozzillo N, Elashoff R, Essner R, et al. Sentinel-node biopsy or nodal observation in melanoma. *N Engl J Med.* 2006;355:1307-17.
- Balch CM, Soong SJ, Gershenwald JE, Thompson JE, Reintgen DS, Cascinelli N, et al. Prognostic factors analysis of 17,600 melanoma patients. Validation of the American Joint Committee on Cancer melanoma system. *J Clin Oncol.* 2001;19:3622-34.
- Bastiaannet E, Oyen WJ, Meijer S, Hoekstra OS, Wobbles T, Jager PL, et al. Impact of 18F fluorodeoxyglucose positron emission tomography on surgical management of melanoma patients. *Br J Surg.* 2006;93:243-9.
- Bastiaannet E, Wobbles T, Hoekstra OS, van der Jagt EJ, Brouwers AH, Koelemij R, et al. Prospective comparison of FDG-PET and CT in 251 melanoma patients with palpable lymph node metastases; diagnostic accuracy and impact on treatment. *J Clin Oncol.* 2009;27:4774-80.
- Bastiaannet E, Hoekstra OS, Oyen WJ, Jager PL, Wobbles T, Hoekstra HJ. Level of fluorodeoxyglucose uptake predicts risk for recurrence in melanoma patients presenting with lymph node metastases. *Ann Surg Oncol.* 2006;13:919-26.
- Brochez L, Naeyaert JM. Serological markers for melanoma. *Br J Dermatol.* 2000;143:256-68.
- Finck SJ, Giuliano AE, Morton DL. LDH and melanoma. *Cancer.* 1983;51:840-3.
- Karakousis CP, Balch CM, Urist MM, Ross MM, Smith TJ, Bartolucci AA. Local recurrence in malignant melanoma: longterm results of the multiinstitutional randomized surgical trial. *Ann Surg Oncol.* 1996;3:446-52.
- Sirott MN, Bajorin DF, Wong GY, Tao Y, Chapman PB, Templeton MA, et al. Prognostic factors in patients with metastatic malignant melanoma. A multivariate analysis. *Cancer.* 1993;72:3091-8.
- Fagnart OC, Sindic CJ, Laterre C. Particle counting immunoassay of S100 protein in serum. Possible relevance in tumors and ischemic disorders of the central nervous system. *Clin Chem.* 1988;34:1387-91.
- Guo HB, Stoffel-Wagner B, Bierwirth T, Mezger J, Klingmuller D. Clinical significance of serum S100 in metastatic malignant melanoma. *Eur J Cancer.* 1995;31A:924-8.
- Smit LH, Korse CM, Hart AA, Bonfrer JM, Haanen JB, Kerst JM, et al. Normal values of serum S-100B predict prolonged survival for stage IV melanoma patients. *Eur J Cancer.* 2005;41:386-92.
- Jackel A, Deichmann M, Waldmann V, Bock M, Naether H. S-100 beta protein in serum, a tumor marker in malignant melanoma-current state of knowledge and clinical experience. *Hautarzt.* 1999;50:250-6.
- Bastiaannet E, Beukema JC, Hoekstra HJ. Radiation therapy following lymph node dissection in melanoma patients: treatment, outcome and complications. *Cancer Treat Rev.* 2005;31:18-26.
- Henderson MA, Burmeister B, Thompson JE, Di Iulio J, Fisher R, Hong A, et al. Adjuvant radiotherapy and regional lymph node field control in melanoma patients after lymphadenectomy: Results of an intergroup randomized trial. 2009 ASCO Annual meeting, *J Clin Oncol.* 2009;27 Supplement: Abstract No.LBA9084.
- Ghanem G, Loir B, Morandini R, Sales F, Lienard D, Eggermont A, et al. On the release and half life of S100B protein in the peripheral blood of melanoma patients. *Int J Cancer.* 2001; 94:586-90.
- Martenson ED, Hansson LO, Nilsson B, von Schoultz E, Ma'nsson, Brahme E, Ringborg U, et al. Serum S-100b protein as a prognostic marker in malignant cutaneous melanoma. *J Clin Oncol.* 2001;19:824-31.
- Mohammed MQ, Abbraha HD, Sherwood RA, MacRae K, Retsas S. Serum S100beta protein as a marker of disease activity in patients with malignant melanoma. *Med Oncol.* 2001;18:109-20.




22. Curry BJ, Farrelly M, Hersey P. Evaluation of S-100beta assays for the prediction of recurrence and prognosis in patients with AJCC stage I–III melanoma. *Melanoma Res.* 1999;9:557-67.
23. Bonfrer JM, Korse CM, Nieweg OE, Rankin EM. The luminescence immunoassay S-100: a sensitive test to measure circulating S-100B: its prognostic value in malignant melanoma. *Br J Cancer.* 1998;77:2210-4.
24. Buer J, Probst M, Franzke A, Duensing S, Haindl J, Volkenandt M, et al. Elevated serum levels of S100 and survival in metastatic malignant melanoma. *Br J Cancer.* 1997;75:1373-6.
25. Hansson LO, von SE, Djureen E, Hansson J, Nilsson B, Ringborg U. Prognostic value of serum analyses of S-100 protein beta in malignant melanoma. *Anticancer Res.* 1997;17:3071-3.
26. Bonfrer JM, Korse CM, Israels SP. Serum S-100 has prognostic significance in malignant melanoma. *Anticancer Res.* 1997;17:2975-7.
27. Hamberg AP, Korse CM, Bonfrer JM, de Gast GC. Serum S100B is suitable for prediction and monitoring of response to chemoimmunotherapy in metastatic malignant melanoma. *Melanoma Res.* 2003;13:45-9.
28. Hauschild A, Engel G, Brenner W, Glaeser R, Moenig H, Henze E, et al. Predictive value of serum S100B for monitoring patients with metastatic melanoma during chemotherapy and/or immunotherapy. *Br J Dermatol.* 1999;140:1065-71.
29. Henze G, Dummer R, Joller-Jemelka HJ, Boenig H, Burg G. Serum S100-a marker for disease monitoring in metastatic melanoma. *Dermatology.* 1997;194:208-12.
30. Banfalvi T, Boldizsar M, Gergye M, Gilde K, Kremmer T, Otto S. Comparison of prognostic significance of serum 5-S-Cysteinyldopa, LDH and S-100B protein in Stage III–IV malignant melanoma. *Pathol Oncol Res.* 2002;8:183-7.
31. Banfalvi T, Gilde K, Gergye M, Boldizsar M, Kremmer T, Otto S. Use of serum 5-S-CD and S-100B protein levels to monitor the clinical course of malignant melanoma. *Eur J Cancer.* 2003;39:164-9.
32. Garbe C, Leiter U, Ellwanger U, Blaheta HJ, Meier F, Rassner G, et al. Diagnostic value and prognostic significance of protein S-100beta, melanoma-inhibitory activity, and tyrosinase/MART-1 reverse transcription-polymerase chain reaction in the follow-up of high-risk melanoma patients. *Cancer.* 2003;97:1737-45.
33. Domingo-Domenech J, Castel T, Auge JM, Garcia-Albeniz XA, Conill C, Puig S, et al. Prognostic implications of protein S-100beta serum levels in the clinical outcome of high-risk melanoma patients. *Tumour Biol.* 2007;28:264-72.
34. Smit LH, Nieweg OE, Korse CM, Bonfrer JM, Kroon BB. Significance of serum S-100B in melanoma patients before and after sentinel node biopsy. *J Surg Oncol.* 2005;90:66-9.
35. Chun YS, Wang Y, Wang DY, McClain DM, Lucci A, Mansfield PF, et al. S100B levels are more accurate than LDH levels at predicting survival in melanoma patients. *Ann Surg Oncol.* 2008;15:96 (P211).
36. Mocellin S, Zavagno G, Nitti D. The prognostic value of serum S100B in patients with cutaneous melanoma: A meta-analysis. *Int J Cancer.* 2008;123:2370-6.
37. Tarhini AA, Stuckert J, Lee S, Sander C, Kirkwood JM. Prognostic significance of serum S100B surgically resected melanoma Patients Participating in Intergroup Trial ECOG 1694. *J Clin Oncol.* 2009;27:38-44.
38. Miliotes G, Lyman GH, Cruse CW, Puleo C, Albertini PA, Rapaport D, et al. Evaluation of new putative tumor markers for melanoma. *Ann Surg Oncol.* 1996;3:558-63.
39. Abraha HD, Fuller LC, Du Vivier AW, Higgins EM, Sherwood RA. Serum S-100 protein: a potentially useful prognostic marker in cutaneous melanoma. *Br J Dermatol.* 1997;137:381-5.
40. Bosserhoff AK, Kaufmann M, Kaluza B, Bartke I, Zirngibl H, Hein R, et al. Melanoma-inhibiting activity, a novel serum marker for progression of malignant melanoma. *Cancer Res.* 1997;57:3149-53.
41. Tofani A, Cioffi RP, Sciuto R, Rea S, Festa A, Di Filippo F, et al. S-100 and NSE as serum markers in melanoma. *Acta Oncol.* 1997;36:761-4.
42. Schultz ES, Diepgen TL, Von Den DP. Clinical and prognostic relevance of serum S-100 beta protein in malignant melanoma. *Br J Dermatol.* 1998;138:426-30.
43. Bonfrer JM, Korse CM. Monitoring malignant melanoma with the S-100B tumour marker. *Recent Results Cancer Res.* 2001;158:149-57.
44. Seregini E, Massaron S, Martinetti A, Illeni MT, Rovini D, Belli F, et al. S100 protein serum levels in cutaneous malignant melanoma. *Oncol Rep.* 1998;5:601-4.
45. Hauschild A, Engel G, Brenner W, Glaeser R, Moenig H, Henze E, et al. S100B protein detection in serum is a significant prognostic factor in metastatic melanoma. *Oncology.* 1999;56:338-44.

46. Berking C, Schlupen EM, Schrader A, Atzpodien J, Volkenandt M. Tumor markers in peripheral blood of patients with malignant melanoma: multimarker RT-PCR versus a luminoimmunometric assay for S-100. *Arch Dermatol Res.* 1999;291:479-84.
47. Kaskel P, Berking C, Sander S, Volkenandt M, Peter RU, Krahn G. S-100 protein in peripheral blood: a marker for melanoma metastases: a prospective 2-center study of 570 patients with melanoma. *J Am Acad Dermatol.* 1999;41:962-9.
48. Brouard M, Quillien V, Ollivier I, Lesimple T, Adamski H, Chevrant-Breton J. Serum S100B protein and stage of cutaneous melanoma: a prospective study. *Ann Dermatol Venereol.* 2000;127:56-9.
49. Vuoristo MS, Kellokumpu-Lehtinen P, Laine S, Parvinen LM, Hahka-Kemppinen M, Korpela M, et al. The value of serum S-100beta and interleukins as tumour markers in advanced melanoma. *Melanoma Res.* 2000;10:237-41.
50. Jury CS, McAllister EJ, MacKie RM. Rising levels of serum S100 protein precede other evidence of disease progression in patients with malignant melanoma. *Br J Dermatol.* 2000; 143:269-74.
51. Juergensen A, Holzapfel U, Hein R, Stolz W, Buettner R, Bosserhoff A. Comparison of two prognostic markers for malignant melanoma: MIA and S100 beta. *Tumour Biol.* 2001; 22:54-8.
52. Rebmann V, Ugurel S, Tilgen W, Reinhold U, Grosse-Wilde H. Soluble HLA-DR is a potent predictive indicator of disease progression in serum from early-stage melanoma patients. *Int J Cancer.* 2002;100:580-5.
53. Andres R, Mayordomo JI, Zaballos P, Rodino J, Isla D, Escudero P, et al. Prognostic value of serum S-100B in malignant melanoma. *Tumori.* 2004;90:607-10.
54. Ugurel S. Serum markers for melanoma. *Hautarzt.* 2005;56: 173-84.
55. Schmidt H, Sorensen BS, Fode K, Nexø E, von der Maase H. Tyrosinase messenger RNA in peripheral blood is related to poor survival in patients with metastatic melanoma following interleukin-2-based immunotherapy. *Melanoma Res.* 2005;15:409-16.
56. Cao MG, Auge JM, Molina R, Marti R, Carrera C, Castel T, et al. Melanoma inhibiting activity protein (MIA), beta-2 microglobulin and lactate dehydrogenase (LDH) in metastatic melanoma. *Anticancer Res.* 2007;27:595-9

# 6

---



# The value of pre-operative S-100B and SUV in clinically Stage III melanoma patients undergoing therapeutic lymph node dissection | S Kruijff, E Bastiaannet, MJ Speijers, AC Muller

Kobold, AH Brouwers, HJ Hoekstra | Eur J Surg Oncol. 2011;37:225-32

---

## ABSTRACT

**Introduction:** High preoperative serum S-100B values and Standardized Uptake Values (SUV) of Fluorodeoxyglucose (FDG) in PET for clinically stage III melanoma patients could be indicators of recurrence after surgical treatment. Aim was to assess the correlation and the prognostic value of these markers.

**Methods:** All melanoma patients with palpable nodal metastases, without distant metastases, were included from February 2004 to December 2007. Preoperative SUV and S-100B was determined. The correlation between SUV and S-100B and their relations with DFS and DSS were calculated by Cox Proportional Hazard Analysis.

**Results:** 62 Patients, median age 56.9 years, were included in the study. An elevated S-100B was found in 31 patients (50%) and elevated SUV in 24 patients (38.7%). No relation was found between S-100B and SUV. DFS was reduced (31.1%) for patients with an elevated S-100B (HR 3.1;  $p=0.02$ ) in comparison to a normal S-100B (44.6%). The DFS was 42.0% for patients with a SUV below the cut-off point and 29.0% for patients with an elevated SUV (HR 1.1;  $p=0.8$ ). DSS was 60.7% in a normal S-100B and 44.7% for patients with an elevated S-100B (HR 2.2;  $p=0.07$ ). DSS was 59.1% for patients with a normal SUV and 43.5% for patients with elevated SUV (HR 1.1;  $p=0.8$ ).

**Conclusion:** S-100B and SUV in stage III melanoma are not correlated and each have different associations with various histopathological factors. S-100B, in contrast with SUV, is associated with nodal tumor load, and when elevated, predicts a shorter DFS.

## INTRODUCTION

Melanoma causes more than 75% of all deaths related to skin cancer and its incidence has increased dramatically worldwide, especially in Caucasian populations.<sup>1</sup> During the period 1998-2007 the Dutch incidence increased from 13.2 to 20.0 newly diagnosed melanoma patients per 100,000 inhabitants.<sup>2</sup> The past 12 years, melanoma incidence rates have increased rapidly and are expected to keep on rising in the future. The absolute total number of new cases in the Netherlands is estimated to be well over 4,800 in 2015, compared to approximately 2400 cases in 2000.<sup>3</sup>

Despite being diagnosed in an earlier phase of disease, which may be a result of increased awareness and better surveillance, melanoma patients presenting with palpable nodal metastases today still have a poor 5-year survival of 59% and 40% for stage IIIB and IIIC respectively.<sup>4</sup> Staging studies with whole-body FDG-PET and/or spiral CT in patients with stage IIIB melanoma have shown that about 27% of patients are upstaged and treatment was changed for one of five patients.<sup>5</sup> Recently Balch et al analyzed 2,313 patients with AJCC stage III disease and for the complete cohort 5-year overall survival was 63%. However, when focusing on more specific patient groups, a tremendous heterogeneity in 5-year survival rates was observed (23%-87% 5-year survival).<sup>6</sup>

More than 60 years ago, Hill published about increased Lactate Dehydrogenase (LDH) as a prognostic serum marker in melanoma patients.<sup>7</sup> The last decades various biomarkers besides LDH have been studied, such as Melanoma Inhibitory Activity (MIA), S-100B and Standardized Uptake Value (SUV) in FDG-PET.<sup>8,9</sup> Weighing the evidence, LDH has high specificity for melanoma and literature demonstrates this marker to be elevated in advanced disease, predominantly in case of dissemination to the liver. However, today the most extensively studied melanoma biomarker is S-100B. This 21 kDa protein was first isolated from the central nervous system in vertebrates.<sup>10</sup> A preoperatively elevated serum S-100B in FDG-PET and CT evaluated patients with palpable nodal metastases is associated with a significantly worse survival.<sup>10-13</sup> Another prognostic marker was studied by assessment of the degree of Fluorodeoxyglucose (FDG) accumulation during Positron Emission Tomography in a melanoma metastasis; the Standardized Uptake Value (SUV).<sup>14</sup> Bastiaannet et al proved that the uptake of FDG in the lymph node metastases in melanoma patients with clinical stage III melanoma is of prognostic value.<sup>15</sup>

High values of S-100B and FDG Standardized Uptake Value (FDG-SUV) measured preoperatively in stage III melanoma patients could both be highly specific indicators of early recurrence after surgical treatment. Possibly, S-100B values and FDG-SUV form a measurable reflection of the presence of a subclinical process of dissemination. Therefore, the aim of this study was to assess the correlation between both markers and to study their association with Disease Free Survival (DFS) and Disease Specific Survival (DSS).

## MATERIAL AND METHODS

### *Patients*

Melanoma patients, previously staged as AJCC I or II, now presenting with palpable and pathologically proven lymph node metastases (AJCC stage IIIB or IIIC) were prospectively included in this study from February 2004 to December 2007. Patients were staged with whole-body FDG-PET and spiral CT. In case of negative test results for distant metastases, patients were eligible for therapeutic lymph node dissection (TLND) and were included in this study. Sentinel lymph node positive patients were excluded from this study. Patients received adjuvant radiotherapy (20 x 2.4 Gy) in case of a nodal metastasis of  $\geq 3$  cm in diameter,  $\geq 3$  tumor positive lymph nodes or the presence of extranodal growth.<sup>16,17</sup>

### *Patient characteristics and follow up*

Age, sex, treatment strategy of primary melanoma, date of primary melanoma diagnosis, characteristics of the primary melanoma (Breslow thickness, localization, ulceration status, Clark level) and characteristics of the lymph node metastases (localization, number of lymph nodes removed, number of tumor positive nodes, presence of extranodal growth, lymph node size and AJCC stage III B/C) were recorded, as well as the date of recurrence or death. Any form of melanoma recurrence after TLND was scored as recurrent disease.

Follow-up strategy was the same for all patients. Three monthly history and physical examination in the first year after TLND, 4 monthly in year 2, 6 monthly in years 3 to 5 and annual visits and chest radiographs for the subsequent five years.

### *PET protocol*

FDG-PET and CT were performed preoperatively in a random order. FDG was produced on site.<sup>18</sup> Before FDG injection, patients were instructed to fast for at least 6 hours and drink 1 liter of water. After intravenous injection of FDG (Range 220 to 690 MBq), whole body PET-imaging was performed; two/three dimensional mode, emission scans 5 minutes per bed position, starting 90 minutes after injection of FDG. Patients were scanned from scalp to feet, using a Siemens ECAT EXACT HR + scanner (Siemens/CTI, Knoxville, TN). FDG-PET readings were performed by attending staff nuclear medicine physicians.

### *CT protocol*

In the CT protocol a 64-slice spiral CT (Siemens Somatom Sensation) was performed of neck, chest and abdomen. Before starting the procedure oral (800 ml) and intravenous contrast agents were administered to the patient using standard imaging protocols. CT images were interpreted by attending staff radiologists.

### *Standardized Uptake Value (SUV)*

The SUV depends on the amount of injected radioactivity, the patient's weight and the calibration factor of the camera. The value is calculated according to the following formula: SUV mean = radioactivity concentration in tissue (Bq/Kg) / (injected dose [Bq] / patient weight [Kg]). Three dimensional regions of interest were placed semi-automatically over the tumor on multiple slices using a software program and a threshold of 70 % of the maximum pixel value within the tumor. In case of multiple metastases in the lymph node basin, the lesion with the most intense uptake was analyzed. For SUV the median of 8.4 was used as cut-off value in absence of a standardized cut-off value. Currently no cut-off point is described in literature although initiatives have been undertaken to create a new protocol using standardized uptake values.<sup>19</sup>

### *S-100B*

Serum S-100B levels were calculated on the basis of a calibration curve and checked against internal standards with a known concentration of S-100B. In this assay, a cut-off point of 0.15 µg/l was set. The reference values for the S-100B assay (Liason Sangtec 100) were established by analysis of S-100B values of 120 healthy men and women according to the CLSI C28A2 guideline. Concentrations below the cut-off point were considered to be normal, whereas concentration above this point were considered to be elevated. S-100B levels were measured one day before operation.

### *Statistics*

The association between elevated S-100B and elevated SUV was tested with a linear regression model and a scatter plot. Factors correlated with DFS and DSS were analyzed with univariate (Log rank test) and multivariable (Cox Proportional Hazard) tests. Factors associated with elevated S-100B concentrations and elevated SUV were tested using univariate logistic regression. Finally, a stratified survival analysis for DFS and DSS was performed to assess the strata in which the marker S-100B has the highest prognostic value. Data were analyzed with STATA version 10.0. A difference was considered significant when  $p \leq 0.05$ .

## **RESULTS**

### *Patient characteristics*

Overall, 62 patients (30 males and 32 females), with a median age of 56.9 (range 24.7-93.2) years, were included in the study. The majority of lymph node metastases were found in the groin (48.3%), followed by the axilla (35%) and the neck (16.7%). Size of the lymph node metastases was less than 3.0 cm in 45.2% and 3.0 cm or bigger in 54.8% of patients. An extranodal growth pattern was found in 37.1% of cases. Patient and tumor characteristics are summarized in Table 1.

**TABLE 1:** Characteristics of 62 AJCC stage III melanoma patients unthergoing lymph node dissection

		N	%
Sex	Male	30	48.4
	Female	32	51.6
Age (yrs)	≤45	16	25.8
	46-64	28	45.2
	≥65	18	29.0
Breslow thickness	≤2.0	30	48.4
	>2.0	32	51.6
Clark Level	I-III	17	27.4
	IV-V		58.1
	Unknown	9	14.5
Ulceration	No	51	82.3
	Yes	11	17.7
Region	Neck	10	16.7
	Axilla	21	35.0
	Groin	29	48.3
Number removed	<15	27	43.6
	≥15	35	56.4
Positive nodes	<2	26	41.9
	≥2	36	58.1
Lymph node size	<3.0	28	45.2
	≥3.0	34	54.8
Extranodal growth	No	39	62.9
	Yes	23	37.1

### *S-100B and SUV*

Thirty-one patients (50%) had a normal preoperative S-100B level (< 0.15 µg/l) and thirty-one patients (50%) had an elevated S-100B level (≥ 0.15 µg/l). Thirty-eight patients (61.3%) had a low preoperative SUV (<8.4) and twenty-four patients (38.7%) had a high preoperative SUV (≥8.4). No association was found between preoperative S-100B and SUV (p=0.7) in a linear regression model.

### *Disease free survival (DFS)*

For DFS, recurrent disease was interpreted as any form of melanoma recurrence. However, all patients who returned with recurrent disease after TLND had distant metastases and none of the patients presented with local or regional recurrences.



**TABLE 2:** Univariate and multivariate analysis on Disease-Free Survival (DFS) for SUV and S100-B with any recurrence as event.

Characteristic		3-years DFS	Univariate HR	Univariate p-value
Breslow	≤2.0	44.9	Ref	0.09
Thickness	>2.0	30.0	1.7 (0.9-3.3)	
Melanoma	No	43.2	Ref	0.01
Ulceration	Yes	11.4	2.5 (1.2-5.3)	
Lymph node positive	<2	52.7	Ref	0.02
	≥2	26.8	2.3 (1.2-4.6)	
Extranodal growth	No	46.5	Ref	0.01
	Yes	22.4	2.3 (1.2-4.5)	
Region	Neck	72.9	Ref	0.1
	Axilla	24.5	4.9 (1.1-1.7)	
	Groin	25.9	4.8 (1.1-0.7)	
S-100B	Low	44.6	Ref	0.2
	High	31.1	1.5 (0.8-2.9)	
SUV	Low	42.0	Ref	0.1
	High	29.0	1.7 (0.9-3.2)	

All variables with  $p \leq 0.2$  were entered in the multivariable model; the following variables were not entered: sex ( $p=0.4$ ), age ( $p=0.9$ ), clark level ( $p=0.7$ ) localization of the primary tumor ( $p=0.9$ ), number of lymph nodes removed ( $p=0.3$ ) and tumor size of the lymph nodes metastasis ( $p=0.6$ ). \*\*SUV and S100 were entered separately in the model.

As shown in Table 2 and Figure 1 DFS was 31.1% for a elevated S-100B and 44.6% for S-100B below cut-off point (univariate: HR=1.5;  $p=0.2$ ). The DFS was 29% for a elevated SUV and 42% for a SUV below cut-off point (univariate: HR=1.7;  $p=0.1$ ).

In multivariate analysis only variables with  $p \leq 0.2$  were entered in the multivariable model. SUV and S-100B were entered separately. In multivariate analyses the DFS was significantly correlated with S-100B (HR 3.1;  $p=0.02$ ). There was no correlation of DFS found with SUV (HR 1.1;  $p=0.8$ ). Primary melanoma ulceration ( $p=0.008$  and  $p=0.01$ ) and  $\geq 2$  lymph nodes positive ( $p=0.02$  and  $p=0.03$ ) were both associated with significant reduced DFS.


### *Disease Specific Survival (DSS)*

As shown in Table 3 and Figure 1 DSS was 44.7% for an elevated S-100B and 60.7% for a S-100B below the cut-off point (univariate: HR=2.0;  $p=0.06$ ). The DSS was 43.5% for a elevated SUV and 59.1% for a SUV below cut-off point (univariate: HR=1.7;  $p=0.1$ ).


In multivariate analysis an elevated S-100B showed a trend for DSS (HR= 2.4;  $p=0.06$ ), but an elevated SUV was not correlated with DSS at all (HR=1.0;  $P=0.9$ ). Primary melanoma ulceration was also correlated with DSS ( $p=0.05$  and  $p=0.06$ ).

Multivariable HR	Multivariable p-value	Multivariable HR	Multivariable p-value
Ref	0.4	Ref	0.7
1.4 (0.6-3.0)		1.2 (0.5-2.5)	
Ref	0.008	Ref	0.01
2.8 (1.3-6.2)		2.7 (1.2-5.9)	
Ref	0.02	Ref	0.03
2.7 (1.2-5.8)		2.5 (1.1-5.4)	
Ref	0.07	Ref	0.2
2.1 (0.9-4.6)		1.7 (0.8-3.9)	
Ref	0.1		
3.2 (0.7-15.3)			
4.6 (1.1-20.0)			
Ref	0.02		
3.1 (1.2-7.8)			
		Ref	0.8
		1.1 (0.5-2.4)	


DFS for S-100B, p=0.02




DSS for S-100B, p= 0.07



DFS for SUV, p=0.8



DSS for SUV, p= 0.8



**TABLE 3:** Univariate and multivariate analysis on Disease-Specific Survival (DSS) for SUV and S-100B with death due to melanoma as event.

Characteristic		3-years DSS	Univariate HR	Univariate p-value
Sex	Male	39.7	Ref	0.2
	Female	64.2	0.6 (0.3-1.3)	
Breslow thickness	< 2.0	63.0	Ref	0.2
	≥ 2.0	42.8	1.6 (0.8-3.3)	
Melanoma	No	58.9	Ref	0.04
Ulceration	Yes	27.3	2.2 (1.0-4.7)	
Lymph node positive	<2	51.9	Ref	0.6
	≥2	44.5	1.2 (0.6-2.5)	
Region	Neck	66.7	Ref	0.2
	Axilla	29.9	2.4 (0.8-7.3)	
	Groin	51.9	1.3 (0.4-3.9)	
S-100B	Low	60.7	Ref	0.06
	High	44.7	2.0 (1.0-4.0)	
SUV	Low	59.1	Ref	0.1
	High	43.5	1.7 (0.8-3.4)	

All variables with  $p \leq 0.2$  were entered in the multivariable model; the following variables were not entered: age ( $p=0.8$ ), clark level ( $p=0.3$ ), localization of the primary tumor ( $p=0.3$ ), number of lymph nodes removed ( $p=0.3$ ), number of nodes positive ( $p=0.5$ ), tumor size of the lymph nodes metastasis ( $p=0.7$ ) and extranodal growth ( $p=0.3$ ). \*\*SUV and S100 were entered separately in the model

**TABLE 4a:** Correlations of different patient and tumor characteristics for prognostic markers S-100B and SUV

		Associated with high S-100B	Associated with high SUV
Sex	Male	0.9	0.8
	Female		
Ulceration primary melanoma	No	0.9	0.04
	Yes		
Tumorsize lymph node metastasis	<3.0	0.04	0.01
	≥3.0		
Extranodal growth	No	0.8	0.03
	Yes		

### *Marker association with histopathological factors*

Table 4a reflects the association between histopathological factors and an elevated S-100B or SUV. For FDG-SUV primary melanoma ulceration ( $p=0.04$ ), size of the lymph node metastases ( $p=0.01$ ) and extranodal growth ( $p=0.03$ ) were associated. For S-100B, a large tumor size in the lymph node metastasis ( $p=0.04$ ) was associated with elevated S-100B levels.

Multivariable HR	Multivariable p-value	Multivariable HR	Multivariable p-value
Ref	0.1	Ref	0.1
0.5 (0.2-1.1)		0.5 (0.2-1.2)	
Ref	0.1	Ref	0.2
2.1 (0.9-5.0)		1.8 (0.8-4.4)	
Ref	0.05	Ref	0.06
2.3 (1.0-5.1)		2.3 (0.9-5.4)	
Ref	0.2	Ref	0.3
1.7 (0.7-3.9)		1.5 (0.7-3.5)	
Ref	0.8	Ref	0.4
1.4 (0.4-4.9)		2.0 (0.6-6.9)	
1.2 (0.4-3.8)		1.0 (0.3-3.3)	
Ref	0.06		
2.4 (0.9-5.9)		Ref	0.9
		1.0 (0.4-2.6)	

### *Stratified analysis of the prognostic values S-100B*

Table 4b reflects the prognostic value of S-100B for DFS in both males ( $p=0.03$ ) and females ( $p=0.01$ ), for patients aged 46-64 yrs ( $p=0.03$ ), patients with groin metastases ( $p=0.02$ ) and with  $\geq 2$  positive lymph nodes ( $p=0.02$ ) or a tumor size  $<3.0$  cm ( $p=0.04$ ) and patients with extranodal growth ( $p=0.01$ ).

For DSS, S-100B was of prognostic value especially in elderly patients ( $\geq 65$  years) ( $p=0.05$ ), in patients with groin metastases ( $p=0.01$ ) and with size of the lymph node metastases  $<3.0$  cm ( $p=0.02$ ). A trend towards significance was seen in patients with presence of  $\geq 2$  positive lymph nodes ( $p=0.05$ ).

## **DISCUSSION**

### *General*

No relation was found between the prognostic markers S-100B and SUV in melanoma patients with palpable lymph nodes. A preoperative elevated S-100B predicts a reduced DFS in stage III melanoma and is a prognostic marker in case of high lymph node tumour burden. An elevated SUV, according to the present data, has no correlation with survival. Therefore, evaluating these data, S-100B could be used as a prognostic marker in the stratification for new adjuvant trials, in order to better stage and select clinically AJCC stage III melanoma patients for systemic treatment.

**TABLE 4b:** Prognostic value of S-100B stratified for different groups

		DFS*	DSS**
		HR (95%CI); p-value	HR(95%CI); p-value
Sex	Male	4.6 (1.1-19.1); p=0.03	2.4 (0.7-8.4); p=0.2
	Female	6.0 (1.5-24.8); p=0.01	2.6 (0.7-9.5); p=0.1
Age (years)	≤45	0.8 (0.1-7.2); p=0.8	3.0 (0.1-75); p=0.5
	46-64	9.1 (1.6-52.1); p=0.03	1.0 (0.2-3.9); p=0.9
	≥65	2.9 (0.2-42.1); p=0.2	22.9 (1.0-520); p=0.05
Localization lymph node metastases	Neck	p=0.9***	p=0.8***
	Axilla	0.4 (0.1-2.7); p=0.3	0.7 (0.1-5.4); p=0.7
	Groin	3.4 (1.2-10.0); p=0.02	5.2 (1.5-18.9); p=0.01
Lymph nodes positive	<2	4.2 (0.9-20.3) ; p=0.07	3.4 (0.7-15.7) ; p=0.1
	≥2	3.6 (1.3-10.0) ; p=0.02	2.7 (1.0-7.6) ; p=0.05
Tumorsize lymph node metastasis	<3.0	4.7 (1.1-21.0); p=0.04	7.4 (1.4-38.3); p=0.02
	≥3.0	2.6 (0.8-8.0); p=0.1	1.7 (0.5-5.2); p=0.4
Extranodal growth	No	2.3 (0.8-6.3); p=0.1	2.6 (0.9-7.8); p=0.1
	Yes	6.0 (1.5-24.3); p=0.01	2.9 (0.8-10.0); p=0.1

\*DFS adjusted for ulceration, Breslow, number of positive nodes, localization and extranodal growth

\*\*DSS adjusted for sex, ulceration, Breslow, number of positive nodes and localization.

\*\*\*no estimation of the HR due to small numbers.

### *Prognostic markers*

This is the first study, to our knowledge, evaluating both prognostic markers S-100B and SUV in melanoma patients before undergoing a therapeutic lymph node dissection. After a lymph node dissection, most melanoma patients die within three years as a consequence of distant metastases. Prediction of survival in AJCC stage III has traditionally been based on the number and size of lymph nodes, the presence of extranodal growth and ulceration of the primary tumor.<sup>20</sup> In the search for effective systemic therapy in stage III melanoma, the development of prognostic markers and staging will be of increasing interest in order to improve patient-tailored adjuvant therapy.

### *Staging*

Accurate staging before treatment is essential for all melanoma patients. However, for the evaluation of early regional lymphatic dissemination in AJCC stage I and II melanoma, determining S-100B or performing a PET/CT scan is unsuitable.<sup>21</sup> Fortunately, nowadays the technique of lymphatic mapping and sentinel lymph node biopsy (SLNB) can be used as a standard method of selecting patients with micro-metastases. The technique has undisputedly proven to be a reliable staging method and the fourth interim analysis of the MSLT I study reveals that SLNB indeed leads to improved melanoma specific survival.<sup>22,23</sup> In staging studies analyzing patients with palpable lymph nodes (AJCC stage III), whole-body FDG-PET and/or spiral CT have proved their value with 27% upstaging to stage IV disease. A sensitivity

of 79-92% and a specificity of 86-90% to detect distant metastases was found.<sup>5,24</sup> The combined FDG PET/CT leads to a change in the planned dissection in 37% of the patients.<sup>25</sup> Acknowledging the heterogeneity of stage III melanoma, the suspicion rises that numerous patients have a form of subclinical dissemination, which can even remain undetected by standard FDG-PET and CT imaging.<sup>6</sup>

### *SUV and S-100B not correlated*

SUV can be easily calculated after a FDG-PET in the diagnostic work-up and might recognize, when elevated, high risk stage IIIB patients. By assessing the degree of FDG accumulation SUV was recorded earlier as a prognostic marker by other researchers.<sup>15</sup> If treatment response can be predicted by measurement of increased glucose metabolism, this would be of particular value in choosing an adequate patient-tailored adjuvant therapy.

Earlier, several other studies have recommended the use of S-100B. Successful treatment with lymph node dissection, chemotherapy or immunotherapy was associated with decreased S-100B concentrations whereas increased concentrations were an expression of disease progression.<sup>26</sup> Also, elevated S-100B was thought to be an expression of early dissemination and an independent prognostic marker of risk for mortality.<sup>11-13</sup>

Our results show that the tumour markers SUV and S-100B are not related. Both tumour markers are correlated with tumor size in the lymph node and SUV is also correlated with extranodal growth and primary ulceration (Table 4a).

### *Consideration*

Interpreting these results leads to the conclusion that serum S-100B elevation seems correlated with lymphatic dissemination and nodal tumor burden. Elevation of S-100B might detect a process of haematogenic and lymphogenic dissemination in melanoma patients. Because S-100B is located in the cytoplasm of melanoma cells, elevated serum S-100B probably occurs as a result of loss of melanoma cell integrity.<sup>13</sup> Earlier evidence concerning FDG uptake suggests that the intensity of uptake does not correlate with lymph node tumor burden, but more with biological aggressiveness and is associated with cell viability and particularly cell proliferative activity.<sup>28</sup>

In the current study SUV did not prove its role as prognostic marker despite earlier findings by Bastiaannet et al.<sup>15</sup> However, the cohorts differed significantly in localization of the primary melanoma, number of positive nodes and extra nodal growth. Furthermore, earlier Bastiaannet et al. used a cut-off point of 6.14 and in the current study 8.4. The cut-off point for SUV is data driven and therefore depends on the patient group. In literature no consensus for a cut-off point for SUV has been formed yet.<sup>19</sup> In general, SUV values are often subject to different sources of variability, such as the timing of PET acquisition after tracer injection, patient size, region of interest definition, partial volume and image reconstruction.<sup>29,30</sup>

## CONCLUSION

Preoperative elevation of S-100B is a prognostic marker for AJCC stage III melanoma patients. S-100B, although not significantly correlated with DSS ( $p=0.06$ ), is a prognostic marker for DFS. The results of the current study, although with a relatively small group, again confirm the value of preoperative S-100B in stage III melanoma.<sup>11</sup> Furthermore, Standardised Uptake Value (SUV) is not associated with S-100B and the marker does not reconfirm its status as independent prognostic factor.

For the future we predominantly recommend the use of S-100B in combination with the established prognostic parameters for better stratification in new adjuvant therapeutic trials. Pre-operative S-100B levels provide information for stage III melanoma patients who want to be optimally informed about their disease prognosis. Ongoing research for prognostic biomarkers in larger prospective, randomized, high-quality methodological studies is essential in establishing treatment better tailored to the characteristics of individual tumor biology in stage III melanoma.

## REFERENCES


- Rinne D, Baum RP, Hor G, Kaufman R. Primary staging and follow-up of high risk melanoma patients with whole-body 18F-fluorodeoxyglucose positron emission tomography: results of a prospective study of 100 patients. *Cancer* 1999;85:1199-201.
- [Http://www.ikcnet.nl/page.php?id=2755&nav\\_id=97](http://www.ikcnet.nl/page.php?id=2755&nav_id=97)
- De Vries E, van de Poll-Franse LV, Louwman WJ, de Gruijl FR, Coebergh JW. Predictions of skin cancer incidence in the Netherlands up to 2015. *Br J Dermatol*. 2005;152:481-8.
- Balch CM, Gershenwald JE, Soong SJ et al. Final version of 2009 AJCC melanoma staging and classification. *J Clin Oncol* 2009;27:6199-206.
- Bastiaannet E, Wobbes T, Hoekstra OS et al. Prospective comparison of [18F] fluorodeoxyglucose positron emission tomography and computed tomography in patients with melanoma with palpable lymph node metastases: diagnostic accuracy and impact on treatment. *J Clin Oncol* 2009;27:4774-80.
- Balch CM, Gershenwald JE, Soong S et al. Multivariate Analysis of Prognostic Factors Among 2,313 Patients With Stage III Melanoma: Comparison of Nodal Micrometastases Versus Macrometastases. *J Clin Oncol*. 2010;28:2452-9.
- Hill BR, Levi C. Elevation of a serum component in neoplastic disease. *Cancer Res* 1954;14: 513-15.
- Cogas H, Eggermont AMM, Hauschild A et al. Biomarkers in melanoma. *Ann Oncol* 2009; 20:8-13.
- Krahn G, Kaskel P, Sander S et al. S100 beta is a more reliable tumor marker in peripheral for patients with newly occurred melanoma metastases compared with MIA, albumine and lactaat-dehydrogenase. *Anticancer Res* 2001;21:1311-16.
- Smit LH, Korse CM, Hart AA et al. Normal values of S-100B predict prolonged survival for stage IV melanoma patients. *Eur J Cancer* 2005;41:386-92.
- Kruijff S, Bastiaannet E, Muller-Kobold AC et al. S-100B concentrations predict disease-free survival in stage III melanoma patients. *Ann Surg Oncol* 2009;16:3455-62.
- Tarhini AA, Stuckert J, Lee S, Sander C, Kirkwood JM. Prognostic significance of serum S-100B surgically resected melanoma patients participating in Intergroup Trial ECOG 1694. *J.Clin.Oncol*. 2008 1;27:38-44.
- Ghanem G, Loir B, Morandini R et al. On the release and half-life of S100B protein in the peripheral blood of melanoma patients. *Int J Cancer* 2001;94:586-90.
- Mochiki E, Kuwano H, Katoh H, Asao T, Oriuchi N, Endo K. Evaluation of 18F-2-deoxy-2-fluoro-D-glucose positron emission tomography for gastric cancer. *World J Surg* 2004;28:247-53.
- Bastiaannet E, Hoekstra OS, Oyen WJ, Jager PL, Wobbes T, Hoekstra HJ. Level of Fluorodeoxyglucose Uptake Predicts Risk for Recurrence in Melanoma Patients Presenting With Lymph Node Metastases. *Ann Surg Oncol* 2006;13:919-26.
- Bastiaannet E, Beukema JC, Hoekstra HJ. Radiation therapy following lymph node dissection in melanoma patients: treatment, outcome and complications. *Cancer Treat Rev*. 2005;31:18-26.
- Henderson MA, Burmeister B, Thompson JE, Di Iulio J, Fisher R, Hong A et al. Adjuvant radiotherapy and regional lymph node field control in melanoma patients after lymphadenectomy: Results of an intergroup randomized trial. 2009 ASCO Annual meeting, *J Clin Oncol*. 2009;27 Supplement:Abstract No.LBA9084.
- Hamacher K, Coenen HH, Stocklin G. Efficient stereospecific synthesis of no-carrier-added 2-[18F]-fluoro-2-deoxy-D-glucose using aminopolyether supported nucleophilic substitution. *J Nucl Med* 1986;27:235-8.
- Westerterp M, Pruim J, Oyen W et al. Quantification of FDG PET studies using standardised uptake values in multi-centre trials: effects of image reconstruction, resolution and ROI definition parameters. *Eur J Nucl Med Mol Imaging* 2007;34:392-404.
- Soong SJ, Ding S, Coit D et al. [www.melanomaprognosis.org](http://www.melanomaprognosis.org). Predicting Survival Outcome of Localized Melanoma: An Electronic Prediction Tool Based on the AJCC Melanoma Database. *Ann Surg Oncol* 2010;17:2006-14.



21. Klode J, Dissmond J, Grabbe S, Hillen U, Poeppel T, Boeing C. Sentinel lymph node excision and PET-CT in the initial stage of malignant melanoma: A retrospective analysis of 61 patients with malignant melanoma in American Joint Committee on cancer stages I and II. *Dermatol Surg* 2010;36:439-45.
22. Morton DL, Thompson JE, Cochran AJ, et al. Sentinel-node biopsy or nodal observation in melanoma. *N Engl J Med* 2006;355:1307-17.
23. Morton DL, Thompson JE, Cochran AJ, et al. Sentinel node biopsy and immediate lymphadenectomy for occult metastases versus nodal observation and delayed lymphadenectomy for nodal recurrence. *Ann Surg Oncol* 2010, 17 Suppl 1, S22-S23 (Abstract 53)
24. Mijnhout GS, Hoekstra OS, van Tulder MW, Teule GJ, Deville WL. Systematic review of the diagnostic accuracy of (18)F-fluorodeoxyglucose positron emission tomography in melanoma patients. *Cancer* 2001;91:1530-42.
25. Aukema TS, Valdés Olmos RA, Wouters MW et al. Utility of Preoperative 18F-FDG PET/CT and Brain MRI in Melanoma Patients with Palpable Lymph Node Metastases. *Ann Surg oncol* 2010;17:2773-8.
26. Hamberg AP, Korse CM, Bonfrer JM, de Gast GCl. Serum S-100B is suitable for prediction and monitoring of response to chemoimmunotherapy in metastatic malignant melanoma. *Melanoma Res* 2003;13:45-49.
27. Minn H, Clavo AC, Grenman R, Wahl RL. In vitro comparison of cell proliferation kinetics and uptake of tritiated fluorodeoxyglucose and L-methionine in squamouscell carcinoma of the head and neck. *J Nucl Med* 1995;36:252-8.
28. Sperti C, Pasquali C, Chierichetti F, Ferronato A, Decet G, Pedrazzoli S. 18 Fluorodeoxyglucose positron emission tomography in predicting survival of patients with pancreatic carcinoma. *J Gastrointest Surg* 2003;7:953-9.
29. Sugawara Y, Zasadny KR, Neuhoff AW, Wahl RL. Reevaluation of the standardized uptake value for FDG: variations with body weight and methods for correction. *Radiology* 1999;213:521-5.
30. Boellaard R, Krak NC, Hoekstra OS, Lammertsma AA. Effects of noise, image resolution and ROI definition on the accuracy of standard uptake values: a simulation study. *J Nucl Med* 2004;45:1519-27.



# 7



# The use of S-100B to evaluate therapy effects during bevacizumab induction treatment in AJCC stage III melanoma

| S Kruijff, E Bastiaannet, AH Brouwers, WB Nagengast, MJ, Speijers, AJH Suurmeijer, GA Hospers, HJ Hoekstra | Ann Surg Oncol. 2011 Aug 23. [Epub ahead of print]

---

## ABSTRACT

**Aim:** To investigate the feasibility of using bevacizumab to improve the survival of AJCC stage III melanoma patients, we investigated how a single bevacizumab treatment affected nodal disease and a panel of biomarkers in clinically FDG-PET/CT staged, stage III melanoma patients, prior to therapeutic lymph node dissection (TLND).

**Methods:** Four weeks before TLND, nine patients (median age, 50 (range 28.8-62.1); two male, seven female) with palpable lymph node metastases received 7.5 mg/kg bevacizumab. Before and after this treatment, all patients were assessed by measurements of the maximum standardized uptake value (SUVmax) by FDG-PET scan, and serum S-100B and LDH. After TLND, the dissection specimen was analyzed for the number of removed lymph nodes, the amount of metastatic lymph nodes, and tumor necrosis.

**Results:** Median follow-up was 15.5 (2.2-32.9) months. Histopathological analysis revealed tumor necrosis in six patients, of which, five had an S-100B decline and one had an unchanged S-100B level after bevacizumab. The other three patients showed an S-100B increase and no necrosis. Tumor necrosis was correlated with S-100B decrease ( $p = 0.048$ ). No association was found between necrosis and the markers SUVmax and LDH. No wound healing disturbances were encountered.

**Conclusion:** Tumor necrosis in dissection specimens was associated with declining S-100B levels, while elevated S-100B was only found in cases with no necrosis. Bevacizumab might be useful in treating AJCC stage III melanoma patients prior to TLND, and S-100B appears to be a useful marker for assessment of treatment effects.

## INTRODUCTION

Melanoma is an aggressive and highly metastatic disease, which can be fatal with a rapid systemic dissemination. Approximately one third of all melanoma patients will experience disease recurrence.<sup>1,2</sup> While almost all organs can be involved, the most frequent target sites are the liver, bone, and the brain. Treatment results for advanced melanoma remain unsatisfactory. No systemic therapy has been demonstrated to affect overall survival, although the recent studies of immunotherapy with ipilimumab and the introduction of BRAF pathway inhibitors have shown promising results.<sup>3,4,5</sup> For melanoma patients with nodal disease, therapeutic lymph node dissection (TLND), with or without adjuvant radiation, remains the only curative therapy, with 5-year survival rates of 78%, 59%, and 40%, respectively, for patients with AJCC stage IIIA, IIIB, and IIIC disease.<sup>6,7</sup> As a consequence of shortages in health care resources, the growing elderly population in the Western world, and the increasing incidence of cancer, the wait time for surgery at some cancer centers has lengthened to an average of four to five weeks. This period before TLND offers a unique opportunity to test novel induction treatments before surgery.

Tumor angiogenesis is a continuous process that allows cancer cells to grow by supplying the tumor with nutrients and oxygen, disposing of metabolic waste products, and providing a route for metastatic spread.<sup>8,9</sup> Vascular endothelial growth factor A (VEGF-A) is a key growth factor involved in the development and maintenance of tumor angiogenesis.<sup>10</sup> Bevacizumab, a fully humanized monoclonal antibody, binds to all VEGF isoforms with high affinity, thereby blocking ligand-receptor signaling.<sup>11</sup> It is currently used in patients with metastatic colon cancer, non-small-cell lung cancer, and renal cell cancer.<sup>12-14</sup> Bevacizumab was previously evaluated in a randomized phase II trial (BEAM trial) in metastatic melanoma, which compared the effects of the combination of carboplatin and paclitaxel, with and without bevacizumab. The addition of bevacizumab had a significant impact on progression-free survival, and some impact on overall survival, although this effect was not significant.<sup>15</sup>

The time spent waiting for a TLND for regional metastatic disease could be used more effectively if an induction therapy could be safely administered to reduce tumor load before surgery. S-100B and SUV are known to be of prognostic value in stage III melanoma; elevated S-100B and SUV in stage III melanoma patients can be specific indicators of disease progression.<sup>16-22</sup> Therefore, we hypothesized that monitoring S-100B and SUV before and after a single bevacizumab treatment might provide a 'measurable reflection' of the response to this angiogenetic treatment.

Here, we investigated the feasibility of using serum biomarkers, S-100B and LDH, and the Standardized Uptake Value (SUV) from FDG-PET to evaluate effects of an induction treatment with a single dose bevacizumab in stage IIIB/C melanoma patients, prior to TLND. We assessed the perioperative changes in biomarker levels following bevacizumab treatment, as well as the induction of tumor necrosis based on final histopathology of the resected lymph nodes.

## PATIENTS AND METHODS

### *Patients*

All consecutive melanoma patients presenting with palpable and cytology-proven lymph node metastases (AJCC IIIB/C) at the UMCG between January and July 2008 were evaluated with FDG-PET and spiral CT in addition to the routine staging, which included a complete medical history, physical examination, and blood chemistry profile. If the FDG-PET and CT were negative for distant metastases, patients were offered participation in the bevacizumab trial bridging the waiting time between the diagnosis and the TLND. Eligible patients were  $\geq 18$  years of age with AJCC stage IIIB melanoma and a WHO performance status of 0-2. Exclusion criteria included a history of radiotherapy on the involved lymph node basin, major surgery within 28 days of start of the study, administration of any investigational drug within 30 days before start of the study, and clinical evidence of brain metastases. Patients who presented with a positive sentinel lymph node biopsy, local recurrence, and/or in-transit disease were also excluded from this study. Examined data included patient demographics, primary tumor characteristics, date and type of operation, and status at last follow-up. Patients with nodal metastases of  $\geq 3$  cm and/or  $\geq 3$  positive lymph nodes and/or extranodal (EN) disease received adjuvant radiotherapy ( $20 \times 2.4$  Gy).

This study was approved by the local medical ethics committee and written informed consent was obtained from all participants. The study was registered under trial number NTR1941 and was organized as a corollary study in concurrence with another feasibility study investigating the presence of VEGF in melanoma lesions by VEGF-SPECT with  $^{111}\text{In}$ -bevacizumab.<sup>23</sup>

### *Study design*

On day 0, all selected patients were evaluated with a FDG-PET scan and on day 7, they received a single dose of 7.5 mg/kg bevacizumab intravenously. On day 40, the patients underwent a second FDG-PET scan, after which TLND was performed on day 42 (five weeks after bevacizumab). S-100B and LDH levels were measured on days 0, 41, and 43 (first visit to outpatient clinic, one day preoperatively, and one day postoperatively). After lymph node dissection, the pathological results were analyzed.

### *S-100B*

Concentrations of serum S-100B were measured using the Liason Sangtec 100 immunoassay on the Advantage (Nichols). The half-life of S-100B is estimated to be about 30 minutes.<sup>24</sup> Levels were calculated using a calibration curve and checked against internal standards with known concentrations of S-100B. In this assay, concentrations were considered normal if they were below a cut-off point of 0.15  $\mu\text{g/L}$ . Concentrations above this range were considered to be elevated. For all patients, the change in S-100B concentration from day 0 to day 41 was determined.

### *LDH*

LDH was routinely analyzed by enzymatic activity measurement with a Roche Modular analyzer (Hitachi). LDH values <250 U/L were considered to be normal.

### *Standardized uptake value (SUV)*

To assess the degree of FDG accumulation before and after bevacizumab treatment, the maximum standardized uptake value (SUV<sub>max</sub>) was calculated from a FDG-PET scan. The SUV depends on the amount of injected radioactivity, the patient's weight, and the calibration factor of the camera, and it is calculated according to the following formula: SUV<sub>max</sub> = radioactivity concentration in tissue (Bq/Kg)/(injected dose [Bq]/patient weight [Kg]). If multiple metastases were present in the lymph node basin, the lesion with the most intense uptake was chosen. The cut-off point for SUV is data driven and therefore depends on the patient group; there is no consensus in the literature regarding a cut-off point for SUV.<sup>25-27</sup> In the present study, any decrease/increase of SUV<sub>max</sub> was recorded simply as the level of decrease/increase.

### *Surgery*

The therapeutic axillary lymph node dissection included a level III resection with resection of the pectoralis minor muscle. The groin lymph node dissection included a superficial (inguinal) and deep (iliac-obturator) groin dissection. The latter procedure was performed as extensively described by Baas et al.<sup>28</sup> Perioperative surgical complications, the number of harvested lymph nodes, number of metastatic lymph nodes, maximum metastasis diameter, and degree of tumor necrosis were scored. The pathological results after lymph node dissection were analyzed by a pathologist.

### *Statistics*

Fischer's exact test was used to assess associations between the SUV, S-100B and LDH levels, and pathology data.

## **RESULTS**

### *Patient characteristics*

Table 1 shows the characteristics of the nine AJCC stage III melanoma patients, which included two males and seven females, with a median age of 50 (range 28.8-62.1) years. The median follow-up time was 15.5 (range 2.2-32.9) months. In these nine patients, metastatic lymph nodes were detected with spiral CT and FDG-PET imaging. After TLND, the resection specimens displayed a median of 16 nodes (range 8-29). No perioperative or wound healing disturbances were encountered.

**TABLE 1** Characteristics of the primary melanoma and localization recurrence.

		Number	%
Sex	Male	2	22.2
	Female	7	77.8
Age yrs (median range)		50.0	(28.8-62.1)
Localization of primary melanoma	Trunk	5	55.6
	Upper extremities Lower extremities	0	0
	Unknown	3	33.3
Clark level	II	1	11.1
	III	2	22.2
	IV	5	55.6
	Unknown primary	1	11.1
Breslow thickness	≤1.0	4	44.4
	>1.0	4	44.4
	Unknown primary	1	11.1
Mitotic index	≤5	4	44.4
	≥6	4	44.4
	Unknown primary	1	11.2
Ulceration	Yes	3	33.3
	No	5	55.6
	Unknown primary	1	11.1
Localization of lymph node metastases	Axilla	5	55.6
	Groin	4	44.4

**TABLE 2** Association between pathology data and the changes in SUV value, S-100B, and LDH.

		SUV		S-100B		LDH	
		Increase	Decrease	Increase	Decrease	Increase	Decrease
Size	≤4.0 cm	3 (75%)	1 (20%)	2 (50%)	2 (40%)	1 (50%)	3 (43%)
	>4.0 cm	1 (25%)	4 (80%)	2 (50%)	3 (60%)	1 (50%)	4 (57%)
Necrosis	Yes	3 (75%)	3 (60%)	1 (25%)*	5 (100%)*	0 (0%)	6 (86%)
	No	1 (25%)	2 (40%)	3 (75%)	0 (0%)	2 (100%)	1 (14%)
Ratio positive / removed nodes	≤0.5	2 (50%)	2 (40%)	2 (50%)	2 (40%)	1 (50%)	3 (43%)
	>0.5	2 (50%)	3 (60%)	2 (50%)	3 (60%)	1 (50%)	4 (57%)
Extranodal growth	No	2 (50%)	3 (60%)	3 (75%)	2 (40%)	1 (50%)	4 (57%)
	Yes	2 (50%)	2 (40%)	1 (25%)	3 (60%)	1 (50%)	3 (43%)
Total		4 (44%)	5 (56%)	4 (44%)	5 (56%)	2 (22%)	7 (78%)

\*p < 0.05 (Fisher's exact test)




### *SUV and S-100B and LDH*

In five of the nine patients, the SUVmax decreased after bevacizumab treatment. In one patient, SUVmax was not influenced, and in the remaining three patients, the SUVmax increased. Between day 0 and day 41, an LDH decrease was noted in seven patients and an elevation in two patients.

### *Pathology*

The pathologic analysis reported tumor necrosis in six TLND specimens after bevacizumab treatment. In five of these six patients, a decrease in S-100B after bevacizumab was documented on day 41; in the remaining one patient, S-100B was unchanged but was comparatively decreased on day 43. Tumor necrosis was significantly associated with declining S-100B levels ( $p = 0.048$ ). These data are shown in Table 2 and Figure 1. No association was found between the markers SUVmax and S-100B ( $p = 0.6$ ) or LDH, nor was a correlation found between LDH or SUVmax and necrosis.



**FIGURE 1** S-100B before and after bevacizumab.

## **DISCUSSION**

This is the first study to assess the use of biomarkers to monitor the response following an induction treatment with bevacizumab, a VEGF-specific antibody, prior to TLND in AJCC stage III melanoma patients with nodal disease. Our data suggest that neo-adjuvant bevacizumab may be useful for preoperative tumor load reduction before TLND in AJCC stage IIIB/C melanoma patients.

Neo-adjuvant treatment with bevacizumab might have induced tumor necrosis in the metastatic lymph nodes in six of the nine patients. In all patients with necrosis in the dissection specimen, a significant decline of serum S-100B levels was observed. In contrast, in patients whose S-100B levels continued to rise following treatment, no necrosis was found in the dissection preparations. No relationship was found between SUV and S-100B or LDH and S-100B or any of the other pathological data.

The data from this feasibility study should only be considered preliminary and it is clear that no firm conclusions can be drawn from such a small group of samples. However, the observations certainly indicate that further investigation is warranted. The data from the present study might indicate that a single neo-adjuvant bevacizumab treatment induced tumor necrosis in the tumor bearing lymph nodes. Even more importantly, a decline of S-100B was observed in all patients who exhibited tumor necrosis after induction therapy.

Unfortunately, there are no data available revealing how often necrosis is discovered after lymph node dissection without prior administration of any induction therapy. However, the relatively high necrosis rate within our small sample of patients and the related S-100B decline might suggest a drug-induced tumor necrosis in the lymph nodes. The associated significant decrease of tumor load might explain the S-100B decline in all six patients.

To date, the most extensively studied biomarker in melanoma is S-100B, a 21-kDa protein that was first isolated from the central nervous system in vertebrates. S-100B is located in the cytoplasm and nucleus in cells with a neuro-ectodermal origin, e.g. melanocytes. S-100B has various intra-cellular functions, mainly in cytoskeleton integrity, cell cycle regulation, and apoptosis. Elevated serum S-100B occurs as the result of loss of melanoma cell integrity.<sup>24</sup> The mechanism by which this protein is released into circulation remains uncertain, but it is probably caused by cell damage or apoptosis. The hypothesis that elevation of S-100B should be interpreted as indication of ongoing metastatic disease has been suggested by earlier data.<sup>16</sup> Proportions of patients with elevated S-100B concentrations are 0-9% in stage I/II, 5-98% in stage III, and 40-100% in stage IV melanoma.<sup>29</sup> For clinically FDG-PET and CT-staged stage III patients, a pre-operatively elevated S-100B level is correlated with decreased survival.<sup>16-19</sup> In stage IV, improved survival is observed for patients with low S-100B levels; increased concentrations of S-100B during systemic treatment are associated with disease progression.<sup>30-34</sup> Recently, Bouwhuis et al. revealed that an elevation of S-100B levels throughout serial serum measurements in stage III is of very significant prognostic value, which is even stronger when combined with disease stage and number of positive lymph nodes.<sup>35</sup> However, not only should elevated S-100B be seen as a prognostic factor for survival and a mere reflection of disease progression alone; the presence of the protein S-100B itself induces disease progression as well. S-100B interacts with p53 and thereby down-regulates its function as a tumor suppressor protein by preventing induction of apoptosis of potential melanoma cells.<sup>36,37</sup>


Taking these results into account, an induction therapy before lymph node dissection might interrupt the melanoma proliferation cycle in two ways: first, by reducing tumor load and second, by temporarily suppressing the serum S-100B concentrations and thus preventing down regulation of tumor suppressor protein p53. Bevacizumab as an induction therapy might

temporarily suppress metastatic proliferation and eventually, in combination with surgery, improve outcome.

Anti-angiogenic therapy is a promising strategy in the treatment of cancer. Angiogenesis, or remodeling of an existing network of blood vessels, performs an essential role in diverse pathophysiological processes.<sup>38,39</sup> Because the neo-vascularization process that supports tumor growth may be similar to that involved in physiological wound healing, a delay in wound repair has been a concern. This may be one of the reasons that bevacizumab has not been widely used as induction therapy before major surgical cancer treatment.<sup>40</sup> None of the patients in our study showed any wound healing disturbances during the course of the study.

The currently in-progress AVAST-M trial is the only known randomized trial evaluating the VEGF inhibitor, bevacizumab, for adjuvant therapy following resection of AJCC stage IIB, IIC, and III cutaneous melanoma. Many randomized trials of adjuvant therapy have been performed in patients after resection of melanoma, but to date, no treatment has convincingly improved overall survival. The AVAST-M trial is a cancer research UK funded, NCRN trial in which patients with previously resected melanoma at high risk of disease recurrence will be randomized to receive either standard observation or bevacizumab administration for 1 year.<sup>41</sup>

In one of the few examples in which induction therapy effects were monitored by biomarkers, Willet et al. performed a phase I-II trial to test the efficacy of adding bevacizumab to standard chemoradiotherapy as an induction therapy before rectal surgery. The phase II clinical trial involved administration of neoadjuvant bevacizumab, in combination with 5-fluorouracil (5-FU) and radiation therapy. The response of one patient's tumor to this regimen with bevacizumab was dramatic, with no residual cancer in the surgical specimen. More surprisingly, the plasma CEA dropped substantially from a pretreatment value of 122.8 ng/ml to 63.6 ng/ml at 12 days after the first bevacizumab infusion, and declined to 4.9 ng/ml at 1 month after completion of neoadjuvant therapy before surgery.<sup>42</sup>



**FIGURE 2** Patient (61 years, female), melanoma lower extremities (Breslow 2.1 mm, ulceration). Inguinal lymph nodes metastases with a SUVmax of 9.2 first scan and 10.5 second scan

In addition to S-100B, we also monitored SUVmax and LDH levels in the patients in this study, but did not find any significant correlation between these values and S-100B or pathology findings. In an earlier study, no relation was found between the prognostic markers S-100B and SUV in melanoma patients with palpable lymph nodes.<sup>43</sup> Evidence concerning FDG uptake suggests that the intensity of uptake does not correlate with lymph node tumor burden, but more with biological aggressiveness.<sup>44</sup> Serum lactate dehydrogenase (LDH) has high specificity for melanoma, but low sensitivity; it is an independent prognostic factor in stage IV melanoma.<sup>16</sup> Taking these facts into consideration, it is not surprising that we observed a lack of correlation of these markers with tumor response in this study.


In conclusion, this feasibility study has shown that induction treatment with bevacizumab in stage III melanoma four weeks prior to surgery may be related to induction of tumor necrosis without any observed incidence of wound healing disturbances. More importantly, when tumor necrosis was found in the dissection preparation, a 100% decline of S-100B was noted. Correspondingly, absence of tumor necrosis was correlated with ongoing S-100B elevation. The ability to target therapy towards well-selected subgroups of patients with the use of biomarkers, such as S-100B, could increase the likelihood of benefit and might improve therapeutic outcomes for future melanoma treatments.<sup>45</sup> Induction treatment with bevacizumab followed by therapeutic lymph node dissection could be a potentially successful combined treatment and might improve locoregional control, and therefore disease-free survival, for lymphogenic disseminated melanoma patients.

## REFERENCES

1. Francken AB, Bastiaannet E, Hoekstra HJ. Follow-up in patients with localised primary cutaneous melanoma. *Lancet Oncol.* 2005;8:608-21.
2. Speijers MJ, Francken AB, Hoekstra-Weebers JEHM, et al. Optimal Follow-up for Melanoma. *Expert Review of Dermatology.* 2010;5:461-478.
3. Hodi FS, O'Day SJ, McDermott DF, et al. Improved survival with ipilimumab in patients with metastatic melanoma. *New Engl J Med.* 2010;363:711-23.
4. Flaherty KT, Puzanov I, Kim KB, et al. Inhibition of mutated, activated BRAF in metastatic melanoma. *N Engl J Med.* 2010;363:809-19.
5. Chapman PB, Hauschild A, Robert C, et al. Improved survival with vemurafenib in melanoma with BRAF V600E mutation. *N Engl J Med.* 2011;364:2507-16.
6. Balch CM, Greshenwald JE, Soong SJ, et al. Final version of 2009 AJCC melanoma staging and classification. *J Clin Oncol.* 2009; 27:6199-206.
7. Bastiaannet E, Beukema JC, Hoekstra HJ. Radiation therapy following lymph node dissection in melanoma patients: treatment, outcome and complications. *Cancer Treat Rev.* 2005;31:18-26.
8. Kerbel RS. Tumor angiogenesis. *N Engl J Med.* 2008; 358:2039-49.
9. Folkman J, Klagsbrun M. Angiogenic factors. *Science* 1987;235:442-7.
10. Gerber HP, Ferrara N. Pharmacology and pharmacodynamics of bevacizumab as monotherapy or in combination with cytotoxic therapy in preclinical studies. *Cancer Res.* 2005;65:671-80.
11. Gaudreault J, Fei D, Rusit J, Suboc P, Shiu V. Preclinical pharmacokinetics of Ranibizumab (rhuFabV2) after a single intravitreal administration. *Invest Ophthalmol Vis Sci.* 2005;46:726-33.
12. Hurwitz H, Fehrenbacher L, Novotny W, et al. Bevacizumab plus irinotecan, fluorouracil and leucovorin for metastatic colorectal cancer. *N Engl J Med* 2004; 350:2335-42.
13. Sandler A, Gray R, Perry MC, et al. Paclitaxel-carboplatin alone or with bevacizumab for non-small-cell lung cancer. *N Engl J Med* 2006;355:2542-50.
14. Yang JC, Haworth L, Sherry RM, et al. A randomized trial of bevacizumab, an anti-vascular endothelial growth factor antibody, for metastatic renal cancer. *N Engl J Med.* 2003; 349:427-34.
15. O'Day SJ, Kim KB, Sosman JA, et al. BEAM: a randomized phase II study evaluating the activity of bevacizumab in combination with carboplatin plus paclitaxel in patients with previously untreated advanced melanoma. *Eur J Cancer Suppl.* 2009;7:Abstr 13.
16. Kruijff S, Bastiaannet E, Muller Kobold AC, et al. S-100B concentrations predict disease-free survival in stage III melanoma patients. *Ann Surg Oncol* 2009; 16:3455-62.
17. Tarhini AA, Stuckert J, Lee S, et al. Prognostic significance of serum S100B surgically resected melanoma patients participating in Intergroup Trial ECOG 1694. *J Clin Oncol.* 2008;27:38-44.
18. Mocellin S, Zavagno G, Nitti D. The prognostic value of serum S100B in patients with cutaneous melanoma: A meta-analysis. *Int J Cancer.* 2008;123:2370-6.
19. Guo HB, Stoffel-Wagner B, Bierwirth T, et al. Clinical significance of serum S-100 in metastatic malignant melanoma. *Eur J Cancer* 1995;31A: 924-928.
20. Mochiki E, Kuwano H, Katoh H, et al. Evaluation of 18F-2-deoxy-2-fluoro-D-glucose positron emission tomography for gastric cancer. *World J Surg.* 2004;28:247-53.
21. Wong RJ, Lin DT, Schoder H, et al. Diagnostic and prognostic value of [(18)F] fluorodeoxyglucose positron emission tomography for recurrent head and neck squamous cell carcinoma. *J Clin Oncol.* 2002;20:4199-208.
22. Bastiaannet E, Hoekstra OS, Oyen WJ, et al. Level of fluorodeoxyglucose uptake predicts risk for recurrence in melanoma patients presenting with lymph node metastases. *Ann Surg Oncol.* 2006;13:919-26.
23. Nagengast WB, Hooge MN, van Straten EM, et al. VEGF-SPECT with <sup>111</sup>In-bevacizumab in stage III/IV melanoma patients. *Eur J Cancer.* 2011;47:1595-602.
24. Ghanem G, Loir B, Morandini R, et al. On the release and half-life of S100B protein in the peripheral blood of melanoma patients. *Int J. Cancer;*2001;94:586-90.
25. Hamacher K, Coenen HH, Stocklin G. Efficient stereospecific synthesis of no-carrier-added 2-[18F]-fluoro-2-deoxy-D-glucose using aminopolyether supported nucleophilic substitution. *J Nucl Med* 1986;27:235-8.
26. Keyes JW Jr. SUV: standard uptake or silly useless value? *J Nucl Med* 1995;36:1836-9.

27. Sugawara Y, Zasadny KR, Neuhoff AW, et al. Re-evaluation of the standardized uptake value for FDG: variations with body weight and methods for correction. *Radiology* 1999;213:521-5.
28. Baas PC, Schraffordt Koops H, Hoekstra HJ, et al. Groin dissection in the treatment of lower-extremity melanoma. Short-term and long-term morbidity. *Arch Surg*. 1992; 127:281-6.
29. Martenson ED, Hansson LO, Nilsson B, et al. Serum S-100b protein as a prognostic marker in malignant cutaneous melanoma. *J Clin Oncol*. 2001;19:824-31.
30. Smit LH, Korse CM, Hart AA, et al. Normal values of S-100B predict prolonged survival for stage IV melanoma patients. *Eur J Cancer*. 2005;41:386-92.
31. Mohammed MQ, Abraha HD, Sherwood RA, et al. S. Serum S100 beta protein as a marker of disease activity in patients with malignant melanoma. *Med Oncol*. 2001; 18:109-20.
32. Bonfrer JM, Korse CM, Israels SP. Serum S-100 has prognostic significance in malignant melanoma. *Anticancer Res*. 1997; 17:2975-7.
33. Hamberg AP, Korse CM, Bonfrer JM, et al. Serum S-100B is suitable for prediction and monitoring of response to chemoimmunotherapy in metastatic malignant melanoma. *Melanoma Res*. 2003;13:45-9.
34. Hauschild A, Engel G, Brenner W, Glaser R, et al. Predictive value of serum S-100B for monitoring patients with metastatic melanoma during chemotherapy and/or immunotherapy. *Br J Dermatol*. 1999;140:1065-71.
35. Bouwhuis MG, Suci S, Kruit W, et al. Prognostic value of serial blood S-100B determinations in stage IIB-III melanoma patients: A corollary study to EORTC trial 18952. *Eur J Cancer*. 2011;47:361-8.
36. Jing Lin, Qingyuan Yang, Wilder PT, et al. The calcium-binding protein S-100B Down-regulates p53 and apoptosis in malignant melanoma. *J Biol Chem* 2010;27: 27487-98.
37. Markowitz J, Mackerell AD Jr, Carrier F, et al Design of inhibitors for S-100B. *Curr Top Med Chem*. 2005;5:1093-108.
38. Ferrara N, Davis-Smyth T. The biology of vascular endothelial growth factor. *Endocr Rev*. 1997; 18:4-25.
39. Ferrara N. Role of vascular endothelial growth factor in physiologic and pathologic angiogenesis: Therapeutic implications. *Semin Oncol*. 2002; 29:10-14.
40. van der Bilt JDW, Borrel Rinkes IHM. Surgery and angiogenesis; *Biochemica et Biophysica Acta (BBA) Rev Cancer*. 2004;1654:95-104.
41. [Http://public.ukcrn.org.uk/search/StudyDetail.aspx?StudyID=1751](http://public.ukcrn.org.uk/search/StudyDetail.aspx?StudyID=1751).
42. Willett CG, Duda DG, di Tomaso E, et al. Complete pathological response to bevacizumab and chemoradiation in advanced rectal cancer. *Nat Clin Pract Oncol*. 2007; 4:316-21.
43. The value of pre operative S-100B and SUV in clinically stage III melanoma patients undergoing therapeutic lymph node dissection. Kruijff S, Bastiaannet E, Speijers MJ, et al. *Eur J Surg Oncol*. 2011;37:225-32.
44. Sperti C, Pasquali C, Chierichetti F, et al. 18 Fluorodeoxyglucose positron emission tomography in predicting survival of patients with pancreatic carcinoma. *J Gastrointest Surg*. 2003;7:953-9.
45. Jubb AM, Harris AL. Biomarkers to predict the clinical efficacy of bevacizumab in cancer. *Lancet Oncol*. 2010;11:1172-83.

# 8



# The current status of S-100B as a biomarker in melanoma | S Kruijff, HJ Hoekstra | Submitted

---

## INTRODUCTION

The incidence rate of cutaneous malignant melanoma has increased over the past decades and causes the majority of all skin cancer mortalities.<sup>1</sup> From 1989 to 2008, an annual 4.1% increase in thin melanomas ( $\leq 1.0$  mm) was reported in The Netherlands, suggesting a true melanoma epidemic.<sup>1</sup> The incidence in the Netherlands is expected to continue to rise from 2,400 new melanoma patients in the year 2000 to a level of 4,800 patients in 2015.<sup>2</sup> Melanoma is increasingly detected earlier, which might explain the growing incidence of thin melanomas.<sup>1,3</sup> When a primary cutaneous melanoma is diagnosed, a curative result can be achieved by means of a simple primary excision of the melanoma with a 1- or 2-centimeter margin according to the Breslow thickness.<sup>4</sup> Disseminated melanoma, on the contrary, is still difficult to treat and almost impossible to cure. The most important prognostic factors for mortality in melanoma remain Breslow thickness, ulceration, mitosis, and the presence of metastases, although the Breslow thickness continues to be the strongest prognostic marker for survival.<sup>5</sup>

About 90% of all melanoma patients have a primary melanoma (stage I or II).<sup>5</sup> Still, after primary excision, there is a 20-28% risk of loco regional recurrence or in-transit metastases, 26-60% risk of regional metastases, and 15-50% risk of distant metastases.<sup>6</sup> Despite campaigns and earlier detection, melanoma mortality increased in the Netherlands from 2.5 to 3.6 per 100,000 in males and from 2.0 to 2.4 per 100,000 in females between 1989-2003.<sup>7</sup> Survival rates for lymphogenic disseminated melanoma (AJCC stage III) remain unsatisfactory. Melanoma is an aggressive and highly metastatic disease, which can be fatal, with a rapid systemic dissemination. Approximately one third of all melanoma patients will eventually experience disease recurrence. While almost all organs can be involved, the most frequent target sites are the liver, bone, and brain. Today, for melanoma patients with palpable lymph nodes, a lymph node dissection is still the only available curative treatment with a 5-year survival of 40-78%.<sup>5</sup> Patients with distant metastases are almost incurable, with a mean survival of only 8-9 months and a 2-year survival of 18-41%.<sup>5</sup>



## TUMOUR MARKERS AND SYSTEMIC THERAPY

Currently, no systemic therapy has demonstrated an overall survival benefit, although recent studies of immunotherapy with ipilimumab and the introduction of BRAF pathway inhibitors have shown promising results.<sup>8-10</sup>

Tumour markers in general can be used for several purposes; for screening, as diagnostic instruments, for staging, as a prognostic tool, to detect a recurrence, and finally as a quality control assessment after therapy. Tumour markers are particularly suitable for assessing treatment results or to detect early recurrences.<sup>11</sup>

The ability to target the new types of therapy described above towards well-selected subgroups of patients with the use of biomarkers such as S-100B could increase the likelihood of benefit, and might improve therapeutic outcomes for future melanoma treatments. Additionally, monitoring S-100B after systemic therapy could provide a 'measurable reflection' of the response to treatment; therefore it might be used as a quality assessment.<sup>12</sup>

### *LDH and S-100B*

More than 60 years ago, Hill reported that lactate dehydrogenase (LDH) could be a prognostic marker for melanoma patients.<sup>13</sup> LDH as a tumour marker is very useful for detecting distant metastases, but has a very low sensitivity and seems to be correlated with other malignancies as well.<sup>14</sup> Nevertheless, LDH is still the most prominent tumour marker in melanoma today, and is used in the American Joint Committee on Cancer (AJCC) staging system for stage IV.<sup>5</sup>

S-100B is a 21 kilo Dalton protein that was first isolated by Moore from the central nervous system of vertebrates.<sup>15</sup> S-100B is chiefly found in glial- and Schwann cells. The name is derived from the protein's 100% solvency in saturated ammonium sulphate at neutral pH.<sup>15</sup> S-100B has calcium-binding properties and, as a dimer, consists of two isomers, a and b. All possible combinations can occur (S-100aa, S-100ab, and S-100bb).<sup>16</sup> The S-100B protein is of neuroectodermal and mesodermal origin, and is expressed in various parts of the body.<sup>17</sup>

S100B is found diffusely in the cell cytoplasm, and a substantial fraction is bound to membranes. Moreover, there is strong evidence suggesting that it is involved in cytoskeletal regulation, and has a possible role in cell cycle progression.<sup>17</sup> However, the mechanism by which this protein is shed into the blood by malignant melanocytes is unknown.

### *S-100B concentrations related to tumour stage*

An increasing number of studies found serum S-100B protein levels to be a useful serum marker in melanoma, particularly in the late stages of the disease. In the past decade, serum S-100B is increasingly analyzed during melanoma treatment in various AJCC melanoma stages. The mean serum concentration of S-100B protein is significantly related to clinical melanoma stage, with the lowest levels in stages I and II (0-9%), elevated levels in stage III (5-98%), and high levels in stage IV (40-100%).<sup>18</sup> The highest S-100B concentrations are found in patients with bone or liver metastases.<sup>19</sup>

The evident correlation, which was found with both tumour burden and survival, suggests that S-100B in melanoma patients' blood is not a result of over expression or increased release, but a result of a loss of cell integrity and proteolytic degradation associated with apoptosis and cell death.<sup>17</sup>

Biomarkers are increasingly important in staging and defining prognosis in patients with metastasized melanoma. In the future, biomarkers could be increasingly used to target therapy towards well-selected subgroups of patients to increase the benefit of and improve therapeutic outcomes for future melanoma treatments.

S-100B is a promising diagnostic marker, which is correlated with most stages of melanoma.<sup>18</sup> Therefore, the current status of the applicability of S-100B as a tumour marker in the treatment of melanoma is discussed for the various melanoma AJCC tumour stages.

### *S-100B in AJCC stage I and II*

Primary melanoma is accompanied by an average 10-year survival in 93-97% of patients with T1aN0M0 melanoma. In the case of poor prognostic factors like thick melanoma (Breslow > 4 mm) with ulceration (T4bNoMo, AJCC stage IIC), the expected 10 year survival decreases to 39%.<sup>4</sup> The detection of micro-metastases by means of a sentinel node biopsy (SLNB) has a great influence on prognosis.<sup>4</sup> Recently, the results of SLNB followed by immediate lymph adenectomy, when compared with nodal observation and delayed lymph adenectomy for nodal recurrence, provided a significant 10-year survival advantage of 60.9% versus 41.8% ( $p = 0.01$ ).<sup>20</sup>


Less invasive diagnostics like FDG-PET/CT or the determination of S-100B in AJCC stage I and II seem inadequate for detecting micro metastases. FDG-PET/CT has a 5% sensitivity and a negative predictive value of 78% for appropriate prediction of lymph node status.<sup>21</sup> Neither S-100B nor LDH are suitable for predicting sentinel node status.<sup>22</sup> Therefore, there is no place for assessing serum LDH or S-100B nor the use of FDG-PET CT in AJCC stage I and II melanoma.<sup>21-23</sup>

### *S-100B in AJCC stage III melanoma*

Melanoma patients with palpable nodes, AJCC stage IIIb, have an average 5 year survival of 39-70%.<sup>4</sup> A therapeutic lymph node dissection, with or without postoperative radiotherapy, is the only available possible curative treatment.<sup>24,25</sup> Survival of this stage of melanoma is related to the number of lymph nodes with metastases, and the size of the malignant nodes.<sup>4</sup>

Balch et al. analyzed 2313 patients with AJCC stage III disease, and the 5-year overall survival was 63% for the cohort. However, when focusing on more specific patient groups, a tremendous heterogeneity in 5-year survival rates was observed (23-87% 5-year survival).<sup>26</sup> Preoperative elevated S-100B in patients with palpable nodes in optimally-staged (i.e. PET/CT) clinical stage III melanoma is associated with decreased disease-free survival (Figure 1).<sup>14, 26-29</sup> S-100B seems to be a strong prognostic marker for lymphogenic metastasized melanoma.<sup>14, 26-29</sup> S-100B is located in the cytoplasm of melanoma cells, and elevated S-100B probably occurs as the result of lost integrity of the melanoma cells.<sup>15,17</sup> Therefore,

elevation of S-100B should be interpreted as a process of subclinical microscopic metastatic disease, and preoperative elevated S-100B could be an expression of early dissemination not detected by today's most advanced imaging techniques.<sup>14, 26-29</sup>



**FIGURE 1.** Disease-specific survival and disease-free survival in patients with elevated S-100B and melanoma AJCC stage IIIB disease.<sup>29</sup>

Recently at this centre, a phase 1 study was published in which S-100B was evaluated to measure therapy effects after bevacizumab-induction treatment in AJCC stage III melanoma in advance of lymph node dissection.<sup>11</sup> This feasibility study showed that induction treatment with bevacizumab 4 weeks prior to surgery may be related to induction of tumour necrosis, but more importantly, when tumour necrosis was found in the dissection preparation, a 100% decline in S-100B was noted.<sup>12</sup> Correspondingly, the absence of tumour necrosis was correlated with ongoing S-100B elevation. The ability to target therapy towards well-selected subgroups of patients with the use of biomarkers, such as S-100B, could increase the likelihood of benefits, and might improve therapeutic outcomes for future melanoma treatments and strategies.<sup>11,14</sup>

### *S-100B in AJCC stage IV*

Patients with distant metastases (AJCC stage IV) have a short 2-year survival of 18-40%.<sup>5</sup>

For a select group of patients with 1 to 3 metastases, who could be operated on with curative intent, a 5-year survival of 40% can be achieved.<sup>5</sup> Treatment results for advanced melanoma have been very unsatisfactory. Patients with stage IV melanoma have traditionally been managed with various systemic treatments; however, overall survival using this approach has been disappointing. Most melanoma patients in this staging group receive palliative systemic treatment or palliative radiotherapy. In melanoma patients with generalized metastases, S-100B is elevated in 40-100% of cases and represents tumour load.<sup>14,18</sup> When S-100B is not elevated earlier in stage IV melanoma, survival was significantly longer.<sup>30</sup> On the contrary, continuing elevation of S-100B during treatment with systemic chemotherapy or immunotherapy is a sign of progressive disseminating disease.<sup>30,31</sup> An effective therapy measured by the RECIST criteria is often correlated with declining S-100B concentrations.<sup>32</sup>

For decades, the progressive treatment of metastasized melanoma has been disappointing with no survival benefit. No systemic therapy has been shown to affect overall survival. Currently however, two promising treatment concepts have arisen. Immunotherapy with ipilimumab, and the introduction of B-Raf proto-oncogene serine/threonine-protein kinase (BRAF) pathway inhibitors have shown spectacular results.<sup>8-10</sup> First, on March 25, 2011, the Food and Drug Administration approved the cytotoxic T-lymphocyte antigen-4 monoclonal antibody ipilimumab (Yervoy), making it the first agent indicated for non-resectable or metastatic melanoma in more than a decade.<sup>7</sup> Second, since the identification in 2002 of a mutation in the BRAF gene in a subset of melanoma patients, research focused on developing inhibitors of the mutated BRAF protein as a therapeutic target in disseminated melanoma.<sup>9,10</sup> Recently, Phase 1 and 2 clinical trials of the BRAF kinase inhibitor vemurafenib (PLX4032) showed response rates of more than 50% in patients with metastatic melanoma with the BRAF V600E mutation.<sup>10</sup> On August 17, the FDA approved vemurafenib (Zelboraf) as a treatment for late-stage melanoma, since it was shown to extend patients' survival. Now that an effective (adjuvant) therapy for loco-regional metastatic and disseminated melanoma seems to have been discovered, the role of S100-B in the treatment of melanoma disease may change dramatically.<sup>33</sup>

## DISCUSSION

Various studies have reported on the use of S-100B as a tumour marker during melanoma treatment. Effective therapy by means of a lymph node dissection, systemic therapy, or immunotherapy is associated with declining S-100B levels, whereas elevated S-100B concentrations correlated with disease progression.<sup>14,27-29</sup> The prognostic biochemical marker, S-100B may be of value in the design of future trials on adjuvant systemic chemotherapy or immunotherapy.

Presently, LDH is still the most prominent serum parameter in stage IV melanoma, and is still part of the AJCC staging system.<sup>4</sup> For optimally-staged clinical stage III melanoma patients, it is clear that preoperative S-100B elevation is associated with significantly poorer survival.<sup>14,29</sup> These results are also supported by an ECOG study in high-risk surgically-resected melanoma by Kirkwood.<sup>27</sup> In this study, postoperative (baseline) S-100B was indeed a prognostic marker for disease-free survival and overall survival in stage III melanoma patients. Recently, Bouwhuis et al. showed that elevated S-100B levels in serial serum measurements in stage III melanoma is of very significant prognostic value, which was even stronger than disease stage and number of positive lymph nodes.<sup>28</sup>

Therefore, S-100B could be used for patient selection for adjuvant systematic treatment in the stratification for new adjuvant therapeutic trials as well as to provide information for stage III melanoma patients who want to be informed about disease prognosis.

Also, in the near future, serum S-100B can also be used to assess response to induction therapy with systemic treatment for nodal disease.<sup>12</sup> The ability to target therapy towards well-selected patient subgroups with the use of biomarkers such as S-100B could increase the likelihood of treatment benefits, and might improve therapeutic outcomes for future melanoma treatments.<sup>12</sup> Not only should elevated S-100B be seen as a prognostic factor for survival and a reflection of disease progression, the presence of the S-100B protein itself induces disease progression as well. S-100B interacts with p53, thereby down-regulating its function as a tumour suppressor protein by preventing induction of apoptosis of potential melanoma cells.<sup>34,35</sup> Taking these results into account, an induction therapy might slow down melanoma proliferation in stage III in two ways; first, by inducing tumour necrosis and second, by temporarily suppressing serum S-100B concentrations, thus preventing down regulation of the tumour suppressor protein p53.

## CONCLUSION

S-100B can be used to perform a individualized risk assessment before administering adjuvant treatment on the one hand, and to measure response to induction therapy for nodal disease on the other.<sup>12,14</sup> Despite these prognostic values, S-100B is not suitable for routine screening purposes in stage I-III melanoma patients, or for the early detection of recurrences during follow-up of melanoma patients. The results of these measurements do not yet translate into

an adequate therapeutic intervention or survival benefit because of limited treatment options in advanced melanoma. However, now that an effective (adjuvant) therapy for loco-regional metastatic and disseminated melanoma seems to have been discovered, the routine use of biomarkers like S-100B may change significantly in the future.<sup>33</sup>

## REFERENCES


1. Hollestein LM, van den Akker SA, Nijsten T, et al. Trends of cutaneous melanoma in The Netherlands: increasing incidence rates among all Breslow thickness categories and rising mortality rates since 1989. *Ann Oncol*. 2011 May 4, Epub ahead of print.
2. De Vries E, Houterman S, Janssen-Heijnen ML, et al. Up-to-date survival estimates and historical trends of cutaneous malignant melanoma in the south-east of the Netherlands. *Ann Oncol* 2007;18:1110-6.
3. Reinhardt MJ. Cutaneous melanoma. *Recent Results Cancer Res* 2008;170:151-7.
4. Sladden MJ, Balch C, Barzilai DA, et al. Surgical excision margins for primary cutaneous melanoma. *Cochrane Database Syst Rev* 2009;4:CD004835.
5. Balch CM, Greshenwald JE, Soong SJ, et al. Final version of 2009 AJCC melanoma staging and classification. *J Clin Oncol*.2009;27:6199-206.
6. Garbe C, Eigentler TK. Diagnosis and treatment of cutaneous melanoma: state of the art 2006. *Melanoma Res* 2007;17:117-127.
7. Siesling S, Visser O, van Dijk JA, et al. Trends in the incidence and death from cancer from 1989-2003 in The Netherlands. *Ned Tijdschr Geneeskd* 2006;150:2490-96.
8. Hodi FS, O'Day SJ, McDermott DF, et al. Improved survival with ipilimumab in patients with metastatic melanoma. *New Engl J Med* 2010;363:711-23.
9. Flaherty KT, Puzanov I, Kim KB, et al. Inhibition of mutated, activated BRAF in metastatic melanoma. *N Engl J Med*. 2010;363:809-19.
10. Chapman PB, Hauschild A, Robert C, et al. Improved survival with vemurafenib in melanoma with BRAF V600E mutation. *N Engl J Med*. 2011;364:2507-16.
11. Keuren JF, Thomas CM, Bonfrère JM, Sweep CG, Boonstra JG. Assessment of tumor markers in clinical practice. *Ned Tijdschr Geneeskd*. 2009;153:A617.
12. S. Kruijff, E Bastiaannet, A.H. Brouwers, et al. The use of S-100B to evaluate therapy effects during bevacizumab induction treatment in AJCC stage III melanoma. *Ann Surg Oncol*. 2011; published online 23 august 2011.
13. Hill BR, Levi C. Elevation of a serum component in neoplastic disease. *Cancer Res* 1954;14:513-15.
14. Kruijff S, Bastiaannet E, Muller-Kobold AC, et al. S-100B concentrations predict disease-free survival in stage III melanoma patients. *Ann Surg Oncol* 2009;16:3455-62.
15. Moore BW. A soluble protein characteristic of the nervous system. *Biochem Biophys Res Commun* 1965;19:739-44.
16. Smit LH, Korse CM, Hart AA, et al. Normal values of S-100B predict prolonged survival for stage IV melanoma patients. *Eur J Cancer*. 2005;41:386-92.
17. Ghanem G, Loir B, Morandini R, et al. On the release and half-life of S-100B protein in the peripheral blood of melanoma patients. *Int J Cancer* 2001;94:586-90.
18. Martenson ED, Hansson LO, Nilsson B, et al. Serum S-100b protein as a prognostic marker in malignant cutaneous melanoma. *J Clin Oncol*. 2001;19:824-31.
19. Mohammed MQ, Abraha HD, Sherwood RA, et al. S. Serum S100 beta protein as a marker of disease activity in patients with malignant melanoma. *Med Oncol*. 2001; 18:109-20.
20. Morton DL, Thompson JF, Cochran AJ, et al. Sentinel node biopsy and immediate lymphadenectomy for occult metastases versus nodal observation and delayed lymphadenectomy for nodal recurrence. *Ann Surg Oncol* 2010, 17 Suppl 1, S22-S23 (Abstract 53)
21. Klode J, Dissmond J, Grabbe S, et al. Sentinel lymph node excision and PET-CT in the initial stage of malignant melanoma: A retrospective analysis of 61 patients with malignant melanoma in American Joint Committee on cancer stages I and II. *Dermatol Surg* 2010;36:439-45.
22. Egberts F, Momkvist A, Egberts JH, Kaehler KC, Hauschild A. Serum S-100B and LDH are not useful in predicting the sentinel node status in melanoma patients. *Anticancer Res*. 2010;30:1799-805.
23. Havenga K, Cobben DC, Oyen WJ, et al. Fluorodeoxyglucose-positron Emission tomography and sentinel lymph node biopsy in staging primary cutaneous melanoma. *Eur J Surg Oncol*. 2003;29:662-4.
24. Bastiaannet E, Beukema JC, Hoekstra HJ. Radiation therapy following lymph node dissection in melanoma patients: treatment, outcome and complications. *Cancer Treat Rev*. 2005;31:18-26.

25. M. A. Henderson, B. Burmeister, J. F. Thompson, et al. Adjuvant radiotherapy and regional lymph node field control in melanoma patients after lymphadenectomy; Results of an intergroup randomized trial (ANZMTG 01.02/TROG 02.01. *J Clin Oncol* 2009; 27:18s, 2009 (suppl; abstr LBA9084)
26. Balch CM, Gershenwald JE, Soong S, et al. Multivariate Analysis of Prognostic Factors Among 2,313 Patients With Stage III Melanoma: Comparison of Nodal Micrometastases Versus Macrometastases. *J Clin Oncol.* 2010;28:2452-9.
27. Tarhini AA, Stuckert J, Lee S et al. Prognostic significance of serum S-100B surgically resected melanoma Patients Participating in Intergroup Trial ECOG 1694. *J.Clin.Oncol.* 2008; 27:38-44.
28. Bouwhuis MG, Suci S, Kruit W, et al. Prognostic value of serial blood S100B determinations in stage IIB-III melanoma patients: A corollary study to EORTC trial 18952. *Eur J Cancer.* 2011;47: 361-8.
29. Kruijff S, Bastiaannet E, Speijers MJ, Kema IP, Van Ginkel RJ, Hoekstra HJ. S-100B Concentrations Predict Disease Specific Survival in AJCC Stage IIIMelanoma Patients. *Ann Surg Oncol* 2011; 18 Suppl 1, S1-S144, P227.
30. Hamberg AP, Korse CM, Bonfrer JM, de Gast GC. Serum S-100B is suitable for prediction and monitoring of response to chemoimmunotherapy in metastatic malignant melanoma. *Melanoma Res.* 2003;13:45-9.
31. Kluger HM, Hoyt K, Bacchiocchi A, et al. Plasma markers for identifying patients with metastatic melanoma. *Clin Cancer Res.* 2011;15:2417-25.
32. Eisenhauer EA, Therasse P, Bogaerts J, et al. New response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1). *Eur J Cancer.*2009;45:228-47.
33. Garbe C, Peris K, Hauschild A, et al. Diagnosis and treatment of melanoma: European consensus-based interdisciplinary guideline. *Eur. J. Cancer* 2010; 46:270-283.
34. Jing Lin, Qingyuan Yang, Wilder PT, et al. The calcium-binding protein S-100B Down-regulates p53 and apoptosis in malignant melanoma. *J Biol Chem* 2010; 27: 27487-98.
35. Markowitz J, Mackerell AD Jr, Carrier F, Charpentier TH, Weber DJ. Design of inhibitors for S-100B. *Curr Top Med Chem.* 2005; 5:1093-108.



# 9

---



## General conclusion and future perspectives

---

As described in this thesis the protein S-100B is a strong prognostic marker in stage III and IV melanoma and is currently the best studied biomarker in melanoma disease. It has the potential to identify high risk stage III melanoma patients who may benefit from adjuvant systematic treatment in the future. The following paragraph describes aspects of currently ongoing research and future research in the development of follow up, elderly, biomarkers in stage III disease and future strategies of metastatic melanoma treatment.

### **FOLLOW-UP**

There is no international consensus on the follow-up of melanoma patients and in many countries high-frequency follow-up after treatment for melanoma is still standard care.<sup>1</sup>

The main purpose of follow-up is early detection of recurrent disease and second primary melanomas that could be treated successfully by surgery or other treatment modalities. However, it is still difficult to predict the behavior of melanoma in the individual patient although three quarters of the melanoma recurrences are still patient detected.<sup>2</sup> Generally, overall treatment results of metastatic or recurrent disease have been disappointing.

In the first chapter in which we evaluated the role of the method of detection in nodal disease for the prognosis, almost 50% of lymph node metastases in Stage I-II melanoma patients are patient detected.<sup>2</sup> Besides, the method of detection of lymph node recurrence by patient or physician didn't have any impact on survival. The data of this study have added to the controversy about the value of high frequency follow-up regimens in melanoma.<sup>3,4</sup> A prospective, randomized, high quality methodological research has been started in order to develop meaningful applicable guidelines (MELFO).<sup>5,6,7</sup> This study, currently conducted in the Netherlands and the UK, randomizes stage IB–IIC patients between the follow-up schedule proposed by the national guideline and an evidence-based schedule with a reduced follow-up

frequency.<sup>5</sup> With its primary end point being health-related quality of life, these results should clarify if reduced follow-up frequency has any effect on patient well-being. Although melanoma can recur more than 10 years after diagnosis, this concerns a very small percentage of patients. We conclude that it is not necessary to keep patients in life-long follow-up.<sup>6,7</sup> During follow-up, doctors are recommended to emphasize educating the patient in detecting recurrences and second primary melanoma, in order to promote early detection. Self-examination of nodes basins and skin changes seems suitable for many patients, but could be improved for optimal benefit, for example, by using the internet.<sup>8,9</sup>

To date, there is no evidence for routine imaging studies, for example PET scanning, or routine measurement of any available serum biomarkers, such as S-100B and LDH, in follow-up for melanoma. Only clinical stage III melanoma patients should be staged with whole-body FDG-PET, PET-CT or spiral-CT as well as using the serum biomarker S-100B.<sup>10,11</sup> Regional treatment, curative or palliative, should be based on the extent of the disease. A randomized clinical trial showed that adjuvant radiotherapy improved regional control in melanoma patients at high risk of regional relapse after lymphadenectomy without affecting survival.<sup>12</sup>

## ELDERLY

For the elderly patient various studies have already shown that the incidence and mortality of melanoma in the older population (>65 yrs) is growing rapidly.<sup>13-17</sup> As the geriatric population increases melanoma will become an important health issue for the elderly age group in this century.<sup>18-21</sup> As was described in the third chapter the incidence for the elderly increased much more rapidly than for the younger melanoma patients. Besides, in contrast to younger patients, elderly patients are still diagnosed with the thickest melanomas. Especially in elderly men the percentage thick melanoma's declined only minimally the last decades. This finding probably explains a worse relative survival for the elderly melanoma patient when compared to the younger population. Literature reveals that awareness in Europe is not at such a high level despite an increasing incidence and focus has been mainly on the young patient.<sup>15,21,22</sup> This might explain the high percentage of thick melanoma's in the elderly who, looking at the rising incidence, seem to have started to pay the costs for earlier sun abuse in the era before melanoma became a health priority. For the future, campaigns in the Netherlands should focus on early detection of melanoma in general and especially on early detection in the elderly.

## BIOMARKERS IN MELANOMA

### *LDH*

Although LDH has a low sensitivity and therefore doesn't seem to be the adequate parameter in staging and follow-up for the earlier stages of melanoma (AJCC stage I, II and III), it is an

independent prognostic factor in disseminated disease. (AJCC stage IV) Almost 60 years ago Hill et al. already reported on increased LDH serum parameters in melanoma patients.<sup>23</sup> In the 1970s, LDH was reported to be an indicator for liver metastases with a specificity of 95% and a sensitivity of 83% in AJCC stage II patients and, respectively, 87 and 57% in stage III patients.<sup>24</sup> Even stage I disease patients with elevated serum LDH seemed to have a significant decreased survival.<sup>25</sup> Weighing today's evidence, LDH has high specificity for melanoma and literature demonstrates that LDH is especially elevated in advanced disease, predominantly when disease has progressed to the liver.<sup>26</sup> Deichmann et al. claimed LDH to be the only significant marker for disease progression when analyzing the marker in combination with S-100B and MIA in stage IV disease.<sup>27</sup> LDH can indeed be used as a prognostic marker for detecting distant metastases (AJCC stage IV) but has a poor sensitivity and is not melanoma specific. Despite these facts, today LDH is still part of the AJCC staging manual for patients with distant metastases (stage IV M1c).<sup>28</sup>

### *S- 100B*

Increased S-100B levels were first detected in melanoma patients in 1980.<sup>29</sup> S-100B is a 21-kDa protein that was first isolated from the CNS in vertebrates. Later, S-100B was found to be a serological tumor marker for melanoma and seemed to be mostly increased in stage III and IV.<sup>30,31</sup> According to the literature to date, the proportions of patients with elevated S-100B concentrations are 0-9% in stage I/II, 5-98% in stage III and 40-100% in stage IV.<sup>11</sup> The highest S-100B concentrations were found in patients with liver and bone metastases.<sup>32,33</sup> It can, therefore, be concluded that S-100B concentrations are correlated with the clinical stage of the disease. Several studies have recommended the use of S-100B as a tumor marker to monitor the course of the disease in stage III melanoma or to evaluate the effect of therapy in stage IV disease. Successful treatment with lymph node dissection, chemotherapy or immunotherapy was associated with decreased S-100B concentrations, whereas increased concentrations were an expression of disease progression.<sup>11,34,35</sup> Today, for AJCC stage III melanoma patients, a therapeutic lymph node dissection is still the first choice of treatment. The lymphogenic disseminated AJCC stage III melanoma patient group is known to be a patient group with a remarkable heterogeneity which was published by Balch in a multivariate analysis of prognostic factors among 2,313 patients with stage III melanoma. The five year overall survival was 63% varying from 67% for patients with nodal micro-metastases to 43% for nodal macrometastases.<sup>36</sup> Recently it was also reported that S-100B can be used as a prognostic marker for disease-free survival and overall survival directly after therapeutic lymph node dissection in stage III melanoma.<sup>37-40</sup> Data from our centre were not only in concordance with the above described findings, but also revealed that S-100B can be used as a prognostic marker for disease-free survival when determined before dissection after staged with whole-body FDG-PET, PET-CT or spiral-CT. Multivariate analysis showed that preoperative elevated S-100B in PET, PET-CT or spiral-CT negative patients in stage III melanoma was an important predictor for a significantly worse survival.<sup>11</sup> The hypothesis that elevating S-100B should be interpreted

as a process of active hematogenic and lymphogenic dissemination seems to have been confirmed by these data. Thus, clinical stage III melanoma patients should be staged with whole-body FDG-PET, PET-CT or spiral-CT as well as the use of serum biomarker S-100B.<sup>11</sup> Further regional treatment, curative or palliative, should be based on the extent of the disease.

In the future, S-100B can thus be used to perform a more refined individualized risk assessment before administering adjuvant treatment. Above all, it provides information to stage III melanoma patients who want to be optimally informed about their disease prognosis. Despite the described prognostic values, this biomarker is not suitable for routine screening purposes in stage I-III melanoma patients or for the early detection of recurrence in the follow-up of melanoma patients; that is, early detection of recurrences in follow-up as the results of these measurements do not yet translate into an adequate therapeutic intervention nor survival benefit due to limited treatment options in advanced melanoma.

## **COMPLETE LYMPH NODE DISSECTION FOR REGIONAL NODAL METASTASIS**

The primary management of lymph nodes involved with metastatic melanoma is regional lymphadenectomy. Axillary or inguinal node complete lymph node dissection (CLND) is performed after occult metastases are found by sentinel lymph node biopsy, or after a clinically apparent regional lymph node metastasis. CLND completely removes all lymph-node-bearing tissue in a nodal basin. Lymph node dissections are known for a considerable morbidity. Whereas a reduction through technical modifications seems to be difficult, efforts to limit the procedures to defined patient subgroups may prove to be a more promising .

SLNB has helped spare the majority of melanoma patients, who are node-negative, from the morbidity of ELND.

Whether or not CLND in patients with positive sentinel lymph nodes increases overall melanoma specific survival remains unclear. The procedure continues to be controversial. No randomized prospective studies have yet determined the survival advantage of CLND. Most national guidelines recommend that all patients with stage III melanoma still have a CLND.<sup>41,42</sup>

Underway is the second MSLT (MSLT-II), a randomized trial that will accrue 4,500 patients from more than 30 countries.<sup>43</sup> Results of MSLT-II will indicate whether removal of the SN alone is adequate nodal surgery in certain patients with regional metastatic melanoma and this trial is therefore expected to clarify this issue.<sup>43</sup>

## STAGE IV DISEASE

### *Surgery in patients with stage IV metastatic melanoma*

Treatment results for advanced melanoma have been very unsatisfactory. Patients with stage IV melanoma have traditionally been managed with various systemic treatments; however, overall survival with this approach has been disappointing. Surprisingly, findings of many retrospective, single-institution, and multicentre studies suggest that participants treated with complete metastasectomy for stage IV metastases have an enhanced overall 5-year survival.<sup>38</sup> Systemic treatment alone for patients with stage IV melanoma have not yet conferred the survival advantage of a complete metastasectomy.<sup>44</sup>

One of the strong believers of metastasectomy in patients with stage IV melanoma, Donald Morton, launched an international trial.<sup>45,46</sup> To be eligible for the study, all participants had to undergo complete metastasectomy for stage IV metastases with tumor free surgical margins. After surgery, patients were randomized to either adjuvant immunotherapy with onmelatucel-L or placebo. Although the main aim of the trial-to assess onmelatucel-L as an adjuvant treatment after surgery-did not show a survival benefit, two important conclusions were made. First, correct selection of patients is vital. Only people with three or fewer visceral sites of stage IV disease who could be rendered clinically and radio graphically free of disease were eligible for this trial. Second, uniformity of resection can be accomplished. In an international, randomized controlled trial surgeons from around the world undertook similar complete metastasectomy procedures for various anatomical sites and achieved tumor-free surgical margins. This similarity in resections led to an 40% 5-year survival for the entire study cohort.<sup>46</sup> Such results for 5-year survival have not yet been achieved in any study of chemotherapy or biological treatment for patients with stage IV melanoma.<sup>45,46</sup> Considering the evidence, the conclusion has to be drawn that complete surgical resection of metastatic disease to stage IV sites-including skin, soft tissue, distant lymph nodes, lungs, or other non-CNS visceral regions-offers the best chance for prolonged survival. Therefore, it seems the time has come to reposition surgery as the first option for properly selected patients with stage IV disease, as long as complete metastasectomy can be undertaken and there is no better alternative systemic treatment.<sup>47</sup>

Of course the use of this aggressive surgical approach should be tempered with the knowledge that incomplete resections put patients at increased risk and without any proven survival benefit these should be reserved only for palliation of symptoms. Thus, the first step for people with newly diagnosed stage IV disease should be assessment of resectability by a skilled surgical oncologist.<sup>44</sup>

### *Adjuvant therapy*

Adjuvant systemic chemotherapy or immunotherapy with various nonspecific immunostimulatory agents in the treatment of high-risk melanoma is not indicated outside the context of a clinical trial.<sup>48</sup> However, adjuvant immunotherapy with high intermediate- and low-dose

IFN- $\alpha$  was extensively studied during the last decade. No systemic therapy has demonstrated to affect overall survival. Currently however, two promising treatment concepts are being explored. The recent immunotherapy with ipilimumab and the introduction of the BRAF pathway inhibitors have shown spectacular results. First, on March 25, 2011, the Food and Drug Administration approved cytotoxic T-lymphocyte antigen-4 monoclonal antibody (Ipilimumab), making it the first agent indicated for unresectable or metastatic melanoma in more than a decade.<sup>49,50</sup> Second, since the identification in 2002 of a mutation in the B-Raf proto-oncogene serine/threonine-protein kinase (BRAF) gene in melanoma, research is focused on developing inhibitors of the mutated BRAF protein as a therapeutic target in disseminated melanoma.<sup>51</sup> Recently, Phase 1 and 2 clinical trials of the BRAF kinase inhibitor vemurafenib (PLX4032) have shown response rates of more than 50% in patients with metastatic melanoma with the BRAF V600E mutation. At 6 months, overall survival was 84% in the vemurafenib group and 64% in the dacarbazine group. In the interim analysis for overall survival and final analysis for progression-free survival, vemurafenib was associated with a relative reduction of 63% in the risk of death and of 74% in the risk of either death or disease progression, as compared with dacarbazine ( $p < 0.001$  for both comparisons).<sup>52</sup>

Maybe, since an effective (adjuvant) therapy for loco-regional metastatic and disseminated melanoma seems to be discovered, the entire discussion with respect to follow-up intensity and routine imaging and use of biomarkers will be significantly altered.<sup>53</sup>

### *The use of biomarkers to evaluate therapy effects*

S-100B has the potential to identify high risk stage III melanoma patients who may benefit from adjuvant systematic treatment. As described in the sixth chapter, in the near future, with the aid of the biomarkers serum S-100B and LDH and Standardized Uptake Value (SUV) in FDG-PET, a response to induction therapy with systemic treatment might be observed for nodal disease. S-100B and SUV are known to be of prognostic value in stage III melanoma and elevated S-100B and SUV in stage III melanoma patients can be specific indicators of disease progression.<sup>33-40,54,55</sup> Recently Bouwhuis et al revealed that an elevation of S-100B levels in serial serum measurements in stage III is a very significant prognostic value, which is even stronger when compared with disease stage and number of positive lymph nodes.<sup>56</sup> However, not only should elevated S-100B be seen as a prognostic factor for survival and a mere reflection of disease progression alone, the presence of the protein S-100B itself induces disease progression as well. S-100B interacts with p53 and thereby down-regulates its function as a tumor suppressor protein by preventing induction of apoptosis of potential melanoma cells.<sup>57,58</sup> Taking these results into account, an induction therapy might slow down melanoma proliferation in stage III in two ways. First, by inducing tumor necrosis and second by temporarily suppressing the serum S-100B concentrations and thus preventing down regulation of tumor suppressor protein p53. The ability to target therapy towards well selected subgroups of patients with help of biomarkers as S-100B in melanoma could increase the likelihood of benefit and might improve therapeutic outcome in the future.<sup>59</sup>

## MELANOMA CLINICS

The challenge for medicine in general and for treating melanoma is to create the right kind of focus in the health care delivery process. This can be achieved by creating special treatment centers in hospitals; units that excel in delivering one type of care.<sup>60</sup> For melanoma this implicates a need for creating integrated multidisciplinary practice units within focused melanoma clinics.<sup>61</sup> So in the coming years these dedicated multidisciplinary clinics are the next step in melanoma patient care. It is a crucial step towards higher treatment efficiency and efficacy; by increasing quality of care and reducing costs, higher Phase I, II and III trial participation and optimizing follow-up. The ultimate goal of melanoma clinics is to improve the survival of patients diagnosed with melanoma. Consequently, the University Medical Center Groningen has recently started a melanoma clinic in which dedicated nurse practitioners participate in the routine follow-up of melanoma patients and clinical trials to reduce the workload of specialists and improve patient satisfaction.<sup>62</sup> Additionally a part of melanoma care may possibly be transferred to GPs.<sup>62,63</sup> Individual patients prognostic models for stage I, II and III were recently developed, made available online and can be used for patient-tailoring initial management and subsequent follow-up.<sup>64,65</sup>

## RISING COSTS

The cost of cancer care is on the rise, from \$104 billion in 2006 to over \$173 billion in 2020. The rise in cost is driven by both the increasing cost in therapy and the extent of care.<sup>66,67</sup>

Efficacy of these new drugs often do not outweigh their cost-effectiveness, resulting in an increasing cost curve that is not sustainable. The new melanoma systemic therapies will definitely not improve this situation. For example, earlier this year, the Food and Drug Administration (FDA) approved the recent developed immunotherapy with ipilimumab which cost about \$120,000 per treatment. The definite price of Vemurafenib is not yet defined but will be likely to cost tens of thousands of dollars per year.

Medical oncologists directly or indirectly control or influence the majority of cancer care costs, including the use and choice of drugs, the types of supportive care, the frequency of imaging, and the number and extent of hospitalizations. Smith and Hillner recently suggested suggesting five changes in medical oncologists behavior and five changes in their attitudes and practice that would bend the cancer-cost curve downward.<sup>68</sup>

In conclusion, some areas of oncology - for example, clinical trials and curative as well as proven adjuvant treatments - should become off limits when it comes to primary cost considerations. This will probably be extraordinarily difficult as patients facing death have a very different and important perspective on risk.<sup>69</sup>



## CONCLUSION

Melanoma is the most aggressive form of skin cancer and its incidence is rising worldwide. While early stages of melanoma can be curatively treated by surgical excision, advanced stages have been uniquely resistant to current therapies. However, we now recognize that melanomas are far more variable at a molecular level than they appear under the microscope.

Therefore, melanoma is to be included in The Cancer Genome Atlas project and this will, in the future, provide more insight into molecular-based profiling studies in melanoma patients.<sup>70</sup> Therefore, rather than treating melanoma as a single disease, eventually molecular biology will unravel the mystery by which melanoma has been surrounded purely by stratifying tumors into molecular subtypes and treating each with the most appropriate therapies.

## REFERENCES


1. Francken AB, Bastiaannet E, Hoekstra HJ. Follow-up in patients with localised primary cutaneous melanoma. *Lancet Oncol* 2005; 6, 608–621.
2. Francken AB, Thompson JF, Bastiaannet E, Hoekstra HJ. Detection of the first recurrence in patients with melanoma: three quarters by the patient, one quarter during outpatient follow-up. *Ned Tijdschr Geneeskd*. 2008;152:557-62.
3. Kruijff S, Bastiaannet E, Suurmeijer AJ, Hoekstra HJ. Detection of melanoma nodal metastases; differences in detection between elderly and younger patients do not affect survival. *Ann Surg Oncol*. 2010;17:3008-14.
4. Speijers MJ, Francken AB, Hoekstra-Weebers JEHM, Bastiaannet E, Kruijff S; Hoekstra HJ. Optimal Follow-up for Melanoma. *Expert Review of Dermatology*, 2010;5:461-478.
5. [Http://dissertations.ub.rug.nl/FILES/faculties/medicine/2007/a.b.francken/melfo.pdf](http://dissertations.ub.rug.nl/FILES/faculties/medicine/2007/a.b.francken/melfo.pdf).
6. Francken AB, Hoekstra HJ. Follow-up of melanoma patients: the need for evidence-based protocols. *Ann Surg Oncol*. 2009;16:804-5.
7. Francken AB, Accortt NA, Shaw HM, Colman MH et al. Follow-up schedules after treatment for malignant melanoma. *Br J Surg*. 2008;1401-7.
8. [Http://www.youtube.com/watch?v=IFv16l-0Ztk](http://www.youtube.com/watch?v=IFv16l-0Ztk).
9. [Http://www.youtube.com/watch?v=P5qKpCjXaLA](http://www.youtube.com/watch?v=P5qKpCjXaLA).
10. Bastiaannet E, Wobbes T, Hoekstra OS et al. Prospective comparison of [18F] fluorodeoxyglucose positron emissiontomography and computed tomography in patients with melanoma with palpable lymph node metastases: diagnostic accuracy and impact on treatment. *J. Clin. Oncol*. 2009; 27: 4774-4780.
11. Kruijff S, Bastiaannet E, Muller Kobold AC et al. S-100B concentrations predict disease free survival in stage III melanoma patients. *Ann. Surg. Oncol*. 2009;16: 3455–3462.
12. Henderson MA, Burmeister B, Thompson JF et al. Adjuvant radiotherapy and regional lymph node control in melanoma patients after lymphadenectomy: results of an intergroup randomized trail (ANZMTG 01.02/TROG 02.01). Presented at: American Society of Clinical Oncology Annual meeting. Orlando, FL, USA, 29 May-2 June 2009.
13. Ulmer MJ, Tonita JM, Hull PR. Trends in invasive cutaneous melanoma in Saskatchewan 1970-1999. *J Cutan. Med Surg*. 2003;7:433-42.
14. Venzon DJ, Moolgavkar SH. Cohort analysis of malignant melanoma in five countries. *Am.J Epidemiol* 1984;119:62-70.
15. Kricke A, Armstrong BK, Goumas C, Litchfield M, Begg CB, Hummer AJ et al. Ambient UV, personal sun exposure and risk of multiple primary melanomas. *Cancer Causes Control* 2007;18:295-304.
16. Armstrong BK, Kricke A. Cutaneous melanoma. *Cancer Surv*. 1994;19-20:219-40.
17. Gaudette LA, Gao RN. Changing trends in melanoma incidence and mortality. *Health Rep*. 1998;10:29-41.
18. Testori A, Sotello J, Sances D et al. Cutaneous melanoma in the elderly. *Melanoma Res*. 2009;19:125-34.
19. Chang CK, Jacobs IA, Vizgirda VM, Salti GI. Melanoma in the elderly patient. *Arch Surg*. 2003 Oct;138:1135-8.
20. Kelly JW. Melanoma in the elderly- a neglected public health challenge. *Med J Aust*. 1998;169:403-4.
21. Criscione VD, Weinstock MA. Melanoma thickness trends in the United States, 1988-2006. *J Invest Dermatol*. 2010 Mar;130(3):793-7.
22. Levi F, Franceschi S, Te VC, Randimbison L, La VC. Trends of skin cancer in the Canton of Vaud, 1976-92. *Br.J Cancer* 1995;72:1047-53.
23. Hill BR, Levi C. Elevation of a serum component in neoplastic disease. *Cancer Res*. 1954; 14: 513–515.
24. Finck SJ, Guillianio AE, Morton DL. LDH and melanoma. *Cancer* 1983; 51: 840-843.
25. Campora E, Repetto L, Giuntini P et al. LDH in the follow up of stage I malignant melanoma. *Eur. J. Cancer Clin. Oncol*. 1988; 24: 277-278.
26. Cogas H, Eggermont AM, Hauschild A et al. Biomarkers in melanoma. *Ann. Oncol*. 2009; 20: 8-13.

27. Deichmann M, Benner A, Bock M et al. S-100B, melanoma-inhibiting activity, and lactate dehydrogenase discriminate progressive from nonprogressive American Joint Committee on Cancer stage IV melanoma. *J. Clin. Oncol.* 1999;17:1891-1896.
28. Balch CM, Greshenwald JE, Soong SJ, and et al. Final version of 2009 AJC melanoma staging and classification. *J Clin Oncol* 2009;27:6199-206.
29. Fagnart OC, Sindic CJ, Laterre C. Particle counting immunoassay of S100 protein in serum. Possible relevance in tumors and ischemic disorders of the central nervous system. *Clin. Chem.* 1988;34:1387-1391.
30. Smit LH, Korse CM, Hart AA et al. Normal values of S-100B predict prolonged survival for stage IV melanoma patients. *Eur. J. Cancer* 2005;41:386-392.
31. Guo HB, Stoffel-Wagner B, Bierwirth T et al. Clinical significance of serum S100 in metastatic malignant melanoma. *Eur. J. Cancer* 1995;31: 924-928.
32. Martenson ED, Hansson LO, Nilsson B et al. Serum S-100b protein as a prognostic marker in malignant cutaneous melanoma. *J. Clin. Oncol* 2001;19:824-831.
33. Mohammed MQ, Abraha HD, Sherwood RA et al. Serum S100b protein as a marker of disease activity in patients with malignant melanoma. *Med. Oncol.* 2001;18, 109-120.
34. Hamberg AP, Korse CM, Bonfrer JM, de Gast CG. Serum S100B is suitable for prediction and monitoring of response to chemoimmunotherapy in metastatic malignant melanoma. *Melanoma Res.*2003;13:45-49.
35. Hauschild AP, Engel G, Brenner JM et al. Predictive value of serum S100B for monitoring patients with metastatic melanoma during chemotherapy and/or immunotherapy. *Br. J. Dermatol.* 1999 140;1065-1071.
36. Balch CM, Gershenwald JE, Soong S et al. Multivariate Analysis of Prognostic Factors Among 2,313 Patients With Stage III Melanoma: Comparison of Nodal Micrometastases Versus Macrometastases. *J Clin Oncol.* 2010;28:2452-9.
37. Domingo-Domenech J, Castel T, Auge JM et al. Prognostic implications of protein S-100b serum levels in the clinical outcome of high-risk melanoma patients. *Tumour Biol.*2007;28: 264-272.
38. Chun YS, Wag Y, Wang DY et al. S100B levels are more accurate than LDH levels at predicting survival in melanoma patients. *Ann. Surg. Oncol.* 2008; 15: 96.
39. Mocellin S, Zavango G, Nitti D. The prognostic value of serum S100B in patients with cutaneous melanoma: a meta-analysis. *Int. J. Cancer* 2008;123: 2370-2376.
40. Tarhini AA, Stuckert J, Lee S, Sander C, Kirkwood JM. Prognostic significance of serum S100B protein in high-risk surgically resected melanoma patients participating in Intergroup Trial ECOG 1694. *J. Clin. Oncol.*2009; 27:38-44.
41. Dzwierzynski WW. Complete lymph node dissection for regional nodal metastasis. *Clin Plast Surg.* 2010 ;37:113-25.
42. Guggenheim MM, Hug U, Jung FJ, Rousson V et al. Morbidity and recurrence after completion lymph node dissection following sentinel lymph node biopsy in cutaneous malignant melanoma. *Ann Surg.* 2008;4:687-93.
43. [Http://clinicaltrials.gov/ct/show/NCT00297895.](http://clinicaltrials.gov/ct/show/NCT00297895)
44. Olilla DW. Complete metasectomy in patients with stage IV metastatic melanoma. *Lancet Oncol.* 2006;7:919-24.
45. Hsueh EC, Essner R, Foshag LJ, et al. Prolonged survival after complete resection of disseminated melanoma and active immunotherapy with a therapeutic cancer vaccine. *J Clin Oncol.* 2002;20:4549-54.
46. Morton DL, Mozzillo N, Thompson JF, et al. An international, randomized, double-blind, phase 3 study of the specific active immunotherapy agent, onmelatucel-L (Canvaxin), compared to placebo as a post-surgical adjuvant in AJCC stage IV melanoma: plenary talk. Society of Surgical Oncology Cancer Symposium, 59th Annual Meeting; March 24, 2006; San Diego, CA, USA.
47. Wasif N, Bagaria SP, Ray P, Morton DL. Does metastasectomy improve survival in patients with stage IV melanoma? a cancer registry analysis of outcomes. *J Surg Oncol.* 2011 Mar 4. [Epub ahead of print].
48. Garbe C, Hauschild A, Volkenandt M et al. Evidence-based and interdisciplinary consensus-based German guidelines: systemic medical treatment of melanoma in the adjuvant and palliative setting. *Melanoma Res.*2008;18: 152-160.
49. Hodi FS, O'Day SJ, McDermott DF et al. Improved survival with ipilimumab in patients with metastatic melanoma. *New Engl J Med* 2010; 363:711-23.
50. Culver ME, Gatesman ML, Mancl EE, Lowe DK. Ipilimumab: a novel treatment for metastatic melanoma. *Ann Pharmacother.* 2011;45:510-9.

51. Flaherty KT, Puzanov I, Kim KB et al. Inhibition of mutated, activated BRAF in metastatic melanoma. *N Engl J Med*. 2010;26:9:809-19.
52. Chapman PB, Hauschild A, Robert C et al. Improved survival with vemurafenib in melanoma with BRAF V600E mutation. *N Engl J Med* 2011;364:2507-16.
53. Garbe C, Peris K, Hauschild A et al. Diagnosis and treatment of melanoma: European consensus-based interdisciplinary guideline. *Eur. J. Cancer* 2010; 46:270-283.
54. Wong RJ, Lin DT, Schoder H et al. Diagnostic and prognostic value of [(18)F]fluorodeoxyglucose positron emission tomography for recurrent head and neck squamous cell carcinoma. *J Clin Oncol* 2002; 20:4199-208.
55. Bastiaannet E, Hoekstra OS, Oyen WJ et al. Level of fluorodeoxyglucose uptake predicts risk for recurrence in melanoma patients presenting with lymph node metastases. *Ann Surg Oncol* 2006; 13:919-26.
56. Bouwhuis MG, Suci S, Kruit W et al. Prognostic value of serial blood S-100B determinations in stage IIB-III melanoma patients: A corollary study to EORTC trial 18952. *Eur J Cancer*, 2011;47:361-8.
57. Jing Lin, Qingyuan Yang, Wilder PT et al. The calcium-binding protein S-100B Down-regulates p53 and apoptosis in malignant melanoma. *J Biol Chem* 2010; 27: 27487-98.
58. Markowitz J, Mackerell AD Jr, Carrier F, Charpentier TH, Weber DJ. Design of inhibitors for S-100B. *Curr Top Med Chem*. 2005; 5:1093-108.
59. Jubb AM, Harris AL. Biomarkers to predict the clinical efficacy of bevacizumab in cancer. *Lancet Oncol*. 2010;11:1172-83.
60. Christensen CM, Grossman JH, Hwang J. *The Innovator's Prescription: A Disruptive Solution For Health Care*. McGraw-Hill, OH, USA, 2008.
61. Porter ME, Teisberg EO. How physicians can change the future of health care. *JAMA* 297, 1103–1111 2007;297:1103-1111
62. Murchie P, Nicolson MC, Hannaford PC et al. Patient satisfaction with GP-led melanoma follow-up: a randomised controlled trial. *Br. J. Cancer* 2010;10:1447–1455.
63. Francken AB, Hoekstra-Weebers JW, Hoekstra HJ. Is GP-led follow-up feasible? *Br J Cancer*. 2010 May 11;102(10):1445-6.
64. Soong SJ, Ding S, Coit D et al. Predicting survival outcome of localized melanoma: an electronic prediction tool based on the AJCC melanoma database. *Ann. Surg. oncol*. 2010;17: 2006-2014.
65. CancerMath.net, Melanoma Outcomes Calculator. <http://cancer.lifemath.net/melanoma/outcome>
66. National Cancer Institute. Cancer trends progress report - 2009/2010 update. ([http://progressreport.cancer.gov/doc\\_detail.asp?pid=1&did=2009&chid=95&coid=926&mid=](http://progressreport.cancer.gov/doc_detail.asp?pid=1&did=2009&chid=95&coid=926&mid=)).
67. Mariotto AB, Yabroff KR, Shao Y et al. Projections of the cost of cancer care in the United States: 2010-2020. *J Natl Cancer Inst* 2011;103:117-128.
68. Smith TJ, Hillner BE. Bending the cost curve in cancer care. *N Engl J Med*. 2011;26;21:2060-5. 69. Matsuyama R, Reddy SN, Smith TJ. Why do patients choose chemotherapy near the end of life? *J Clin Oncol* 2006;24:3490-6.
70. National Cancer Institute, The Cancer Genome Atlas <http://cancergenome.nih.gov>.

# 10

---



**FIGURE 1** : Structure of the protein S-100B

## Summary


---

Melanoma causes more than 75% of all deaths related to skin cancer and melanoma incidence rates have increased rapidly worldwide and are expected to keep on rising in the future. The absolute total number of new cases in the Netherlands is estimated to be well over 4,800 in 2015.<sup>1</sup> Despite being diagnosed in an earlier phase of disease as a result of increased awareness and better surveillance, melanoma patients presenting with palpable nodal metastases today still have a poor 5-year survival (59% and 40% for stage IIIB and IIIC respectively) but a tremendous heterogeneity in 5-year survival rates is observed (23%-87% 5-year survival).<sup>2,3</sup> Biomarkers measured preoperatively in stage III melanoma patients could both be highly specific indicators of early recurrence after surgical treatment and could optimize and refine the staging of melanoma patients with palpable nodes.<sup>4</sup>

In this thesis the main focus is on stage III melanoma and how to identify stage III melanoma patients who may benefit from adjuvant systematic treatment with the aid of biomarkers as S-100B (figure 1) and FDG- SUV in order to improve, in the long run, survival.

### **PART I : DETECTION AND TREATMENT OF NODAL MELANOMA METASTASES**

In chapter two the role of the method of detection in nodal disease in the prognosis of melanoma patients who underwent therapeutic lymph node dissection (TLND) was evaluated. International consensus on the follow-up of melanoma patients has not been achieved and high-frequency follow-up is still practiced in many countries around the world. It was found that the method of detection of lymph node recurrence in melanoma patients by patient or physician has no impact on survival. 45% of lymph node metastases in Stage I-II melanoma patients are physician detected. Younger patients detect their own lymph node metastases significantly more often than elderly patients. However, neither the method of detection nor




**FIGURE 2:** Bed rest and flexion of the hip and knee in flexion using a Bohler splint

age correlates with DSS. More frequent follow-up would not alter DFS and DSS significantly. The data of this study will add to the controversy about the value of high frequency follow-up regimens in melanoma. A prospective, randomized, high quality methodological research has been started in order to develop meaningful applicable guidelines (MELFO). In the third chapter we conducted an extensive retrospective study to assess the present morbidity of therapeutic or completion lymph node dissection to determine risk factors for short term morbidity and to evaluate recommendations on peri-operative treatment. Therapeutic groin dissection for melanoma remains a surgical procedure with a high morbidity. Presently, more complications occur after therapeutic ilio-inguinal lymph node dissection than 20 years ago. Analysing our data it was found that a high BMI was significantly correlated with the occurrence of wound infections. Bed rest and flexion of the hip and knee in flexion using a Bohler splint (figure 2) improved wound healing after therapeutic ilio-inguinal lymph node dissection.


## **PART II : PROGNOSTIC MARKERS IN MELANOMA**

When compared with other markers and tumor characteristics as ulceration, Clark level and Mitosis index, Breslow thickness is the most important biomarker and predictor for mortality. In chapter four, patients diagnosed with invasive melanoma between 1994 and 2008 were selected from the Netherlands Cancer Registry (NCR). Incidence (per 100,000) over time was calculated for young (<65 years) and elderly patients ( $\geq 65$  years). Distribution of Breslow thickness for young and elderly males and females was assessed. Regression analysis of the log-transformed Breslow thickness was used to assess changes over time. Relative survival was calculated as the ratio of observed survival to expected survival.

We found that melanoma incidence increased more rapidly among the elderly (5.4% Estimated Annual Percent Change (EAPC),  $p < 0.0001$ ) than among younger patients (3.9% EAPC,  $p < 0.0001$ ) (figure 3) and the overall Breslow thickness declined significantly over time ( $p < 0.001$ ). However, the decline in Breslow thickness is less prominent among elderly patients than among young patients with a relative survival of elderly patients that was worse compared with that of younger patients ( $p < 0.001$ ). Campaigns in the Netherlands must focus more on early melanoma detection in the elderly.



**FIGURE 3** Incidence of invasive melanoma among young and elderly patients



**FIGURE 4** : Disease specific survival (DSS) correlated to S-100B in AJCC stadium IIIB.

Up to today the role of S-100B for patients with palpable lymph node metastases before undergoing therapeutic lymph node dissection was still unknown. In chapter five the role of S-100B for disease-free survival (DFS) was evaluated and compared with the tumor marker



LDH. Preoperative S-100B elevation in patients with optimally staged clinical stage III melanoma was correlated with shortened disease-free survival (figure 4).

The new serum tumour marker S-100B has the potential to identify stage III melanoma patients who may benefit from adjuvant systematic treatment. In the stratification of new adjuvant therapeutic trials in patients with loco regional recurrence stage III melanoma, we would recommend the incorporation of S-100B in the stratification instead of the more commonly used LDH tumour marker. Pre-operative S-100B levels provide information for stage III melanoma patients who want to be optimal informed with respect to their disease prognosis.

In chapter six S-100B values and Standardized Uptake Value (SUV) in FDG-PET for clinically stage III melanoma patients were analysed as indicators of early recurrence in stage III disease. S-100B and SUV in stage III melanoma seemed to have no correlation and both markers have different associations with various histo-pathological factors. However, as concluded in chapter five, S-100B, in contrast with SUV, is associated with nodal tumor load, and when elevated, predicts a shorter DFS.

In chapter seven to investigate the feasibility of using bevacizumab to improve the survival of AJCC stage III melanoma patients, we analyzed how a single bevacizumab treatment affected nodal disease and a panel of biomarkers in clinically FDG-PET/CT staged, stage III melanoma patients, prior to therapeutic lymph node dissection (TLND). Four weeks before TLND, nine patients with palpable lymph node metastases received 7.5 mg/kg bevacizumab. Before and after this treatment, all patients were assessed by measurements of the maximum standardized uptake value (SUVmax) by FDG-PET scan, and serum S-100B and LDH. After TLND, the dissection specimen was analyzed for the number of removed lymph nodes, the amount of metastatic lymph nodes, and tumor necrosis. Tumor necrosis in dissection specimens was associated with declining S-100B levels, while elevated S-100B was only found in cases with no necrosis. Bevacizumab might be useful in treating AJCC stage III melanoma patients prior to TLND, and S-100B appears to be a useful marker for assessment of treatment effects. Future research, preferably in a randomized controlled trial using serum S-100B for response monitoring, should determine the effect of (neoadjuvant) bevacizumab, and adjuvant Ipilimumab and/or BRAF pathway inhibitors as a new combined treatment strategy of systemic treatment with TLND to improve disease free and overall survival in clinically stage III melanoma patients.

The current status of S-100B in treatment of melanoma is described in chapter eight. A review was composed describing the role of S-100B for the different tumor stages in melanoma and how the protein could be used by a practicing surgical oncologist. S-100B was already known as a marker in stage IV palliative melanoma treatment, however in the future it could be used for high risk stage III melanoma to perform better staging. S-100B can be used to perform a individualized risk assessment before administering adjuvant treatment on the one hand and on the other hand to measure response to induction therapy for nodal disease.<sup>4,5</sup> Despite these described prognostic values, S-100B is not suitable for routine screening purposes in stage I–III melanoma patients or for the early detection of recurrence in the follow-up of melanoma patients; that is, early detection of recurrences in follow-up do

not yet translate into an adequate therapeutic intervention nor survival benefit due to limited treatment options in advanced melanoma. However, maybe, since an effective (adjuvant) therapy for loco-regional metastatic and disseminated melanoma seems to be discovered, the routine use of biomarkers as S-100B seems about to alter dramatically in the future.<sup>6-9</sup>


**REFERENCES**

1. De Vries E, van de Poll-Franse LV, Louwman WJ et al. Predictions of skin cancer incidence in the Netherlands up to 2015. *Br J Dermatol* 2005; 152:481-488.
2. Balch CM, Greshenwald JE, Soong SJ et al. Final version of 2009 AJCC melanoma staging and classification. *J Clin Oncol* 2009; 27:6199-206.
3. Balch CM, Gershenwald JE, Soong S et al. Multivariate Analysis of Prognostic Factors Among 2,313 Patients With Stage III Melanoma: Comparison of Nodal Micrometastases Versus Macrometastases. *J Clin Oncol*. 2010;28:2452-9.
4. Kruijff S, Bastiaannet E, Muller-Kobold AC et al. S-100B concentrations predict disease-free survival in stage III melanoma patients. *Ann Surg Oncol* 2009;16:3455-62.
5. Kruijff S, Bastiaannet E, Brouwers AH, et al. The use of S-100B to evaluate therapy effects during bevacizumab induction treatment in AJCC stage III melanoma. *Ann Surg Oncol*. 2011; accepted for publication.
6. Hodi FS, O'Day SJ, McDermott DF, et al. Improved survival with ipilimumab in patients with metastatic melanoma. *New Engl J Med*. 2010;363:711-23.
7. Flaherty KT, Puzanov I, Kim KB, et al. Inhibition of mutated, activated BRAF in metastatic melanoma. *N Engl J Med*. 2010;363:809-19.
8. Chapman PB, Hauschild A, Robert C, et al. Improved survival with vemurafenib in melanoma with BRAF V600E mutation. *N Engl J Med*. 2011;364:2507-16.
9. Garbe C, Peris K, Hauschild A, et al. Diagnosis and treatment of melanoma: European consensus-based interdisciplinary guideline. *Eur. J. Cancer* 2010; 46:270-283.



# 10

---



**FIGUUR 1** : Structuur van het eiwit S-100B

## Samenvatting

---


Het melanoom veroorzaakt meer dan 75% van alle huid kanker gerelateerde mortaliteit en de incidentie van deze maligne huidaandoening heeft wereldwijd een grote vlucht genomen. De verwachting is dat die opwaartse trend zich zal voortzetten in de toekomst en dat het absolute aantal nieuwe patiënten dat zich in 2015 zal presenteren met een melanoom in Nederland rond de 4800 is.<sup>1</sup> Ondanks dat melanomen eerder vastgesteld lijken te worden met een toename aan dunnere melanomen, mogelijk als een gevolg van een toegenomen patiënt bewustzijn en toezicht, hebben patiënten die zich presenteren met palpabele klieren (AJCC stadium III) nog altijd een zeer matige prognose. Opvallend is met name de heterogeniteit in 5 jaar overleving uiteenlopend van 23% tot 87%.<sup>2,3</sup> Voor deze patiënten zou daarom een strategie gehanteerd kunnen worden waarbij in eerste instantie biomarkers bepaald worden. Enerzijds met het doel in een vroege fase een recidief te kunnen vaststellen maar met name om het stadierings proces te verfijnen voor deze heterogene groep.<sup>4</sup> Dit proefschrift richt de aandacht op de melanoom patiënt met lymfogene metastasen en op de mogelijkheid om met behulp van biomarkers zoals S-100B (figuur 1) die specifieke hoog risico patiënt te identificeren die in de toekomst overlevingsvoordeel zou kunnen hebben van systemische therapie.

### **DEEL I : HET VASTSTELLEN EN BEHANDELEN VAN LYMFKLIER METASTASEN BIJ HET MELANOOM**

In hoofdstuk 2 werd de prognostische invloed van de wijze van vaststellen van lymfklier metastasen (door de dokter of door de patiënt zelf) bij patiënten die later een lymfklier dissectie ondergingen, geëvalueerd. Er bestaat nog steeds geen algemene internationale consensus over de follow-up van melanoom patiënten en in verscheidene landen over de gehele wereld wordt nog altijd een hoog frequente follow-up strategie toegepast. Vastgesteld werd dat de wijze van vaststellen van palpabele klieren dan wel door een dokter tijdens de follow-up dan wel door een

patiënt zelf geen invloed had op de overleving van deze patiënten groep. 45 % van de lymfklier metastasen in stadium I en II melanoom patiënten worden door een dokter vastgesteld tijdens follow-up. Jonge patiënten ontdekken hun lymfklier metastasen significant sneller dan oudere patiënten. Echter, noch de wijze van vaststellen noch de leeftijd van de patiënt lijken gecorreleerd met ziekte specifieke overleving. Het intensiveren van de follow-up frequentie zou de ziekte specifieke- of ziekte vrije-overleving niet beïnvloeden. De data van deze studie dragen bij aan de controverse die bestaat over de waarde van het uitvoeren van hoog frequente follow-up bij het melanoom. Een prospectieve gerandomiseerde methodologische hoog kwalitatieve studie is derhalve reeds gestart met als doel uiteindelijke waardevolle goed toepasbare follow-up richtlijnen te creëren (MELFO).

In het derde hoofdstuk voerden we een zeer uitgebreide retrospectieve studie uit om de huidige morbiditeit van therapeutische ilio-inguinale liesklier dissecties verricht in ons



**FIGUUR 2:** Flecteren van de heup door middel van een Bohler splint

centrum te evalueren. Het doel was de risicofactoren voor korte termijn morbiditeit in kaart te brengen en daarmee aanbevelingen te doen voor het peri-operatieve beleid. Een therapeutische liesklierdissectie is een chirurgische procedure met een hoge morbiditeit. Vandaag de dag lijken meer complicaties op te treden na een therapeutische ilio-inguinale lies klierdissectie dan 20 jaar geleden. Uitgebreide analyse van de data uit dit centrum leverde onder meer de conclusie op dat bij patiënten met een hoog body mass index (BMI) een significante correlatie bestond met het optreden van wondinfecties. Daarnaast ging bij die patiënten bij wie postoperatieve bedrust en het flecteren van de heup door middel van het gebruik van een Bohler splint was gehandhaafd (figuur 2) de wond genezing gepaard met minder problemen.


## DEEL II : PROGNOSTISCHE MARKERS VOOR HET MELANOOM

Wanneer de Breslow dikte vergeleken wordt met andere markers en tumor kenmerken als ulceratie, Clark niveau en Mitose index , blijft de Breslow dikte van het melanoom de meest belangrijke biomarker en voorspeller voor mortaliteit. In het vierde hoofdstuk werden alle patiënten bij wie tussen 1994 en 2008 de diagnose melanoom gesteld werd, geselecteerd vanuit de Nederlandse integrale kanker centra (IKC). De incidentie (per 100 000) over de tijd werd

uitgerekend voor de jongere (<65 jaar) en oudere patiënten (≥65 jaar). De verdeling van Breslow dikte van jongere en oudere mannen en vrouwen werd in kaart gebracht. Er werd een regressie analyse gebruikt om de veranderingen van de Breslow dikte over de tijd in beeld te brengen. De relatieve overleving brachten we in kaart als de ratio van de geobserveerde overleving en de verwachte overleving. De resultaten lieten zien dat de incidentie van het melanoom veel sneller steeg onder ouderen (5.4% Estimated Annual Percent Change (EAPC),  $p < 0.0001$ ) dan onder jongere patiënten (3.9% EAPC,  $p < 0.0001$ ) (figuur 3). Weliswaar daalde als verwacht de Breslow dikte over de tijd ( $p < 0.001$ ) maar deze daling was opvallend minder onder ouderen dan onder jongere melanoom patiënten met een relatieve overleving die eveneens slechter was voor de oudere patiënt ( $p < 0.001$ ). Het lijkt daarom toenemend van belang in Nederland meer aandacht te gaan vestigen op het vroeger vaststellen van het melanoom bij oudere patiënten.


In het vijfde hoofdstuk werd de rol van het eiwit S-100B en ziekte vrije overleving geëvalueerd en vergeleken met de klassiek gebruikte tumor marker LDH die nu nog deel uitmaakt van het AJCC stadierings systeem. Tot op heden was de rol van S-100B voor patiënten met lymfogeen gemetastaseerde ziekte, voordat zij een klierdissectie ondergingen, nog volledig onbekend. Uit onze resultaten bleek dat een preoperatief verhoogd S-100B bij AJCC bij stadium III patiënten die eerder optimaal geanalyseerd waren door middel van FDG PET en CT scan, significant gecorreleerd was met een verkorte ziekte vrije overleving (figuur 4).

De nieuwe tumor marker S-100B heeft de potentie om hoog risico stadium III patiënten te identificeren die zouden kunnen profiteren van adjuvante therapie. Bij het selecteren van patiënten met een loco-regionaal lymfogeen recidief voor nieuwe therapeutische trials, zouden we het toevoegen van S-100B aan het AJCC stadierings systeem sterk willen aanbevelen in plaats van de klassieke tumor marker LDH. In stadium III melanoom geeft het preoperatief



**FIGUUR 3** De incidentie van het melanoom onder jonge en oudere patiënten





**FIGUUR 4 :** Ziekte specifieke overleving (DSS) gecorreleerd met S-100B in AJCC stadium IIIB.

bepalen van serum S-100B informatie aan melanoom patiënten die optimaal geïnformeerd willen worden over hun prognose.

In het zesde hoofdstuk werd zowel serum S-100B als de marker Standardized Uptake Value (SUV) geanalyseerd bij de klinisch stadium III melanoom patiënt. S-100B en SUV leken geen correlatie te hebben met elkaar en beide markers bleken beiden tevens verschillende relaties met histo- pathologische factoren te hebben. Duidelijk was, zoals ook al eerder geconcludeerd in hoofdstuk vijf, dat S-100B evident gecorreleerd was met de lymfogene tumor massa bij patiënten en wanneer verhoogd een goede voorspeller voor een verkorte ziekte vrije overleving. De Standardized Uptake Value (SUV) bleek geen relatie te hebben met ziekte vrije overleving.

In hoofdstuk zeven werd een pilot studie verricht om de mogelijkheid te onderzoeken voor het gebruik van bevacizumab bij stadium III patiënten om de algehele overleving te verbeteren.

We analyseerden hoe een eenmalige toediening van bevacizumab de behandeling van lymfogene ziekte en het verloop van de biomarkers S-100B en LDH kon beïnvloeden voorafgaand aan een therapeutische lymfklierdissectie. Vier weken voor de klierdissectie kregen negen stadium III patiënten 7.5 mg/kg bevacizumab toegediend waarbij hiervoor en erna bij alle negen patiënten de maximale standardized uptake value (SUVmax) en S-100B en LDH werd afgenomen. Na de dissectie werd het klier preparaat pathologisch geanalyseerd op het aantal verwijderde klieren, de hoeveelheid metastatische lymfklieren en de hoeveelheid tumornecrose. De hoeveelheid tumor necrose in het dissectie preparaat bleek gecorreleerd aan een dalend S-100B daar waar een doorstijgend S-100B alleen werd gezien wanneer de tumornecrose ontbrak. Concluderend zou bevacizumab dus van waarde kunnen zijn bij de behandeling van AJCC stadium III melanoom patiënten voorafgaand aan een klierdissectie, en daarnaast zou

S-100B een waardevolle marker kunnen zijn voor het evalueren van het behandelresultaat. Toekomstig onderzoek, bij voorkeur in de vorm van een gerandomiseerde trial waarbij serum S-100B toegepast wordt om behandelrespons te meten, zal moeten gaan uitmaken of het effect van (neoadjuvante) bevacizumab, eventueel in samenwerking met de veelbelovende adjuvante middelen als Ipilimumab en/of BRAF remmers als nieuwe gecombineerde strategie, de ziekte vrije en algemene overleving zal verbeteren in stadium III melanoom patiënten.

In hoofdstuk 8 werd de huidige status van het gebruik van het eiwit S-100B beschreven bij de behandeling van het melanoom in een overzichtartikel. Hier wordt de rol van het S-100B eiwit beschreven volgens de verschillende tumor stadia van het AJCC classificatie model en hoe het gebruikt zou kunnen bij de behandeling van het melanoom nu en in de toekomst.

S-100B was reeds bekend als een marker bij de palliatieve behandeling van stadium IV melanoom maar lijkt dus in de toekomst ook gebruikt te kunnen gaan worden bij de behandeling van hoog risico stadium III melanoom om beter te kunnen stadieren.

Daarnaast kan het eiwit gebruikt worden om een betere individuele risico analyse te maken voordat adjuvante behandeling toegediend wordt alsmede voor het meten van response op een inductie behandeling voorafgaand aan een klierdissectie bij lymfogene metastasen.<sup>4,5</sup>

Ondanks deze prognostische kwaliteiten, lijkt S-100B niet geschikt voor routine screenings doeleinden in stadium I en II patiënten of voor het gebruik bij vroeg opsporing van recidieven in de follow up; dat wil zeggen, vroege opsporing van recidieven tijdens de follow-up vertaalt zich nog niet in overlevingsvoordeel doordat er nog te weinig behandelopties zijn voor het gemetastaseerd melanoom. Echter, misschien, doordat er een serie aan veel belovende effectieve adjuvante middelen voor loco-regionaal en gegeneraliseerd gemetastaseerd melanoom op de markt zijn gekomen, zou het gebruik van biomarkers als S-100B in de nabije toekomst drastisch kunnen gaan veranderen.<sup>6-9</sup>

**REFERENCES**

1. de Vries E, van de Poll-Franse LV, Louwman WJ et al. Predictions of skin cancer incidence in the Netherlands up to 2015. *Br J Dermatol* 2005; 152:481-488.
2. Balch CM, Greshenwald JE, Soong SJ et al. Final version of 2009 AJCC melanoma staging and classification. *J Clin Oncol* 2009; 27:6199-206.
3. Balch CM, Gershenwald JE, Soong S et al. Multivariate Analysis of Prognostic Factors Among 2,313 Patients With Stage III Melanoma: Comparison of Nodal Micrometastases Versus Macrometastases. *J Clin Oncol*. 2010;28:2452-9.
4. Kruijff S, Bastiaannet E, Muller-Kobold AC et al. S-100B concentrations predict disease-free survival in stage III melanoma patients. *Ann Surg Oncol* 2009;16:3455-62.
5. Kruijff S, Bastiaannet E, Brouwers AH, et al. The use of S-100B to evaluate therapy effects during bevacizumab induction treatment in AJCC stage III melanoma. *Ann Surg Oncol*. 2011; accepted for publication.
6. Hodi FS, O'Day SJ, McDermott DF, et al. Improved survival with ipilimumab in patients with metastatic melanoma. *New Engl J Med*. 2010;363:711-23.
7. Flaherty KT, Puzanov I, Kim KB, et al. Inhibition of mutated, activated BRAF in metastatic melanoma. *N Engl J Med*. 2010;363:809-19.
8. Chapman PB, Hauschild A, Robert C, et al. Improved survival with vemurafenib in melanoma with BRAF V600E mutation. *N Engl J Med*. 2011;364:2507-16.
9. Garbe C, Peris K, Hauschild A, et al. Diagnosis and treatment of melanoma: European consensus-based interdisciplinary guideline. *Eur. J. Cancer* 2010; 46:270-283.

## List of contributing authors

---

Prof. dr. H.J. Hoekstra, Division of Surgical Oncology, University Medical Centre Groningen, University of Groningen.

Dr. R.J. van Ginkel, Division of Surgical Oncology, University Medical Centre Groningen, University of Groningen.

Dr. A.B. Francken, Surgical trainee, NKI/AvL, Amsterdam.

Ir. E. Bastiaannet, Leiden University Medical Center, Leiden.

Drs H.P.A.M. Poos, Surgical trainee, Medisch Spectrum Twente, Enschede.

Drs M.J. Speijers, Surgical trainee, Isala Klinieken, Zwolle.

Dr. A.C. Muller Kobold, Nuclear Medicine and Molecular Imaging, University Medical Centre Groningen, University of Groningen.

Prof. dr. G.A.P. Hospers, Medical Oncology, University Medical Centre Groningen, University of Groningen.

Dr. W.B. Nagengast, resident gastro-enterology, Medisch Spectrum Twente, Enschede.

Dr. M. van der Aa, Comprehensive Cancer Centre the Netherlands, Utrecht.

Dr. M. Schaaapveld, NKI/AvL, Amsterdam, the Netherlands.



# DANK!

Een proefschrift is een proeve van bekwaamheid in de wetenschap. Mijn proeve van bekwaamheid beschouw ik als een zeer leerzame periode waarin mij bijgebracht is hoe een onderzoek op te zetten, te analyseren, op te schrijven en vervolgens te publiceren. Ik zie dit proefschrift als een begin; een begin van het bedrijven van wetenschap met een zekere kwaliteit.

De totstandkoming van dit werk heb ik beschouwd als een opleiding. Het was voor mij een voorrecht deze opleiding te kunnen combineren met mijn klinisch chirurgische opleiding. Het was namelijk een project van mijzelf, waar alleen ik eindverantwoordelijk voor was. Ik heb het als prettig ervaren om naast de dagelijkse patiëntenzorg aan dit project te werken. Want patiëntenzorg is prachtig werk - maar nooit af. Een artikel is op een dag wel af. En het voelt goed voor een mens om iets af te ronden.

Het doorlopen van deze opleiding heb ik gedaan volgens een piramidemodel. Gek genoeg vormt de promotie zelf pas de top van deze piramide.

Het fundament van de piramide werd thuis gevormd door Annelies, Otto, Alexa en Eelco, zonder wier liefde en steun dit project niet had kunnen worden voltooid. Het middelste deel van de piramide werd gevormd door mijn chirurgische opleider Dr. W.J. Mastboom en de gehele opleidingsgroep, zonder wier flexibiliteit en steun ik nooit twee opleidingen tegelijk had kunnen doen. Deze belangrijke fundamenten brachten mij drie jaar lang aan de top van de piramide: eenmaal daar aangekomen was de wetenschappelijke opleiding zonder Prof. dr. H. J. Hoekstra en Ir. E. Bastiaannet onmogelijk geweest. Zij inspireerden en lanceerden mij als beginnend onderzoeker en boden vertrouwen. De laagdrempelige begeleiding en de vruchtbare samenwerking resulteerden in deze promotie. Het was eervol deel uit te maken van dit driemanschap.

Enschede , 29 augustus 2011  
Schelto Kruijff

## Mijn

## d a n k

gaat eveneens

uit naar de leden

van de promotiecommissie:

Prof. dr. G.A.P. Hospers, Prof. dr.

E. Heineman en Prof. dr. I.H.M. Borel

Rinkes; naar de paranimfen Drs H. Frima en Dr.

P.R. de Reuver alsmede de ceremoniemeester Dr. J.E.

Jansen en naar de mede-auteurs: Drs H.P.A.M. Poos, Dr.

A.C. Muller Kobold, Dr. R.J. van Ginkel, Prof. dr. G.A.P. Hospers,

Dr. W.B. Nagengast, Drs M.J. Speijers, Dr. A.B. Francken, Dr. M. van

der Aa, Dr. M. Schaapveld, Prof. dr. I.P. Kema, Prof. dr. A.J.H. Suurmeijer,

Dr. A.H. Brouwers en naar alle mensen met goed advies: Dr. M. de Vries, Dr. I.

Grossmann, Dr. M.F. Lutke Holzik, Dr. D. Daryanani en natuurlijk naar de ondersteuners:

Melchior Spies, Tom Letteboer, Sjoerd Kruijff en tenslotte naar de mensen die er altijd zijn:

Denia Kruijff-Stoffelsma, Hajo Kruijff, Letta van Walsem-van der Griendt en Arnoud van Walsem.

# Curriculum vitae

Schelto Kruijff werd geboren op 18 mei 1977 in Rijpwetering, als laatste uit een gezin van drie kinderen. Vanaf zijn zevende jaar ging hij in het naburige Leiderdorp (heen en weer 14 kilometer fietsen) naar zijn lagere school. Zijn eindexamen VWO behaalde hij in 1996 hij aan het Stedelijk Gymnasium in Leiden. Daarna volgde een wereldreis van een jaar.

In 1997 startte Schelto in Groningen met de studie Biologie, maar toen bleek dat hij het ontleden van haaien interessanter vond dan het determineren van planten, stapte hij over naar de Geneeskunde.

Tijdens zijn studietijd in Groningen was Schelto actief op de studentenvereniging aan de Grote Markt 27, wat leidde tot een jaar onderbreking om deel uit te maken van het bestuur van die vereniging. Ook de studiegebonden activiteiten, zoals het leverteam en de stichting Johannes Borgesius, werden met groot plezier uitgevoerd.

Tijdens de co-schappen in Deventer, waar Schelto twee jaar in een boerderij in de bossen achter Diepenveen woonde, raakte hij voorgoed verslingerd aan de Heelkunde. Na het beëindigen van zijn studie werkte Schelto vier maanden in het Nkhoma mission Hospital in Malawi samen met Herman Firma (tropenarts en chirurg). Tijdens deze indrukwekkende periode ontwikkelde Schelto een visie op zijn toekomst als dokter. Vanuit Malawi stuurde hij zijn sollicitatieformulieren voor de Heelkunde op.

Via een Agnio-schap Heelkunde in het MST in Enschede werd Schelto aangenomen voor de opleiding Heelkunde per mei 2006 in het UMCG onder leiding van Prof. dr. H.J. ten Duis. Aldaar begon hij aan zijn oncologisch onderzoek over het eiwit S-100B en de melanoompatiënt met lymfogeen gemetastaseerd melanoom onder begeleiding van Prof. dr. H.J. Hoekstra. Nauwelijks met het onderzoek aangevangen, vertrok hij in 2008 naar het MST. Daar vervolgde hij zijn opleiding onder leiding van Dr. W.J. Mastboom en legde hij zich toenemend toe op zijn promotie.

In Enschede raakte Schelto klinisch steeds meer geïnteresseerd in de chirurgische oncologie en kreeg hij de kans zich hierin verder te ontwikkelen. Daarnaast was Schelto tijdens zijn opleiding bestuurlijk actief in de Jonge orde en het bestuur van de Vagh. Schaatsen is een passie en hij verreed de alternatieve Elfstedentocht op de Weissensee dan ook drie keer.

Schelto zal zijn opleiding voltooien in juli 2012. Daarna zal hij met zijn gezin vertrekken naar Sydney vanwege een fellowship in de endocriene chirurgie.

Schelto is getrouwd met Annelies van Walsem, die werkzaam is in het onderwijs. Zij hebben drie kinderen: Otto, Alexa en Eelco.

## List of publications

---

- 2011      The use of S-100B to evaluate therapy effects during bevacizumab induction treatment in AJCC stage III melanoma. S. Kruijff, E Bastiaannet, A.H. Brouwers, et al. *Ann Surg Oncol*, aug 2011, accepted for publication.
- 2011      VEGF-SPECT with 111In-bevacizumab in stage III/IV melanoma patients  
Nagengast WB, Kruijff S, Hoekstra HJ, Hospers G.A.P  
*Eur J Cancer*. 2011;47:1595-602.
- 2010      Comparison of S-100B and SUV and their relation with survival in stage III melanoma  
Kruijff S, Spijers M, Muller Kobold A, Bastiaannet E, HJ Hoekstra  
*Eur J Surg Oncol*. 2011;37:225-32.
- 2010      Optimal Follow-up for Melanoma  
Speijers MJ, Francken AB, Hoekstra-Weebers JEHM, Bastiaannet E, Kruijff S, Hoekstra HJ. *Expert Rev. of Derm.*2010;5:461.
- 2010      De effectiviteit van beugelbehandeling versus chirurgie bij de ingegroeide teennagel  
Kruijff S, van det RJ, van der Palen J, Gerritsma AC, Geelkerken RH, NtvH, june 2010.
- 2010      Detection of Melanoma Nodal Metastases; Differences in Detection Between Elderly and Younger Patients Do Not Affect Survival.  
Kruijff S, Bastiaannet E, Suurmeijer AJ, Hoekstra HJ.  
*Ann Surg Oncol*. 2010;17:3008-14.



- 2009 S-100B concentrations predict disease-free survival in stage III melanoma patients.  
Kruijff S, Bastiaannet E, Kobold AC, van Ginkel RJ, Suurmeijer AJ, Hoekstra HJ.  
Ann Surg Oncol. 2009;16:3455-62.
- 2008 Therapeutic groin dissection for melanoma: risk factors for short term morbidity.  
Poos HP, Kruijff S, Bastiaannet E, van Ginkel RJ, Hoekstra HJ.  
Eur J Surg Oncol. 2009;35:877-83.
- 2008 Partial matrix excision or orthonyxia for ingrowing toenails.  
Kruijff S, van Det RJ, van der Meer GT, van den Berg IC, van der Palen J, Geelkerken  
RH. Am Coll Surg. 2008;206:148-53.