

University of Groningen

Glutamate transport in thermophilic and mesophilic bacteria

Tolner, Berend

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

1995

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Tolner, B. (1995). *Glutamate transport in thermophilic and mesophilic bacteria*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Summary

So far, it has been difficult to obtain 3-dimensional crystals of integral membrane proteins which are suitable for structural analysis at high resolution. Membrane (associated) proteins from thermophilic organisms may offer the advantage that solubilized proteins are more stable than proteins from mesophilic organisms, which may be very helpful in crystallization attempts.

In *chapter 1* an overview is given of the special properties of membranes and enzymes from thermophilic microorganisms relative to those of mesophiles. Thermophiles are adapted to life at higher temperature by having special membrane lipids and a higher thermo-stability and thermo-activity of the proteins. Other adaptations that have been described include high turnover rates of the energy transducing enzymes and the use of Na^+ rather than H^+ as coupling ion in energy transduction.

The aim of the present project was to characterize bacterial glutamate transporters from related bacterial species which differ with respect to the optimal temperature of growth (T_{opt}). For this purpose, *Bacillus caldotenax* (T_{opt} of 70°C), *Bacillus stearothermophilus* (T_{opt} of 63°C), *Escherichia coli* (T_{opt} of 37°C) and *Bacillus subtilis* (T_{opt} of 37°C) were selected.

The cloning of the genes encoding the $\text{Na}^+/\text{H}^+/\text{L}$ -glutamate symport proteins, and characterization of the respective gene products, of *B. stearothermophilus* (gltT_{BS}), *B. caldotenax* (gltT_{BC}), and the H^+/L -glutamate symport protein of *B. subtilis* (gltP_{BSU}) is described in *chapters 2* and *4*, respectively. The genes were cloned by complementation in *E. coli* JC5412 by selecting for growth on glutamate as sole source of carbon, energy and nitrogen. The gltT_{BS} , gltT_{BC} , and gltP_{BSU} genes encode homologous proteins that are also homologous to transporters for structurally related compounds such as aspartate, fumarate, malate and succinate. Both, sodium- and proton-coupled transporters, belong to this family of dicarboxylate transporter proteins. Hydropathy profiling and multiple alignment of the family of carboxylate transporters suggest that each of the proteins spans the cytoplasmic membrane 12-times with both the amino-acid and carboxy-terminus on the inside (*Chapter 2* and *4*). Comparison of the primary structure of GltT_{BS} and GltT_{BC} , with that of the $\text{H}^+/\text{glutamate}$ transport protein of *E. coli* K-12 (GltP_{EC}) revealed regions of high similarity among regions which differed completely. However, by translating the nucleotide sequence of gltP_{EC} in different reading frames and by comparing the translated sequences with those of GltT_{BS} and GltT_{BC} , it

became apparent
by sequencing
which is de
encodes a
hydropathy
suggested m
The substr
chapter 4.
electrogeni
symported w
studies in
generated b
by artifici
had no (st
glutamate up
L-aspartate
acid, but n
In *chapter*
 $\text{H}^+/\text{glutamate}$
(GltP_{EC} and
symport pr
caldotenax
proteins was
which the p
expressed.
strain (glt
Transport of
electrogenic
cation-symp
translocate
cilitates th
by GltP_{EC} , C
and cysteic
glutamate. C
ions. When t
are express
be electroge
is surprisi
stearotherm
glutamate s
dependency
with temper
ion-specific
transport p
Bacillus me

al crystals
of structural
proteins from
solubilized
organisms,
s.
properties of
ms relative
to life at
and a higher
eins. Other
rnover rates
rather than

the bacterial
pecies which
owth (T_{opt}).
, *Bacillus*
 T_{opt} of 37°C)

ate symport
e products,
 glT_{Bc}), and
 glT_{Bsu}) is
genes were
ecting for
energy and
homologous
rters for
fumarate,
on-coupled
transporter
ent of the
ach of the
h both the
r 2 and 4).
 glT_{Bc} , with
coli K-12
ions which
nucleotide
comparing
 glT_{Bc} , it

became apparent that the divergence of the sequence was caused by sequencing errors. This was confirmed by resequencing *glTP_{EC}* which is described in chapter 3. The correct open reading frame encodes a protein of 437 (instead of 395) amino acids, and hydropathy analysis predicts 12 rather than 4 previously suggested membrane spanning α -helical regions.

The substrate and cation-selectivity of $GltP_{Bsu}$ is described in chapter 4. Transport of acidic amino acids in *B. subtilis* is an electrogenic process in which L-glutamate or L-aspartate is symported with at least two protons. This is shown by transport studies in membrane vesicles in which a proton motive force is generated by oxidation of ascorbate / phenazine methosulfate or by artificial ion-gradients. An inwards-directed sodium gradient had no (stimulatory) effect on proton motive force driven L-glutamate uptake. The transporter is specific for L-glutamate and L-aspartate, and is inhibited by β -hydroxyaspartate and cysteic acid, but not by α -methyl-glutamate.

In chapter 5 the substrate and cation-selectivity of the H^+ /glutamate and Na^+ /glutamate symport proteins of *E. coli* K-12 ($GltP_{EC}$ and $GltS_{EC}$, respectively), and the Na^+/H^+ /glutamate symport proteins of *B. stearothermophilus* ($GltT_{Bs}$) and *B. caldotenax* ($GltT_{Bc}$) are described. L-glutamate transport by these proteins was studied in membrane vesicles derived from cells in which the proteins were either homologously or heterologously expressed. For this purpose, a glutamate transport deficient strain (*gltS⁻, gltP⁻*) of *E. coli* (ECOMUT1) was constructed. Transport of L-glutamate by each of the proteins was shown to be electrogenic and characterized as high affinity L-glutamate cation-symport. The $GltP_{EC}$, $GltT_{Bs}$ and $GltT_{Bc}$ transporters translocate L-glutamate and L-aspartate, whereas $GltS_{EC}$ only facilitates the transport of L-glutamate. Transport of L-glutamate by $GltP_{EC}$, $GltT_{Bs}$ and $GltT_{Bc}$ is inhibited by β -hydroxyaspartate and cysteic acid, but not by the $GltS_{EC}$ inhibitor α -methyl-glutamate. $GltS_{EC}$ symports L-glutamate with (at least) two sodium ions. When the L-glutamate transporters $GltP_{EC}$, $GltT_{Bs}$ and $GltT_{Bc}$ are expressed in *E. coli* transport of L-glutamate was found to be electrogenic and to occur in symport with $\geq 2 H^+$. This finding is surprising since earlier studies in membrane vesicles of *B. stearothermophilus* and *B. caldotenax* indicated a $Na^+/H^+/L$ -glutamate symport for both $GltT_{Bs}$ and $GltT_{Bc}$. The sodium ion dependency of these $GltT$ transporters was observed to increase with temperature. These observations suggest that the coupling ion-specificity is influenced by the conformation of the transport proteins, which is different in *E. coli* and the *Bacillus* membranes.