

University of Groningen

## The terminal oxidases of *Paracoccus denitrificans*

Gier, Jan-Willem L. de; Lübben, Mathias; Reijnders, Willem N.M.; Tipker, Corinne A.; Slotboom, Dirk-Jan; Spanning, Rob J.M. van; Stouthamer, Adriaan H.; Oost, John van der

*Published in:*  
Molecular Microbiology

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
1994

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Gier, J-W. L. D., Lübben, M., Reijnders, W. N. M., Tipker, C. A., Slotboom, D-J., Spanning, R. J. M. V., Stouthamer, A. H., & Oost, J. V. D. (1994). The terminal oxidases of *Paracoccus denitrificans*. *Molecular Microbiology*, 13(2), 183-196.

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

# The terminal oxidases of *Paracoccus denitrificans*

Jan-Willem L. de Gier,<sup>1</sup> Mathias Lübben,<sup>2</sup>  
Willem N. M. Reijnders,<sup>1</sup> Corinne A. Tipker,<sup>1</sup>  
Dirk-Jan Slotboom,<sup>1,2</sup> Rob J. M. van Spanning,<sup>1</sup>  
Adriaan H. Stouthamer<sup>1</sup> and John van der Oost<sup>1\*</sup>

<sup>1</sup>Department of Molecular and Cellular Biology,  
BioCentrum Amsterdam, Vrije Universiteit, De Boelelaan  
1087, 1081 HV Amsterdam, The Netherlands.

<sup>2</sup>European Molecular Biology Laboratory,  
Meyerhofstrasse 1, 69018 Heidelberg, Germany.

## Summary

Three distinct types of terminal oxidases participate in the aerobic respiratory pathways of *Paracoccus denitrificans*. Two alternative genes encoding subunit I of the  $aa_3$ -type cytochrome *c* oxidase have been isolated before, namely *ctaDI* and *ctaDII*. Each of these genes can be expressed separately to complement a double mutant ( $\Delta ctaDI$ ,  $\Delta ctaDII$ ), indicating that they are isoforms of subunit I of the  $aa_3$ -type oxidase. The genomic locus of a quinol oxidase has been isolated: *cyoABC*. This protohaem-containing oxidase, called cytochrome  $bb_3$ , is the only quinol oxidase expressed under the conditions used. In a triple oxidase mutant ( $\Delta ctaDI$ ,  $\Delta ctaDII$ , *cyoB*::Km<sup>R</sup>) an alternative cytochrome *c* oxidase has been characterized; this  $cbb_3$ -type oxidase has been partially purified. Both cytochrome  $aa_3$  and cytochrome  $bb_3$  are redox-driven proton pumps. The proton-pumping capacity of cytochrome  $cbb_3$  has been analysed; arguments for and against the active transport of protons by this novel oxidase complex are discussed.

## Introduction

In mitochondrial respiration the reduction of oxygen to water is catalysed by cytochrome *c* oxidase. This membrane-bound complex is the last component of the respiratory chain, in which electrons from a reductant (NADH, succinate) are transferred in subsequent redox reactions to oxygen (Babcock and Wikström, 1992). In bacteria aerobic respiration is often a more complex branched pathway. For many bacterial species more than one terminal oxidase has been described, catalysing the oxidation of either cytochrome *c* or quinol. The

possession of alternative respiratory branches enables a bacterium to adapt its efficiency of oxidative phosphorylation to changes in the environmental conditions.

An extensively studied prokaryotic cytochrome *c* oxidase is cytochrome  $aa_3$  from *Paracoccus denitrificans* and *Rhodobacter sphaeroides*. The primary sequences of its three subunits show strong homology to the mitochondrion-encoded eukaryotic cytochrome oxidase subunits I, II and III (Raitio *et al.*, 1987; 1990; Steinrücke *et al.*, 1987; Cao *et al.*, 1991; Shapleigh and Gennis 1992; Saraste, 1990). Subunit I of cytochrome  $aa_3$  contains two haems A: one is in the low-spin (6-co-ordinate) configuration and the other is in the high-spin (5-co-ordinate) configuration. The catalytic site of oxygen reduction is formed by a binuclear centre of the high-spin haem A and a copper ion, Cu<sub>B</sub> (Wikström *et al.*, 1981). Subunit II harbours another copper site, Cu<sub>A</sub> (van der Oost *et al.*, 1992; Lappalainen *et al.*, 1993; von Wachenfeldt *et al.*, 1994), which is probably involved in the oxidation of the substrate cytochrome *c* (Hill, 1993). Like the mitochondrial cytochrome *c* oxidase (Wikström, 1977; Krab and Wikström, 1978), the bacterial cytochrome  $aa_3$  is a redox-driven proton pump: the reduction of oxygen is coupled to the translocation of protons across the membrane (van Verseveld *et al.*, 1981; Solioz *et al.*, 1981; Hosler *et al.*, 1993).

The best-characterized prokaryotic quinol oxidase is cytochrome  $bo_3$  from *Escherichia coli*. As the name suggests, a low-spin haem B and a high-spin haem O are located in the major subunit, in addition to Cu<sub>B</sub> (Minghetti *et al.*, 1992). Like the cytochrome *c* oxidases, cytochrome  $bo_3$  functions as a proton pump (Puustinen *et al.*, 1989). Sequence analysis has revealed that cytochrome  $bo_3$  is closely related to cytochrome  $aa_3$  (Saraste *et al.*, 1988; Chepuri *et al.*, 1990). However, in contrast to the  $aa_3$ -type cytochrome *c* oxidases the quinol-oxidizing cytochrome  $bo_3$  does not contain a Cu<sub>A</sub> site (Puustinen *et al.*, 1991; van der Oost *et al.*, 1992). An alternative type of quinol oxidase from *E. coli*, cytochrome *bd*, is not related to the haem-copper oxidases (Anraku and Gennis, 1987; Green *et al.*, 1988).

Apart from the  $aa_3$ -type cytochrome *c* oxidase (John and Whatley, 1975), *P. denitrificans* has been reported to express alternative terminal oxidases. Several ubiquinol oxidases (cytochrome  $b_o$ , Cox *et al.*, 1978; cytochrome *d*, Henry and Vignais, 1979; cytochrome  $a_1$ , van Verseveld *et al.*, 1983; cytochrome  $ba_3$ , Ludwig, 1992) and cytochrome *c* oxidases (cytochrome  $c_o$ , Bosma, 1989; cytochrome  $aa_3$  isoenzyme, Raitio *et al.*, 1990) have

been either demonstrated or suggested. However, in contrast to cytochrome *aa*<sub>3</sub> the alternative oxidases from *Paracoccus* have not been described in detail.

In this paper a molecular genetic approach is used to acquire more detailed information on the terminal oxidases from *Paracoccus*. A new oxidase locus (*cyoABC*) has been cloned and its product identified as a protohaem-containing quinol oxidase. Single and multiple mutants have been generated for cytochrome *aa*<sub>3</sub>, the putative cytochrome *aa*<sub>3</sub> isoenzyme, and the quinol oxidase. For the first time expression of the *ctaDI* gene has been observed and the iso-cytochrome *aa*<sub>3</sub> has been characterized. After mutagenesis of the *cyoABC* locus no alternative quinol oxidase activity is detectable. In a triple oxidase mutant a *cb*-type cytochrome *c* oxidase has been analysed in more detail.

## Results

### Mutagenesis of cytochrome *aa*<sub>3</sub> and iso-cytochrome *aa*<sub>3</sub>

The genomic loci encoding the *aa*<sub>3</sub>-type cytochrome *c* oxidase (*ctaCBGE* and *ctaDII*), as well as the gene encoding a putative subunit I isoenzyme (*ctaDI*) have been isolated previously (Raitio *et al.*, 1987; 1990; Steinrücke *et al.*, 1987; van Spanning *et al.*, 1990). In the course of this study deletions have been generated in both subunits I ( $\Delta$ *ctaDI*,  $\Delta$ *ctaDII*) as shown in Fig. 1A, using a gene-replacement method as described by van Spanning *et al.* (1991). In this method the target gene is first inactivated by insertion of a kanamycin-resistance marker; the subsequent removal of this insertion cassette in a second recombination event results in an unmarked deletion.


### Cloning and mutagenesis of a new oxidase locus

Degenerated oligonucleotides derived from conserved regions in subunit I of haem-copper oxidases have been designed as described by M. Lübben *et al.* (submitted). A polymerase chain reaction (PCR) has been performed on genomic DNA from the cytochrome *aa*<sub>3</sub> double mutant ( $\Delta$ *ctaDI*,  $\Delta$ *ctaDII*), in conditions essentially as before (M. Lübben *et al.*, submitted). A PCR product of the expected size has been cloned into an M13 derivative and sequenced. The predicted amino acid sequence turned out to be very similar to that of subunit I of cytochrome *bo*<sub>3</sub> from *E. coli* (Chepuri *et al.*, 1990). Using the PCR fragment as a probe, a genomic *SphI* fragment of approx. 5 kb has been isolated (Fig. 1B).

### Sequence analysis of the *cyoABC* locus

The genomic *SphI* fragment has been subcloned and sequenced. Independently, the same locus has been isolated by O. Preisig and B. Ludwig (personal communication); their complete sequence has been submitted to the EMBL Data Library and they will publish their data elsewhere. Upstream of the subunit I gene (*cyoB*) a subunit II (*cyoA*) homologue was found, downstream of a *cyoB* subunit III-like gene (*cyoC*) (Fig. 1B). Screening of the data library revealed highest homology with *E. coli cyoABCDE* (Chepuri *et al.*, 1990) and *Bacillus subtilis qox-ABCD* (Santana *et al.*, 1992), encoding the quinol oxidases cytochrome *bo*<sub>3</sub> and *aa*<sub>3</sub>, respectively.

Primary sequences have been analysed in order to obtain information on the metal-binding sites. In subunit I of haem-copper oxidases six invariant histidines are the established ligands for the redox centres, i.e. two haems



**Fig. 1.** Genomic oxidase loci of *P. denitrificans*.

A. Subunit I genes of iso-cytochrome *aa*<sub>3</sub> (*ctaDI*) and cytochrome *aa*<sub>3</sub> (*ctaDII*); the latter is located adjacent to the *cycA* gene, encoding cytochrome *c*<sub>550</sub>.

B. The *cyoABC* locus, encoding the *b*-type quinol oxidase (cytochrome *bb*<sub>3</sub>). Sites of deletion and/or insertion mutagenesis are indicated.

## A subunit I

		helix II	helix VI
P.d. CtaDI	...	VVTY <b>H</b> GILMMFFVVPALFG...	...HILWFFG <b>H</b> PEVYIIILPGFG...
P.d. CtaDII	...	MITY <b>H</b> GVLMFFVVPALFG...	...HILWFFG <b>H</b> PEVYIIILPGFG...
B.s. CtaD	...	VMT <b>H</b> GGTMIFLAAMPLFA...	...HLFWIFG <b>H</b> PEVYILILPAFG...
B.j. FixN	...	LRPL <b>H</b> TSAVIFAFGGNVLIA...	...MFQWWY <b>H</b> NAVGFLLTAGFF...
P.d. CyoB	...	IFTA <b>H</b> GVIMIFFVAMPFITG...	...NLIWIW <b>H</b> PEVYILILPLFG...
E.c. CyoB	...	IFTA <b>H</b> GVIMIFFVAMPFVIG...	...NLIWAW <b>H</b> PEVYILILPVFG...
S.a. SoxA	...	ALT <b>H</b> GWAAAMIAFVPMAAA...	...ILFWFY <b>H</b> PVVVYVFPPLFG...
		102 121	277 296
		helix VII	helix X
P.d. CtaDI	...	MAAIAFLGFIVWA <b>H</b> HMYTAG...	...YHDYYIV <b>A</b> H <b>F</b> HYVMSLGAV...
P.d. CtaDII	...	MAAIGILGFVVWA <b>H</b> HMYTAG...	...YHDYYV <b>V</b> A <b>H</b> FHYVMSLGAV...
B.s. CtaD	...	AIVLGFGLGFVV <b>V</b> H <b>H</b> MFTTG...	...FHDTYF <b>V</b> V <b>A</b> H <b>F</b> HYVVIIGGV...
B.j. FixN	...	FWALIFLYIWAG <b>P</b> H <b>H</b> LHYTA...	...SHYTDWT <b>I</b> G <b>H</b> V <b>H</b> SALGWVG...
P.d. CyoB	...	TVCITVLSYLVL <b>L</b> H <b>H</b> FFTMG...	...LHNSLFL <b>I</b> A <b>H</b> F <b>H</b> NVIIGGV...
E.c. CyoB	...	TVCITVLSFIVL <b>L</b> H <b>H</b> FFTMG...	...LHNSLFL <b>I</b> A <b>H</b> F <b>H</b> NVIIGGV...
S.a. SoxA	...	IYLLAIGTMGV <b>V</b> H <b>L</b> LQTPW...	...FHNSY <b>V</b> V <b>G</b> H <b>F</b> H <b>L</b> MIWTLII...
		320 339	410 429

## B subunit II

P.d. CtaC	...	TATDVI <b>H</b> AWTIPAFVAVKQDAVPGRIAQLWFSVD-----QEGVYFGQCSELGGIN <b>H</b> AYMPIVVK...
B.s. CtaC	...	KASDV <b>K</b> HSFWIPSVGGKLDNTDNDENKFFLTFDSKRSKEAGDMFFGKCAELCGPS <b>H</b> ALMDFKVKT...
P.d. CyoA	...	TSTSVMN <b>A</b> FYIPAMAGMIYAMPGMETKLNGLV-----HPGKYKGIASHYSYSGHGF <b>S</b> GMHFKAHA...
E.c. CyoA	...	TNSVSMNS <b>F</b> IPRLGSQIYAMAGMQTRLHLIAN-----EPGTYDGISASYSYSGP <b>G</b> FMKFKATA...
		166 225


**Fig. 2.** Alignment of metal-binding amino acid sequences of four putative membrane-spanning helices of subunit I (A), and the C-terminal periplasmic domain of subunit II (B). Ligands are in bold. *P. denitrificans* CtaDI, CtaDII, CtaC (Raitio *et al.*, 1987; 1990), *B. subtilis* CtaD, CtaC (Saraste *et al.*, 1990), *B. japonicum* FixN (Preisig *et al.*, 1993), *P. denitrificans* CyoB, CyoA (this study), *E. coli* CyoB, CyoA (Chepuri *et al.*, 1990), *Sulfolobus acidocaldarius* SoxB (Lübben *et al.*, 1992). Sequence numbering refers to *E. coli* proteins.

and a copper ion, Cu<sub>B</sub> (reviewed by Hosler *et al.*, 1993). Alignment of *Paracoccus* CyoB sequences with corresponding fragments from several haem-copper oxidases clearly indicates similar metal-binding sites (Fig. 2A). A fourth redox centre resides in subunit II of cytochrome *c* oxidase, i.e. a binuclear copper centre, Cu<sub>A</sub> (van der Oost *et al.*, 1992; Lappalainen *et al.*, 1993; von Wachenfeldt *et al.*, 1994). Five amino acids in subunit II (2 cysteines, 2 histidines and 1 methionine) have been identified as a copper ligand (Kelly *et al.*, 1993). In quinol-oxidizing members of the haem-copper oxidase family this copper centre is absent (Puustinen *et al.*, 1991; van der Oost *et al.*, 1992). In *Paracoccus* CyoA four of the five Cu<sub>A</sub> ligands are substituted (Fig. 2B), indicating that the *cyoABC* genes code for a quinol oxidase.

## Optical spectra, oxygen consumption

Deletion of the *Paracoccus ctaDI* gene, proposed to encode subunit I of cytochrome *aa*<sub>3</sub> (Raitio *et al.*, 1987), did not affect the cytochrome *aa*<sub>3</sub> biosynthesis. This result has led to the isolation of a second gene, *ctaDII* (Raitio *et al.*, 1990). Deletion of the latter gene ( $\Delta$ *ctaDII*) results in a complete loss of cytochrome *aa*<sub>3</sub> absorption, with typical maxima at 445 nm and 605 nm (de Gier *et al.*, 1992; Raitio and Wikström, 1994).

The single, double, and triple oxidase mutants generated in this study have been analysed spectroscopically using whole cells, grown in batch culture on minimal medium with succinate (Fig. 3). As expected, the cytochrome *aa*<sub>3</sub> double mutant ( $\Delta$ *ctaDI*,  $\Delta$ *ctaDII*) has a



**Fig. 3.** Optical spectra of cells of *Paracoccus*. Cells were grown in batch cultures on minimal medium with succinate, and harvested during late-exponential growth. The figure presents absolute spectra of dithionite-reduced cell suspensions: wild type (Pd1222), cytochrome *bb*<sub>3</sub> mutant (Pd2621), cytochrome *aa*<sub>3</sub> double mutant (Pd9220), and the triple oxidase mutant (Pd9311).

**Table 1.** Oxygen-consumption measurements on whole-cell suspensions of different *P. denitrificans* strains: Pd1222 (wild type), Pd9220 (*aa*<sub>3</sub> double mutant), Pd2621 (*bb*<sub>3</sub> mutant), and Pd9311 (*aa*<sub>3</sub>/*bb*<sub>3</sub> triple mutant).

Strain	Inhibitor	Substrate			Ascorbate/TMPD			
		Endogenous	Succinate	Ascorbate/TMPD	Endogenous	Succinate	Ascorbate/TMPD	
Wild type	-	1.2	0.9	0.9	3.8	1.0	1.0	4.3
<i>aa</i> <sub>3</sub> double mutant	-	1.2	1.2	1.2	3.8	3.2	3.3	3.1
<i>bb</i> <sub>3</sub> mutant	-	1.1	0	0	3.6	0	0	4.3
<i>aa</i> <sub>3</sub> / <i>bb</i> <sub>3</sub> triple mutant	-	1.2	0	0	3.7	0	0	3.4

Values are the means of three independent assays (nmol O<sub>2</sub> mg protein<sup>-1</sup> s<sup>-1</sup>). Inhibitors: antimycin A (AA), myxothiazol (myx).

cytochrome *aa*<sub>3</sub>-minus phenotype. The spectrum of the *cyoB* mutant (*cyoB*::Km<sup>R</sup>), on the other hand, did not differ significantly from that of the wild type. In the optical spectrum of the triple mutant ( $\Delta$ *ctaDI*,  $\Delta$ *ctaDII*, *cyoB*::Km<sup>R</sup>) a clear increase of cytochrome *b* and *c* is observed (Fig. 3).

The oxygen consumption of wild-type and mutant cells has been measured polarographically, both without supplemented substrate (endogenous) and after addition of succinate (Table 1). In separate assays the electron flow via cytochrome *c* reductase has been inhibited by addition of either antimycin A or myxothiazol. The endogenous respiration of the wild type and the *aa*<sub>3</sub> double mutant is only slightly affected by these inhibitors. The inhibition becomes relatively more pronounced when succinate is used as substrate. In contrast, the oxygen consumption is blocked completely in the *cyoB* mutant and in the triple mutant. This is in agreement with the analysis of the primary structure, and implies that CyoABC encodes a quinol oxidase. In addition, the oxidation of the artificial electron-donating couple ascorbate and *N,N,N',N'*-tetramethyl-*p*-phenylenediamine (TMPD) has been assayed; more or less equal values indicate the presence of cytochrome *c* oxidase(s) in all strains.

#### Mutant complementations

All three oxidase mutants have been complemented using an expression vector and the triple oxidase mutant as a host. Although the *lac* promoter is thought to be inactive in *Paracoccus*, the orientation of the oxidase genes in the expression constructs is such that they are encoded by the complementary strand, indicating that transcription has been under the control of the native promoters. An interesting observation was that all three complementation constructs, but not the vector alone, gave rise to a decline of the *bc* peak (Fig. 4).


The plasmid-encoded expression of CtaDII was demonstrated by the restoration of the cytochrome *aa*<sub>3</sub> optical

spectrum, with characteristic maxima at 445 nm and 605 nm (not shown). For the first time the spectrum of cytochrome *aa*<sub>3</sub> with subunit I expressed from *ctaDI* has been observed (Fig. 4); it is obvious that it concerns a cytochrome *aa*<sub>3</sub> isoenzyme, as originally proposed by Raitio *et al.* (1990).

The quinol oxidase has been complemented using the *SphI* fragment shown in Fig. 1B. In this case the effect was most clearly demonstrated by analysis of the oxygen consumption. As stated above, the respiration of the triple mutant was inhibited completely by antimycin A and myxothiazol (Table 1). In the CyoABC-complemented clone, however, the direct oxidation of ubiquinol (independent of cytochrome *c* oxidase) is restored (not shown).

#### Haem analysis

Haems have been extracted from wild-type and mutant cells and subsequently analysed by high-performance liquid chromatography (HPLC) reversed-phase chromatography (Table 2). In extracts of the wild type and the quinol oxidase mutant, haem B (protohaem) and haem A are present abundantly. In the cytochrome *aa*<sub>3</sub> double mutant and in the triple mutant the haem B quantity is not altered significantly; in contrast, haem A is just detectable in the absence of cytochrome *aa*<sub>3</sub> (the concentration decreased 10–15-fold). At least under the cultivation conditions used, no haem O is detected in membranes of *Paracoccus*. This implies that the cloned quinol oxidase is a *b*-type cytochrome, and that the alternative cytochrome *c* oxidase is a *cb*-type cytochrome.



**Fig. 4.** Optical spectra of cells of *Paracoccus*. The figure presents absolute spectra of dithionite-reduced cell suspensions (as described in legend Fig. 3) of the triple oxidase mutant Pd9311, and triple oxidase mutant after complementation with CtaDI: Pd9311(pEG.ctaDI).

**Table 2.** Haem analysis by reversed-phase HPLC (Deltapak C18 column, Waters) of different *P. denitrificans* strains: Pd1222 (wild type), Pd9220 (*aa*<sub>3</sub> double mutant), Pd2621 (*bb*<sub>3</sub> mutant), and Pd9311 (*aa*<sub>3</sub>/*bb*<sub>3</sub> triple mutant).

Strain	Haem		
	B	O	A
Wild type	++	—	++
<i>aa</i> <sub>3</sub> double mutant	++	—	±
<i>bb</i> <sub>3</sub> mutant	++	—	++
<i>aa</i> <sub>3</sub> / <i>bb</i> <sub>3</sub> triple mutant	++	—	±


Haems were extracted from isolated membranes and eluted on an acetonitrile gradient in 0.05% trifluoroacetic acid, as described by Sone and Fujiwara (1991). Relative quantities of extracted haem: ++, abundantly present; ±, just detectable; —, not detectable.

#### Partial purification of alternative cytochrome *c* oxidase

From membranes of the triple mutant ( $\Delta$ *ctaDI*,  $\Delta$ *ctaDII*, *cyoB::Km<sup>R</sup>*), grown on minimal medium with succinate, the alternative cytochrome *c* oxidase has been extracted. Partial purification of the oxidase complex has been accomplished by column chromatography, and the resulting preparation oxidizes both TMPD and horse-heart cytochrome *c* at high rates (M. Saraste and J. van der Oost, unpublished). The optical spectrum shows maxima at 551 nm and 559 nm, indicating the presence of the cytochromes *c* and *b* (Fig. 5).

#### Proton translocation

The coupling of respiratory electron ( $e^-$ ) transfer to the translocation of protons ( $H^+$ ) across the bacterial cytoplasmic membrane drives the oxidative phosphorylation. The efficiency of the latter phenomenon in the different mutants has been studied by determination of the  $H^+/2e^-$  stoichiometry. A cell suspension is supplemented



**Fig. 5.** Absolute absorbance spectrum of the alternative cytochrome *c* oxidase (cytochrome *cbb*<sub>3</sub>) complex after reduction with Na-dithionite; the oxidase has been extracted from membranes of the triple oxidase mutant Pd9311.

with substrate, either ascorbate and TMPD or succinate; a subsequent oxygen pulse induces the proton efflux, which is measured as acidification of the medium (Fig. 6, Table 3).


Ascorbate donates electrons via TMPD to cytochrome *c* (van Verseveld *et al.*, 1981) (Fig. 7). When ascorbate is oxidized to dehydro-ascorbate one 'scalar' proton is released per two electrons; when a proton-pumping cytochrome *c* oxidase is operative, a maximum of two

**Table 3.** Proton-translocation measurements on whole-cell suspensions of different *P. denitrificans* strains.  $H^+/2e^-$  ratios are averages  $\pm$  standard deviation.

Strain	succinate O <sub>2</sub> ( $H^+/2e^-$ )	succinate Fe(CN) <sub>6</sub> <sup>3-</sup> ( $H^+/2e^-$ )	ascorbate O <sub>2</sub> ( $H^+/2e^-$ )
Wild type(s)			2.40 $\pm$ 0.11
<i>aa</i> <sub>3</sub> double mutant(s)			1.21 $\pm$ 0.16
<i>bb</i> <sub>3</sub> mutant(s)	5.15 $\pm$ 0.21	3.99 $\pm$ 0.11	2.01 $\pm$ 0.18
<i>aa</i> <sub>3</sub> / <i>bb</i> <sub>3</sub> triple mutant(s)	4.15 $\pm$ 0.14	4.05 $\pm$ 0.10	1.12 $\pm$ 0.10
<i>aa</i> <sub>3</sub> / <i>bb</i> <sub>3</sub> triple mutant, <i>ctaDI</i> compl.(s)			2.20 $\pm$ 0.36
<i>aa</i> <sub>3</sub> / <i>bb</i> <sub>3</sub> triple mutant, <i>ctaDII</i> compl.(s)			2.12 $\pm$ 0.10
Wild type, aerobic/ methanol-limited (m/ch)			2.81 $\pm$ 0.14
Wild type, aerobic/O <sub>2</sub> - limited (s/ch)			1.21 $\pm$ 0.35
Wild type, anaerobic/ NO <sub>3</sub> -limited (s/ch)			1.19 $\pm$ 0.07

Cells were cultivated as aerobic batch cultures, in minimal medium with either succinate (s) or methanol (m); data are compared with analyses from chemostat-cultured cells (ch) of *P. denitrificans* NCIB 8944 (wild type) with limitations as indicated, reported by van Verseveld *et al.* (1981; 1983). The electron donor was either succinate or ascorbate; anaerobic suspensions were pulsed with oxygen (O<sub>2</sub>) or ferricyanide (Fe(CN)<sub>6</sub><sup>3-</sup>). A control pulse with ferricyanide of the ascorbate mixture routinely resulted in  $H^+/2e^- = 1.1 \pm 0.1$ , both in the presence and in the absence of cells (not shown). The assay was as described in the legend to Fig. 5; for details see the *Experimental procedures*.

'vectorial' protons is translocated to the medium in addition (Wikström *et al.*, 1981). In the wild type and the quinol oxidase mutant an  $H^+/2e^-$  ratio of 2.0–2.4 is measured upon ascorbate oxidation; in the *aa*<sub>3</sub> double mutant and the triple oxidase mutant an  $H^+/2e^-$  ratio of 1.1–1.2 (Table 3) is found. Apparently, vectorial proton translocation is detected only in the presence of cytochrome *aa*<sub>3</sub>, which is an established proton pump (van Verseveld *et al.*, 1981); in its absence, however, only the scalar proton release is measured. This observation is confirmed by analysis of the triple oxidase mutant, in which cytochrome *aa*<sub>3</sub> is reintroduced by complementation with either *CtaDII* or *CtaDI*. The induced increase of the  $H^+/2e^-$  ratios (2.1–2.2) also indicates that both isoforms of cytochrome *aa*<sub>3</sub> are proton-pumping cytochrome *c* oxidases (Table 3).



**Fig. 6.** Respiration-driven proton translocation in cells of *Paracoccus*. Anaerobic cell suspensions were pulsed with oxygen ( $O_2$ ) by injection of air-saturated KCl. Traces were calibrated by injecting an anaerobic solution of 1 mM HCl. The calculated  $H^+/2e^-$  ratio is shown for each oxygen pulse.


Succinate-dehydrogenase catalyses the oxidation of succinate and the reduction of ubiquinone. Subsequently, ubiquinol is oxidized either by a quinol oxidase (cytochrome  $bb_3$ ;  $H^+/2e^- = 4.0$ ), or by cytochrome  $c$  reductase (cytochrome  $bc_1$ ;  $H^+/2e^- = 4.0$ ). When electrons flow from the latter complex, via cytochrome  $c$ , to a proton pumping cytochrome  $c$  oxidase the energy-transducing efficiency increases (cytochrome  $aa_3$ ;  $H^+/2e^- = 6.0$ ). In the actual experiments with succinate the same tendency is observed with cells of the  $aa_3$  double mutant and the triple oxidase mutant: in the presence of cytochrome  $aa_3$  the  $H^+/2e^-$  stoichiometry is 5.0–5.3, but in the cytochrome  $aa_3$  mutants it never exceeds an  $H^+/2e^-$  ratio of 4.0–4.3 (Table 3). As an internal control, the proton efflux associated with the succinate oxidation has been measured with ferricyanide as terminal electron acceptor.

## Discussion

*P. denitrificans* contains a number of alternative respiratory pathways, branching at the level of ubiquinol. One route closely resembles the mitochondrial respiratory chain in which ubiquinol is oxidized by the so-called supercomplex consisting of cytochrome  $c$  reductase (cytochrome  $bc_1$ ), cytochrome  $c_{552}$  and cytochrome  $c$  oxidase (cytochrome  $aa_3$ ) (Berry and Trumpower, 1985). In a commonly expressed alternative pathway ubiquinol is oxidized directly by a quinol oxidase called cytochrome  $b_0$  by Cox *et al.* (1978).

When *Paracoccus* is cultivated under microaerobic or anaerobic conditions, the expression level of cytochrome  $aa_3$  and, to some extent, that of cytochrome  $c_{552}$  appears significantly lower (Bosma *et al.*, 1987a). Instead of the

## *Paracoccus* respiratory pathways: sites of proton-translocation



**Fig. 7.** Respiratory pathways in *Paracoccus*. Indicated here is the theoretical stoichiometry of the respiration-driven proton translocation ( $H^+/2e^-$ ), detectable in the medium, for each of the contributing respiratory complexes: NADH-dehydrogenase (NADH.dh), succinate dehydrogenase (succ.dh), cytochrome  $c$  reductase (cyt. $bc_1$ ), cytochrome  $c_{552}$  (cyt. $c_{552}$ ),  $aa_3$ -type cytochrome  $c$  oxidase (cyt. $aa_3$ ),  $cbb_3$ -type cytochrome  $c$  oxidase (cyt. $cbb_3$ ) and the  $bb_3$ -type quinol oxidase (cyt. $bb_3$ ). Ascorbate (with TMPD) is an artificial reductant of cytochrome  $c$ , and ferricyanide ( $Fe(CN)_6^{3-}$ ) an oxidant. The chemical conversion of ascorbate is indicated (open arrow) to show generation of the 'scalar' proton. For discussion, see text.

aforementioned supercomplex, a quinol-oxidizing complex has been isolated from such oxygen-limited cells in which cytochrome  $bc_1$  is associated with an alternative  $cb$ -type cytochrome  $c$  oxidase, called cytochrome  $c_o$  by Bosma (1989).

#### Ubiquinol oxidase

A new oxidase locus (*cyoABC*) has been isolated from a *P. denitrificans* genomic library. The predicted protein sequences share a high degree of homology with subunits I, II and III of the quinol-oxidizing cytochrome  $bo_3$  from *E. coli* (Fig. 2, Chepuri *et al.*, 1990). In subunit I (CyoB) the six conserved histidines are present, indicating that it concerns a haem-copper oxidase (Saraste, 1990; Hosler *et al.*, 1993). As in *E. coli* CyoA, subunit II of this *Paracoccus* oxidase has substituted four out of five residues that have been identified as  $Cu_A$  ligands (Kelly *et al.*, 1993). This suggests that CyoABC is a quinol oxidase (Saraste *et al.*, 1991). Moreover, the oxygen consumption of the *cyoB* mutant was completely inhibited by either antimycin A or myxothiazol, whereas the wild-type *Paracoccus* strain was hardly affected by these compounds (Table 1). This confirms that the *cyoABC* locus codes for a quinol oxidase.

The *E. coli* *cyo* operon is a cluster of five genes: *cyoABCDE*. The *cyoD* gene product is thought to be a fourth subunit (Minghetti *et al.*, 1992), which may be absent from the *Paracoccus* quinol oxidase complex. The *cyoE* product is proposed to be a protohaem IX farnesyl transferase, involved in the conversion of haem B into haem O (Saiki *et al.*, 1992). Deletion of the latter gene in *E. coli* resulted in an oxidase with two B haems, however, without oxidase activity (Hill *et al.*, 1992; Saiki *et al.*, 1992).

In well-aerated, succinate-grown cells of *Paracoccus* the contribution of the quinol oxidase to the endogenous respiration varies: it is 75% in wild-type cells and 100% in the cytochrome  $aa_3$  double mutant. Apparently, relatively more electrons flow via the cytochrome  $c$  branch when succinate is added as a substrate (Table 1).

Haem analysis indicates that wild-type membranes, with quinol oxidase activity, contain haem B and haem A; in the cytochrome  $aa_3$  double mutant haem B and only traces of haem A could be detected (Table 2). Unlike haem A and haem B, haem O has never been detected in membrane extracts from the *Paracoccus* strains grown under these conditions. We conclude that the CyoABC quinol oxidase is a cytochrome with two protohaems. This implies that the hydroxyethylfarnesyl side-chain, which is present in haems A and O but not in haem B (Wu *et al.*, 1992), is not essential for catalytic activity or for proton translocation. Previously this oxidase has been referred to as cytochrome  $o$  or  $b_o$  (Cox *et al.*, 1978;

Puustinen *et al.*, 1989); we propose to call it cytochrome  $bb_3$ , according to the classical terminology of the mitochondrial cytochrome  $c$  oxidase in which the  $O_2$ -binding haem is denoted by the subscript 3 (Puustinen and Wikström, 1991).

An alternative quinol oxidase complex, cytochrome  $ba_3$ , has been purified from a *Paracoccus* cytochrome  $c$  reductase mutant (Ludwig, 1992). The genomic locus of cytochrome  $ba_3$  has been isolated (O. Richter and B. Ludwig, personal communication), and comparison of these data with the partial sequences as presented in Fig. 2 revealed that cytochrome  $ba_3$  and cytochrome  $bb_3$  are encoded by the same set of genes (*cyoABC*). As far as we are aware, the cytochrome  $bc_1$  mutant is the only *Paracoccus* strain in which this mixed haem oxidase is detectable. Hence, it may be more likely that the genotype rather than the growth conditions gives rise to the observed variable haem incorporation into the CyoB apo-cytochrome. Similar variability has recently been reported for the cytochrome  $c$  oxidase from the thermophilic bacterium PS3 (Sone and Fujiwara, 1991), and for quinol oxidases from *E. coli* and *Acetobacter aceti* as well (Puustinen *et al.*, 1992; Matsushita *et al.*, 1992).

In the quinol oxidase mutant and in the triple mutant, no alternative quinol oxidase activity could be demonstrated (Table 1). Moreover, extensive attempts to generate a cytochrome  $c$  reductase/quinol oxidase double mutant have not succeeded (not shown), which indeed might indicate that the *Paracoccus* genome contains only one quinol oxidase locus. Previously, the existence of two more terminal oxidases was deduced from optical spectra of whole cells. First, a cyanide-tolerant oxidase with a high affinity for oxygen and an absorbance maximum at 629 nm has been called cytochrome  $d$  by Henry and Vignais (1979). However, cytochrome  $bb_3$  might have been responsible for the reported cyanide-tolerant respiration of *Paracoccus*; in a membrane preparation of the cytochrome  $aa_3$  double mutant a  $K_i$  for cyanide of approx. 250  $\mu$ M has been determined, whereas the triple mutant is extremely sensitive to cyanide (J.-W. L. de Gier and J. van der Oost, unpublished). The observed absorbance band may have originated from the nitrite reductase cytochrome  $cd_1$  (Timkovitch *et al.*, 1982). Second, an oxidase that is expressed microaerobically, with an absorbance maximum at 590 nm, has been called cytochrome  $a_1$  by van Verseveld *et al.* (1983). It is not unlikely, however, that the observed cytochrome belongs to another class of enzymes, e.g. the CO-binding catalase-peroxidase cytochrome  $b_{590}$  (Appleby and Poole, 1991), which was originally proposed to be the high-affinity oxidase (cytochrome  $a_1$ ) from nitrogen-fixing *Bradyrhizobium japonicum*. Indeed, a cytochrome  $b_{595}$ -associated peroxidase activity has been observed in periplasmic fractions of *Paracoccus* (J. Ras, unpublished).



*aa<sub>3</sub>-type cytochrome c oxidase*

Expression of the *Paracoccus* cytochrome *aa<sub>3</sub>* has been studied extensively. When *Paracoccus* is growing on succinate, either microaerobically or anaerobically, cytochrome *aa<sub>3</sub>* is undetectable or barely detectable in the optical spectrum (Bosma, 1989; Stouthamer, 1991). In contrast, this cytochrome *c* oxidase plays an active role during autotrophic growth on the C<sub>1</sub> substrates methanol and methylamine (reviewed by Harms and van Spanning 1991). On the other hand, a cytochrome *aa<sub>3</sub>* mutant of *Paracoccus* (Willison *et al.*, 1981) is able to grow on methanol (Harms *et al.*, 1985), and a cytochrome *c* reductase/cytochrome *aa<sub>3</sub>* double mutant of *Paracoccus* is able to grow on methylamine (de Gier *et al.*, 1992). This suggests the presence of an alternative cytochrome *c* oxidase (see below).

The genomic loci encoding the cytochrome *aa<sub>3</sub>* isoenzymes have been isolated previously (reviewed by van der Oost *et al.*, 1991). In the triple oxidase mutant ( $\Delta$ *ctaDI*,  $\Delta$ *ctaDII*, *cyoB::Km<sup>R</sup>*) we show that it is possible to restore the optical spectrum of cytochrome *aa<sub>3</sub>* by complementation with either *CtaDI* or *CtaDII* (Fig. 4). This has been confirmed by Western blotting, using antibodies raised against the three-subunit cytochrome *aa<sub>3</sub>* from *Paracoccus* (not shown). No spectral/immunological evidence has been found so far for expression of *CtaDI* in the  $\Delta$ *ctaDII* mutant. The successful expression *in trans* might indicate that a repression of the *ctaDI* promoter activity has been (partially) overcome by the elevated copy number. The physiological conditions under which the expression of the cytochrome *aa<sub>3</sub>* isoenzyme is induced remain unknown.

*cb-type cytochrome c oxidase*

Cells of wild-type *Paracoccus* grown anaerobically/microaerobically on succinate are still able to oxidize ascorbate/TMPD although no *aa<sub>3</sub>-type* cytochrome *c* oxidase is detectable in the optical spectrum (van Verseveld *et al.*, 1983). The latter authors have reported 'the presence of a cytochrome oxidase with a very high affinity for oxygen and a low affinity for CO'. This high-affinity cytochrome *c* oxidase has been characterized in more detail by Bosma (1989), who partially purified it from anaerobically grown cells. The alternative cytochrome *c* oxidase was isolated as a complex with cytochrome *c* reductase, but without cytochrome *c*<sub>552</sub>. Spectroscopic analysis revealed that it concerns a *cb-type* oxidase complex, consisting of a cytochrome *b*, a 30 kDa cytochrome *c* and perhaps a loosely membrane-associated 45 kDa dihaem cytochrome *c* (Bosma *et al.*, 1987b; Bosma 1989).

In the present study a triple oxidase mutant ( $\Delta$ *ctaDI*,  $\Delta$ *ctaDII*, *cyoB::Km<sup>R</sup>*) has been generated, in which an

alternative cytochrome *c* oxidase has been studied *in vivo*. The optical spectrum of this mutant indicates the absence of cytochrome *aa<sub>3</sub>* and an increase in the *bc* pool (Fig. 3). This increased amount of cytochromes *b* and *c* turned out to be reversible: upon complementation of the triple mutant *in trans* with either one of the three oxidases, the absorbance peak decreased to a wild-type level (Fig. 4). The alternative cytochrome *c* oxidase has recently been partially purified from the membranes of the triple mutant; the optical spectrum indicates the presence of cytochromes *c* and *b* (Fig. 5). It is tempting to assume that an expression induction/repression of this *cb-type* oxidase is responsible for the observed increase/decrease in the absorbance spectra of the triple mutant (Figs 3 and 4). Analysis of the haems extracted from membranes of the triple mutant only indicates the presence of haem B (Table 2). Haem A and haem O are absent; haem C is not extracted, owing to its covalent binding. Oxygen-consumption measurements indicate that respiration in the triple mutant could be inhibited completely at the level of cytochrome *bc<sub>1</sub>* (Table 1). It may be that the alternative cytochrome *c* oxidase is the only remaining terminal oxidase in this mutant.

Recently, we obtained additional genetic information on this *cb-type* oxidase. Using peptide sequences from the aforementioned purified cytochrome *cb* complex, a DNA fragment has been isolated from a *P. denitrificans* genomic library; preliminary sequence analysis indicates the presence of a new oxidase locus (M. Saraste and J. van der Oost, unpublished). The isolated gene cluster is very similar to *fixNOQP* from *B. japonicum* (Preisig *et al.*, 1993). The genes from *Bradyrhizobium* have been demonstrated to encode an alternative oxidase that is expressed microaerobically or anaerobically; this is in good agreement with the finding that the gene cluster is located adjacent to the genes of the oxygen-sensing regulators FixJL. The predicted protein sequences suggest that FixN is the haem- and copper-binding subunit I homologue, FixO a monohaem cytochrome *c*, and FixP a dihaem cytochrome *c* (Preisig *et al.*, 1993). The alignment of the *Bradyrhizobium* FixN sequence (Fig. 2A) suggests that all six invariant histidines, typical of the haem-copper oxidase superfamily, are present in the *cb-type* cytochrome *c* oxidase; it probably concerns a *b-type* haem-copper oxidase (Table 2, see below) with associated cytochrome(s) *c*. In accordance with classical cytochrome oxidase terminology (Puustinen and Wikström, 1991), we propose to name this *cb-type* oxidase cytochrome *cbb<sub>3</sub>*.

In addition, a *b-type* cytochrome *c* oxidase has been described in *Rhodobacter capsulatus* (Hüdig *et al.*, 1987). Regulation has been demonstrated at two levels: the expression as well as the activity of this oxidase respond to oxygen and light. Upon a shift from phototrophic conditions to chemotrophic conditions (microaerobically) the

activity of this oxidase increases fourfold. Garcia-Horsman *et al.* (1994) have purified a *cb*-type cytochrome *c* oxidase from a *Rhodobacter sphaeroides* cytochrome *aa*<sub>3</sub> mutant. The oxidase consists of three subunits: two cytochromes *c* and a cytochrome *b*. Metal and haem analyses indicate that this *cbb*<sub>3</sub>-type oxidase contains haem C, haem B, and copper in a ratio of 3:2:1. The presence of a binuclear centre has been demonstrated, including a high-spin haem B and a Cu<sub>B</sub> (Garcia-Horsman *et al.*, 1994).

#### Proton translocation

Analysis of proton translocation by the generated set of *Paracoccus* oxidase mutants has been performed in order to determine whether or not the alternative cytochrome *c* oxidase is a proton-pumping oxidase. During succinate oxidation, when ubiquinol is oxidized directly by cytochrome *bb*<sub>3</sub>, a minimum H<sup>+</sup>/2e<sup>-</sup> stoichiometry of 4 is expected theoretically; when ubiquinol is oxidized via cytochrome *bc*<sub>1</sub> (H<sup>+</sup>/2e<sup>-</sup> = 4) and cytochrome *aa*<sub>3</sub> (H<sup>+</sup>/2e<sup>-</sup> = 2) the maximal H<sup>+</sup>/2e<sup>-</sup> stoichiometry is 6. An obvious consequence of the deletion of cytochrome *aa*<sub>3</sub> is a decreased H<sup>+</sup>/2e<sup>-</sup> stoichiometry (Table 3). When ferricyanide is used as terminal electron acceptor instead of oxygen, equal values are obtained because ferricyanide is reduced directly by the cytochrome *c* pool (Table 3, Fig. 6). A similar tendency is observed when ascorbate/TMPD is used as substrate, donating electrons at the level of cytochrome *c*. Apart from the scalar proton (H<sup>+</sup>/2e<sup>-</sup> = 1), an additional acidification is expected in the case of a proton-translocating cytochrome *c* oxidase, such as cytochrome *aa*<sub>3</sub> (H<sup>+</sup>/2e<sup>-</sup> = 2); the maximal H<sup>+</sup>/2e<sup>-</sup> stoichiometry would be 3. Again, in the absence of cytochrome *aa*<sub>3</sub> only the scalar ascorbate proton is measured (Table 3). When cytochrome *aa*<sub>3</sub> is present, an intermediate value of 2.0–2.4 is commonly detected. This indicates that, unlike cytochrome *aa*<sub>3</sub> (van Verseveld *et al.*, 1981) and cytochrome *bb*<sub>3</sub> (Puustinen *et al.*, 1989), the *cbb*<sub>3</sub>-type cytochrome *c* oxidase does not couple the reduction of oxygen to the active transport of protons across the membrane, at least under these conditions (see below).

The proton translocation data as presented in this study are consistent with the absence of a proton-pumping oxidase in cytochrome *aa*<sub>3</sub> deletion strains. However, Raitio and Wikström (1994) recently presented experimental data that strongly suggest that the alternative cytochrome *c* oxidase is able to pump protons. These authors report an H<sup>+</sup>/2e<sup>-</sup> ratio of 6, with whole cells (wild-type, cytochrome *aa*<sub>3</sub> double mutant), succinate as substrate, and oxygen as electron acceptor. With ascorbate/TMPD as the electron-donating system, on the other hand, hardly any pumping was observed (H<sup>+</sup>/2e<sup>-</sup> = 1.2–1.5). The buffer composition of the proton translocation assay as presented by the latter authors is the main

difference in the method used in this study (HEPES (0.5 mM) and glycylglycine (1.5 mM), respectively). Whereas the proton translocation linked to the cytochrome *aa*<sub>3</sub> activity is easily detectable when the cell suspension is buffered by either glycylglycine or HEPES, proton translocation by an alternative oxidase, which is likely to be cytochrome *cbb*<sub>3</sub>, is apparently only detectable in the HEPES-buffered cell suspension.

It has been reported previously that maximal rates of oxidative phosphorylation in *Paracoccus* were measured only in the presence of cytochrome *aa*<sub>3</sub>, e.g. after growth on methanol but not after microaerobic growth (Boogerd *et al.*, 1981; van Verseveld *et al.*, 1983) (Table 3). Stouthamer (1991) has recently reviewed the results obtained by his co-workers over the last decade, on *Paracoccus* cultivated in chemostats under a wide variety of growth conditions. Calculations of growth yields and proton-translocation rates (assayed in glycylglycine buffer) led him to suggest that cytochrome *aa*<sub>3</sub> is a proton-translocating cytochrome *c* oxidase, unlike the alternative cytochrome *c* oxidase. In view of the aforementioned variations of the *cbb*<sub>3</sub>-catalysed proton translocation, it can be imagined that the *in situ* energy-transducing efficiency of this oxidase might also vary under different external conditions. Chemostat experiments with the available set of oxidase mutants are intended to address this point.

Understanding the nature of the differences observed in the proton pumping behaviour of the *cbb*<sub>3</sub>-type oxidase as compared to, for example, the *aa*<sub>3</sub>-type oxidase will be the goal of future experiments. With current knowledge, it is anticipated that cytochrome *cbb*<sub>3</sub> is indeed capable of pumping protons, perhaps only under specific external conditions. A detailed study of this 'distant' member of the haem-copper oxidase family will contribute substantially to a model of the redox-driven proton pump.

## Experimental procedures

### *Bacterial strains, plasmids, and growth conditions*

The strains of *P. denitrificans* and *E. coli*, as well as the plasmids, that were used in this study are listed in Table 4. Cells of the wild-type *P. denitrificans* (Pd1222) and the mutant strains were cultivated in aerobic batch cultures (0.5 l bottles with 100 ml of culture, on a rotary shaker at 30°C), either with brain-heart infusion broth (BHI) or minimal medium supplemented with 25 mM succinate, as described previously (van Spanning *et al.*, 1991). *E. coli* strains were cultivated in YT medium at 37°C. When appropriate, antibiotics were added: rifampicin (Rif, 40 mg l<sup>-1</sup>), streptomycin (Sm, 50 mg l<sup>-1</sup>), tetracycline (Tc, 12.5 mg l<sup>-1</sup>), kanamycin (Km, 50 mg l<sup>-1</sup>) and ampicillin (Amp, 100 mg l<sup>-1</sup>).

### *DNA manipulations*

General cloning techniques were carried out as described by

Table 4. Strains and plasmids.

Strain/Plasmid	Relevant characteristics	Reference
<u>Strain</u>		
<i>E. coli</i>		
TG1	<i>supE</i> , <i>hsdD5</i> , <i>thi</i> , $\Delta(lac^- proAB)$ , F', ( <i>traD36 proAB lacI<sup>q</sup> lacZ</i> $\Delta$ M15)	Sambrook et al. (1989)
HB101	F <sup>-</sup> , <i>hsdS20</i> , ( <i>r<sub>B</sub><sup>-</sup> m<sub>B</sub><sup>-</sup></i> ), <i>lacY1</i> , <i>proA2</i> , <i>recA13</i>	Boyer et al. (1969)
S17.1	Sm <sup>R</sup> , <i>pro</i> , ( <i>r<sub>B</sub><sup>-</sup> m<sub>B</sub><sup>+</sup></i> ), RP4-2, integrated (Tc::Mu) (Km::Tn7)	Simon et al. (1983)
<i>P. denitrificans</i>		
Pd1222	Rif <sup>R</sup> , enhanced conjugation frequencies, ( <i>r<sub>B</sub><sup>-</sup> m<sub>B</sub><sup>+</sup></i> )	De Vries et al. (1989)
Pd2521	Pd1222 derivative, <i>ctaDI</i> ::Km <sup>R</sup>	This work
Pd2541	Pd2521 derivative, $\Delta$ <i>ctaDI</i>	This work
Pd9218	Pd2541 derivative, $\Delta$ <i>ctaDI</i> , <i>ctaDI</i> ::Km <sup>R</sup>	This work
Pd9220	Pd9218 derivative, $\Delta$ <i>ctaDI</i> , $\Delta$ <i>ctaDII</i>	This work
Pd2621	Pd1222 derivative, <i>cyoB</i> ::Km <sup>R</sup>	This work
Pd9311	Pd9220 derivative, $\Delta$ <i>ctaDI</i> , $\Delta$ <i>ctaDII</i> , <i>cyoB</i> ::Km <sup>R</sup>	This work
<u>Plasmid</u>		
pUC18	Amp <sup>R</sup> , <i>lacZ'</i>	Yanisch-Perron et al. (1985)
pUC4K	Km <sup>R</sup> (Tn903)	Pharmacia
pUC4KISS	Km <sup>R</sup> (Tn903)	Pharmacia
M13mp18	pUC18 mcs, <i>lacZ'</i>	Sanger et al. (1980)
pEG400/401	<i>IncP</i> , Sm <sup>R</sup> , Spec <sup>R</sup> , pUC12/13 mcs, <i>lacZ'</i>	Gerhus et al. (1990)
pRK2020	Tc <sup>R</sup> , pRK2013 Km::Tn10	Ditta et al. (1985)
pGRPd1	<i>oriV</i> , (colE1), Amp <sup>R</sup> , <i>oriT</i> , Sm <sup>R</sup> , (Tn1831)	Van Spanning et al. (1990)
pRVS1	<i>oriV</i> (colE1), Amp <sup>R</sup> , <i>oriT</i> , Sm <sup>R</sup> , (Tn1831), Tn5p, <i>lacZ</i>	Van Spanning et al. (1991)
pMR3	pUC18 derivative, <i>ctaDI</i>	M. Raitio, unpublished
pMR3.Km <sup>R</sup>	pMR3 derivative, <i>ctaDI</i> ::Km <sup>R</sup>	This work
pMR3D	pMR3 derivative, $\Delta$ <i>ctaDI</i>	This work
pUC.cyoABC	pUC18 derivative, <i>cyoABC</i>	This work
pUC.cyoABC.Km <sup>R</sup>	pUC.cyoABC derivative, <i>cyoB</i> ::Km <sup>R</sup>	This work
pRT2521	pGRPd1 derivative, <i>ctaDI</i> ::Km <sup>R</sup>	This work
pRT2541	pRVS1 derivative, $\Delta$ <i>ctaDI</i>	This work
pRT2321	pGRPd1 derivative, <i>ctaDII</i> ::Km <sup>R</sup>	This work
pRT2341	pRVS1 derivative, $\Delta$ <i>ctaDII</i>	This work
pRT2621	pRVS1 derivative, <i>cyoB</i> ::Km <sup>R</sup>	This work
pEG.ctaDII	pEG400 derivative, <i>ctaDII</i>	This work
pEG.ctaDI	pEG401 derivative, <i>ctaDI</i>	This work
pEG.cyoABC	pEG401 derivative, <i>cyoABC</i>	This work

Ausubel et al. (1992). Conjugations were performed as described previously (de Vries et al., 1989). The matings of *Paracoccus* host strains were performed either directly with *E. coli* S17-1 transformed with the plasmid of interest, or via a triparental mating using any *E. coli* strain transformed with the plasmid of interest, in combination with *E. coli* HB101/ (pRK2020) containing the 'helper plasmid'.

*P. denitrificans* mutant strains were constructed by gene-replacement techniques as described previously (van Spanning et al., 1991). First, a kanamycin-resistance cassette from pKISS (Boehringer) was inserted in the *SmaI* site of pMR3, a pUC18 derivative carrying a *ctaDI*-containing genomic *Bam*HI fragment (a generous gift of Dr M. Raitio, Fig. 1A). From the resulting construct (pMR3.Km) a *Pvu*II fragment was ligated in the *SmaI* site of pGRPd1 (pRT2521), which was transformed to *E. coli* S17.1 and

conjugated to *P. denitrificans* Pd1222 to give Pd2521. Exconjugants were selected for kanamycin and rifampicin resistance and streptomycin sensitivity; genomic Southern blots were used to confirm the mutations (not shown).

From the *ctaDI* gene a *Sa*II-*Sma*I fragment was deleted (Fig. 1A); sticky ends were filled in using Klenow DNA polymerase fragment and religated (pMR3 $\Delta$ ). The in-frame deleted locus was cloned as a *Pvu*II fragment in the *Sma*I site of the suicide vector pRVS1 (pRT2541), which was transformed to *E. coli* S17.1 and conjugated to the insertion mutant Pd2521. *Paracoccus* colonies were initially selected by rifampicin-, kanamycin- and streptomycin resistance. Subsequent restreaking revealed two distinct phenotypes: (i) integrants turned blue on plates supplemented with Xgal (pRVS1 carries the  $\beta$ -galactosidase gene) and are streptomycin- and kanamycin resistant, and (ii) double recombinant colonies stayed

white and had lost streptomycin- and kanamycin resistance. The deletions in the latter clones (Pd2541) were verified by genomic Southern blotting (not shown). From the *ctaDI* deletion mutant, a *ctaDI/ctaDII* double deletion mutant was derived, using the same methods. As an intermediate an insertion mutant (Pd9218) was generated by conjugation with a pGRP1 derivative (pRT2321), carrying a *Bam*HI-*Sph*I fragment of the *ctaDII* locus with the Km-resistance gene from pUC4K inserted in the *Pst*I site; in the subsequently derived *ctaDII* deletion mutant (Pd9220) the *Hinc*II-*Nco*I fragment was removed after a double recombination event with pRT2341 (Fig. 1A).

The degenerated primers used in the PCR experiment were: #151 (5'-GCGCGGAATTC-CAT/CGGG/CGTG/CATT-CATGATT/CTTT/CTT) and #152 (5'-GCGCGGAATTC-TAG/CACT/CTCG/CGGA/GTGG/CCCA/GAAA/GAACCA). Using genomic DNA from the cytochrome *aa*<sub>3</sub> double mutant and these primers a PCR was performed under conditions essentially like those described before (M. Lübben *et al.*, submitted); in this case the annealing temperature was 37°C. A PCR product of the expected size was digested with *Eco*RI (sites introduced in primers, underlined) and cloned into M13mp18. Using the PCR fragment as a probe, a clone was isolated (pUC.cyoABC) that appeared to contain the complete oxidase locus. This plasmid was digested with *Eco*RI, removing the *Pst*I site from the pUC multiple cloning site, and religated. Subsequently the internal *Pst*I fragment was substituted for the kanamycin-resistance box from pUC4KISS. The *Eco*RI-*Sph*I insert was cloned in pRVS1, transformed to *E. coli* S17.1, and conjugated to the wild-type *Paracoccus* Pd1222 and the *ctaDI/ctaDII* double deletion mutant (Pd9220). Conjugants were selected by kanamycin resistance; colonies of integrants turned blue on plates supplemented with Xgal, but colonies of double recombinants remained white. The insertions in the latter clones were verified by genomic Southern blotting (not shown).

#### Optical spectroscopy

Spectra of whole-cell suspensions were recorded at room temperature on an Aminco/SLM DW2 UV/Vis Spectrophotometer, essentially as described previously (de Gier *et al.*, 1992).

#### Oxygen-consumption analysis

Oxygen consumption by bacterial cell suspensions was analysed polarographically at 30°C in a biological oxygen monitor, Model 53 (Yellow Springs Instrument Co.). Succinate (25 mM), or ascorbate (0.4 mM) and TMPD (0.1 mM) were added as external substrates; electron flow via cytochrome *c* reductase was inhibited by antimycin A (6 µM) or myxothiazol (6 µM). Assays were repeated at least three times and were found to be reproducible within 10%. No auto-oxidation of ascorbate/TMPD was detected when solutions were prepared freshly.

#### Proton-translocation analysis

Measurement of H<sup>+</sup> translocation in intact cells was performed essentially as described by Boogerd *et al.* (1981). Cells from

batch cultures (minimal medium with succinate) were harvested, washed, and resuspended in 1.5 mM glycyglycine buffer (pH 7.0), 100 mM KSCN, 50 mM KCl (approx. 25 mg dry weight cells ml<sup>-1</sup>), incubated at 25°C, and bubbled with nitrogen. In the reaction vessel, 150 µl of cell suspension was mixed with 3.1 ml of buffer. With succinate (25 mM) as electron donor, rotenone (30 µM) was added and incubated for 30 min in order to inhibit endogenous respiration. With ascorbate/TMPD (0.4 mM/0.1 mM) antimycin A (6 µM) was added (final concentrations); in the presence of the quinol oxidase, rotenone (30 µM) was added as well. The anaerobic cell suspension was pulsed with oxygen (O<sub>2</sub>) by injecting 20 µl of air-saturated KCl (9.4 ng atoms of oxygen), or with ferricyanide by injecting 20 µl of an anaerobic solution of 1 mM K<sub>3</sub>Fe(CN)<sub>6</sub>. Traces were calibrated by injecting 20 µl of an anaerobic solution of 1 mM HCl.

#### Haem analysis

The analysis of the haem composition after extraction from membranes was performed as described by Sone and Fujiwara (1991). Haems were extracted from isolated membranes of different *P. denitrificans* strains, separated by reversed-phase HPLC (Deltapak C18 column, Waters) on an acetonitrile gradient in 0.05% trifluoroacetic acid.

#### Acknowledgements

We thank Matti Saraste (Heidelberg) for critical reading of the manuscript. Oliver Richter and Bernd Ludwig (Frankfurt), Arthuro Garcia-Horsman and Robert B. Gennis (Urbana), Mirja Raitio and Mårten Wikström (Helsinki) are gratefully acknowledged for communicating data prior to publication. This research was supported by the Netherlands Foundation for Chemical Research, with financial aid from the Netherlands Organization for the Advancement of Science. M.L. had a long-term EMBO Fellowship, and J.v.d.O. has a Fellowship of the Royal Dutch Academy of Science.

#### References

- Anraku, Y., and Gennis, R.B. (1987) The aerobic respiratory chain of *Escherichia coli*. *Trends Biochem Sci* **12**: 262–266.
- Appleby, C.A., and Poole, R.K. (1991) Characterization of a soluble catalase-peroxidase hemoprotein *b*-590, previously identified as 'cytochrome *a*<sub>1</sub>' from *Bradyrhizobium japonicum* bacteroids. *FEMS Microbiol Lett* **78**: 325–332.
- Ausubel, F.M., Brent, R., Kingston, R.E., Moore, D.D., Smith, J.A., Seidman, J.C., and Struhl, K. (1992) *Current Protocols in Molecular Biology*. New York: John Wiley and Sons, Inc.
- Babcock, G.T., and Wikström, M. (1992) Oxygen activation and the conservation of energy in cell respiration. *Nature* **356**: 301–309.
- Berry, E.A., and Trumpower, B.L. (1985) Isolation of ubiquinol oxidase from *Paracoccus denitrificans* and resolution into cytochrome *bc*<sub>1</sub> and cytochrome *c-aa*<sub>3</sub> complexes. *J Biol Chem* **260**: 2458–2467.
- Boogerd, F.C., van Verseveld, H.W., and Stouthamer, A.H. (1981) Respiration driven proton translocation with nitrate

- and nitrous oxide in *Paracoccus denitrificans*. *Biochim Biophys Acta* **638**: 181–191.
- Bosma, G. (1989) Synthesis of electron transfer components in *Paracoccus denitrificans*. Ph.D. Thesis, Vrije Universiteit, Amsterdam, The Netherlands.
- Bosma, G., Braster, M., Stouthamer, A.H., and van Verseveld, H.W. (1987a) Isolation and characterization of ubiquinol oxidase complexes from *Paracoccus denitrificans* cells cultured under various limiting conditions in the chemostat. *Eur J Biochem* **165**: 657–663.
- Bosma, G., Braster, M., Stouthamer, A.H., and van Verseveld, H.W. (1987b) Isolation and characterization of soluble c-type cytochromes from *Paracoccus denitrificans* cultured under various limiting conditions in the chemostat. *Eur J Biochem* **165**: 665–670.
- Boyer, H.W., and Roulland-Dussoix, D. (1969) A complementation analysis of the restriction and modification of DNA in *Escherichia coli*. *J Mol Biol* **41**: 459–472.
- Cao, J., Shapleigh, J., Gennis, R.B., Revzin, A., and Ferguson-Miller, S. (1991) The gene encoding cytochrome c oxidase subunit II from *Rhodobacter sphaeroides*; comparison of the deduced amino acid sequence with sequences of corresponding peptides from other species. *Gene* **101**: 133–137.
- Chepuri, V., Lemieux, L.J., Au, D.C.-T., and Gennis, R.B. (1990) The sequence of the *cyo* operon indicates substantial structural similarities between the cytochrome o ubiquinol oxidase of *Escherichia coli* and the *aa*<sub>3</sub>-type family of the cytochrome c oxidases. *J Biol Chem* **265**: 11185–11192.
- Cox, J.C., Ingledew, W.J., Haddock, B.A., and Lawford, H.G. (1978) The variable cytochrome content of *Paracoccus denitrificans* grown aerobically under different conditions. *FEBS Lett* **93**: 261–265.
- Ditta, G., Stanfield, S., Corbin, D., and Helinski, D.R. (1985) Broad host range DNA cloning system for gram-negative bacteria: construction of a gene bank of *Rhizobium meliloti*. *Proc Natl Acad Sci USA* **77**: 7347–7351.
- Garcia-Horsman, J.A., Berry, E., Shapleigh, J.P., Alben, J.O., and Gennis, R.B. (1994) A novel cytochrome c oxidase from *Rhodobacter sphaeroides* that lacks Cu<sub>A</sub>. *Biochemistry* **33**: 3113–3119.
- Gerhus, E.P., Steinrücke, P., and Ludwig, B. (1990) *Paracoccus denitrificans* cytochrome c<sub>1</sub> gene replacement mutants. *J Bacteriol* **172**: 2392–2400.
- Green, N.G., Fang, H., Lin, R.-J., Newton, G., Mather, M., Georgiou, C., and Gennis, R.B. (1988) The nucleotide sequence of the *cyd* locus encoding the two subunits of the cytochrome d terminal oxidase complex of *Escherichia coli*. *J Biol Chem* **263**: 13138–13143.
- de Gier, J.-W.L., van Spanning, R.J.M., Oltmann, L.F., and Stouthamer, A.H. (1992) Oxidation of methylamine by a *Paracoccus denitrificans* mutant impaired in the synthesis of the bc<sub>1</sub> complex and the *aa*<sub>3</sub>-type oxidase. Evidence for the existence of an alternative cytochrome c oxidase in this bacterium. *FEBS Lett* **306**: 23–26.
- Harms, N., and van Spanning, R.J.M. (1991) C<sub>1</sub> metabolism in *Paracoccus denitrificans*: genetics of *Paracoccus denitrificans*. *J Bioenerg Biomembr* **23**: 187–210.
- Harms, N., de Vries, G.E., Maurer, K., Veltkamp, E., and Stouthamer, A.H. (1985) Isolation and characterization of *Paracoccus denitrificans* with defects in the metabolism of one-carbon compounds. *J Bacteriol* **164**: 1064–1070.
- Henry, M.F., and Vignais, P.M. (1979) Induction by cyanide of cytochrome d in the plasma membrane of *Paracoccus denitrificans*. *FEBS Lett* **100**: 41–46.
- Hill, B.C. (1993) The sequence of electron carriers in the reaction of cytochrome c oxidase with oxygen. *J Bioenerg Biomembr* **25**: 115–120.
- Hill, J., Chepuri Goswitz, V., Calhoun, M., Garcia-Horsman, J.A., Lemieux, L., Alben, J.O., and Gennis, R.B. (1992) Demonstration by FTIR that the bo-type quinol oxidase of *Escherichia coli* contains a heme-copper binuclear center similar to that in cytochrome c oxidase and that proper assembly of the binuclear center requires the *cyoE* gene product. *Biochemistry* **31**: 11435–11440.
- Hosler, J.P., Ferguson-Miller, S., Calhoun, M.W., Thomas, J.W., Hill, J., Lemieux, L.J., Ma, J., Georgiou, C., Fetter, J., Shapleigh, J., Tecklenburg, M.M.J., Babcock, G.T., and Gennis, R.B. (1993) Insight into the active-site structure and function of cytochrome oxidase by analysis of site-directed mutants of bacterial cytochrome *aa*<sub>3</sub> and cytochrome bo. *J Bioenerg Biomembr* **25**: 121–136.
- Hüdig, H., Stark, G., and Drews, G. (1987) The regulation of cytochrome c oxidase of *Rhodobacter capsulatus* by light and oxygen. *Arch Microbiol* **149**: 12–18.
- John, P., and Whatley, F.R. (1975) *Paracoccus denitrificans* and the evolutionary origin of the mitochondrion. *Nature* **254**: 495–498.
- Kelly, M., Lappalainen, P., Talbo, G., Haltia, T., van der Oost, J., and Saraste, M. (1993) Two cysteines, two histidines and one methionine are ligands of a binuclear purple copper center. *J Biol Chem* **268**: 16781–16787.
- Krab, K., and Wikström, M. (1978) Proton-translocating cytochrome c oxidase in artificial phospholipid vesicles. *Biochim Biophys Acta* **504**: 200–214.
- Lappalainen, P., Aasa, R., Malmström, B.G., and Saraste, M. (1993) Soluble Cu<sub>A</sub>-binding domain from the *Paracoccus* cytochrome oxidase. *J Biol Chem* **268**: 26416–26421.
- Lübber, M., Kolmerer, B., and Saraste, M. (1992) An archaeobacterial terminal oxidase combines core structures of two mitochondrial respiratory complexes. *EMBO J* **11**: 805–812.
- Ludwig, B. (1992) Terminal oxidases in *Paracoccus denitrificans*. In *European Bioenergetics Conference Reports*. Vol. 7. Wikström, M. (ed.). Amsterdam: Elsevier Science, pp. 195–197.
- Matsushita, K., Ebisuya, H., and Adachi, O. (1992) Homology in the structure and the prosthetic groups between two different terminal oxidases, cytochrome a<sub>1</sub> and cytochrome o, of *Acetobacter aceti*. *J Biol Chem* **267**: 24748–24753.
- Minghetti, K.C., Goswitz, V.C., Gabriel, N.E., Hill, J., Barassi, C.A., Georgiou, C.D., Chan, S.I., and Gennis, R.B. (1992) A modified, large-scale purification of the cytochrome o complex of *Escherichia coli* yields a two heme/one copper terminal oxidase with high specific activity. *Biochemistry* **31**: 6917–6924.
- van der Oost, J., Haltia, T., Raitio, M., and Saraste, M. (1991) Genes coding for cytochrome c oxidase in *Paracoccus denitrificans*. *J Bioenerg Biomembr* **23**: 257–267.
- van der Oost, J., Lappalainen, P., Musacchio, A., Warne, A.,

- Lemieux, L.J., Rumbley, J., Gennis, R.B., Aasa, R., Pascher, T., Malmström, B.G., and Saraste, M. (1992) Restoration of a lost metal-binding site: construction of two different copper sites into a subunit of the *E. coli* quinol oxidase complex. *EMBO J* **11**: 3209–3217.
- Preisig, O., Antamatten, D., and Hennecke, H. (1993) Genes for a microaerobically induced oxidase complex in *Bradyrhizobium japonicum* are essential for a nitrogen-fixing endosymbiosis. *Proc Natl Acad Sci USA* **90**: 3309–3313.
- Puustinen, A., and Wikström, M. (1991) The heme groups of cytochrome *o* from *Escherichia coli*. *Proc Natl Acad Sci USA* **88**: 6122–6126.
- Puustinen, A., Finel, M., Virkki, M., and Wikström, M. (1989) Cytochrome *o* (*bo*) is a proton pump in *Paracoccus denitrificans* and *Escherichia coli*. *FEBS Lett* **249**: 163–167.
- Puustinen, A., Finel, M., Haltia, T., Gennis, R.B., and Wikström, M. (1991) Properties of the two terminal oxidases of *Escherichia coli*. *Biochemistry* **30**: 3936–3942.
- Puustinen, A., Morgan, J.E., Verkhovski, M., Thomas, J.W., Gennis, R.B., and Wikström, M. (1992) The low-spin heme site of cytochrome *o* from *Escherichia coli* is promiscuous with respect to heme type. *Biochemistry* **31**: 10363–10369.
- Raitio, M., and Wikström, M. (1994) An alternative cytochrome oxidase of *Paracoccus denitrificans* functions as a proton pump. *Biochim Biophys Acta*, in press.
- Raitio, M., Jalli, T., and Saraste, M. (1987) Isolation and analysis of the genes for cytochrome *c* oxidase in *Paracoccus denitrificans*. *EMBO J* **6**: 2825–2833.
- Raitio, M., Pispala, J.M., Metso, T., and Saraste, M. (1990) Are there isoenzymes of cytochrome *c* oxidase in *Paracoccus denitrificans*? *FEBS Lett* **206**: 154–156.
- Saiki, K., Mogi, T., and Anraku, Y. (1992) Heme O biosynthesis in *Escherichia coli*: the *cyoE* gene in the cytochrome *bo* operon encodes a protoheme IX farnesyl-transferase. *Biochem Biophys Res Commun* **189**: 1491–1497.
- Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual*, 2nd edn. Cold Spring Harbor, New York, Cold Spring Harbor Laboratory Press.
- Sanger, F., Coulson, R., Barrel, B.G., Smith, J.H., and Roe, B.A. (1980) Cloning in single-stranded bacteriophage as an aid to rapid DNA sequencing. *J Mol Biol* **143**: 161–178.
- Santana, M., Kunst, F., Hullo, M.F., Raparort, G., Danchin, A., and Glaser, P. (1992) Molecular cloning, sequencing and physiological characterization of the *cox* operon from *Bacillus subtilis* encoding the *aa*<sub>3</sub>-600 quinol oxidase. *J Biol Chem* **267**: 10225–10231.
- Saraste, M. (1990) Structural features of cytochrome oxidase. *Q Rev Biophys* **23**: 331–366.
- Saraste, M., Raitio, M., Jalli, T., Chepuri, V., Lemieux, L.J., and Gennis, R.B. (1988) Cytochrome *o* from *E. coli* is structurally related to cytochrome *aa*<sub>3</sub>. *Ann N Y Acad Sci* **550**: 314–324.
- Saraste, M., Holm, L., Lemieux, L.J., Lübben, M., and van der Oost, J. (1991) The happy family of cytochrome oxidases. *Biochem Soc Trans* **19**: 608–612.
- Shapleigh, J.P., and Gennis, R.B. (1992) Cloning, sequencing and deletion from the chromosome of the gene encoding subunit I of the *aa*<sub>3</sub>-type cytochrome oxidase of *Rhodobacter sphaeroides*. *Mol Microbiol* **6**: 635–642.
- Simon, R., Priefer, U., and Pühler, A. (1983) Vector plasmids for *in vivo* and *in vitro* manipulations of Gram-negative bacteria. In *Molecular Genetics of the Bacteria-Plant interaction*. Pühler, A. (ed.). Berlin: Springer Verlag, pp. 98–106.
- Soloz, M., Carafoli, E., and Ludwig, B. (1981) The cytochrome *c* oxidase of *Paracoccus denitrificans* pumps protons in a reconstituted system. *J Biol Chem* **257**: 1579–1582.
- Sone, N., and Fujiwara, Y. (1991) Heme O can replace heme A in the active site of cytochrome *c* oxidase thermophilic bacterium PS3. *FEBS Lett* **288**: 154–158.
- Van Spanning, R.J.M., Wansell, C.W., Harms, N., Oltmann, L.F., and Stouthamer, A.H. (1990) Mutagenesis of the gene encoding cytochrome *c*<sub>550</sub> of *Paracoccus denitrificans* and analysis of the resultant physiological effects. *J Bacteriol* **172**: 986–996.
- Van Spanning, R.J.M., Wansell, C.W., Reijnders, W.N.M., Oltmann, L.F., and Stouthamer, A.H. (1991) A method for introduction of unmarked mutations in the genome of *Paracoccus denitrificans*: construction of strains with multiple mutations in the genes encoding periplasmic cytochromes *c*<sub>550</sub>, *c*<sub>551i</sub>, and *c*<sub>553i</sub>. *J Bacteriol* **173**: 6962–6970.
- Steinrücke, P., Steffens, G.C., Pankus, G., Buse, G., and Ludwig, B. (1987) Subunit II of cytochrome *c* oxidase from *Paracoccus denitrificans* — DNA sequence, gene expression and the protein. *Eur J Biochem* **167**: 431–439.
- Stouthamer, A.H. (1991) Metabolic regulation including anaerobic metabolism in *Paracoccus denitrificans*. *J Bioenerg Biomembr* **23**: 163–185.
- Timkovich, R., Dhesi, R., Martinkus, K.J., Robinson, N.K., and Rea, T.M. (1982) Isolation of *Paracoccus denitrificans* cytochrome *cd*<sub>1</sub>: comparative kinetics with other nitrite reductases. *Arch Biochem Biophys* **215**: 47–58.
- van Verseveld, H.W., Krab, K., and Stouthamer, A.H. (1981) Proton pump coupled to cytochrome *c* oxidase in *Paracoccus denitrificans*. *Biochim Biophys Acta* **635**: 525–534.
- van Verseveld, H.W., Braster, M., Boogerd, F.C., Chance, B., and Stouthamer, A.H. (1983) Energetic aspects of growth of *Paracoccus denitrificans*: oxygen-limitation and shift from anaerobic nitrate-limitation to aerobic succinate-limitation. Evidence for a new alternative oxidase, cytochrome *a*<sub>1</sub>. *Arch Microbiol* **135**: 229–236.
- de Vries, G.E., Harms, N., Hoogendijk, J., and Stouthamer, A.H. (1989) Isolation and characterization of *Paracoccus denitrificans* mutants with increased conjugation frequencies and pleiotropic loss of a (nGATCn) DNA-modifying property. *Arch Microbiol* **152**: 52–57.
- Von Wachenfeldt, C., de Vries, S., and van der Oost, J. (1994) The Cu<sub>A</sub> site of the *caa*<sub>3</sub>-type oxidase of *Bacillus subtilis* is a mixed-valence binuclear copper centre. *FEBS Lett* **340**: 109–113.
- Wikström, M. (1977) Proton pump coupled to cytochrome *c* oxidase in mitochondria. *Nature* **266**: 271–273.

- Wikström, M., Krab, K., and Saraste, M. (1981) *Cytochrome Oxidase: A Synthesis*. London: Academic Press.
- Willison, J.C., Haddock, B.A., and Boxer, D.H. (1981) A mutant of *Paracoccus denitrificans* deficient in cytochrome  $a+a_3$ . *FEMS Microbiol Lett* **10**: 249–255.
- Wu, W., Chang, C.K., Varotsis, C., Babcock, G.T., Puustinen, A., and Wikström, M. (1992) Structure of the heme O prosthetic group from the terminal quinol oxidase from *Escherichia coli*. *J Am Chem Soc* **114**: 1182–1187.
- Yanisch-Perron, C., Vieira, J., and Messing, J. (1985) Improved M13 phage cloning vectors and host strains: nucleotide sequences of M13mp18 and pUC19 vectors. *Gene* **33**: 103–119.

This document is a scanned copy of a printed document. No warranty is given about the accuracy of the copy. Users should refer to the original published version of the material.