

University of Groningen

Wintering on different continents

Kentie, Rosemarie

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2015

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Kentie, R. (2015). *Wintering on different continents: Effects on breeding parameters of Black-tailed Godwits.*

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Wintering on different continents


Effects on breeding parameters of Black-tailed Godwits

Rosemarie Kentie¹, Rocío Márquez-Ferrando², Jos Hooijmeijer¹, Jordi Figuerola², Theunis Piersma^{1,3}


¹ Conservation Ecology Group, Groningen Institute for Evolutionary Life Sciences (GELIFES), University of Groningen, PO Box 11103, 9700 CC Groningen, The Netherlands, ² Department of Wetland Ecology, Estación Biológica de Doñana-CSIC, Avda. Américo Vespucio s/n, Sevilla 41092, Spain, ³ NIOZ Department of Marine Ecology, Royal Netherlands Institute for Sea Research, PO Box 59, 1790 AB Den Burg, Texel, The Netherlands

Hypothesis: Godwits that winter in Doñana benefit from shorter migration distances, which will be reflected in their breeding parameters


First resighting date in breeding area




Laying date



Egg volume (females)



Nest success



Black-tailed Godwits wintering in Doñana seem to have similar breeding parameters compared to godwits wintering 3000 km further south, in West Africa.

- No difference in arrival date in their breeding area
- Laying dates of Doñana winterers were 3 days later than those wintering in W-Africa
- No difference in egg volume
- No difference in nest success

Methods:

Since 2004 we have colour-ringed more than 9000 Black-tailed Godwits (*Limosa limosa limosa*). We have more than 75.000 resightings throughout the flyway. If an individual was seen at least once during the non-breeding period in West Africa, we classified it as a West Africa winterer. If an individual was seen in Doñana in Sept-Nov, we classified it to winter there (see Marquez *et al.* 2014). We assumed that they were faithful to their wintering location, which unpublished data on satellite tags and geolocators confirms. Breeding parameters were collected in 8500 ha field site in The Netherlands (see Groen *et al.* 2012 and Kentie *et al.* 2015 for more details).

1. Márquez-Ferrando, R., Figuerola, J., Hooijmeijer, J.C.E.W. & Piersma, T. (2014) Recently created man-made habitats in Doñana provide alternative wintering space for the threatened Continental European black-tailed godwit population. *Biological Conservation*, **171**, 127-135.
2. Groen, N.M., Kentie, R., de Goeij, P., Verheijen, B., Hooijmeijer, J.C.E.W. & Piersma, T. (2012) A modern landscape ecology of Black-tailed Godwits: habitat selection in southwest Friesland, The Netherlands. *Ardea*, **100**, 19-28.
3. Kentie, R., Both, C., Hooijmeijer, J.C.E.W. & Piersma, T. (2015) Management of modern agricultural landscapes increases nest predation rates in Black-tailed Godwits (*Limosa limosa limosa*). *Ibis*, **157**, 614-625.



The fieldwork was funded by the FP7 Regpot project "ECOGENES: Adapting to Global Change in the Mediterranean hotspot: from genes to ecosystems" 2011-2013, by ExpeER Ecosystem Research 2013-2014, 'Kenniskringing Weidevogels' of the former Dutch Ministry of Agriculture, Nature Management and Food Safety, and Provincie Fryslân. We thank all the volunteer bird observers and field assistants for helping collecting so much data, It Fryske Gea, Staatsbosbeheer and private landowners for allowing us on their land.

