

University of Groningen

Outbreak of NDM-1-Producing *Klebsiella pneumoniae* in a Dutch Hospital, with Interspecies Transfer of the Resistance Plasmid and Unexpected Occurrence in Unrelated Health Care Centers

Bosch, Thijs; Lutgens, Suzanne P M; Hermans, Mirjam H A; Wever, Peter C; Schneeberger, Peter M.; Renders, Nicole H M; Leenders, Alexander C. A. P.; Kluytmans, Jan A. J. W.; Schoffelen, Annelot F; Notermans, Daan

Published in:
Journal of Clinical Microbiology

DOI:
[10.1128/JCM.00535-17](https://doi.org/10.1128/JCM.00535-17)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2017

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Bosch, T., Lutgens, S. P. M., Hermans, M. H. A., Wever, P. C., Schneeberger, P. M., Renders, N. H. M., Leenders, A. C. A. P., Kluytmans, J. A. J. W., Schoffelen, A. F., Notermans, D., Witteveen, S., Bathoorn, E., & Schouls, L. M. (2017). Outbreak of NDM-1-Producing *Klebsiella pneumoniae* in a Dutch Hospital, with Interspecies Transfer of the Resistance Plasmid and Unexpected Occurrence in Unrelated Health Care Centers. *Journal of Clinical Microbiology*, 55(8), 2380-2390. <https://doi.org/10.1128/JCM.00535-17>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Outbreak of NDM-1-Producing *Klebsiella pneumoniae* in a Dutch Hospital, with Interspecies Transfer of the Resistance Plasmid and Unexpected Occurrence in Unrelated Health Care Centers

Thijs Bosch,^a Suzanne P. M. Lutgens,^b Mirjam H. A. Hermans,^b Peter C. Wever,^b Peter M. Schneeberger,^b Nicole H. M. Renders,^b Alexander C. A. P. Leenders,^b Jan A. J. W. Kluytmans,^{c,d} Annelot Schoffelen,^a Daan Notermans,^a Sandra Witteveen,^a Erik Bathoorn,^e Leo M. Schouls^a

Centre for Infectious Disease Control, National Institute for Public Health and the Environment, Bilthoven, the Netherlands^a; Department of Medical Microbiology and Infection Control, Jeroen Bosch Hospital, 's-Hertogenbosch, the Netherlands^b; Laboratory for Microbiology and Infection Control, Amphia Hospital Breda, Breda, the Netherlands^c; Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht, the Netherlands^d; Department of Medical Microbiology, University of Groningen, University Medical Center Groningen, Groningen, the Netherlands^e

ABSTRACT In the Netherlands, the number of cases of infection with New Delhi metallo-beta-lactamase (NDM)-positive *Enterobacteriaceae* is low. Here, we report an outbreak of NDM-1-producing *Klebsiella pneumoniae* infection in a Dutch hospital with interspecies transfer of the resistance plasmid and unexpected occurrence in other unrelated health care centers (HCCs). Next-generation sequencing was performed on 250 carbapenemase-producing *Enterobacteriaceae* isolates, including 42 NDM-positive isolates obtained from 29 persons at the outbreak site. Most outbreak isolates were *K. pneumoniae* ($n = 26$) and *Escherichia coli* ($n = 11$), but 5 isolates comprising three other *Enterobacteriaceae* species were also cultured. The 26 *K. pneumoniae* isolates had sequence type 873 (ST873), as did 7 unrelated *K. pneumoniae* isolates originating from five geographically dispersed HCCs. The 33 ST873 isolates that clustered closely together using whole-genome multilocus sequence typing (wgMLST) carried the same plasmids and had limited differences in the resistome. The 11 *E. coli* outbreak isolates showed great variety in STs, did not cluster using wgMLST, and showed considerable diversity in resistome and plasmid profiles. The *bla*_{NDM-1} gene-carrying plasmid present in the ST873 *K. pneumoniae* isolates was found in all the other *Enterobacteriaceae* species cultured at the outbreak location and in a single *E. coli* isolate from another HCC. We describe a hospital outbreak with an NDM-1-producing *K. pneumoniae* strain from an unknown source that was also found in patients from five other Dutch HCCs in the same time frame without an epidemiological link. Interspecies transfer of the resistance plasmid was observed in other *Enterobacteriaceae* species isolated at the outbreak location and in another HCC.

KEYWORDS *Klebsiella*, NDM, carbapenems, outbreak

In 2008, a new carbapenemase, designated New Delhi metallo-beta-lactamase 1 (NDM-1), was detected in a *Klebsiella pneumoniae* isolate originating from a Swedish patient who had been hospitalized in India (1). NDM-1 belongs to the Ambler class B beta-lactamases and hydrolyzes virtually all beta-lactam antibiotics, including carbapenems (2). Besides NDM-1, 13 other variants of this metallo-beta-lactamase (NDM-2 to NDM-14) have currently been described, and most of the variants contain either single-

Received 30 March 2017 Returned for modification 28 April 2017 Accepted 10 May 2017

Accepted manuscript posted online 17 May 2017

Citation Bosch T, Lutgens SPM, Hermans MHA, Wever PC, Schneeberger PM, Renders NHM, Leenders ACAP, Kluytmans JAJW, Schoffelen A, Notermans D, Witteveen S, Bathoorn E, Schouls LM. 2017. Outbreak of NDM-1-producing *Klebsiella pneumoniae* in a Dutch hospital, with interspecies transfer of the resistance plasmid and unexpected occurrence in unrelated health care centers. *J Clin Microbiol* 55:2380–2390. <https://doi.org/10.1128/JCM.00535-17>.

Editor Betty A. Forbes, Virginia Commonwealth University Medical Center

Copyright © 2017 American Society for Microbiology. All Rights Reserved.

Address correspondence to Thijs Bosch, thijs.bosch@rivm.nl.

or two-amino-acid substitutions compared to NDM-1 (3). The NDM gene has been isolated from a wide range of bacterial hosts, including *Vibrio cholerae*, and has been located on a great variety of plasmids (4, 5).

After the first report, NDM-positive bacteria were shown to be widely present among hospitalized patients in many countries of the Indian subcontinent and in the United Kingdom in 2008 and 2009 (6). At present, NDM-positive bacteria have been isolated from many countries worldwide, but many isolates have travel-related epidemiological links, with or without hospitalization, to the Indian subcontinent (7). However, some reports showed the presence of NDM-positive isolates in patients with no history of foreign travel, suggesting that the bacteria or the NDM gene had been acquired through local transmission (8, 9).

In the Netherlands, until now, the number of NDM-positive cases has been low, with only sporadic cases reported (10). The first NDM-positive bacteria were detected in 2010 in two Dutch patients who had recently traveled to India (11). More recently, in 2015, an NDM-5-producing *K. pneumoniae* isolate was isolated from a Dutch patient without a history of travel abroad (12).

Here, we describe a large outbreak of NDM-1-producing *Enterobacteriaceae* in the Netherlands that occurred in a large teaching hospital. Unexpectedly, the same strain was also found in other health care centers with no detectable epidemiological link.

RESULTS

Transmission of NDM-1-producing *K. pneumoniae*. The *K. pneumoniae* isolates collected from 26 cases at the outbreak location (Table 1) were NDM positive, and analysis of the next-generation sequencing (NGS) data showed that they all harbored the *bla*_{NDM-1} gene. Three NDM-positive *K. pneumoniae* isolates originating from the outbreak location were not epidemiologically linked to the outbreak but also carried the *bla*_{NDM-1} gene. The top three carbapenemase genes carried by the 133 *K. pneumoniae* isolates from the Dutch carbapenemase-producing *Enterobacteriaceae* (CPE) surveillance, which were used as context in this study, were OXA-48-like ($n = 75$), KPC ($n = 27$), and NDM ($n = 26$) genes (Table 2). Twenty-three of the NDM-positive surveillance isolates carried the NDM-1 variant. Two other isolates carried the *bla*_{NDM-5} gene, and the remaining isolate harbored the *bla*_{NDM-7} gene.

The 26 presumed outbreak cases had meropenem MICs ranging from 0.75 to 12 $\mu\text{g/ml}$, but the majority of the isolates ($n = 20$) had MICs below 2 $\mu\text{g/ml}$. In contrast, 21 of the 26 NDM-positive context isolates from the CPE surveillance had MICs greater than 6 $\mu\text{g/ml}$, and only 2 isolates had MICs below 2 $\mu\text{g/ml}$.

Based on the NGS data on the *K. pneumoniae* isolates, all 26 presumed outbreak cases had sequence type 873 (ST873). Seven of the 133 *K. pneumoniae* context isolates from the Dutch CPE surveillance also were ST873, and these isolates originated from five other health care centers (HCCs). The HCCs were geographically dispersed throughout the Netherlands, and the ST873 isolates were submitted between December 2015 and October 2016. Six of the seven isolates were submitted after the outbreak period, but a single isolate was submitted during the outbreak period (Fig. 1). The other 126 isolates and the 3 unrelated isolates from the outbreak site submitted between 2015 and 2016 had various other STs.

The whole-genome multilocus sequence typing (wgMLST) was based on 4,978 genes, and the resulting minimum spanning tree (MST) showed that the NDM-positive isolates were scattered over the tree, with the exception of a group of 33 ST873 isolates that clustered closely together with a maximum of five genes difference between neighboring isolates (Fig. 2). The distances between this group and the nearest neighbors ranged from 2,926 to 3,666 genes. The distance from the other three epidemiologically unrelated *K. pneumoniae* isolates that were isolated from the outbreak location ranged from 3,781 to 3,837 genes, indicating that they did not belong to the outbreak.

Resistome analysis of the 33 ST873 NDM-1 *K. pneumoniae* isolates that clustered closely together in the wgMLST analysis showed that 25 isolates had identical resis-

TABLE 1 Species isolated from the outbreak location during the NDM-1 outbreak

Species isolated during the outbreak	No. of cases
<i>K. pneumoniae</i>	17
<i>E. coli</i>	2
<i>C. freundii</i>	1
<i>K. pneumoniae</i> and <i>E. coli</i>	7
<i>K. pneumoniae</i> , <i>E. coli</i> , <i>K. oxytoca</i> , and <i>C. freundii</i>	1
<i>K. pneumoniae</i> , <i>E. coli</i> , <i>K. oxytoca</i> , and <i>P. mirabilis</i>	1
Total no. of cases	29

tomes (Table 3). Of these, 20 were obtained from the outbreak location and 5 isolates originated from four other HCCs. The remaining 8 isolates differed in their resistance gene profiles, each lacking one or more resistance genes compared to the group of 25 isolates.

Plasmid analyses showed that the cluster of 33 ST873 *K. pneumoniae* isolates all had identical plasmid compositions. The *bla*_{NDM-1} gene was located on a variant of the pNDM-US plasmid (CP006661), a 140-kb plasmid belonging to the IncA/C2 group (13). Compared to the pNDM-US reference genome, the plasmid in the 33 ST873 *K. pneumoniae* isolates lacked a 740-bp segment that carried the *aac*(6')-1B-cr (fluoroquinolone and aminoglycoside resistance) gene. This finding was corroborated by electroporation experiments, since the transformants originating from two ST873 *K. pneumoniae* isolates, one from the outbreak location and one from another HCC, did not carry the gene. Besides *bla*_{NDM-1}, the plasmid also harbored the beta-lactam resistance gene *bla*_{CMY-6} and the resistance genes *rmtC* (aminoglycoside resistance) and *sul-1* (sulfonamide resistance). In addition to the variant of the pNDM-US plasmid, all 33 ST873 *K. pneumoniae* isolates also carried two plasmids belonging to the IncFIB(K) and IncFII(K) groups. The three unrelated *K. pneumoniae* isolates from the outbreak location and the NDM-positive *K. pneumoniae* context isolates obtained from the national CPE surveillance did not carry the plasmid found in the ST873 isolates.

TABLE 2 Isolates collected from January 2015 to December 2016, used as context in the study (one isolate per case per year)

Species	Carbapenemase allele	No. of cases
<i>K. pneumoniae</i>	OXA-48	69
	OXA-232	4
	OXA-181	1
	OXA-162	1
	KPC-2	19
	KPC-3	8
	NDM-1	23
	NDM-5	2
	NDM-7	1
	NDM-1, OXA-48	3
	NDM-1, OXA-232	1
	VIM-1	1
	Total	133
	<i>E. coli</i>	OXA-48
OXA-181		7
OXA-244		4
KPC-2		1
NDM-5		15
NDM-1		7
NDM-7		2
NDM-6		1
NDM-5, OXA-181		1
VIM-2		1
Total		70




FIG 1 Geographic origins of the 26 *K. pneumoniae* isolates from outbreak cases and locations of the other five health care centers in the Netherlands. The outbreak cases were plotted based on the postal codes of the NDM carriers, and the sizes of the circles indicate the numbers of carriers with the same postal code.

Transmission of the pNDM-US variant to *E. coli*. During the outbreak, *Escherichia coli* isolates producing NDM-1 carbapenemase were cultured from 11 patients at the outbreak location. Nine of these patients also carried an ST873 *K. pneumoniae* isolate. An epidemiologically unrelated *E. coli* isolate submitted from the outbreak location in June 2015, 5 months prior to the outbreak, also produced NDM-1, whereas the other unrelated isolate carried a *bla*_{NDM-5} gene. PCR of the 70 *E. coli* isolates originating from the national CPE surveillance that served as context showed that they carried the *bla*_{OXA-48} (*n* = 42), *bla*_{NDM} (*n* = 26), *bla*_{KPC} (*n* = 1), and *bla*_{VIM} (*n* = 1) carbapenemase genes (Table 2). Only 7 of the 26 *E. coli* NDM-positive context isolates produced the NDM-1 variant, whereas 15 other isolates produced NDM-5 and the remaining isolates produced NDM-7 (*n* = 2) and NDM-6 (*n* = 1). Similar to the *K. pneumoniae* outbreak isolates, the meropenem MICs of the 11 *E. coli* isolates ranged from 0.75 to 6 μg/ml and were considerably lower than those of the *E. coli* context isolates.

The STs, inferred from the NGS data, revealed that each of the 11 *E. coli* isolates from the outbreak location had a unique ST. The two unrelated *E. coli* isolates submitted from the outbreak location and the 70 *E. coli* isolates obtained from the national CPE surveillance also showed great diversity in STs. The wgMLST analysis, based on 4,503 genes, showed that the 11 *E. coli* outbreak isolates were scattered over the tree and that no clustering occurred (Fig. 3). The average distance from one of the outbreak *E. coli* isolates to the closest neighboring *E. coli* isolate was 1,091 genes, with a range between




FIG 2 Minimum spanning tree of the 162 *K. pneumoniae* isolates. The tree was based on 4,978 genes of the *K. pneumoniae* whole-genome MLST scheme. Clustering was done using a categorical coefficient, and the sizes of the circles indicate the numbers of isolates with identical wgMLST profiles. The lengths of the lines between isolates represent the numbers of different alleles.

304 and 2,319 genes. The unrelated bla_{NDM-1} -positive *E. coli* isolate from the outbreak location did not cluster with any other isolate, and the distance from the closest neighbor was 3,016 genes. The other unrelated bla_{NDM-5} -positive *E. coli* isolate clustered closely with another *E. coli* isolate carrying $bla_{OXA-181}$, with a distance of 87 alleles.

Compared to the *K. pneumoniae* isolates, the *E. coli* isolates from the outbreak location showed more diversity in the resistome (Table 3). Plasmid analysis showed that all 11 *E. coli* outbreak isolates and the NDM-1-positive *E. coli* isolate, submitted in June 2015, harbored the same plasmid as the *K. pneumoniae* isolates, namely, the Inc A/C2 group plasmid variant of pNDM-US. A transformant of an *E. coli* isolate from the outbreak location confirmed that the same plasmid was present in both species. Apart from this plasmid, the *E. coli* isolates showed considerable diversity in their plasmid profiles, and the compositions were different in all the isolates (data not shown).

Four of the seven NDM-1-positive *E. coli* isolates from the national CPE surveillance also carried a plasmid belonging to the Inc A/C2 group. A single isolate carried the same plasmid variant of pNDM-US as the *K. pneumoniae* and *E. coli* outbreak isolates. This isolate originated from a patient from another HCC, and the patient was one of the

TABLE 3 Resistance genes among the outbreak isolates identified by ResFinder software

Phenotype	Resistance gene	Presence ^a											
		<i>K. pneumoniae</i> (n = 33)											
		V1 (n = 25)	V2 (n = 2)	V3 (n = 2)	V4 (n = 1)	V5 (n = 1)	V6 (n = 1)	V7 (n = 1)	<i>E. coli</i> (n = 11)				
									V1 (n = 5)	V2 (n = 3)	V3 (n = 1)	V4 (n = 1)	V5 (n = 1)
Beta-lactam resistance	<i>bla</i> _{NDM-1}	+	+	+	+	+	+	+	+	+	+	+	+
	<i>bla</i> _{CMV-6}	+	+	+	+	+	+	+	+	+	+	+	+
	<i>bla</i> _{OXA-1}	+	+	+	+	+	+	+	+	+	+	+	+
	<i>bla</i> _{SHV-27}	+	+	+	+	+	+	+	+	+	+	+	+
	<i>bla</i> _{CTX-M-15}	+	+	+	+	+	+	+	+	+	+	+	+
	<i>bla</i> _{TEM-1B}	+							+	+	+	+	+
Aminoglycoside resistance	<i>aac</i> (3)-II/d												+
	<i>aadA1</i>												+
	<i>aadA2</i>												+
	<i>aadA5</i>												+
Fluoroquinolone and aminoglycoside resistance	<i>aac</i> (6')IIb-cr	+	+	+	+	+	+	+	+	+	+	+	+
Aminoglycoside resistance	<i>aph</i> (3')-Ia	+	+	+	+	+	+	+	+	+	+	+	+
	<i>rmtC</i>	+	+	+	+	+	+	+	+	+	+	+	+
	<i>strA</i>	+	+	+	+	+	+	+	+	+	+	+	+
	<i>strB</i>	+	+	+	+	+	+	+	+	+	+	+	+
	<i>aph</i> (3')-VIa	+	+	+	+	+	+	+	+	+	+	+	+
Rifampin resistance	<i>arr-2</i>	+	+	+	+	+	+	+	+	+	+	+	+
Phenicol resistance	<i>catB3</i>	+	+	+	+	+	+	+	+	+	+	+	+
	<i>cmiA1</i>	+	+	+	+	+	+	+	+	+	+	+	+
Trimethoprim resistance	<i>dhfrA12</i>	+	+	+	+	+	+	+	+	+	+	+	+
	<i>dhfrA14</i>	+	+	+	+	+	+	+	+	+	+	+	+
	<i>dhfrA17</i>	+	+	+	+	+	+	+	+	+	+	+	+
Fosfomycin resistance	<i>fosA</i>	+	+	+	+	+	+	+	+	+	+	+	+
	<i>mph</i> (A)	+	+	+	+	+	+	+	+	+	+	+	+
Macrolide resistance													
Quinolone resistance	<i>oqx4</i>	+	+	+	+	+	+	+	+	+	+	+	+
	<i>oqx8</i>	+	+	+	+	+	+	+	+	+	+	+	+
	<i>qnrB66</i>	+	+	+	+	+	+	+	+	+	+	+	+
	<i>qnrS1</i>	+	+	+	+	+	+	+	+	+	+	+	+
Sulfonamide resistance	<i>sul1</i>	+	+	+	+	+	+	+	+	+	+	+	+
	<i>sul2</i>	+	+	+	+	+	+	+	+	+	+	+	+
	<i>sul3</i>	+	+	+	+	+	+	+	+	+	+	+	+
Tetracycline resistance	<i>tet</i> (A)	+	+	+	+	+	+	+	+	+	+	+	+

^a+, present.




FIG 3 Minimum spanning tree based on whole-genome MLST of 83 *E. coli* isolates. The tree was based on 4,503 genes of the *E. coli* scheme, and clustering was done using a categorical coefficient. Each isolate in the tree is displayed as a circle. The lines between the isolates denote the distance in numbers of alleles.

seven persons who also carried an ST873 *K. pneumoniae* isolate (Fig. 1 and 2). The plasmid compositions of the other three NDM-1-positive *E. coli* surveillance isolates differed from the plasmid present in the *K. pneumoniae* and *E. coli* outbreak isolates, indicating that another plasmid from the Inc A/C2 group carried the NDM-1 gene.

Multiple NDM-1-producing species in the same patient. From 9 of the 31 outbreak patients, two to four different NDM-1-positive *Enterobacteriaceae* species were isolated during the outbreak period. Seven patients carried *K. pneumoniae* and *E. coli*, one also carried *Citrobacter freundii* and *Klebsiella oxytoca*, and one patient carried *Proteus mirabilis* and *K. oxytoca*, in addition to *K. pneumoniae* and *E. coli* (Table 1 and Fig. 4). In all nine cases, the *K. pneumoniae* isolate was cultured before or on the same day as the other species. However, from a patient not related to the outbreak whose *E. coli* isolate was submitted in June 2015, prior to the outbreak, a ST873 *K. pneumoniae* isolate was also cultured after the outbreak in March 2016. Similar to *K. pneumoniae* and *E. coli*, all the other species carried the variant of plasmid pNDM-US lacking the *aac(6′)-1B-cr* gene. An NDM-1-positive *C. freundii* isolate was




FIG 4 Timeline of the outbreak for 10 patients who carried multiple NDM-1-producing *Enterobacteriaceae* species. The colors indicate isolates originating from the same patient. The dates below the isolates represent the sampling dates.

cultured from a single patient, and this isolate had the same ST and resistome as the *C. freundii* isolate cultured from the patient who also carried NDM-1-positive *K. pneumoniae* and *E. coli* isolates.

DISCUSSION

To date, only a few NDM-positive isolates have been described in the Netherlands, indicating a limited presence of strains producing this metallo-beta-lactamase (11, 12). Here, we report a large outbreak in a Dutch hospital comprising 26 patients colonized with NDM-1-producing *K. pneumoniae*. In addition, 16 NDM-1-positive isolates comprising five other *Enterobacteriaceae* species harboring the same plasmid were also cultured at the outbreak location.

All *K. pneumoniae* outbreak isolates had ST873, were identical or very closely related in the wgMLST, and contained a variant of the IncA/C2 plasmid pNDM-US (CP006661) that harbors the *bla_{NDM-1}* gene. This indicates that the outbreak was caused by transmission of the same NDM-1-producing *K. pneumoniae* strain. Three epidemiologically unrelated *bla_{NDM-1}*-positive *K. pneumoniae* isolates collected at the outbreak location were also genetically unrelated according to wgMLST, had a completely different resistome, and did not carry the variant of plasmid pNDM-US. Consequently, they did not belong to the outbreak.

The NDM-1-positive *K. pneumoniae* outbreak strain was also isolated from seven other persons originating from five other HCCs geographically dispersed throughout the Netherlands. These seven isolates were submitted for the Dutch CPE surveillance between December 2015 and October 2016. Despite extensive epidemiological investigations, none of the seven cases could be connected to the cases from the hospital where the outbreak occurred. Considering their very limited genetic variation, these strains most likely come from a common source. This conclusion was supported when the observed genetic variation was compared to a recently defined cutoff for relatedness between *K. pneumoniae* isolates (14). Although no evidence was found for this outbreak, NDM-1-producing *K. pneumoniae* strains could have been introduced via food supplied to various HCCs, as exemplified by the recent study from Egypt, where

12 of 100 tested carcasses were contaminated with CPE isolates, 11 of which were *K. pneumoniae* (15). The contamination of meat with CPE has not been described in the Netherlands, but several reports have shown high prevalence of extended-spectrum beta-lactamase (ESBL)-producing *Enterobacteriaceae* (16, 17). Furthermore, several of the health care centers, including the hospital with the outbreak, received poultry from the same supplier. Unfortunately, an attempt to reconstruct the diets of the patients involved in the outbreak was unsuccessful at the time that food was considered a possible source.

Another scenario could be a prolonged, unnoticed circulation of this *K. pneumoniae* strain or of other NDM-1 strain-carrying *Enterobacteriaceae* in the Dutch population or in patients or health care workers at the outbreak location. This scenario is supported by the fact that an *E. coli* strain carrying the same plasmid as the outbreak isolates was isolated at the outbreak location in June 2015, 5 months prior to the outbreak. Moreover, an NDM-1-positive ST873 *K. pneumoniae* isolate was cultured from the same patient after the outbreak in March 2016. However, given the short outbreak period and the fact that no further transmission was noticed after the outbreak, the latter suggestion seems less likely.

The *bla*_{NDM-1} gene was also present in 11 *E. coli* isolates collected at the outbreak location during the outbreak. The wgMLST and resistome analyses showed that the isolates were genetically unrelated. Like the *K. pneumoniae* outbreak isolates, all 11 *E. coli* isolates carried the same variant of plasmid pNDM-US, indicating multiple occurrences of horizontal gene transfer rather than transmission of the same *E. coli* strain. This was further supported by the fact that the transfer of the plasmid was also observed in five other isolates, comprising three other *Enterobacteriaceae* species, at the outbreak location. In addition, the transfer of the plasmid was also seen in another HCC where an ST873 *K. pneumoniae* isolate and an *E. coli* isolate from the same patient harbored the same plasmid as the outbreak isolates.

Our study has a number of limitations. First, our study has fully focused on bacterial isolates that harbored carbapenemase genes. The genetic background and potential spread of noncarbapenemase isolates from the different HCCs is therefore unknown. Second, our analysis comprised only isolates from the Dutch CPE surveillance. NGS data from publicly available databases containing international carbapenemase-producing *K. pneumoniae* isolates were not assessed and used. However, we did notice that ST873 *K. pneumoniae* isolates were occasionally found in recent studies from Switzerland and China, but these isolates did not produce carbapenemase (18, 19).

In conclusion, we report a hospital outbreak with an NDM-1-producing *K. pneumoniae* strain from an unknown source that was also found in patients from five other HCCs in the Netherlands in the same time frame. The plasmid carrying the *bla*_{NDM-1} gene was also found in other *Enterobacteriaceae* species isolated at the outbreak location and in another HCC. The fact that the outbreak strain was also found in other HCCs without a clear epidemiological link shows that analysis of the NGS data on CPE isolates collected during national CPE surveillance is important to reveal the spread of carbapenemase-producing pathogens in the Netherlands.

MATERIALS AND METHODS

Hospital setting and outbreak description. The outbreak was first noted during screening for a possible ESBL transmission in the surgical ward of a 683-bed tertiary teaching hospital in the southern Netherlands. On 23 November 2015, a carbapenemase-producing *bla*_{NDM-1}-positive, ESBL-positive *K. pneumoniae* strain was isolated during this screening. At the time of identification, the patient had already been discharged. Shortly after, ESBL screening of cultures from long-term-admitted surgical patients revealed two additional patients with NDM-producing *K. pneumoniae*. Contact tracing and weekly screening rounds identified additional NDM carriers. There were two wards with a single NDM carrier and seven wards with two or more NDM carriers. All the patients admitted to one of these wards from the 1 October 2015 on were defined as at risk for carrying NDM. The outbreak period was defined as the date 2 weeks prior (1 October) to the admission of the initial patient until the date when all suspected NDM carriers were identified and put into contact isolation (30 December). Weekly screening rounds were continued during the first 4 weeks of January 2016 on risk wards to confirm that the outbreak was successfully controlled. All 2,964 patients admitted to one of the risk wards during the

outbreak period were tested for NDM carriage. Six months after the end of the outbreak, over 95% of the patients at risk had been screened, and in total, 29 NDM carriers (1%) had been identified.

Bacterial isolates. The first three isolates were obtained from an ESBL screening using an ESwab on ChromID ESBL agar (bioMérieux Inc., Marcy l'Etoile, France). All subsequent isolates were obtained by screening for carbapenem resistance using enrichment in tryptone soya broth supplemented with ertapenem and vancomycin. During the outbreak period, 42 isolates carrying a *bla*_{NDM} gene detected by PCR were obtained from 29 persons at the outbreak site and included in the study (Table 1). In most cases, a single NDM-producing species was found per patient. However, in nine patients, two to four distinct NDM-producing species were isolated. Most isolates were *K. pneumoniae* ($n = 26$) and *E. coli* ($n = 11$), but two *C. freundii* isolates, two *K. oxytoca* isolates, and one *P. mirabilis* isolate were also cultured. Retrospective analysis showed that five more isolates carrying a *bla*_{NDM} gene had been isolated at the outbreak site between 2015 and 2016 and subsequently submitted to the national reference center. These isolates (three *K. pneumoniae* and two *E. coli*) were also included in the study. For the Dutch national surveillance of CPE, medical microbiology laboratories are requested to submit *Enterobacteriaceae* isolates with a meropenem MIC of >0.25 $\mu\text{g/ml}$ and/or an imipenem MIC of >1 $\mu\text{g/ml}$ to the National Institute for Public Health and the Environment for further molecular characterization. From the Dutch CPE surveillance, NGS data from all *K. pneumoniae* ($n = 133$) and all *E. coli* ($n = 70$) carbapenemase-producing isolates obtained between January 2014 and December 2016 were included for analysis to provide genetic context. All isolates were tested for the production of carbapenemase using the carbapenem inactivation method (CIM) and subjected to PCR to detect the presence of carbapenemase-encoding genes, as previously described (20). Isolates that were shown by PCR to carry a *bla*_{NDM} gene and by CIM to produce carbapenemase are referred to as NDM positive.

Antimicrobial susceptibility testing. At the outbreak location, all isolates were cultured and analyzed for resistance using Vitek (bioMérieux Inc., Marcy l'Etoile, France). Resistance to carbapenem was confirmed by assessing the meropenem MICs for all 250 isolates using Etest (bioMérieux Inc., Marcy l'Etoile, France).

Next-generation sequencing. All 250 isolates were subjected to NGS using the Illumina HiSeq 2500 (BaseClear, Leiden, the Netherlands). The NGS data on the *K. pneumoniae* and *E. coli* isolates were used for MLST and wgMLST analyses using the available wgMLST schemes in SeqSphere software version 2.3.0 (Ridom GmbH, Münster, Germany). The resulting data were imported into Bionumerics version 7.6 for comparative analyses (Applied Maths, Sint-Martens-Latem, Belgium).

The resistomes and plasmid compositions of all the isolates were determined by uploading the assembled contigs into the online tools ResFinder (version 2.1) and PlasmidFinder (version 1.3) via the Center for Genomic Epidemiology website (21, 22). For ResFinder, a 90% identity threshold and a minimum length of 60% were used as criteria, whereas for PlasmidFinder, an identity of 95% was utilized. The NGS data on the presumed outbreak isolates were mapped against known NDM plasmids present in the NCBI database in May 2016.

Electroporation experiments. To confirm that the same plasmid was transferred between species, an NDM-positive *K. pneumoniae* isolate and an NDM-positive *E. coli* isolate from the outbreak location were used to transfer the plasmid carrying the *bla*_{NDM} gene to a susceptible *E. coli* (DH10B) isolate by electroporation. The plasmid of another NDM-positive *K. pneumoniae* isolate from another health care center was also transferred to show that the same plasmid was present in multiple locations. For this, the electroporation protocol of the manufacturer was used (Invitrogen, Carlsbad, CA, USA). Transformants were subjected to PCR to confirm the presence of the *bla*_{NDM} gene, and NGS was performed on each transformant. The resulting NGS data were analyzed in CLC Genomics Workbench version 9.5.3 (Qiagen, Hilden, Germany).

ACKNOWLEDGMENTS

We thank all the Dutch medical microbiology laboratories for submitting their CPE isolates for the national CPE surveillance. We particularly thank Karen Heemstra for searching for epidemiological links between the related NDM-1-positive *K. pneumoniae* cases from another health care center and the outbreak cases.

REFERENCES

1. Yong D, Toleman MA, Giske CG, Cho HS, Sundman K, Lee K, Walsh TR. 2009. Characterization of a new metallo-beta-lactamase gene, *bla*_{NDM-1}, and a novel erythromycin esterase gene carried on a unique genetic structure in *Klebsiella pneumoniae* sequence type 14 from India. *Antimicrob Agents Chemother* 53:5046–5054. <https://doi.org/10.1128/AAC.00774-09>.
2. Nordmann P. 2014. Carbapenemase-producing *Enterobacteriaceae*: overview of a major public health challenge. *Med Mal Infect* 44: 51–56. <https://doi.org/10.1016/j.medmal.2013.11.007>.
3. Zou D, Huang Y, Zhao X, Liu W, Dong D, Li H, Wang X, Huang S, Wei X, Yan X, Yang Z, Tong Y, Huang L, Yuan J. 2015. A novel New Delhi metallo-beta-lactamase variant, NDM-14, isolated in a Chinese hospital possesses increased enzymatic activity against carbapenems. *Antimicrob Agents Chemother* 59:2450–2453. <https://doi.org/10.1128/AAC.05168-14>.
4. Khong WX, Xia E, Marimuthu K, Xu W, Teo YY, Tan EL, Neo S, Krishnan PU, Ang BS, Lye DC, Chow AL, Ong RT, Ng OT. 2016. Local transmission and global dissemination of New Delhi metallo-beta-lactamase (NDM): a whole genome analysis. *BMC Genomics* 17:452. <https://doi.org/10.1186/s12864-016-2740-0>.
5. Walsh TR, Weeks J, Livermore DM, Toleman MA. 2011. Dissemination of NDM-1 positive bacteria in the New Delhi environment and its implications for human health: an environmental point prevalence study. *Lancet Infect Dis* 11:355–362. [https://doi.org/10.1016/S1473-3099\(11\)70059-7](https://doi.org/10.1016/S1473-3099(11)70059-7).
6. Kumarasamy KK, Toleman MA, Walsh TR, Bagaria J, Butt F, Balakrishnan R, Chaudhary U, Doumith M, Giske CG, Irfan S, Krishnan P, Kumar AV, Maharjan S, Mushtaq S, Noorie T, Paterson DL, Pearson A, Perry C, Pike R, Rao B, Ray U, Sarma JB, Sharma M, Sheridan E, Thirunarayan

- MA, Turton J, Upadhyay S, Warner M, Welfare W, Livermore DM, Woodford N. 2010. Emergence of a new antibiotic resistance mechanism in India, Pakistan, and the UK: a molecular, biological, and epidemiological study. *Lancet Infect Dis* 10:597–602. [https://doi.org/10.1016/S1473-3099\(10\)70143-2](https://doi.org/10.1016/S1473-3099(10)70143-2).
7. Johnson AP, Woodford N. 2013. Global spread of antibiotic resistance: the example of New Delhi metallo-beta-lactamase (NDM)-mediated carbapenem resistance. *J Med Microbiol* 62:499–513. <https://doi.org/10.1099/jmm.0.052555-0>.
 8. Denis C, Poirel L, Carricajo A, Grattard F, Fascia P, Verhoeven P, Gay P, Nuti C, Nordmann P, Pozzetto B, Berthelot P. 2012. Nosocomial transmission of NDM-1-producing *Escherichia coli* within a non-endemic area in France. *Clin Microbiol Infect* 18:E128–E130. <https://doi.org/10.1111/j.1469-0691.2012.03761.x>.
 9. Jovicic B, Lepsanovic Z, Suljagic V, Rackov G, Begovic J, Topisirovic L, Kojic M. 2011. Emergence of NDM-1 metallo-beta-lactamase in *Pseudomonas aeruginosa* clinical isolates from Serbia. *Antimicrob Agents Chemother* 55:3929–3931. <https://doi.org/10.1128/AAC.00226-11>.
 10. European Centre for Disease Prevention and Control. 2013. Carbapenemase-producing bacteria in Europe: interim results from the European Survey on carbapenemase-producing *Enterobacteriaceae* (EuSCAPE) project. European Centre for Disease Prevention and Control, Stockholm, Sweden.
 11. Leverstein-van Hall MA, Stuart JC, Voets GM, Versteeg D, Roelofsen E, Fluit AC. 2010. Carbapenem-resistant *Klebsiella pneumoniae* following foreign travel. *Ned Tijdschr Geneesk* 154:A2013. (In Dutch.)
 12. Bathoorn E, Rossen JW, Lokate M, Friedrich AW, Hammerum AM. 2015. Isolation of an NDM-5-producing ST16 *Klebsiella pneumoniae* from a Dutch patient without travel history abroad, August 2015. *Euro Surveill* 20. <https://doi.org/10.2807/1560-7917.ES.2015.20.41.30040>.
 13. Hudson CM, Bent ZW, Meagher RJ, Williams KP. 2014. Resistance determinants and mobile genetic elements of an NDM-1-encoding *Klebsiella pneumoniae* strain. *PLoS One* 9:e99209. <https://doi.org/10.1371/journal.pone.0099209>.
 14. Kluytmans-van den Bergh MF, Rossen JW, Bruijning-Verhagen PC, Bonten MJ, Friedrich AW, Vandenbroucke-Grauls CM, Willems RJ, Kluytmans JA. 2016. Whole-genome multilocus sequence typing of extended-spectrum-beta-lactamase-producing *Enterobacteriaceae*. *J Clin Microbiol* 54:2919–2927. <https://doi.org/10.1128/JCM.01648-16>.
 15. Abdallah HM, Reuland EA, Wintermans BB, Al Naiemi N, Koek A, Abdelwahab AM, Ammar AM, Mohamed AA, Vandenbroucke-Grauls CM. 2015. Extended-spectrum beta-lactamases and/or carbapenemases-producing *Enterobacteriaceae* isolated from retail chicken meat in Zagazig, Egypt. *PLoS One* 10:e0136052. <https://doi.org/10.1371/journal.pone.0136052>.
 16. Leverstein-van Hall MA, Dierikx CM, Cohen Stuart J, Voets GM, van den Munckhof MP, van Essen-Zandbergen A, Platteel T, Fluit AC, van de Sande-Bruinsma N, Scharinga J, Bonten MJ, Mevius DJ, National ESBL Surveillance Group. 2011. Dutch patients, retail chicken meat and poultry share the same ESBL genes, plasmids and strains. *Clin Microbiol Infect* 17:873–880. <https://doi.org/10.1111/j.1469-0691.2011.03497.x>.
 17. Overdevest I, Willemsen I, Rijnsburger M, Eustace A, Xu L, Hawkey P, Heck M, Savelkoul P, Vandenbroucke-Grauls C, van der Zwaluw K, Huijsdens X, Kluytmans J. 2011. Extended-spectrum beta-lactamase genes of *Escherichia coli* in chicken meat and humans, The Netherlands. *Emerg Infect Dis* 17:1216–1222. <https://doi.org/10.3201/eid1707.110209>.
 18. Wohlwend N, Endimiani A, Francey T, Perreten V. 2015. Third-generation-cephalosporin-resistant *Klebsiella pneumoniae* isolates from humans and companion animals in Switzerland: spread of a DHA-producing sequence type 11 clone in a veterinary setting. *Antimicrob Agents Chemother* 59:2949–2955. <https://doi.org/10.1128/AAC.04408-14>.
 19. Zhang J, Zhou K, Zheng B, Zhao L, Shen P, Ji J, Wei Z, Li L, Zhou J, Xiao Y. 2016. High prevalence of ESBL-producing *Klebsiella pneumoniae* causing community-onset infections in China. *Front Microbiol* 7:1830.
 20. van der Zwaluw K, de Haan A, Pluister GN, Bootsma HJ, de Neeling AJ, Schouls LM. 2015. The carbapenem inactivation method (CIM), a simple and low-cost alternative for the Carba NP test to assess phenotypic carbapenemase activity in gram-negative rods. *PLoS One* 10:e0123690. <https://doi.org/10.1371/journal.pone.0123690>.
 21. Carattoli A, Zankari E, Garcia-Fernandez A, Voldby Larsen M, Lund O, Villa L, Moller Aarestrup F, Hasman H. 2014. In silico detection and typing of plasmids using PlasmidFinder and plasmid multilocus sequence typing. *Antimicrob Agents Chemother* 58:3895–3903. <https://doi.org/10.1128/AAC.02412-14>.
 22. Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother* 67:2640–2644. <https://doi.org/10.1093/jac/dks261>.