

University of Groningen

Campylobacter blaseri sp. nov., isolated from common seals (*Phoca vitulina*)

Gilbert, Maarten J; Zomer, Aldert L; Timmerman, Arjen J; Spaninks, Mirlin P; Rubio-García, Ana; Rossen, John W; Duim, Birgitta; Wagenaar, Jaap A

Published in:
International Journal of Systematic and Evolutionary Microbiology

DOI:
[10.1099/ijsem.0.002742](https://doi.org/10.1099/ijsem.0.002742)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Gilbert, M. J., Zomer, A. L., Timmerman, A. J., Spaninks, M. P., Rubio-García, A., Rossen, J. W., Duim, B., & Wagenaar, J. A. (2018). *Campylobacter blaseri* sp. nov., isolated from common seals (*Phoca vitulina*). *International Journal of Systematic and Evolutionary Microbiology*, 68(5), 1787-1794.
<https://doi.org/10.1099/ijsem.0.002742>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Campylobacter blaseri sp. nov., isolated from common seals (*Phoca vitulina*)

Maarten J. Gilbert,^{1,2,*} Aldert L. Zomer,^{1,3} Arjen J. Timmerman,¹ Mirlin P. Spaninks,¹ Ana Rubio-García,^{1,4,5} John W. Rossen,⁴ Birgitta Duim^{1,3} and Jaap A. Wagenaar^{1,3,6}

Abstract

During a study to assess the faecal microbiome of common seals (*Phoca vitulina*) in a Dutch seal rehabilitation centre, 16S rRNA gene sequences of an unknown *Campylobacter* taxon were identified. *Campylobacter* isolates, which differed from the established *Campylobacter* taxa, were cultured and their taxonomic position was determined by a polyphasic study based on ten isolates. The isolates were characterized by 16S rRNA and *atpA* gene sequence analyses and by conventional phenotypic testing. Based on the whole genome sequences, the average nucleotide identity and core genome phylogeny were determined. The isolates formed a separate phylogenetic clade, divergent from all other *Campylobacter* taxa and most closely related to *Campylobacter corcagiensis*, *Campylobacter geochelonis* and *Campylobacter ureolyticus*. The isolates can be distinguished phenotypically from all other *Campylobacter* taxa based on their lack of motility, growth at 25 °C and growth on MacConkey agar. This study shows that these isolates represent a novel species within the genus *Campylobacter*, for which the name *Campylobacter blaseri* sp. nov. is proposed. The type strain for this novel species is 17S00004-5^T (=LMG 30333^T=CCUG 71276^T).

A growing number of *Campylobacter* species have been isolated from marine mammals [1–3]. The species *Campylobacter insulaenigrae* and *Campylobacter pinnipediorum* have been shown to be associated with pinnipeds [1, 3]. These species belong to separate divergent clades within the *Campylobacter* genus; *C. insulaenigrae* belongs to the clade which includes the clinically relevant species *Campylobacter jejuni*, while *C. pinnipediorum* belongs to the clade which includes *Campylobacter concisus*. Both *Campylobacter* species have also been isolated from the widely distributed common seal (*Phoca vitulina*), suggesting that this species serves as an important host reservoir for *Campylobacter*. Furthermore, seals may harbour a larger *Campylobacter* diversity than currently known, which is supported by microbiome data [4]. Here we describe a novel, urease-positive, non-motile *Campylobacter* species isolated from common seals, which is most closely related to *Campylobacter corcagiensis*, *Campylobacter geochelonis* and *Campylobacter ureolyticus*.

During a study to assess the faecal microbiome of common seals in a Dutch seal rehabilitation centre, partial 16S rRNA gene sequences (450 bp of the V3 and V4 regions) of an unknown *Campylobacter* taxon were obtained, which showed 95 % BLAST identity to *C. jejuni*, *C. corcagiensis* and *Campylobacter sputorum* (Rubio-García, unpublished work). Noteworthy, this *Campylobacter* taxon was the tenth most common species observed in the seal microbiome and comprised up to 20 % of all sequences in some seals, with loads increasing with age (3–6 months). The 16S rRNA gene sequences of this *Campylobacter* taxon were identified in 35 out of 98 animals, which were unrelated and did not have prior contact with each other, already at uptake in the rehabilitation centre, indicating that it is a widespread microbiome constituent of wild common seals outside a captive setting as well.

In an attempt to isolate this *Campylobacter* taxon for further characterization, faecal swab samples were collected in 2017

Author affiliations: ¹Department of Infectious Diseases and Immunology, Faculty of Veterinary Medicine, Utrecht University, Utrecht, The Netherlands; ²Reptile, Amphibian and Fish Conservation Netherlands (RAVON), Nijmegen, The Netherlands; ³WHO Collaborating Centre for Campylobacter/OIE Reference Laboratory for Campylobacteriosis, Utrecht, The Netherlands; ⁴Sealcentre, Pieterburen, The Netherlands; ⁵Department of Medical Microbiology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands; ⁶Wageningen Bioveterinary Research, Lelystad, The Netherlands.

*Correspondence: Maarten J. Gilbert, m.j.gilbert@uu.nl

Keywords: *Campylobacter*; novel species; common seal; microbiome; core genome phylogeny; average nucleotide identity.

Abbreviations: ANI, average nucleotide identity; CCDA, charcoal cefoperazone deoxycholate agar; DDH, DNA–DNA hybridization; TSI agar, triple sugar iron agar; TTE broth, Thomann transport and enrichment broth.

The GenBank accession numbers for the whole genome sequences of isolates 17S00004-5 and 17S00008-12 are PDHH00000000 and PDHI00000000, respectively. The GenBank accession numbers for the *atpA* sequences of isolates 17S00004-5 and 17S00008-12 are MG958595 and MG958596, respectively. The GenBank accession numbers for the 16S rRNA sequences of isolates 17S00004-5, 17S00008-12, 17S01633-4, 17S01636-3, 17S01637-3, 17S01639-3, 17S01643-3, 17S01644-3, 17S01646-3, and 17S01649-5 are MG013475, MG013476, MG013477, MG013478, MG013479, MG013480, MG013481, MG013483, MG013485 and MG013486, respectively.

One supplementary figure is available with the online version of this article.

from 40 common seals in January ($n=20$) and in September ($n=20$). Swabs were extracted in 1 ml PBS (Gibco) prior to direct culturing. Specimens from the first sampling event were cultured directly on Columbia agar with 5 % sheep blood (Oxoid) with a filter membrane (0.65 μm ; Millipore) using 300 μl of the suspension, followed by incubation under aerobic conditions at 37 °C for 30 min and subsequent removal of the filter, and on CCD, Preston, and Skirrow agar (Oxoid) using 10 μl of the suspension. Samples were also cultured stagnant in three different selective pre-enrichment media (Bolton, Preston, and Thomann transport and enrichment (TTE) medium [5]; Oxoid), 14 ml of each medium in a 15 ml tube (Sarstedt) with a closed screwcap under aerobic conditions at 37 °C for 48 h, followed by incubation on Columbia agar with 5 % sheep blood with a filter membrane. All agar plates were incubated under microaerobic conditions (83.3 % N_2 , 7.1 % CO_2 , 3.6 % H_2 , and 6 % O_2) at 37 °C for 48–72 h. Additionally, all samples were also cultured both directly and after pre-enrichment on blood agar plates with and without a filter membrane under anaerobic conditions at 37 °C for 48–72 h. Based on results from the first sampling event, samples of the second sampling event were cultured on Preston and Skirrow agar, with and without a selective pre-enrichment in Preston and TTE broth, and incubated in a microaerobic atmosphere at 37 °C for 48–72 h. For the pre-selective enrichment, samples were cultured shaking in 5 ml of each medium in a vertically positioned 25 cm^2 cell culture flask (Corning) with a loose screwcap under microaerobic conditions at 37 °C for 48 h. All isolates of the unknown *Campylobacter* taxon were obtained by direct culturing on Preston and Skirrow agar. In addition, *C. insulaenigrae* was isolated from one animal using the filtration method under microaerobic conditions. In total, 12 isolates of the unknown *Campylobacter* taxon were obtained from ten different animals, of which ten isolates were used for further taxonomic characterization (Table 1).

A polyphasic approach was used to determine the taxonomic position of the ten isolates [6]. Comparisons based on 16S rRNA and *atpA* gene sequences were made to determine the taxonomic position of all isolates. Whole genome sequencing was performed on two isolates on the basis of which the average nucleotide identity (ANI) and phylogenies based on the

full *AtpA* protein and core genome could be determined. Phenotypic characteristics were determined by conventional biochemical testing for all isolates.

Complete genome sequences for isolates 17S00004-5 and 17S00008-12 were obtained by Illumina MiSeq sequencing (Illumina), with a 2 \times 250 bp paired-end output. Sequencing reads were assembled using SPAdes version 3.1.1, resulting in 1 869 677 and 1 869 950 bp genomes, which consisted of 30 and 33 contigs for isolates 17S00004-5 and 17S00008-12, respectively. The GenBank accession numbers for the genome sequences of isolates 17S00004-5 and 17S00008-12 are PDHH00000000 and PDHI00000000, respectively (Table 1).

The taxonomic position of all isolates was determined by comparison of 16S rRNA gene sequences. The 16S rRNA gene sequences (≥ 1211 bp) were extracted from the whole genome sequences or obtained by PCR and Sanger sequencing [7]. Accession numbers are listed in Table 1. The 16S rRNA gene sequences of the other *Campylobacter* taxa were obtained from EzTaxon [8]. Sequence alignment and dendrogram reconstruction were performed using MEGA version 6.05 [9]. A neighbour-joining dendrogram containing all *Campylobacter* taxa was reconstructed, with bootstrap values based on 500 repetitions (Fig. 1). The 16S rRNA gene sequence of *Arcobacter butzleri* strain ATCC 49616^T was used as an outgroup to root the tree. The 16S rRNA gene sequence similarity between the isolates was 99–100 %, while the sequence similarity between these isolates and the most closely related species, *C. geochelonis*, was 94 %.

For improved taxonomic resolution [10], full *AtpA* protein sequences were extracted from the whole genome sequences or obtained from GenBank. Alignment and dendrogram reconstruction were performed as described for 16S rRNA gene phylogeny; the *AtpA* sequence of *Arcobacter butzleri* strain ATCC 49616^T was used as an outgroup to root the tree. Consistent with the 16S rRNA phylogeny, the isolates formed a clade distinct from other *Campylobacter* taxa, with *C. geochelonis* being the most closely related species (Fig. 2).

Additional intraspecific variation amongst all isolates was explored based on partial *atpA* sequences (489 nt), extracted from the whole genome sequences or obtained by PCR and

Table 1. Features of the *C. blaseri* sp. nov. isolates used in this study

Isolate	Host species	Isolation date	16S rRNA gene accession no.	<i>atpA</i> accession no.	Whole genome
17S00004-5 ^T	<i>Phoca vitulina</i>	6 January 2017	MG013475	MG958595	PDHH00000000
17S00008-12	<i>Phoca vitulina</i>	6 January 2017	MG013476	MG958596	PDHI00000000
17S01633-4	<i>Phoca vitulina</i>	12 September 2017	MG013477	N/A	N/A
17S01636-3	<i>Phoca vitulina</i>	11 September 2017	MG013478	N/A	N/A
17S01637-3	<i>Phoca vitulina</i>	10 September 2017	MG013479	N/A	N/A
17S01639-3	<i>Phoca vitulina</i>	12 September 2017	MG013480	N/A	N/A
17S01643-3	<i>Phoca vitulina</i>	10 September 2017	MG013481	N/A	N/A
17S01644-3	<i>Phoca vitulina</i>	10 September 2017	MG013483	N/A	N/A
17S01646-3	<i>Phoca vitulina</i>	10 September 2017	MG013485	N/A	N/A
17S01649-5	<i>Phoca vitulina</i>	14 September 2017	MG013486	N/A	N/A




Fig. 1. Neighbour-joining phylogenetic dendrogram based on 16S rRNA gene sequences. *Arcobacter butzleri* ATCC 49616^T is used as an outgroup and root. Bootstrap values (≥75%) based on 500 replications are indicated at the nodes. Bar, 0.01 substitutions per nucleotide position.




Fig. 2. Neighbour-joining phylogenetic dendrogram based on AtpA protein sequences. Bootstrap values ($\geq 75\%$) based on 500 replications are indicated at the nodes. *Arcobacter butzleri* ATCC 49616^T is used as an outgroup and root. Bar, 0.02 substitutions per amino acid position.

Sanger sequencing. Primers and PCR conditions were as described previously [10], with primers adapted as follows: forward primer 5' GGT CAA GAT GTT GTA TGT GTG TAT GTT GC 3' and reverse primer 5' TTT AAT ATT TCA ACC ATT TTT TCA CC 3'. Alignment and dendrogram reconstruction were performed as described for the 16S rRNA gene phylogeny. The intraspecific variation was similar as observed for the 16S rRNA gene phylogeny, with the exception that isolate 17S01649-5 formed an additional genetic variant (Fig. S1, available in the online version of this article).

In support of the 16S rRNA gene and AtpA phylogeny, a core genome phylogeny without correction for recombination was reconstructed using Fasttree [11], based on a 407 260 bp core gene superalignment of 382 core genes of which the protein sequences had at least 35% sequence identity as determined by Roary [12], from genomes re-annotated using Prokka [13]. The genome of *Helicobacter pylori* strain J99 was used as an outgroup to root the tree and to find orthologs of known virulence factors using VFDB [14]. Consistent with the 16S rRNA gene phylogeny, the isolates formed a clade distinct from other *Campylobacter* taxa (Fig. 3). The most closely




Fig. 3. Phylogenetic dendrogram based on the core genome. *Helicobacter pylori* J99 is used as an outgroup and root. Local support values, calculated with the Shimodaira–Hasegawa test [25], for all branches were 1. Bar, 0.1 substitutions per nucleotide position in the core gene superalignment.

related species were *C. corcagiensis*, *C. ureolyticus* and *C. geochelonis*.

Notable virulence factors, such as flagella encoding genes, *kps* genes coding for capsule production, S-layer protein encoding genes, but also the cytolethal distending toxin encoding *cdtABC* genes, which are widely conserved in *Campylobacter*, were absent from the genomes of both isolates. The majority of the virulence factors absent were related to the flagella and glycosylation of the flagella.

As an alternative for DNA–DNA hybridization (DDH), ANI has been suggested [15, 16]. A DDH species delineation of 70 % corresponds to about 95 % ANI [17]. Using

JSpecies version 1.2.1 [18], pair-wise ANI values based on whole genome sequences were calculated for the seal-associated *Campylobacter* isolates and other taxa of the *Campylobacter* genus. ANI values of the genomes of the isolates were 100 % amongst each other (Table 2). The ANI between the genomes of the isolates and most closely related species *C. ureolyticus* and *C. geochelonis* was maximally 72.2 and 71.9 %, respectively, which is well below the 95 % species cut-off suggested by Goris and colleagues [17].

The G+C content was determined based on the whole genome sequences using Artemis version 13.2 [19]. Both isolates had a G+C content of 29.2 %, which is within the lower end of the range observed for *Campylobacter* (Table 3).

Table 2. Average nucleotide identity (ANI) values (%) based on BLAST for *C. blaseri* sp. nov. and other *Campylobacter* taxa

Strains: 1, *Campylobacter blaseri* sp. nov. 17S00004-5^T; 2, *C. blaseri* sp. nov. 17S00008-12; 3, *C. ureolyticus* RIGS 9880; 4, *C. corcagiensis* CIT 045^T; 5, *C. geochelonis* RC20^T; 6, *C. hominis* ATCC BAA-381^T; 7, *C. sputorum* RM8705; 8, *C. lanienae* NCTC 13004^T; 9, *C. iguaniorum* 1485E^T; 10, *C. hyointestinalis* subsp. *lawsonii* LMG 15993; 11, *C. hyointestinalis* subsp. *hyointestinalis* LMG 9260; 12, *C. fetus* subsp. *testudinum* 03-427; 13, *C. fetus* subsp. *fetus* 82-40; 14, *C. fetus* subsp. *venerealis* 97-608; 15, *C. pinnipediorum* subsp. *pinnipediorum* RM17260^T; 16, *C. pinnipediorum* subsp. *caledonicus* M302/10/6; 17, *C. mucosalis* DSM 21682^T; 18, *C. concisus* ATCC 33237^T; 19, *C. curvus* 525-92; 20, *C. gracilis* ATCC 33236^T; 21, *C. showae* CC57C; 22, *C. rectus* RM3267; 23, *C. upsaliensis* DSM 5365^T; 24, *C. helveticus* ATCC 51209^T; 25, *C. cuniculorum* LMG 24588^T; 26, *C. jejuni* RM1285; 27, *C. hepaticus* HV10^T; 28, *C. coli* FB1; 29, *C. ornithocola* WBE38^T; 30, *C. lari* Slaughter Beach; 31, *C. subantarcticus* LMG 24374; 32, *C. peloridis* LMG 23910^T; 33, *C. insulaeni-grae* NCTC 12927^T; 34, *C. volucris* LMG 24379; 35, *C. avium* LMG 24591^T.

1	---	100.0	72.1	71.0	71.9	68.4	70.4	68.2	68.2	68.4	68.6	68.6	68.5	68.6	70.7	69.6	67.4	67.0	64.8	64.5	64.8	64.4	66.3	66.3	66.5	67.4	67.5	67.2	67.9	68.0	67.6	68.2	68.0	68.1	66.4
2	100.0	---	72.1	71.0	71.8	68.4	70.3	68.1	68.2	68.4	68.6	68.6	68.5	68.6	70.6	69.5	67.4	67.0	64.8	64.5	64.8	64.4	66.3	66.3	66.5	67.3	67.5	67.2	67.8	68.0	67.6	68.2	67.9	68.1	66.4
3	72.2	72.2	---	72.5	71.2	70.2	70.3	68.0	68.4	68.4	68.7	68.6	68.4	68.5	68.9	68.8	67.7	67.2	65.0	64.8	65.0	64.7	66.5	66.6	66.8	67.8	67.5	67.4	68.2	68.1	68.0	68.3	68.0	68.4	66.7
4	70.9	70.9	72.4	---	70.5	68.2	69.8	68.0	68.1	68.0	68.3	68.1	67.9	68.1	68.2	67.8	67.3	67.1	65.0	64.7	65.1	64.6	65.9	65.9	66.3	67.0	66.7	66.7	67.1	67.3	67.1	67.2	66.9	67.4	66.1
5	71.9	71.9	71.0	70.4	---	68.7	69.7	68.2	69.2	68.9	69.0	69.2	68.8	68.8	68.5	68.4	68.0	68.2	66.4	66.7	66.5	66.0	66.4	66.4	66.0	66.9	66.4	66.8	67.0	67.2	66.8	67.2	66.5	67.1	66.0
6	68.6	68.6	70.2	68.3	69.0	---	69.0	66.9	67.7	67.9	67.9	68.0	67.9	67.9	67.6	67.6	67.0	66.9	65.6	66.9	65.9	65.6	66.2	66.0	66.3	67.1	66.9	67.0	67.0	66.8	66.9	66.9	66.7	67.1	65.7
7	70.3	70.3	70.3	69.8	69.6	68.9	---	67.9	68.2	68.5	68.8	68.8	68.7	68.7	69.1	69.1	67.6	67.2	65.1	64.6	65.0	64.6	66.4	66.4	66.9	67.5	67.6	67.3	68.0	67.9	67.5	68.2	67.8	68.2	66.5
8	68.2	68.2	68.1	68.1	68.5	68.9	---	71.5	71.9	73.9	72.0	71.7	71.7	71.8	67.9	68.0	67.6	66.2	65.9	66.2	65.9	66.5	66.5	66.6	66.5	67.1	67.0	67.3	68.0	67.3	67.2	67.5	67.3	67.6	66.6
9	68.6	68.6	68.4	68.4	69.8	67.8	68.6	71.6	---	75.4	75.5	75.5	75.0	75.0	68.4	68.4	68.2	68.5	67.1	66.5	67.2	66.9	66.1	66.1	66.1	67.1	66.7	67.0	67.1	67.3	67.1	67.3	66.7	67.0	66.2
10	68.5	68.5	68.5	68.1	69.2	67.9	68.6	71.9	75.4	---	94.4	79.2	78.0	78.0	68.4	68.4	67.9	68.1	66.6	66.1	66.9	66.5	66.0	66.3	66.1	67.1	66.7	66.9	67.0	67.2	66.9	67.3	66.9	67.2	66.4
11	68.9	68.9	68.9	68.5	69.3	67.9	69.1	73.8	75.5	94.3	---	78.2	77.3	77.4	68.7	68.6	68.0	68.2	66.8	66.1	66.8	66.4	66.6	66.5	66.4	67.3	67.1	67.3	67.5	67.6	67.2	67.7	67.3	67.6	66.4
12	68.7	68.7	68.6	68.2	69.3	67.7	69.1	71.9	75.4	79.2	78.0	---	91.9	92.0	68.8	68.6	67.9	67.8	66.4	65.7	66.6	66.1	66.1	66.3	66.1	67.0	66.9	67.0	67.3	67.2	67.1	67.3	67.0	67.3	66.2
13	68.5	68.5	68.4	68.0	68.8	67.7	68.7	71.8	74.9	77.9	77.2	92.0	---	99.7	68.3	68.3	67.6	67.5	66.5	65.7	66.4	66.1	66.0	66.1	66.2	66.9	66.7	66.9	67.3	67.2	67.1	67.3	66.9	67.2	66.3
14	68.9	68.9	68.7	68.2	69.1	68.1	69.0	71.9	75.0	78.2	77.4	92.1	99.6	---	68.7	68.5	67.6	67.5	66.4	65.8	66.6	66.3	66.0	66.2	66.3	67.0	66.8	67.1	67.3	67.3	67.2	67.4	66.9	67.3	66.3
15	70.8	70.8	69.2	68.6	68.9	67.7	69.5	68.2	68.5	68.6	68.7	68.9	68.5	68.7	---	94.3	71.1	70.1	68.0	64.9	67.0	66.5	66.5	66.8	66.8	67.9	67.7	67.7	68.4	68.4	68.1	68.4	68.2	68.5	67.0
16	69.7	69.7	68.9	68.1	68.9	67.6	69.2	68.1	68.3	68.3	68.6	68.6	68.4	68.4	94.3	---	71.1	70.2	68.0	65.0	66.8	66.4	66.6	66.7	66.8	67.7	67.8	67.6	68.2	68.3	68.0	68.4	68.1	68.4	66.9
17	67.3	67.3	67.2	67.3	68.1	66.7	67.6	67.8	67.6	67.7	67.4	67.4	71.0	71.0	---	71.0	70.1	70.0	68.3	66.3	66.8	68.5	65.9	66.0	65.4	66.0	65.0	65.4	66.0	66.2	66.2	65.7	66.0	65.8	
18	67.1	67.1	67.4	67.2	68.5	66.9	67.7	67.4	68.4	67.9	68.1	67.9	67.6	67.6	70.0	70.1	71.1	---	73.1	67.5	71.3	70.4	66.0	66.0	65.4	66.4	65.8	66.4	66.4	66.3	66.4	66.4	66.0	66.3	
19	64.9	64.9	65.0	65.3	66.8	65.8	65.2	67.1	66.8	66.8	66.4	66.4	66.4	67.9	67.9	70.3	73.0	---	68.8	71.9	71.6	65.3	64.5	64.9	64.1	64.9	64.6	64.8	64.7	64.6	64.4	64.6	64.4	65.0	
20	64.5	64.5	64.7	65.1	66.9	66.9	64.7	65.7	66.1	66.1	65.9	65.8	65.8	65.9	65.1	64.9	66.8	67.6	68.9	---	71.4	71.5	64.9	64.8	64.3	64.5	63.7	64.4	64.1	64.5	64.2	64.3	64.0	64.2	64.5
21	64.8	64.8	64.7	65.0	66.6	65.6	64.8	65.8	66.7	66.4	66.4	66.1	66.0	66.2	66.5	66.5	68.5	71.0	71.5	71.2	---	89.0	64.8	64.7	64.0	64.4	63.9	64.7	64.4	64.5	64.5	64.5	64.0	64.4	64.5
22	64.4	64.4	64.4	64.7	66.2	65.4	64.6	65.5	66.6	66.2	66.0	65.7	65.8	66.0	66.3	66.0	68.4	70.3	71.5	71.3	89.1	---	64.9	64.7	64.0	64.6	64.0	64.5	64.4	64.4	64.2	64.3	63.9	64.1	64.5
23	66.3	66.3	66.1	65.9	66.2	65.9	66.2	66.1	65.9	65.9	66.0	66.0	65.9	65.9	66.1	66.1	65.9	65.6	65.0	64.4	65.1	64.9	---	85.1	73.7	74.3	73.4	74.2	70.5	70.9	70.3	70.7	70.1	70.6	69.2
24	66.6	66.6	66.7	66.4	66.7	66.2	66.8	66.7	66.3	66.4	66.5	66.5	66.4	66.3	66.5	66.5	66.7	66.2	65.5	65.0	65.7	65.6	85.4	---	73.8	74.6	73.6	74.1	70.9	70.6	71.0	70.2	70.7	69.4	
25	66.9	66.9	67.0	66.5	66.5	66.7	67.1	66.4	66.1	66.0	66.3	66.3	66.3	66.3	66.7	66.8	65.8	65.5	64.3	64.3	64.3	64.1	73.8	73.8	---	75.8	75.3	75.6	71.7	71.5	71.3	72.2	71.2	71.7	69.3
26	67.6	67.6	67.8	67.1	67.3	67.2	67.6	66.9	67.0	67.1	67.3	67.1	67.1	67.5	67.4	66.8	66.4	65.3	64.4	65.1	64.9	74.3	74.5	75.7	---	83.9	84.1	73.3	73.3	72.8	73.3	72.6	73.3	69.8	
27	67.3	67.3	67.5	66.7	66.6	66.6	67.6	66.6	66.2	66.5	66.8	66.8	66.6	66.6	67.5	67.5	65.5	63.9	63.3	64.2	63.9	73.4	73.4	75.2	84.0	---	80.5	72.9	72.9	72.7	73.3	72.7	73.1	69.4	
28	67.3	67.3	67.4	66.9	67.2	67.0	67.4	67.1	66.9	66.9	67.1	67.1	67.0	67.1	67.5	67.4	66.4	66.2	65.0	64.6	65.1	65.0	74.2	74.1	75.3	84.1	80.5	---	72.9	73.0	72.7	73.1	72.2	72.8	69.8
29	67.9	67.9	67.9	67.3	67.2	66.8	67.9	67.0	66.8	67.0	67.2	67.4	67.2	67.2	68.0	67.9	66.5	66.0	64.6	64.0	64.6	64.3	70.5	70.7	71.3	73.1	72.8	72.8	---	91.7	88.4	86.0	80.9	82.0	69.2
30	68.3	68.3	68.2	67.5	67.6	66.8	68.0	67.1	67.2	67.5	67.4	67.4	67.4	67.3	67.9	68.0	66.7	66.4	65.0	64.5	65.0	64.5	70.7	70.7	71.3	73.1	73.0	72.9	92.0	---	90.0	86.3	81.0	82.3	69.5
31	67.9	67.9	68.0	67.2	67.5	66.8	67.6	67.1	66.9	67.1	67.1	67.4	67.2	67.2	67.9	67.8	66.4	66.3	64.7	64.4	65.0	64.4	70.4	70.5	71.4	73.0	72.6	72.6	88.4	89.8	---	85.4	80.3	81.9	69.0
32	68.0	68.0	68.0	67.4	67.4	66.8	68.0	67.3	67.1	67.1	67.4	67.3	67.3	67.3	67.9	68.0	66.4	66.1	64.4	64.0	64.5	64.2	70.7	71.2	72.5	73.3	73.2	72.9	86.5	86.4	85.5	---	81.1	82.5	69.4
33	68.0	68.0	68.2	67.1	67.1	67.0	67.9	66.9	66.7	66.8	67.1	67.1	67.0	66.9	67.9	67.7	66.4	65.9	64.3	64.3	64.6	64.2	70.0	70.0	71.0	72.6	72.7	72.2	81.0	81.0	80.2	81.1	---	82.9	68.8
34	68.2	68.2	68.4	67.5	67.4	67.1	68.2	67.3	66.9	67.4	67.5	67.6	67.2	67.2	68.2	68.1	66.6	66.3	64.9	64.4	65.0	64.5	70.7	70.6	71.5	73.2	73.0	72.9	82.2	82.3	81.8	82.4	82.9	---	69.5
35	66.7	66.7	66.6	66.4	66.5	65.9	66.6	66.3	66.2	66.5	66.5	66.4	66.3	66.8	66.7	66.3	65.9	65.0	64.5	65.0	64.8														

Conflicts of interest

The authors declare that there are no conflicts of interest.

Ethical statement

The procedures conducted on the animals in the study were combined with routine veterinary diagnostic and therapeutic acts and were not considered to cause any additional discomfort. Consequently, the study was not considered an animal experiment under the Dutch 'Experiments on Animals Act', making assessment by an animal ethics committee unnecessary.

References

- Foster G, Holmes B, Steigerwalt AG, Lawson PA, Thorne P et al. *Campylobacter insulaenigrae* sp. nov., isolated from marine mammals. *Int J Syst Evol Microbiol* 2004;54:2369–2373.
- Goldman CG, Matteo MJ, Loureiro JD, Almuzara M, Barberis C et al. Novel gastric helicobacters and oral campylobacters are present in captive and wild cetaceans. *Vet Microbiol* 2011;152:138–145.
- Gilbert MJ, Miller WG, Leger JS, Chapman MH, Timmerman AJ et al. *Campylobacter pinnipediorum* sp. nov., isolated from pinnipeds, comprising *Campylobacter pinnipediorum* subsp. *pinnipediorum* subsp. nov. and *Campylobacter pinnipediorum* subsp. *caledonicus* subsp. nov. *Int J Syst Evol Microbiol* 2017;67:1961–1968.
- Bik EM, Costello EK, Switzer AD, Callahan BJ, Holmes SP et al. Marine mammals harbor unique microbiotas shaped by and yet distinct from the sea. *Nat Commun* 2016;7:10516.
- Harwood LJ, Thomann A, Brodard I, Makaya PV, Perreten V. *Campylobacter fetus* subspecies *venerealis* transport medium for enrichment and PCR. *Vet Rec* 2009;165:507–508.
- On SLW, Miller WG, Houf K, Fox JG, Vandamme P. Minimal standards for describing new species belonging to the families *Campylobacteraceae* and *Helicobacteraceae*: *Campylobacter*, *Arcobacter*, *Helicobacter* and *Wolinella* spp. *Int J Syst Evol Microbiol* 2017;67:5296–5311.
- Maiwald M. Broad-range PCR for detection and identification of bacteria. In: Persing DH, Tenover FC, Versalovic J, Tang YW, Relman D et al. (editors). *Molecular Microbiology: Diagnostics Principles and Practice*. Washington, DC: ASM Press; 2004. pp. 379–390.
- Kim OS, Cho YJ, Lee K, Yoon SH, Kim M et al. Introducing EzTaxon-e: a prokaryotic 16S rRNA gene sequence database with phylotypes that represent uncultured species. *Int J Syst Evol Microbiol* 2012;62:716–721.
- Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol* 2013;30:2725–2729.
- Miller WG, Yee E, Jolley KA, Chapman MH. Use of an improved *atpA* amplification and sequencing method to identify members of the *Campylobacteraceae* and *Helicobacteraceae*. *Lett Appl Microbiol* 2014;58:582–590.
- Price MN, Dehal PS, Arkin AP. FastTree: computing large minimum evolution trees with profiles instead of a distance matrix. *Mol Biol Evol* 2009;26:1641–1650.
- Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S et al. Roary: rapid large-scale prokaryote pan genome analysis. *Bioinformatics* 2015;31:3691–3693.
- Seemann T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 2014;30:2068–2069.
- Chen L, Yang J, Yu J, Yao Z, Sun L et al. VFDB: a reference database for bacterial virulence factors. *Nucleic Acids Res* 2005;33:D325–D328.
- Konstantinidis KT, Tiedje JM. Genomic insights that advance the species definition for prokaryotes. *Proc Natl Acad Sci USA* 2005;102:2567–2572.
- Konstantinidis KT, Ramette A, Tiedje JM. The bacterial species definition in the genomic era. *Philos Trans R Soc Lond B Biol Sci* 2006;361:1929–1940.
- Goris J, Konstantinidis KT, Klappenbach JA, Coenye T, Vandamme P et al. DNA-DNA hybridization values and their relationship to whole-genome sequence similarities. *Int J Syst Evol Microbiol* 2007;57:81–91.
- Richter M, Rosselló-Móra R. Shifting the genomic gold standard for the prokaryotic species definition. *Proc Natl Acad Sci USA* 2009;106:19126–19131.
- Rutherford K, Parkhill J, Crook J, Horsnell T, Rice P et al. Artemis: sequence visualization and annotation. *Bioinformatics* 2000;16:944–945.
- Elharrif Z, Mégraud F. Characterization of thermophilic *Campylobacter*: I. Carbon-substrate utilization tests. *Curr Microbiol* 1986;13:117–122.
- Elharrif Z, Mégraud F. Characterization of thermophilic *Campylobacter*. II. Enzymatic profiles. *Curr Microbiol* 1986;13:317–322.
- On SL, Holmes B. Effect of inoculum size on the phenotypic characterization of *Campylobacter* species. *J Clin Microbiol* 1991;29:923–926.
- On SL, Holmes B. Reproducibility of tolerance tests that are useful in the identification of campylobacteria. *J Clin Microbiol* 1991;29:1785–1788.
- On SL, Holmes B. Assessment of enzyme detection tests useful in identification of campylobacteria. *J Clin Microbiol* 1992;30:746–749.
- Shimodaira H, Hasegawa M. Multiple comparisons of log-likelihoods with applications to phylogenetic inference. *Mol Biol Evol* 1999;16:1114–1116.
- On SL, Holmes B, Sackin MJ. A probability matrix for the identification of campylobacters, helicobacters and allied taxa. *J Appl Bacteriol* 1996;81:425–432.
- Rossi M, Debruyne L, Zanoni RG, Manfreda G, Revez J et al. *Campylobacter avium* sp. nov., a hippurate-positive species isolated from poultry. *Int J Syst Evol Microbiol* 2009;59:2364–2369.
- Debruyne L, Broman T, Bergström S, Olsen B, On SL et al. *Campylobacter subantarcticus* sp. nov., isolated from birds in the sub-Antarctic region. *Int J Syst Evol Microbiol* 2010;60:815–819.
- Debruyne L, Broman T, Bergström S, Olsen B, On SL et al. *Campylobacter volucris* sp. nov., isolated from black-headed gulls (*Larus ridibundus*). *Int J Syst Evol Microbiol* 2010;60:1870–1875.
- Koziel M, O'Doherty P, Vandamme P, Corcoran GD, Sleator RD et al. *Campylobacter corcagiensis* sp. nov., isolated from faeces of captive lion-tailed macaques (*Macaca silenus*). *Int J Syst Evol Microbiol* 2014;64:2878–2883.
- Piccirillo A, Niero G, Calleros L, Pérez R, Naya H et al. *Campylobacter geochelonis* sp. nov. isolated from the western Hermann's tortoise (*Testudo hermanni hermanni*). *Int J Syst Evol Microbiol* 2016;66:3468–3476.
- van TT, Elshagmani E, Gor MC, Scott PC, Moore RJ. *Campylobacter hepaticus* sp. nov., isolated from chickens with spotty liver disease. *Int J Syst Evol Microbiol* 2016;66:4518–4524.
- Cáceres A, Muñoz I, Iraola G, Díaz-Viraqué F, Collado L. *Campylobacter ornithocola* sp. nov., a novel member of the *Campylobacter lari* group isolated from wild bird faecal samples. *Int J Syst Evol Microbiol* 2017;67:1643–1649.